九州大学学術情報リポジトリ Kyushu University Institutional Repository

# Identification of genes involved in fluoride resistance in oral streptococci

門, 暁晨

https://doi.org/10.15017/1785374

出版情報:九州大学,2016,博士(歯学),課程博士 バージョン: 権利関係:全文ファイル公表済

### Identification of genes involved in fluoride resistance in oral streptococci

2016.9 Xiaochen Men

Section of Preventive and Public Health Dentistry Division of Oral Health, Growth and Development, Graduate School of Dental Science Kyushu University

Supervisors

Professor Yoshihisa Yamashita Section of Preventive and Public Health Dentistry Division of Oral Health, Growth and Development, Faculty of Dental Science Kyushu University

#### A part of this thesis has been submitted to Journal of Dental Research:

Anion channels involved in fluoride resistance in oral streptococci

Xiaochen Men, Yukie Shibata, Toru Takeshita, and Yoshihisa Yamashita

#### CONTENTS

ABSTRACT · · · · · · · · · · · · · · · · · · ·	4
INTRODUCTION · · · · · · · · · · · · · · · · · · ·	5
MATERIALS AND METHODS • • • • • • • • • • • • • • • • • • •	7
RESULTS · · · · · · · · · · · · · · · · · · ·	12
DISCUSSION · · · · · · · · · · · · · · · · · · ·	27
ACKNOWLEDGMENTS · · · · · · · · · · · · · · · · · · ·	32
REFERENCES · · · · · · · · · · · · · · · · · · ·	33

#### ABSTRACT

Recently, it has been reported that *eriC* and *crcB* are involved in bacterial fluoride resistance (Baker et al. 2012). However, the fluoride-resistance mechanism in oral streptococci remains unclear. BLAST studies showed that two types of eriCs (eriCl and eriC2) and two types of crcBs (crcB1 and crcB2) are present across 18 oral streptococci, which were selected based on the following criteria: identification in  $\ge 10\%$  of 166 orally healthy subjects and  $\ge 0.01\%$ of the mean relative abundance. They were divided into three groups based on the distribution of these four genes: group I, only *eriC1*; group II, *eriC1* and *eriC2*; and group III, *eriC2*, crcB1, and crcB2. Group I consisted of Streptococcus mutans, in which one of the two eriC1s predominantly affected fluoride resistance. Group II consisted of eight species, in which eriCl was involved in fluoride resistance, but eriC2 was not, in Streptococcus anginosus as a representative species. Group III consisted of nine species, in which both crcB1 and crcB2 were crucial for fluoride resistance, but eriC2 was not, in Streptococcus sanguinis as a representative species. Based on these results, either EriC1 or CrcB play a role in fluoride resistance in oral streptococci. Complementation between S. mutans EriC1 and S. sanguinis CrcB1/B2 was confirmed in both S. mutans and S. sanguinis. However, neither transfer of S. sanguinis CrcB1/B2 into wild-type S. mutans nor S. mutans EriC1 into wild-type S. sanguinis increased the fluoride resistance of the wild-type strain. It is possible that EriC1 and CrcB1/B2 are responsible for fluoride resistance in oral streptococci by sharing specific pathways.

#### **INTRODUCTION**

Fluoride is commonly used as an effective caries-preventive agent in many countries. Organisms present in the oral cavity are frequently exposed to fluoride ions from drinking water or from the use of fluoride dentifrices and fluoride mouth rinses. Fluoride is known to interfere with metabolic processes in many organisms (Marquis et al. 2003), and high-concentration fluoride shows a bactericidal effect (Mayhew and Brown 1981) (Maltz and Emilson 1982). These anti-microbial effects contribute to the anti-caries effect of fluoride. However, the anti-microbial effects of fluoride may have an unexpected effect on the bacterial composition of oral microflora.

The widespread, long-term use of fluoride can result in the emergence of fluoride-resistant oral *Streptococcus* species, including *Streptococcus mutans*, a cariogenic bacterium (Streckfuss et al. 1980) (Hamilton 1969) (Bunick and Kashket 1981) (Brown et al. 1983) (Sheng and Liu 2000). These fluoride-resistant strains show clear phenotypic differences in growth, adherence, and metabolic activity compared to the fluoride-sensitive strains (Van Loveren et al. 1991) (Van Loveren et al. 1991) (Hoelscher and Hudson 1996). The emergence of fluoride-resistant oral *Streptococcus* species may not only decrease the anti-caries effects of fluoride, but also disrupt the composition of oral streptococci in the oral cavity, followed by the deterioration of oral health.

More than 700 bacterial species are present in the oral cavity, and oral streptococci are predominant. Oral streptococci account for approximately 20% of all bacteria in saliva (Moritani et al. 2015) and approximately 50% of those during the early stages of dental plaque formation (Takeshita et al. 2015). Oral streptococci pioneer early dental plaque

 $\mathbf{5}$ 

formation and have a specific temporal and spatial distribution that is crucial for the development of oral biofilms (Rickard et al. 2003).

Bacteria have evolved numerous strategies to alleviate the toxic effects of ions other than fluoride, and yet analogous systems for fluoride toxicity mitigation were notably absent (Silver 1996). Recently, it has been reported that the *eriC* gene of *Pseudomonas syringae* and the *crcB* gene of *Escherichia coli* are involved in bacterial fluoride resistance (Baker et al. 2012). The *eriC* gene is described as a CIC chloride channel in many bacteria, and the *crcB* gene has previously been implicated in resistance to camphor-induced chromosome decondensation. Furthermore, Liao et al. (2015) identified two single nucleotide polymorphisms (SNPs) in the gene cluster, including two *eriC* genes in the genome of the fluoride-resistant mutant *S. mutans* C180-2FR. These results suggested that *eriC* may be related to the response to fluoride in *S. mutans*. However, it remains unclear whether *eriC* and/or *crcB* play a role in fluoride resistance in oral streptococci. To maintain the appropriate bacterial composition of oral microflora, it is important to investigate the fluoride resistance mechanism in oral streptococci. In this study, we attempted to identify and characterize fluoride-resistance-related genes in oral streptococci.

#### MATERALS AND METHODS

#### **Bacterial strains and culture conditions**

The following bacterial strains were used in this study: *Streptococcus mutans* UA159, *Streptococcus sobrinus* OMZ175, *Streptococcus mitis* ATCC 49456, *Streptococcus oralis* ATCC 10557, *Streptococcus gordonii* ATCC 10558, *Streptococcus sanguinis* SK36, *Streptococcus parasanguinis* ATCC 15912, *Streptococcus tigurinus* ATCC 15914, *Streptococcus australis* ATCC 700641, *Streptococcus infantis* ATCC 700779, *Streptococcus salivarius* HHT, *Streptococcus anginosus* NCTC 10707, *Streptococcus intermedius* ATCC 27335, and *Escherichia coli* DH5 $\alpha$ . *E. coli* strains were grown in 2×YT Broth (Becton Dickinson, Franklin Lakes, NJ, USA). Oral streptococci strains were grown in brain heart infusion (BHI) broth (Becton Dickinson) at 37°C in 5% CO<sub>2</sub>. Antibiotics were used at the following concentrations: 300 µg/mL erythromycin and 50 µg/mL ampicillin for *E. coli*, 20 µg/mL erythromycin for oral streptococci, 600 µg/mL spectinomycin for *S. mutans* and *S. anginosus*, and 150 µg/mL spectinomycin for *S. sanguinis*.

#### Transformation

Transformation of oral streptococci was performed as described previously (Perry et al. 1983). Briefly, *Streptococcus* cells were grown for 18 h in BHI broth containing 10% heat-inactivated (56°C, 30 min) horse serum (BHI-HS), diluted 1:10 into fresh BHI-HS, and incubated at 37°C until an optical density of about 0.2 (at 550 nm) was attained. Competent cells were then exposed to DNA for 30 min, diluted 1:2 into fresh BHI-HS, and incubated for 2 h. An aliquot of cell suspensions was spread on BHI plates containing appropriate antibiotics.

Transformation of *E. coli* was performed as described previously (Hanahan 1983). Briefly, *E. coli* cells were grown in SOB medium (Becton Dickinson) until the cell density reaches 0.3 (at 550 nm). Cell cultures (50 mL) were placed on ice for 10 min, pelleted at 5000 rpm for 10 min at 4°C, and resuspended in 17 mL of cool CCMB solution [80 mM CaCl<sub>2</sub>, 20 mM MnCl<sub>2</sub>, 10 mM MgCl<sub>2</sub>, 10 mM potassium acetate, and 10% (v/v) glycerol]. Cell suspensions were again placed on ice for 20 min, harvested at 5000 rpm for 10 min at 4°C, and resuspended in 5mL of cool CCMB solution. Competent cells and DNA were mixed and placed on ice for 45 min. The mixture was heat-pulsed without agitation at 42°C for 60 sec and placed on ice for 2 min. After addition of 1 mL of fresh SOB medium, the cell suspension was incubated at 37°C for 1~4 h to allow the bacteria to recover and to express the antibiotic resistance marker encoded by the plasmid. The appropriate volume of transformed competent cells was spread onto 2×YT agar plate with appropriate antibiotic.

#### Construction of mutant forms of the genes in Streptococcus species

Various deletion mutants were constructed by replacing the target gene with an erythromycin resistance or spectinomycin resistance gene using double cross-over homologous recombination, as described previously (Li et al. 2012). As an example, we describe a strategy for construction of the *eriCla/eriClb* double mutant (Fig. 1). An 869-bp fragment upstream of *eriCla* and a 767-bp fragment downstream of *eriClb* were amplified from *S. mutans* UA159 genomic DNA and inserted upstream and downstream, respectively, of the Em<sup>r</sup> gene in pBSSKII-Em<sup>r</sup> (Kawada-Matsuo et al. 2009), in which the Em<sup>r</sup> fragment was cloned into HindIII- and EcoRV-digested pBluescript SK II (+). The resultant plasmid

(pBSSKII-Em<sup>r</sup>-*eriC1a/eriC1b* -UD) was digested with KpnI and SacII, and the assembled fragment was transformed into *S. mutans* UA159. Correct insertions or replacements of transformants were confirmed by PCR.

Complementation was performed using the shuttle vector, pSEP, which consisted of an erythromycin resistance gene, pC194ori for replication in Gram-positive organisms, pUCori for replication in *E. coli*, and the promoter region of the erythromycin resistance gene derived from Gram-positive organisms. Fig. 2 shows a strategy for complementation of *eriC1a/eriC1b* by *eriC1b* in the *eriC1a/eriC1b* double mutant of *S. mutans*.

S. mutans UA159



Fig. 1. Construction of the *eriCla/eriClb* double mutant



Fig. 2. Construction of the mutant complemented by EriC1b

#### Evaluation of the fluoride sensitivity of strains

The mutant strains and wild-type control strain cells were grown in BHI broth overnight at  $37^{\circ}$ C in 5% CO<sub>2</sub>. The cultures were then diluted 1:10 into fresh BHI and grown to an OD<sub>550</sub> of ~0.5. Aliquots (20  $\mu$ L) of cell suspensions with the same turbidity were inoculated into wells containing 200  $\mu$ L of fresh BHI medium with several different sodium fluoride (NaF) concentrations. Growth was monitored by measuring OD<sub>550</sub> using a SpectraMax 340PC384 microplate spectrophotometer (Molecular Devices, Sunnyvale, CA, USA). Wells containing only BHI were used as controls. Growth yields were estimated based on the means of the data obtained from three independent experiments.

#### RESULTS

#### Distribution of *eriC* and *crcB* genes in oral streptococci

To examine the distribution of *eriC* and *crcB* genes in oral streptococci, we performed sequence homology analysis for the complete genome sequences of 18 types of oral streptococci with the BLAST program via the National Center for Biotechnology Information server. These species were selected based on the following criteria: identification in  $\ge 10\%$  of 166 orally healthy subjects and  $\geq 0.01\%$  of the mean relative abundance. There were two types of *eriC* genes (*eriC1* and *eriC2*) and two types of *crcB* genes (*crcB1* and *crcB2*) in oral streptococci (Fig. 3). The *eriC1* gene product showed about 50% similarity with EriC of *P*. syringae DC3000, which was identified as a fluoride channel protein in a previous report (Baker et al. 2012). On the other hand, another EriC2 had no similarity with *P. syringae* EriC. Both crcB products showed about 50% similarity with CrcB of E. coli K-12, which was involved in fluoride resistance in a previous report (Baker et al. 2012). As shown in Fig. 3, these oral streptococci were divided into three groups based on the distribution of eriC and crcB genes: group I with only eriCl (S. mutans), group II with eriCl and eriC2 (S. sobrinus, S. salivarius, S. vestibularis, S. australis, S. parasanguinis, S. anginosus, S. constellatus, and S. intermedius), and group III with eriC2, crcB1, and crcB2 (S. cristatus, S. gordonii, S. infantis, S. mitis, S. oralis, S. peroris, S. pneumonia, S. sanguinis, and S. tigurinus). The eriC1 and *crcB* genes were flanked by highly similar gene arrangements.



**Fig. 3.** Locations of *eriC* and *crcB* genes in 18 oral streptococci. These species were selected based on the following criteria: identification in  $\ge 10\%$  of 166 orally healthy subjects and  $\ge 0.01\%$  of the mean relative abundance.

#### Fluoride resistance of oral streptococci

To examine the fluoride resistance of oral streptococci, 13 available species among the above 18 oral Streptococcus species were grown in BHI broth with various concentrations of NaF. As shown in Fig. 4, these oral streptococci showed various resistances to fluoride. S. parasanguinis, S. sanguinis, S. sobrinus, S. salivarius, and S. anginosus grew fairly well, even in the presence of 300 ppm NaF. S. australis, S. oralis, S. infantis, S. mitis, and S. tigurinus grew poorly under the same conditions. In the presence of 600 ppm NaF, all 13 oral streptococci did not grow. Among species with the higher fluoride-resistance, S. parasanguinis, S. sobrinus, S. salivarius, and S. anginosus belong to group II and S. sanguinis belongs to group III. On the other hand, among species with the lower fluoride-resistance, S. oralis, S. infantis, S. mitis, and S. tigurinus belonged to group III and S. australis belonged to group II. Streptococcus mutans, which belonged to group I, showed medium fluoride-resistance. No strong relationship was observed between the distribution of *eriC* and *crcB* genes and fluoride resistance. Next, we explored the related gene(s) for fluoride resistance in S. mutans, S. anginosus, and S. sanguinis, which possess a fair fluoride resistance, as representative species of the three groups, respectively.



**Fig. 4.** Growth yields of 13 oral streptococci at various NaF concentrations. The y-axis represents the  $OD_{550}$  after incubation for 16 h. The data represent the mean standard deviations of three independent experiments.

#### Identification of fluoride-resistance-related genes in S. mutans

Sequence homology analysis revealed two eriCl genes [SMU 1290 (eriCla) and SMU 1289 (eriClb)] in S. mutans UA159. eriCla and eriClb exist in tandem and in the same orientation. Both eriCla and eriClb encode a protein of 406 amino acids, named the chloride channel permease. The amino acid sequences deduced from eriCla and eriClb showed 52.6 and 52.5% similarity, respectively, to *P. syringae* EriC. The amino acid sequence similarity between EriC1a and EriC1b was 74.4%. To examine whether inactivation of *eriC1a* or/and eriClb resulted in the loss of fluoride resistance, eriCla and eriClb single mutants, and the eriCla/eriClb double mutant were constructed using double-crossover homologous recombination, as described in the Materials and Methods. As shown in Fig. 5, the *eriClb* mutant showed extremely limited growth in the presence of 75 ppm NaF, while it grew similarly to the wild-type UA159 in the absence of NaF. In contrast, the *eriCla* mutant showed a growth rate similar to wild-type UA159 in both the presence and absence of NaF, and inactivation of eriCla did not result in a loss of fluoride resistance. Furthermore, the growth rates of the *eriCla/eriClb* double mutant and the *eriClb* single mutant were similar in the presence of NaF. Complementation of *eriC1b* was performed using the shuttle vector, pSEP, in the *eriCla/eriClb* double mutant. The complemented strain showed rescued fluoride resistance (Fig. 5). On the other hand, introduction of *eriCla* into the *eriCla/eriClb* double mutant did not restore fluoride resistance. These results demonstrated that EriC1b was responsible for the fluoride resistance in S. mutans.

16



**Fig. 5.** Growth yields of the parent strain, eriC1a mutant, eriC1b mutant, eriC1a/eriC1b double mutant and complemented strains in *S. mutans* at various NaF concentrations. The y-axis represents the OD<sub>550</sub> after incubation for 16 h. Data represent the mean standard deviations of three independent experiments.

#### Identification of fluoride-resistance-related genes in S. anginosus

*S. anginosus* possesses two types of *eriC* genes. *eriC1* encodes a protein of 405 amino acids with amino acid sequence similarity of 52% to the *P. syringae* EriC, and is named the voltage-gated chloride channel protein. *eriC2* encodes a protein of 518 amino acids with no amino acid sequence similarity to the *P. syringae* EriC. To examine whether the inactivation of *eriC1* or *eriC2* leads to the loss of fluoride resistance, the *eriC1* and *eriC2* single mutants were constructed in *S. anginosus* NCTC 10707 in the same manner as the *eriC1a/eriC1b* double mutant. The *eriC1* mutant hardly grew in the presence of 75 ppm NaF, while it grew similarly to wild-type NCTC 10707 in the absence of NaF (Fig. 6). On the other hand, the *eriC2* mutant showed a growth rate similar to that of wild-type NCTC 10707 in both the absence and presence of NaF. Furthermore, as shown in Fig. 6, complementation of *eriC1* rescued fluoride resistance, while the introduction of *eriC2* did not. These results demonstrated that EriC1 was responsible for the fluoride resistance in *S. anginosus*.



**Fig. 6.** Growth yields of the parent strain, eriC1 mutant, eriC2 mutant, and complemented strains in *S. anginosus* at various NaF concentrations. The y-axis represents the OD<sub>550</sub> after incubation for 16 h. Data represent the mean standard deviations of three independent experiments.

#### Identification of fluoride-resistance-related genes in S. sanguinis

The eriC2, crcB1, and crcB2 genes are present in S. sanguinis. The eriC2 gene product is a protein of 518 amino acids with no similarity to P. syringae EriC, but with a similarity of 71% to S. anginosus EriC2, which is not involved in fluoride resistance in S. anginosus. The crcB1 gene encodes a protein of 124 amino acids with amino acid sequence similarity of 51% to E. coli CrcB (127-aa protein), which is involved in fluoride resistance in E. coli, and the crcB2 gene product encodes a protein of 116 amino acids with a similarity of 62.2% to E. coli CrcB. Amino acid sequence similarity between CrcB1 and CrcB2 was 58.4%. The crcB1 and crcB2 genes exist in tandem and in the same orientation. To examine whether the inactivation of *eriC2*, *crcB1*, or *crcB2* leads to the loss of fluoride resistance, *eriC2*, *crcB1*, and *crcB2* single mutants were constructed in S. sanguinis SK36 in the same manner as the eriCla/eriClb double mutant. As shown in Fig. 7, inactivation of eriC2 did not affect fluoride resistance of wild-type SK36. On the other hand, both crcB1 and crcB2 single mutants showed extremely limited growth in the presence of 75 ppm NaF, while they grew similarly to wild-type SK36 in the absence of NaF. The crcB1/crcB2 double mutant showed a growth rate similar to that of the crcB1 or crcB2 single mutants in both the absence and presence of NaF. At a lower concentration of 25 and 50 ppm NaF, the growth rates of these three mutants were similar (Fig. 8). Furthermore, as shown in Fig. 7, complementation of *crcB1/crcB2* by crcB1/crcB2 restored fluoride resistance to wild-type levels. However, the fluoride resistance of the crcB1/crcB2 mutant complemented by either crcB1 or crcB2 did not reach the level of the mutant complemented by crcB1/crcB2. These results demonstrated that both crcB1 and crcB2 genes were critical for fluoride resistance in S. sanguinis.



**Fig. 7.** Growth yields of the parent strain, *crcB1* mutant, *crcB2* mutant, *crcB1/crcB2* double mutant, *eriC2* mutant, and complemented strains in *S. sanguinis* at various NaF concentrations. The y-axis represents the  $OD_{550}$  after incubation for 16 h. Data represent the mean standard deviations of three independent experiments.



**Fig. 8.** Growth yields of the parent strains, *crcB1* mutant, *crcB2* mutant, and *crcB1/crcB2* double mutant at a lower concentration of NaF (25 and 50 ppm). The y-axis represents the  $OD_{550}$  after incubation for 16 h. Data represent the mean standard deviations of three independent experiments.

#### Complementation of the fluoride-resistance-related genes between S. mutans and S.

#### anginosus or S. mutans and S. sanguinis

We explored whether fluoride-resistance-related genes were able to complement each other between S. mutans and S. anginosus or S. mutans and S. sanguinis. As shown in Fig. 9A, complementation of eriCla/eriClb by S. anginosus eriCl in the eriCla/eriClb double mutant rescued fluoride resistance, and the fluoride resistance ability of this complemented strain was higher than that complemented by *eriC1b*. Complementation of *eriC1* by *eriC1b* in S. anginosus also restored fluoride resistance, and the fluoride resistance ability of this transformant was lower than that complemented by *eriCl* (Fig. 9B). These results may reflect the fact that the fluoride resistance ability of S. anginosus wild-type NCTC 10707 tended to be higher than that of S. mutans wild-type UA159. Next, we performed complementation of eriC and crcB between S. mutans and S. sanguinis. Both the eriCla/eriClb mutant complemented by crcB1/crcB2 and the crcB1/crcB2 mutant complemented by eriC1b showed restored fluoride resistance (Fig 9A, C), as well as that E. coli crcB knock-out mutant rescued fluoride resistance by introduction of the fluoride-resistance- related EriCs of other bacteria (Stockbridge et al. 2012). The fluoride resistance ability of the eriCla/eriClb mutant complemented by crcB1/crcB2 was lower than that complemented by its own eriC1b, while the fluoride resistance ability of S. sanguinis wild-type SK36 tended to be higher than that of S. mutans wild-type UA159. Furthermore, we examined whether introductions of eriClb and crcB1/crcB2 into wild-type S. sanguinis and S. mutans, respectively, affects fluoride resistance in the wild-type strains. Neither introduction of *eriC1b* nor *crcB1/crcB2* increased the fluoride resistance of the wild-type strain (Fig. 9 A, C).

23



**Fig. 9.** Growth yields of the strains obtained by complementation between *S. mutans* EriC1b and *S. anginosus* EriC1 and between *S. mutans* EriC1b and *S. sanguinus* CrcB1/B2. The y-axis represents the  $OD_{550}$  after incubation for 16 h. The data represent the mean standard deviations of three independent experiments.

## Responses of the *eriC1a* mutant, the *eriC1b* mutant, and the *eriC1a / eriC1b* double mutant to bacitracin stress and environmental stressors other than fluoride

The *eriC1a* mutant, the *eriC1b* mutant, and the *eriC1a / eriC1b* double mutant of *S. mutans* were tested for its ability to grow at a low pH (5.5), an elevated temperature (44°C), and high osmotic pressure (NaCl at a final concentration of 0.585 M). As shown in Fig. 10, these mutants showed growth rates similar to that of wild-type UA159 under the above stress conditions.



**Fig. 10.** Growth yields of the parent strain, UA159, the *eriC1a* mutant, the *eriC1b* mutant, and the *eriC1a/eriC1b* double mutant at pH 7.5 or 5.5 (A), at 37°C and 44°C (B), in BHI broth supplemented with 0.5 M NaCl (C), and in BHI broth supplemented with 5U/mL bacitracin (D). The y-axis represents the  $OD_{550}$  after incubation for 16 h. The data represent the mean standard deviations of three independent experiments.

#### DISCUSSION

ClC-type anion-selective channels are widespread throughout eukaryotic organisms and play a crucial role in controlling the ionic composition of the cytoplasm and the volume of cells, as well as the regulation of membrane electrical excitability. Although these channels may conduct other anions (e.g., I<sup>-</sup> or NO<sub>3</sub><sup>-</sup>) better than Cl<sup>-</sup>, they are often called chloride channels because Cl<sup>-</sup> is the most abundant anion in organisms (Jentsch et al. 2002). BLAST homology searches revealed that many microbial genomes contain members of the CIC family, and the prokaryotic CIC channel was designated an E. coli-derived CIC chloride channel homolog (EriC) (Maduke et al. 1999). E. coli EriC has been confirmed experimentally to function as a CIC CI<sup>-</sup> channel. On the other hand, P. syringae EriC and Clostridium difficile EriC have been reported to be associated with fluoride resistance (Baker et al. 2012). The crcB (confers resistance to camphor B) gene has previously been implicated in resistance to camphor-induced chromosome decondensation. However, it has recently been reported that an E. coli crcB knock-out mutant became sensitive to fluoride, and crcB is predicted to code for membrane proteins belonging to a superfamily composed predominantly of transporters (Baker et al. 2012). Furthermore, it was shown that the eukaryotic crcB homolog [renamed FEX (fluoride exporter)] encodes a previously unrecognized class of fluoride exporters required for survival under standard environmental conditions (Li et al. 2013).

Oral streptococci are classified into four groups; namely, the mutans, the salivarius, the mitis, and the anginosus groups, according to their 16S rRNA sequences (Kawamura et al. 1995). The present study suggests that oral streptococci possess either *eriC1* or *crcB*, which are associated with fluoride resistance. Among oral streptococci examined in this study, all

*Streptococcus* species belonging to the mutans and the salivarius groups contained EriC1. Nine of 11 *Streptococcus* species belonging to the mitis group possessed CrcB and the other 2 species possessed EriC1. Although a distinct relationship between the type of genes (*eriC1* or *crcB*) and the fluoride-resistance level was not observed, oral streptococci with *eriC1* were prone to higher fluoride resistance than those with *crcB*. The difference in the fluoride-resistance level related to the gene types suggests that the mechanisms through EriC1 and CrcB are not identical.

Baker et al. (2012) showed that P. syringae EriC and C. difficile EriC associated with fluoride resistance commonly carry a specific set of amino acids in the conserved anion selectivity filter region (Dutzler, et al., 2002), which are distinct from validated chloride-specific EriC proteins (Table). In this study, two novel genes encoding EriC1b of S. mutans and EriC1 of S. anginosus were shown to be involved in fluoride-resistance, and subsequently seven EriC1s in other oral streptococci were predicted to be responsible for fluoride resistance of these organisms based on amino acid sequence homology. Our table shows amino acid sequences of the conserved anion selectivity filter region of the 18 EriC homologs; 3 EriCs are Cl<sup>-</sup> channels and 15 EriCs are predicted to be F<sup>-</sup> channels. All EriC1s of oral streptococci possessed a similar set of amino acid sequences to those of the F<sup>-</sup> channel EriCs, but they were distinct from those of Cl<sup>-</sup> channel EriCs. Although it is possible that EriC1s in oral streptococci may be generally responsible for the fluoride resistance, the role of EriC1s from oral streptococci other than S. mutans and S. anginosus in fluoride resistance should be examined. The amino acid sequences of the region in the streptococci were more similar to those of C. difficile, Eubacterium ventriosum, and Lactococcus lactis EriCs than those of *P. syringae*, *Pirellula staleyi*, and *Ralstonia picketti* EriCs, possibly due to the

difference between Gram-positive and Gram-negative bacteria. On the other hand, EriC2s identified in all oral streptococci examined in this study, except for *S. mutans*, possessed amino acid sequences with conserved Cl<sup>-</sup> channels (data not shown), suggesting that EriC2 may be related to Cl<sup>-</sup> channels.

Complementation between S. mutans EriC1b and S. anginosus EriC1 and that between S. mutans EriC1b and S. sanguinus CrcB1/B2 were confirmed in this study. Introduction of S. anginosus EriC1 into an EriC1s-knockout mutant of S. mutans increased fluoride resistance when compared with the same complementation experiment with EriC1b. On the other hand, when EriC1b was transferred into the EriC1-knockout mutant of S. anginosus, fluoride resistance of the complemented strain was lower than that complemented with its own EriC1. The fluoride resistance level of S. anginosus NCTC10707 was higher than that of S. mutans UA159, and the differences in fluoride resistance levels between the above species may be attributed to differences in the kinetics of ion transport between EriC1s. Moreover, the EriC1s-knockout mutant of S. mutans complemented by S. sanguinis CrcB1/B2 and the CrcB1/CrcB2-knockout mutant complemented by S. mutans EriC1b showed restored fluoride resistance. However, neither introduction of S. sanguinis CrcB1/B2 into wild-type S. mutans nor S. mutans EriC1b into wild-type S. sanguinis increased the fluoride resistance level of the wild-type strain. It is possible that EriC1 and CrcB1/B2 may be responsible for fluoride resistance in oral streptococci by sharing pathways.

Fluoride application is highly effective in preventing dental caries. On the other hand, we must consider that fluoride has anti-bacterial effects. The frequent application of fluoride results in the emergence of fluoride resistant strains, leading to the dysbiosis of oral microflora. This disturbance may affect not only oral health conditions, but also general

29

health conditions. Exploring the mechanism of fluoride resistance may contribute to countermeasures against the risk of emergence of the fluoride-resistant strains in oral microflora.

Liao et al. (Liao, *et al.*, 2015) reported that two SNPs related to fluoride resistance were identified in the genome of the fluoride resistant strain *S. mutans* C180-2FR. These were located in the region of the gene cluster composed of SMU\_1291, *eriC1a*, and *eriC1b*; one in its promoter region and the other in *eriC1b*. Expression of the cluster was approximately 10-fold higher in C180-2FR than in the parent strain C180-2. However, we demonstrated that *eriC1b* is involved in fluoride resistance of *S. mutans*, while *eriC1a* is not. It remains unknown why *S. mutans* possesses two EriC1s, unlike other oral streptococci, and why *eriC1a* is not involved in fluoride resistance in *S. mutans*, even though the two EriC1s showed amino acid sequence similarities of 74%.

Inactivation of *eriC1a* or/and *eriC1b* did not affect the resistance to other stress conditions, such as low pH and high osmotic pressure, in *S. mutans*. On the other hand, the function of EriC1a is not clear, while EriC1b is obviously involved in fluoride resistance. EriC1a might be responsible for the transport of other molecules. Anyhow, it is interesting to investigate, including the additional role of EriC1b, what kind of molecules EriC1a transports in order to response to environmental stressors other than those examined in the present study.

**Table.** Amino acid sequences of the conserved anion selectivity filter region of several EriCs. Eight EriCs (shadowed) were shown to be involved in fluoride resistance (Baker et al. 2012) (Stockbridge et al. 2012). Boxes show the amino acid sequences associated with fluoride resistance. Related EriCs commonly contain a specific set of amino acids, which are distinct from validated chloride-specific EriCs.

Substrate	Organism	CΓ selectivity filter residues
Cl-	human (ClC-1)	G S G I P G K E G P G G F M P Y
Cl⁻	Escherichia coli	G S G I P G R E G P G I F A P Y
Cl⁻	Salmonella typhimurium	G S G I P G R E G P G I F A P Y
F-	Pseudomonas syringae	G N N L I G R E G T G E V T P Y
F-	Pirellula staleyi	G N N L L G R E G T G E V T P Y
F-	Ralstonia picketti	G N N L L G R E G T G E V T P Y
F	Clostridium difficile	GMNLIGREGVGEVTPY
F	Eubacterium ventriosum	G M N L V G R E G V G E V T P Y
F	Lactococcus lactis	GMGLIGREGVGEVTPY
F⁻	Streptococcus mutans	GMGLIGREGVGEVTPY
F-	Streptococcus anginosus	GMTLIGREGVGEVTPY
F⁻	Streptococcus sobrinus	G M G L V G R E G V G E V T P Y
F	Streptococcus australis	G M E L L G R E G V G E V T P Y
F	Streptococcus parasanguinis	GMGLIGREGVGEVTPY
F	Streptococcus salivarius	GMGLIGREGVGEVTPY
F	Streptococcus vestibularis	GMGLIGREGVGEVTPY
F	Streptococcus constellatus	GMTLIGREGVGEVTPY
F	Streptococcus intermedius	GMTLIGREGVGEVTPY

#### ACKNOLEDGMENTS

On the completion of this thesis, I would like to express my deepest gratitude to Professor Yoshihisa Yamashita of his invaluable assistance, resourceful counsel and enlightening advice guided me in every stage of the writing of this work. I am also tremendously indebted to my supervisor, Dr. Yukie Shibata, who is Assistant Professor of Section of Preventive and Public Health Dentistry, Division of Oral Health, Growth and Development, Faculty of Dental Science Kyushu University, for the huge amount of time and work she spent throughout.

My gratefulness also goes t to Dr. Toru Takeshita, who is Associate Professor of Section of Preventive and Public Health Dentistry, Division of Oral Health, Growth and Development, Faculty of Dental Science Kyushu University; Dr. Michiko Furuta and Dr. Kenji Takeuchi who are Assistant Professors from Section of Preventive and Public Health Dentistry, Division of Oral Health, Growth and Development, Faculty of Dental Science Kyushu University, for their warm-hearted help from which I benefited enormously.

Finally, I thank all the other laboratory members who supported and brightened my life and studies here.

32

#### REFERENCES

- Baker JL, Sudarsan N, Weinberg Z, Roth A, Stockbridge RB, Breaker RR. 2012. Widespread genetic switches and toxicity resistance proteins for fluoride. Science. 335(13):233-235.
- Brown LR, White JO, Horton IM, Dreizen S, Streckfuss JL. 1983. Effect of continuous fluoride gel use on plaque fluoride retention and microbial activity. J Dent Res. 62(6):746-751.
- Bunick, Kashket S. 1981. Enolases from fluoride-sensitive and fluoride-resistant streptococci. Infect Immun. 34(3):856-863.
- Dutzler R, Campbell EB, Cadene M, Chait BT, MacKinnon R. 2002. X-ray structure of a CIC chloride channel at 3.0 Å reveals the molecular basis of anion selectivity. Nature. 415(17): 287-294.
- Hamilton IR. 1969. Growth characteristics of adapted and ultraviolet-induced mutants of *Streptococcus salivarius* resistant to sodium fluoride. Can J Microbiol. 15(3):287-295.
- Hanahan D. 1983. Studies on transformation of *Escherichia coli* with plasmids. J Mol Biol. 166(4):557-580.
- Hoelscher GL, Hudson MC. 1996 Characterization of an unusual fluoride-resistant *Streptococcus mutans* isolate. Curr microbiol. 32(3):156-161.
- Jentsch TJ, Stein V, Weinreich F, Zdebik AA. 2002. Molecular structure and physiological function of chloride channels. Physiol rev. 82(2):503-568.
- Kawada-Matsuo M, Shibata Y, Yamashita Y. 2009. Role of two component signaling response regulators in acid tolerance of *Streptococcus mutans*. Oral Microbiol Immunol 24(2):173-176.

- Kawamura Y, Hou XG, Sultana F, Miura H, Ezaki T. 1995. Determination of 16S rRNA sequences of *Streptococcus mitis* and *Streptococcus gordonii* and phylogenetic relationships among members of the genus *Streptococcus*. Int J Sys Bacteriol. 45(2):406-408.
- Li S, Smith KD, Davis JH, Gordon PB, Breaker RR, Strobel SA. 2013. Eukaryotic resistance to fluoride toxicity mediated by a widespread family of fluoride export proteins. Proc Natl Acad Sci USA. 110(47):19018-19023.
- Li D, Shibata Y, Takeshita T, Yamashita Y. 2012. A novel gene involved in the survival of *Streptococcus mutans* under stress conditions. Appl Environ Microbiol. 80(1):97-103.
- Liao Y, Chen J, Brandt BW, Zhu Y, Li J, van Loveren C, Deng DM. 2015. Identification and functional analysis of genome mutations in a fluoride-resistant *Streptococcus mutans* strain. PloS One. 10(4):e0122630.
- Maduke M, Pheasant DJ, Miller C. 1999. High-level expression, functional reconstitution, and quaternary structure of a prokaryotic ClC-type chloride channel. J Gen Physiol. 114(5):713-722.
- Maltz M, Emilson CG. 1982. Susceptibility of oral bacteria to various fluoride salts. J Dent Res. 61(6):786-790.
- Marquis RE, Clock SA, Mota-Meira M. 2003. Fluoride and organic weak acids as modulators of microbial physiology. FEMS Microbiol Rev. 26(5):493-510.
- Mayhew RR, Brown LR. 1981. Comparative effect of SnF<sub>2</sub>, NaF, and SnCl<sub>2</sub> on the growth of *Streptococcus mutans*. J Dent Res. 60(10):1809-1814.
- Moritani K, Takeshita T, Shibata Y, Ninomiya T, Kiyohara Y, Yamashita Y. 2015. Acetaldehyde production by major oral microbes. Oral Dis. 21(6):748-754.

- Perry D, Wondrack LM, Kuramitsu HK. 1983. Genetic transformation of putative cariogenic properties in *Streptococcus mutans*. Infect Immun. 41: 722-727.
- Rickard AH, Gilbert P, High NJ, Kolenbrander PE, Handley PS. 2003. Bacterial coaggregation: an integral process in the development of multi-species biofilms. Trends Microbiol. 11(2): 94-100.
- Sheng JY, Liu Z. 2000. Acidogenicity and acidurance of fluoride-resistant *Streptococcus sobrinus* in vitro. Chin J Dent Res. 3(2):7-14.

Silver S. 1996. Bacterial resistances to toxic metal ions-a review. Gene. 179(1):9-19.

- Stockbridge RB, Lim HH, Otten R, Williams C, Shane T, Weinberg Z, Miller C. 2012. Fluoride resistance and transport by riboswitch-controlled CLC antiporters. Proc Natl Acad Sci USA. 109(38):15289-15294.
- Streckfuss JL, Perkins D, Horton IM, Brown LR, Dreizen S, Graves L. 1980. Fluoride resistance and adherence of selected strains of *Streptococcus mutans* to smooth surfaces after exposure to fluoride. J Dent Res. 59(2):151-158.
- Takeshita T, Yasui M, Shibata Y, Furuta M, Saeki Y, Eshima N, Yamashita Y. 2015. Dental plaque development on a hydroxyapatite disk in young adults observed by using a barcoded pyrosequencing approach. Sci Rep. 5:8136.
- Van Loveren C, Spitz LM, Buijs JF, Ten Cate JM, Eisenberg AD. 1991. In vitro demineralization of enamel by F-sensitive and F-resistant mutans streptococci in the presence of 0, 0.05, or 0.5 mmol/L NaF. J Dent Res. 70(12):1491-1496.
- Van Loveren C, Van de Plassche-Simons YM, De Soet JJ, De Graaff J, Ten Cate JM. 1991. Acidogenesis in relation to fluoride resistance of *Streptococcus mutans*. Oral Microbiol Immunol. 6(5):288-291.