

# A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

Takaesu, Toshimitsu  
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/17859>

---

出版情報 : MI Preprint Series. 24, 2010-07-06. 九州大学大学院数理学研究院  
バージョン :  
権利関係 :



# **MI Preprint Series**

**Kyushu University  
The Global COE Program  
Math-for-Industry Education & Research Hub**

## **A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra**

**Toshimitsu Takaesu**

**MI 2010-24**

( Received July 6, 2010 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

Toshimitsu TAKAESU

*Faculty of Mathematics, Kyushu University,  
Fukuoka, 812-8581, Japan*

**Abstract.** In this article we consider linear operators satisfying a generalized commutation relation of a type of the Heisenberg-Lie algebra. It is proven that a generalized inequality of the Hardy's uncertainty principle lemma follows. Its applications to time operators and abstract Dirac operators are also investigated.

**Key words :** weak commutation relations, Heisenberg-Lie algebra, time operators, Hamiltonians, time-energy uncertainty relation, Dirac operators, essential self-adjointness.

**MSC 2010 :** 81Q10, 47B25, 46L60.

## 1 Introduction and Results

In this article we investigate a norm-inequality of the linear operators which obey a generalized weak commutation relation of a type of the Heisenberg-Lie algebra, and consider its application to the theory of the time operator [7, 2], and an abstract Dirac operator. Let  $\mathbf{X} = \{X_j\}_{j=1}^N$ ,  $\mathbf{Y} = \{Y_j\}_{j=1}^N$  and  $\mathbf{Z} = \{Z_j\}_{j=1}^N$  be symmetric operators on a Hilbert space  $\mathcal{H}$ . The weak commutator of operators  $A$  and  $B$  is defined for  $\psi \in \mathcal{D}(A) \cap \mathcal{D}(B)$  and  $\phi \in \mathcal{D}(A^*) \cap \mathcal{D}(B^*)$  by

$$[A, B]^w(\phi, \psi) = (A^* \phi, B\psi) - (B^* \phi, A\psi).$$

Here the inner product has a linearity of  $(\eta, \alpha\psi + \beta\phi) = \alpha(\eta, \psi) + \beta(\eta, \phi)$  for  $\alpha, \beta \in \mathbf{C}$ . We assume that  $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$  satisfies the following conditions.

(A.1)  $Z_j$ ,  $1 \leq j \leq N$ , is bounded operator.

(A.2) Let  $\mathcal{D}_{\mathbf{X}} = \cap_{j=1}^N \mathcal{D}(X_j)$  and  $\mathcal{D}_{\mathbf{Y}} = \cap_{j=1}^N \mathcal{D}(Y_j)$ . It follows that for  $\phi, \psi \in \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$ ,

$$\begin{aligned} [X_j, Y_l]^w(\phi, \psi) &= \delta_{j,l}(\phi, iZ_j\psi), \\ [X_j, Z_l]^w(\phi, \psi) &= [Y_j, Z_l]^w(\phi, \psi) = 0 \\ [X_j, X_l]^w(\phi, \psi) &= [Y_j, Y_l]^w(\phi, \psi) = [Z_j, Z_l]^w(\phi, \psi) = 0. \end{aligned}$$

Note that  $[Z_j, Z_l]\psi = 0$  follows for  $\psi \in \mathcal{H}$ , since  $Z_j$ ,  $j = 1, \dots, N$ , is bounded. In this article we consider an generalization of the inequality

$$\int_{\mathbf{R}^N} \frac{1}{|\mathbf{r}|^2} |u(\mathbf{r})|^2 d\mathbf{r} \leq \frac{4}{(N-2)^2} \int_{\mathbf{R}^N} |\nabla u(\mathbf{r})|^2 d\mathbf{r}, \quad N \geq 3.$$

This inequality is a basic one of Hardy's uncertainty principle inequalities. For Hardy's uncertainty inequalities, refer to e.g. [5, 6, 13].

Let us introduce the additional conditions.

**(A.3)**  $X_j$  is self-adjoint for all  $1 \leq j \leq N$ .

**(A.4)**  $X_i$  and  $Z_l$  strongly commutes for all  $1 \leq j \leq N$  and  $1 \leq l \leq N$ .

Since  $Z_j$ ,  $j = 1, \dots, N$ , is bounded self-adjoint operator, we can set  $\lambda_{\min}(\mathbf{Z})$  and  $\lambda_{\max}(\mathbf{Z})$  by

$$\begin{aligned}\lambda_{\min}(\mathbf{Z}) &= \min_{1 \leq j \leq N} \inf \sigma(Z_j), \\ \lambda_{\max}(\mathbf{Z}) &= \max_{1 \leq j \leq N} \sup \sigma(Z_j),\end{aligned}$$

where  $\sigma(O)$  denotes the spectrum of the operator  $O$ .

**Theorem 1** Assume **(A.1)**-**(A.4)**. Let  $\Psi \in \mathcal{D}(|\mathbf{X}|^{-1}) \cap \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$ . Then the following **(1)** and **(2)** hold  
**(1)** If  $N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z}) > 0$ , it follows that

$$\left\| |\mathbf{X}|^{-1} \Psi \right\|^2 \leq \frac{4}{(N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z}))^2} \sum_{j=1}^N \left\| Y_j \Psi \right\|^2. \quad (1)$$

**(2)** If  $2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}) > 0$ , it follows that

$$\left\| |\mathbf{X}|^{-1} \Psi \right\|^2 \leq \frac{4}{(2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}))^2} \sum_{j=1}^N \left\| Y_j \Psi \right\|^2. \quad (2)$$

Before proving Theorem 1, let us consider the replacement of  $\mathbf{X}$  and  $\mathbf{Y}$  in Theorem 1. Let us introduce the following conditions substitute for **(A.3)** and **(A.4)**.

**(A.5)**  $Y_j$  is self-adjoint for all  $1 \leq j \leq N$ .

**(A.6)**  $Y_i$  and  $Z_l$  strongly commutes for all  $1 \leq j \leq N$  and  $1 \leq l \leq N$ .

It is seen from **(A.2)**, that

$$[Y_j, X_l]^w(\phi, \psi) = \delta_{j,l}(\phi, i(-Z_j)\psi), \quad \phi, \psi \in \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}. \quad (3)$$

Note that  $\inf \sigma(-Z_j) = -\sup \sigma(Z_j)$  and  $\sup \sigma(-Z_j) = -\inf \sigma(Z_j)$  follow. Then we obtain a following corollary :

**Corollary 2** Assume (A.1)-(A.2) and (A.5)-(A.6). Let  $\Psi \in \mathcal{D}(|\mathbf{Y}|^{-1}) \cap \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$ . Then the following (1) and (2) hold.

(1) If  $2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}) > 0$ , it follows that

$$\left\| |\mathbf{Y}|^{-1}\Psi \right\| \leq \frac{4}{(2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}))^2} \sum_{j=1}^N \left\| X_j \Psi \right\|^2. \quad (4)$$

(2) If  $N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z}) > 0$ , it follows that

$$\left\| |\mathbf{Y}|^{-1}\Psi \right\| \leq \frac{4}{(N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z}))^2} \sum_{j=1}^N \left\| X_j \Psi \right\|^2. \quad (5)$$

**(Proof of Theorem 1)**

(1) Let  $\Psi \in \mathcal{D}(|\mathbf{X}|^{-1}) \cap \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$ . For  $\varepsilon > 0$  and  $t > 0$ , it is seen that

$$\left\| (Y_j - itX_j(\mathbf{X}^2 + \varepsilon)^{-1}) \Psi \right\|^2 = \|Y_j \Psi\|^2 - it[Y_j, X_j(\mathbf{X}^2 + \varepsilon)^{-1}]^w(\Psi, \Psi) + t^2 \|X_j(\mathbf{X}^2 + \varepsilon)^{-1} \Psi\|^2. \quad (6)$$

We see that

$$[Y_j, X_j(\mathbf{X}^2 + \varepsilon)^{-1}]^w(\Psi, \Psi) = [Y_j, X_j]^w(\Psi, (\mathbf{X}^2 + \varepsilon)^{-1}\Psi) + [Y_j, (\mathbf{X}^2 + \varepsilon)^{-1}]^w(X_j \Psi, \Psi). \quad (7)$$

From (A.2) and (A.4), we obtain that

$$[Y_j, X_j]^w(\Psi, (\mathbf{X}^2 + \varepsilon)^{-1}\Psi) = -i((\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi, Z_j(\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi). \quad (8)$$

Note that for a symmetric operator  $A$  and the non-negative symmetric operator  $B$ , the resolvent formula  $[A, (B + \lambda)^{-1}]^w(v, u) = [B, A]^w((B + \lambda)^{-1}v, (B + \lambda)^{-1}u)$  for  $\lambda > 0$  follows. Then by using this formula, (A.2) and (A.4) yield that

$$[Y_j, (\mathbf{X}^2 + \varepsilon)^{-1}]^w(X_j \Psi, \Psi) = 2i(X_j(\mathbf{X}^2 + \varepsilon)^{-1}u, Z_j X_j(\mathbf{X}^2 + \varepsilon)^{-1}u) \quad (9)$$

Since  $\left\| (Y_j - itX_j(\mathbf{X}^2 + \varepsilon)^{-1}) u \right\|^2 \geq 0$  and  $t > 0$ , we see from (7), (8) and (9) that

$$\begin{aligned} & \|Y_j \Psi\|^2 \\ & \geq -t^2 \|X_j(\mathbf{X}^2 + \varepsilon)^{-1}u\|^2 + t((\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi, Z_j(\mathbf{X}^2 + \varepsilon)^{-1/2}u) - 2t(X_j(\mathbf{X}^2 + \varepsilon)^{-1}u, Z_j X_j(\mathbf{X}^2 + \varepsilon)^{-1}\Psi) \\ & \geq \left( -t^2 - 2t\lambda_{\max}(\mathbf{Z}) \right) \|X_j(\mathbf{X}^2 + \varepsilon)^{-1}u\|^2 + t\lambda_{\min}(\mathbf{Z}) \|(\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi\|. \end{aligned} \quad (10)$$

Then we have that

$$\sum_{j=1}^N \|Y_j \Psi\|^2 \geq \left( -t^2 - 2t\lambda_{\max}(\mathbf{Z}) \right) \left\| |\mathbf{X}|(\mathbf{X}^2 + \varepsilon)^{-1}\Psi \right\|^2 + tN\lambda_{\min}(\mathbf{Z}) \|(\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi\|. \quad (11)$$

Note that  $\lim_{\varepsilon \rightarrow 0} \left\| |\mathbf{X}|(\mathbf{X}^2 + \varepsilon)^{-1}\Psi \right\|^2 = \left\| |\mathbf{X}|^{-1}\Psi \right\|^2$  and  $\lim_{\varepsilon \rightarrow 0} \|(\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi\| = \left\| |\mathbf{X}|^{-1}\Psi \right\| = 0$  follow from the spectral decomposition theorem. Then we have

$$\sum_{j=1}^N \|Y_j \Psi\|^2 \geq (-t^2 + (N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z}))t) \left\| |\mathbf{X}|^{-1}\Psi \right\|. \quad (12)$$

By taking  $t = \frac{N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z})}{2} > 0$  in the right side of (12), we obtain **(1)**.

**(2)** By computing  $\| (Y_j + itX_j(\mathbf{X}^2 + \varepsilon)^{-1}) \Psi \|^2$  for  $t > 0$  and  $\varepsilon > 0$ , in a similar way of **(1)**, we see that

$$\begin{aligned} & \|Y_j \Psi\|^2 \\ & \geq -t^2 \|X_j(\mathbf{X}^2 + \varepsilon)^{-1}u\|^2 - t((\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi, Z_j(\mathbf{X}^2 + \varepsilon)^{-1/2}u) + 2t(X_j(\mathbf{X}^2 + \varepsilon)^{-1}u, Z_jX_j(\mathbf{X}^2 + \varepsilon)^{-1}\Psi) \\ & \geq \left( -t^2 + 2t\lambda_{\min}(\mathbf{Z}) \right) \|X_j(\mathbf{X}^2 + \varepsilon)^{-1}u\|^2 - t\lambda_{\max}(\mathbf{Z}) \|(\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi\|. \end{aligned} \quad (13)$$

Then by taking  $\varepsilon \rightarrow 0$  in the right side of (13), it follows that

$$\sum_{j=1}^N \|Y_j \Psi\|^2 \geq \left( -t^2 + (2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}))t \right) \| |\mathbf{X}|^{-1}\Psi \|. \quad (14)$$

By taking  $t = \frac{(2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}))}{2} > 0$  in (14), we obtain **(2)**. ■

## 2 Applications

### 2.1 Time-Energy Uncertainty inequality

In this subsection we consider an applicaion to the theory of time operators [2, 7]. Let  $H$ ,  $T$ , and  $C$  be linear operators on a Hilbert space  $H$ . It is said that  $H$  has the weak time operator  $T$  with the uncommutative factor  $C$  if  $(H, T, C)$  satisfy the following conditions.

**(T.1)**  $H$  and  $T$  are symmetric.

**(T.2)**  $C$  is bounded and self-adjoint.

**(T.3)** It follows that for  $\phi, \psi \in \mathcal{D}(H) \cap \mathcal{D}(T)$ ,

$$[T, H]^w(\phi, \psi) = (\phi, C\psi).$$

**(T.4)**

$$\delta_C := \inf_{\psi \in (\ker C)^\perp \setminus \{0\}} \frac{|(\Psi, C\Psi)|}{\|\Psi\|^2} > 0.$$

Assume that  $(H, T, C)$  satisfies **(T.1)-(T.4)**. Then by using  $\|Au\| \|Bu\| \geq |\operatorname{Im}(Au, Bu)| \geq \frac{1}{2} |[A, B]^w(u, u)|$ , it is seen that  $(H, T, C)$  satisfies the time-energy uncertainty inequality ([2], Proposition 4.1):

$$\frac{\| (H - \langle H \rangle_\psi) \psi \| \| (T - \langle T \rangle_\psi) \psi \|}{\|\psi\|^2} \geq \frac{\delta_C}{2}, \quad \psi \in \mathcal{D}(H) \cap \mathcal{D}(T), \quad (15)$$

where  $\langle O \rangle_\psi = (\psi, O\psi)$ . From **(2)** in Theorem 1 and **(1)** in Corollary 2, we obtain another type of the inequality between  $T$  and  $H$  :

**Corollary 3 (Time-Energy Uncertainty Inequalities)**

Assume (T.1)-(T.3). Then the following (i) and (ii) hold.

(i) If  $T$  is self-adjoint,  $C$  and  $T$  strongly commute, and  $\sup \sigma(C) < 2 \inf \sigma(C)$ , it follows that for  $\psi \in \mathcal{D}(|T|^{-1}) \cap \mathcal{D}(T) \cap \mathcal{D}(H)$ ,

$$\left\| |T|^{-1} \psi \right\| \leq \frac{2}{2 \inf \sigma(C) - \sup \sigma(C)} \left\| H \Psi \right\|. \quad (16)$$

(ii) If  $H$  is self-adjoint,  $C$  and  $H$  strongly commute, and  $\sup \sigma(C) < 2 \inf \sigma(C)$ , it follows that for  $\psi \in \mathcal{D}(|H|^{-1}) \cap \mathcal{D}(H) \cap \mathcal{D}(T)$ ,

$$\left\| |H|^{-1} \psi \right\| \leq \frac{2}{2 \inf \sigma(C) - \sup \sigma(C)} \left\| T \Psi \right\|. \quad (17)$$

## 2.2 Abstract Dirac Operators with Coulomb Potential

Next let us consider the application to abstract Dirac operators. We consider the self-adjoint operators  $\mathbf{P} = \{P_j\}_{j=1}^N$  and  $\mathbf{Q} = \{Q_j\}_{j=1}^N$  on a Hilbert space  $\mathcal{H}$ . Let us set a subspace  $\mathcal{D} \subset \cap_{j,l} (\mathcal{D}(P_j) \cap \mathcal{D}(Q_l))$ . It is said that  $(\mathcal{H}, \mathcal{D}, \mathbf{P}, \mathbf{Q})_N$  is the weak representation of the CCR with degree  $N$ , if  $\mathcal{D}$  is dense in  $\mathcal{H}$  and it follows that for  $\phi, \psi \in \mathcal{D}$ ,

$$\begin{aligned} [P_j, Q_l]^w(\phi, \psi) &= i\delta_{j,l}(\phi, \psi), \\ [P_j, P_l]^w(\phi, \psi) &= [Q_j, Q_l]^w(\phi, \psi) = 0. \end{aligned}$$

Let us define an abstract Dirac operator as follows. Let  $(\mathcal{H}, \mathcal{D}, \mathbf{P}, \mathbf{Q})_3$  be the weak representation of the CCR with degree three. Let  $\mathbf{A} = \{A_j\}_{j=1}^3$  and  $B$  be the bounded self-adjoint operators on a Hilbert space  $\mathcal{K}$ . Here  $\mathbf{A} = \{A_j\}_{j=1}^3$  and  $B$  satisfy the canonical anti-commutation relations  $\{A_j, A_l\} = 2\delta_{j,l}$ ,  $\{A_j, B\} = 0$ ,  $B^2 = I_{\mathcal{K}}$  where  $I_{\mathcal{K}}$  is the identity operator on  $\mathcal{K}$ . The state Hilbert space is defined by  $\mathcal{H}_{\text{Dirac}} = \mathcal{K} \otimes \mathcal{H}$ . The free abstract Dirac operator is defined by

$$H_0 = \sum_{j=1}^3 A_j \otimes P_j + B \otimes M.$$

Here we assume the following condition.

**(D.1)**  $P_j$  and  $P_l$  strongly commute for  $1 \leq j \leq 3$ ,  $1 \leq l \leq 3$ .  $P_j$ ,  $1 \leq j \leq 3$ , and  $M$  strongly commute.

Then it is seen that  $H_0^2 \Psi = (\mathbf{P}^2 + M^2) \Psi$  for  $\Psi \in \mathcal{D}$ . The abstract Dirac Operator with the Coulomb potential is defined by

$$H(\kappa) = H_0 + \kappa I_{\mathcal{K}} \otimes |\mathbf{Q}|^{-1},$$

where  $\kappa \in \mathbf{R}$  is a parameter called the coupling constant. We assume that the following condition

**(D.2)** It follows that  $\mathcal{D} \subset \mathcal{D}(|\mathbf{Q}|^{-1})$ .

Then it follows from (1) in Theorem 1 that for  $\psi \in \mathcal{D}$ ,

$$\|I_{\mathcal{K}} \otimes |\mathbf{Q}|^{-1} \psi\|^2 \leq 4 \sum_{j=1}^3 \|P_j \Psi\|^2 \leq 4 \|H_0 \Psi\|^2.$$

Hence by the Kato-Rellich theorem, we obtain the following corollary.

**Corollary 4** Assume (D.1) and (D.2). Then for  $|\kappa| < \frac{1}{2}$ ,  $H(\kappa)$  is essentially self-adjoint on  $\mathcal{D}$ .

## Acknowledgments

It is a pleasure to thank assistant professor Akito Suzuki and associate professor Fumio Hiroshima for their advice and comments.

## References

- [1] A.Arai, *Mathematical principles of quantum phenomena*, Asakura-shoten, 2005. (in japanese)
- [2] A.Arai, Generalized weak Weyl relation and decay of quantum dynamics, *Rev. Math. Phys.* **17** (2005) 1071-1109.
- [3] A.Arai, Heisenberg Operators, invariant domains and Heisenberg equations of motion, *Rev. Math. Phys* **19** (2007) 1045-1069.
- [4] A.Arai, Spectrum of time operators. *Lett. Math. Phys.* **80** 211-221 (2007).
- [5] G.B.Folland and A.Sitaram, The uncertainty principle : A mathematical survey, *J. Fourier Anal. Appl.* **3** (1997) 207-238.
- [6] V.Havin and B.Jorjick, *The uncertainty principle in harmonic analysis*, Springer 1994.
- [7] M.Miyamoto, A generalized Weyl relation approach to the time operator and its connection to the survival probability, *J. Math. Phys* **42** (2001) 1038-1052.
- [8] J.G.Muga, R.S.Mayato, and I.L.Egusquiza (eds.), *Time in quantum mechanics*. Springer 2002.
- [9] P.Pfeifer and J. Frölich, Generalized time-energy uncertainty relations and bounds on lifetimes of resonances, *Rev. Mod. Phys.* **67** (1995) 759-779.
- [10] M.Reed and B.Simon, *Methods of Modern Mathematical Physics Vol.II*, Academic Press, 1979.
- [11] K.Schmüdgen, On the Heisenberg commutation relation. I, *J. Funct. Anal.* **50** (1983) 8-49.
- [12] B. Thaller, *The Dirac equation*, Springer, 1992.
- [13] S.Thangavelu, *An introduction to the uncertainty principle : Hardy's theorem on Lie groups*, Birkhäuser, 2004.



# List of MI Preprint Series, Kyushu University

The Global COE Program  
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA  
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO  
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU  
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI  
Torsion points of abelian varieties with values in infinite extensions over a  $p$ -adic field
- MI2008-6 Yoshiyuki TOMIYAMA  
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI  
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA  
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA  
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA  
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA  
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO  
On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA  
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA  
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA  
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO  
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI  
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI  
Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI  
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI  
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA  
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI  
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI  
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA  
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO  
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA  
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA  
Hypergeometric  $q$ -functions of the  $q$ -Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA  
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA  
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI  
Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE  
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO  
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI  
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI  
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO  
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA  
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO  
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI  
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO  
On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection

- MI2009-26 Manabu YOSHIDA  
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI  
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA  
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA  
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA  
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI  
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA  
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA  
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA  
Projective reduction of the discrete Painlevé system of type  $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI  
Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA  
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA  
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA  
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA  
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI  
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI  
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI  
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA  
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO  
Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA  
An algebraic approach to underdetermined experiments

- MI2010-10 Kei HIROSE & Sadanori KONISHI  
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA  
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU  
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA  
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI  
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO  
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE  
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU  
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA  
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE  
Lagrangian approach to weakly nonlinear stability of an elliptical flow

MI2010-21 Hiroki MASUDA

Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test

MI2010-22 Toshimitsu TAKAESU

A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs

MI2010-23 Takahiro ITO, Mitsuhiko FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI

Composition, union and division of cellular automata on groups

MI2010-24 Toshimitsu TAKAESU

A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra