Assessment of Cell Membrane Thermostability and Silicon Supplement on Dendrobium Lucky Girl

Wu, Shing-Kuan

Department of Horticulture, National Chung Hsing University | Institute of Tropical Agriculture, Kyushu University

Miyajima, Ikuo

Institute of Tropical Agriculture, Kyushu University

Huang, Kuang-Liang

Department of Horticultural Science, National Chiayi University | Institute of Tropical Agriculture, Kyushu University

Kuo, Ya-Chin

Department of Horticulture, National Chung Hsing University | Institute of Tropical Agriculture, Kyushu University

他

<https://doi.org/10.5109/1833814>

出版情報：九州大学大学院農学研究院紀要. 62 (2), pp.337-343, 2017-09-08. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Assessment of Cell Membrane Thermostability and Silicon Supplement on *Dendrobium* Lucky Girl

Shing-Kuan WU¹, Ikuo MIYAJIMA², Kuang-Liang HUANG³,
Ya-Chin KUO¹ and Ruey-Song LIN^{1*}

Institute of Tropical Agriculture, Kyushu University,
Fukuoka 812–8581, Japan

(Received April 25, 2017 and accepted May 10, 2017)

Dendrobium Lucky Girl leaves position presented different cell membrane thermostability, respectively. It has enhancing the relative injury value when temperature exceeded 50°C. In terms of silicon dioxide treatments decreased leaves ion leakage, malondialdehyde content, and existed higher chlorophyll content which evaluated by water bath temperature at 50°C. Result not only presented higher root activity and chlorophyll SPAD value at 150 mg·L⁻¹ silicon dioxide treatment, but also showed more leaves at 300–600 mg·L⁻¹ silicon dioxide treatments during vegetative stage. Even though, silicon dioxide treatments had no significant difference among some horticultural characteristic, however it decreased flower bud abortion rate at 150 mg·L⁻¹ treatment.

Key words: cell membrane thermostability, heat-tolerance, ion leakage, Nobile *Dendrobium*, silicon dioxide

INTRODUCTION

Temperature is an important factor for growth and development of plant, high temperature cause morpho-anatomical, physiological and biochemical changes of plant during the growing season (Wahid *et al.*, 2007). The mesophyll cell of thermo-sensitive cultivar cabbage (*Brassica oleracea* var. *capitata* L.) showed significant ultrastructural change than thermo-resistant cultivar, such as chloroplasts membrane rupture, structural alteration of thylakoid membrane and disruption of nuclear envelope (Miao *et al.*, 1994). High temperature lead to roll up of leaves in Chinese cabbage (*Brassica campestris* ssp. *L. pekinensis*) (Wu *et al.*, 1995). The photosynthesis or photosynthetic rate was inhibited by high temperature in common bean and salvia taxa (Chaisompongpan *et al.*, 1990; Lasseigne *et al.*, 2007). Even Liu and Huang (2000a) reported that carbohydrate availability was associated with heat stress in creeping bentgrass. Moreover, high temperature stress also caused to affect chlorophyll content, chlorophyll fluorescence parameter and antioxidant systems in early cauliflower (Wang *et al.*, 2004).

Silicon is not an essential element on plant growth and development. It has absorbed, translocated and accumulated of plant by silicic acid and silica gel form (Epstein, 1999; Yoshida, 1975). In fact, silicon has accelerated growth, development and enhanced stresses tolerance of plant, such as inhibited serious extra transpiration and increased water utility and photosynthetic activity of rice (Agarie *et al.*, 1992, 1998), it also

changed morphological and density of stomata in maize (Gao *et al.*, 2006), increased chlorophyll and iron content, decreased proline content in cowpea (Mail and Aery, 2009), appeared thicker flower stalk, bigger diameter and blooming earlier affected leaf macro and micro element content in gerbera (Kamenidou *et al.*, 2010), previous research indicated silicon also enhanced stresses tolerance such as increased cell wall polysaccharide content, decreased leakage at drought and heat stresses in rice (Agarie *et al.*, 1998). Moreover, silicon applied increased superoxide dismutase, root H⁺-ATPase activity, decreased lipid peroxidation and malondialdehyde content at salt stress in barley (Liang, 1999; Liang *et al.*, 2003, 2006). Silicon could alleviate cell damage in norway spruce during aluminum stress (Prabagar *et al.*, 2011).

Cell membrane structure, integrity and stability are most important of plant during heat stress, because it caused lipid liquefaction and electrolytes leakage of cell membrane (Ke, 2006; Yao *et al.*, 2000). Cell membrane thermostability (CMT) has been becoming an efficiency method for evaluating heat tolerance in plants. The method measured as ion leakage from leaves over a range of temperature, it is rapid, inexpensive, require little space and highly correlation (Marcum, 1998; Chen *et al.*, 2014; Yang and Hu, 2015; Yeh and Lin, 2003).

Nobile *Dendrobium* is belong *Dendrobium nobile* type and it has been found from native habitats of Himalayas, Thailand, Laos and South China (Kamemoto *et al.*, 1999). Kuo (2011) indicated that high temperature has an influenced on development of nobile *Dendrobium*. In aspect of micro-propagation and breeding among *Doritaenopsis* and *Phalaenopsis* had difficultly problem to solve (Chuang *et al.*, 2014; Liao *et al.*, 2015), similarly to nobile *Dendrobium* orchid usually do not get used to higher temperature during vegetative growth around June to September (Wu, 2016).

¹ Department of Horticulture, National Chung Hsing University, Taichung 40227, Taiwan, ROC.

² Institute of Tropical Agriculture, Kyushu University, Fukuoka 812–8581, Japan

³ Department of Horticultural Science, National Chiayi University, Chiayi 60004, Taiwan, ROC.

* Corresponding author (E-mail: rsolin@dragon.nchu.edu.tw)

Therefore, it should be a limited factor during summer growing season in Taiwan. The purpose of this studies were determined cell membrane stability by CMT technique and silicon dioxide supply on development of nobile *Dendrobium*.

MATERIALS AND METHODS

Plant materials

Dendrobium Lucky Girl plants were purchased in Mar. 2014 from an orchid nursery in Douliu, Taiwan. Those plants were brought up several pseudobulb with 10–17 nodes and grown in 9-cm diameter soft black plastic pot in greenhouse of National Chung Hsing University horticultural research station. The light intensity was $185\text{--}216\ \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ and average temperature about $23.6\text{--}25.3^\circ\text{C}$ of experimental locations. The plants with two pseudobulb and vegetative buds were transplanted that switched old sphagnum moss medium, then pruned part of root and reporting with new sphagnum moss in 13-cm diameter pots. Those plants were watered as needed with tap water and no fertilizer applied for control. The plants were selected for experiment in Jan. 2015.

Experiment processes

On Mar. to Oct. 2015, the plants were supplied 2 g per pot of Hi-Control No.1 ($14\text{N}\text{--}11\text{P}_2\text{O}_5\text{--}13\text{K}_2\text{O}$, 180 type, JCAM Inc., Japan), while the pseudobulb terminal leaf had already formed and matured after that measured leaves cell membrane thermostability. In terms of silicon dioxide experiment, the plants were supplied nutrient solution ($16.0\ \text{g}\cdot\text{L}^{-1}\ \text{NH}_4\text{NO}_3$, $11.5\ \text{g}\cdot\text{L}^{-1}\ \text{NH}_4\text{H}_2\text{PO}_4$, $48.6\ \text{g}\cdot\text{L}^{-1}\ \text{MgSO}_4\cdot 7\text{H}_2\text{O}$, $47.2\ \text{g}\cdot\text{L}^{-1}\ \text{Ca}(\text{NO}_3)_2\cdot 4\text{H}_2\text{O}$, $60.6\ \text{g}\cdot\text{L}^{-1}\ \text{KNO}_3$, $81.4\ \text{g}\cdot\text{L}^{-1}\ \text{CaCl}_2\cdot 2\text{H}_2\text{O}$, $47.8\ \text{g}\cdot\text{L}^{-1}\ \text{P}_2\text{O}_5$, $2.86\ \text{g}\cdot\text{L}^{-1}\ \text{H}_3\text{BO}_3$, $0.22\ \text{g}\cdot\text{L}^{-1}\ \text{ZnSO}_4\cdot 7\text{H}_2\text{O}$, $0.08\ \text{g}\cdot\text{L}^{-1}\ \text{CuSO}_4\cdot 5\text{H}_2\text{O}$, $1.54\ \text{g}\cdot\text{L}^{-1}\ \text{MnSO}_4\cdot \text{H}_2\text{O}$, $0.017\ \text{g}\cdot\text{L}^{-1}\ \text{H}_2\text{MoO}_4\cdot \text{H}_2\text{O}$) with 0, 75, 150, 300 and $600\ \text{mg}\cdot\text{L}^{-1}$ silicon dioxide (Sigma-Aldrich, USA), 200 mL per pot twice a weeks. Furthermore, we mentioned the leaves chlorophyll SPAD value during vegetative stage, when pseudobulb terminal leaf has already formed and matured after that measured leaves ion leakage, chlorophyll content, root activity and malondialdehyde concentration. The horticultural characteristics were also investigated.

Leaves cell membrane thermostability

The leaves for analysis were harvested from the 3–5 nodes (the location downstream from terminal leaf), and were evaluated for CMT (Cell membrane thermostability) following procedures described by Yeh and Lin (2003) with some modification. Each samples for assay consisted of five leaf discs (6 mm diameter), has rinsed three times thoroughly tap water and distilled water before assay. Leaf discs were placed in 25 mL flask containing 10 mL distilled water then was measured at 25, 30, 35, 40, 45, 50, 55, 60, 65, and 70°C water bath temperature for 30 min, after that the solution was measured with conductivity meter (Isfet, IQ-180, USA). The flask were capped with foil autoclaved (HY-300SV,

121°C , $1.05\ \text{kg}\cdot\text{cm}^{-2}$, Taiwan) for 20 min. The calibrated relative injury (RI) was calculated as follows:

$\text{RI}(\%) = \{1 - [1 - (T_i/T_F)] / [1 - (C_i/C_F)]\} \times 100$, where T and C refer to conductance value for treatment and control, respectively, and subscript I and F refer to initial and final conductance value, respectively.

Leaves ion leakage

The leaves for analysis were harvested from the 3–5 nodes (the location downstream from terminal leaf), and were evaluated for ion leakage following procedures described by Martineau *et al.* (1979) with some modification. Each samples for assay consisted of five leaf discs at 6 mm diameter, has rinsed three times thoroughly tap water and distilled water before assay. Leaf discs were placed in 25 mL flask containing 10 mL distilled water then was measured at 50°C water bath temperature for 30 min, after that the solution was measured with conductivity meter (Isfet, IQ-180, USA). The flask were capped with foil autoclaved (HY-300SV, 121°C , $1.05\ \text{kg}\cdot\text{cm}^{-2}$, Taiwan) for 20 min. The calibrated ion leakage was calculated as follows:

$\text{Leakage}(\%) = (R_i/R_f) \times 100$, where R_i and R_f refer to conductance value for initial and final conductance value, respectively.

Malondialdehyde concentration (MDA)

MDA concentration was measured by using Heath and Packer (1968) with some modification. The sample of 0.3 g leaves tissue were harvested from the 3–5 nodes (the location downstream from terminal leaf), then homogenized with sand and 1.8 mL TCA (trichloroacetic acid 5%), after that was added to 10 mL tube and centrifuged (Kubata, KN-70, Japan) 3500 rpm for 15 min. A 1 mL supernatant solution was added to tube containing 4 mL of 20% (v/v) TCA and 0.5% (v/v) thiobarbituric acid. The sample was heated in a water bath at 95°C for 30 min then quickly cooled in an ice bath followed by centrifugation at 3500 rpm for 15 min. The absorbance of supernatant was determined with a spectrophotometer at 532–600 nm (Hitachi, U-2001, Japan). The MDA concentration was calculated as follow: $\text{MDA}(\text{nmol}\cdot\text{g}^{-1}) = (A_{532} - A_{600}) \div 155 (\text{extinction coefficient, } \text{mM}^{-1}\cdot\text{cm}^{-1}) \times 5 (\text{reaction volume, mL}) \times 1.8 (\text{dilution factor}) \times 1000 \div \text{sample fresh weight (g)}$.

Chlorophyll content

The leaves sample were harvested from the 3–5 nodes (the location downstream from terminal leaf), and were measured by using the method of Arnon *et al.* (1954) and Marker (1972) with some modification. Sample of 0.1 g leaf tissue added 10 mL mixture solution (acetone: methanol=80: 20) soaked in darkness overnight, then determined with a spectrophotometer (Hitachi, U-2001, Japan) at 645, 652 and 663 nm.

Chlorophyll SPAD value

Each leaf sample has been determined on same position, and measured with Chlorophyll Meter (SPAD-502, Minolta Co. Ltd., Japan) during vegetative stage.

The results expressed as SPAD units.

Root activity

Root activity of plants was determined by using Steponkus and Lanphear, (1967) with modified. The sample of 0.01 g root tip added 0.6% TTC solution (triphenyl tetrazolium chloride, 0.05 mM Na_2HPO_4 and KH_2PO_4 buffer pH 7.4) then soaked in darkness over 17 hours. The sample has been rinsed distill water and added 20 mL alcohol (95%) into tube, when it being heated in a water bath at 78°C for 20 min after that quickly cooled in room temperature and determined with a spectrophotometer (Hitachi, U-2001, Japan) at 480 nm.

Statistical analysis

The data were subjected to an analysis of variance using a completely randomized design. Separation of means among treatments was used least significant difference test at $P \leq 0.05$. Statistical analyses were used Costat 6.1 (CoHort Software, Monterey, CA, USA). SigmaPlot software 12.5 (Systat Software Inc., Chicago, CA, USA) was used to assay regression analysis. The figures and tables were made out from the data using Microsoft Excel 2010 (Microsoft Office Software,

Redmond, WA, USA).

RESULTS

The relationship between the relative injury value of leaves and water bath temperature were sigmoidal in *Dendrobium* Lucky Girl. The temperature has been increasing over 50.0°C that leaves (L3–L5) enhancing the relative injury value, the leaf (L3) appeared higher value than others (L4–5). The relative injury value mid-point of leaves position occurred the sigmoid response curve between 54.9–55.2°C (Fig. 1). In terms of silicon experiment, result showed higher root activity and chlorophyll SPAD value at 150 mg·L⁻¹ treatment (Fig. 2 and Fig. 3). The leave ion leakage emerged fluctuation when heating at 50.0°C for 30 min. The leave position (L3–L5) showed lower ion leakage at 150–600 mg·L⁻¹ treatments compared with control (Fig. 4). Silicon dioxide treatments have gradually increasing leave chlorophyll content and decreasing MDA concentration (Table 1). In terms of horticultural characteristic, silicon dioxide treatments did not improve plant height and number of nodes, but it enhanced number of leaves during vegetative growth stage (Table 2), moreover, silicon dioxide treatments had no significant difference among numbers of flower bud, numbers of node with flower bud, flower per inflorescence and flower per pseudobulb, however 150 mg·L⁻¹ treatments occurred lower flower bud abortion rate during reproductive growth stage (Table 3 and Fig. 5).

Fig. 1. Effect of water bath temperature on leaves relative injury of *Dendrobium* Lucky Girl. Formula represent sigmoidal regression, $n=3$. L3, L4, L5: downstream from terminal leaf 3–5 location of node

Fig. 2. Effect of silicon dioxide on root activity of *Dendrobium* Lucky Girl. Bar = $\pm$ SE; $n=3$.

Fig. 3. Effect of silicon dioxide on SPAD value of *Dendrobium* Lucky Girl vegetative bud. Bar = $\pm$ SE; $n=3$.

Fig. 4. Effect of silicon dioxide on leaves ion leakage of *Dendrobium* Lucky Girl. Bar = $\pm$ SE; $n=3$. L3, L4, L5: downstream from terminal leaf 3–5 location of node

Table 1. Effect of silicon dioxide on leaves malondialdehyde and chlorophyll content of *Dendrobium* Lucky Girl

SiO ₂ (mg·L ⁻¹)	L3 ^z		L4		L5	
	MDA (nmol·g ⁻¹)	Total Chl. (mg·g ⁻¹)	MDA (nmol·g ⁻¹)	Total Chl. (mg·g ⁻¹)	MDA (nmol·g ⁻¹)	Total Chl. (mg·g ⁻¹)
0	11.9 a ^y	0.54 c	3.5 a	0.62 c	6.0 a	0.61 b
75	4.1 b	0.70 bc	3.4 ab	0.72 bc	3.9 ab	0.65 b
150	3.2 b	0.81ab	3.6 ab	0.89 ab	2.7 b	0.97 a
300	4.2 b	0.95 a	2.4 b	0.87 ab	2.7 ab	0.87 ab
600	1.8 b	0.98 a	2.1 b	0.98 a	2.8 b	1.01 a

^z L3, L4, L5: downstream from terminal leaf 3–5 location of node^y Mean separation within each column by LSD test at $P \leq 0.05$. n=3**Table 2.** Effect of silicon dioxide on pseudobulb growth of *Dendrobium* Lucky Girl

SiO ₂ (mg·L ⁻¹)	Plant height (cm)	Number of nodes	Number of leaves
0	28.7 a ^z	15.3 a	4.0 b
75	30.0 a	16.7 a	7.5 b
150	29.7 a	17.7 a	9.7 a
300	28.3 a	16.0 a	10.7 a
600	28.0 a	15.7 a	10.7 a

^z Mean separation within each column by LSD test at $P \leq 0.05$. n=3.**Table 3.** Effect of silicon dioxide on blooming of *Dendrobium* Lucky Girl

SiO ₂ (mg·L ⁻¹)	Numbers of flower bud	Numbers of node with flower bud	Flowers per inflorescence	Flowers per pseudobulb	Flower bud abortion (%) ^z
0	7.7 a ^y	5.7 a	1.8 a	10.3 a	25.0 a
75	7.3 a	6.3 a	2.0 a	12.3 a	12.5 ab
150	8.0 a	7.7 a	2.1 a	15.6 a	2.8 b
300	6.3 a	5.0 a	2.1 a	10.0 a	23.1 a
600	6.0 a	5.0 a	2.3 a	11.3 a	16.7 ab

^z Percentage data were logarithmic transformed prior to analysis^y Mean separation within each column by LSD test at $P \leq 0.05$. n=3. **Fig. 5.** Effect of silicon dioxide treatments on flowering of *Dendrobium* Lucky Girl.

DISCUSSION

The cell membrane thermostability technique has been screening crop for heat tolerance in soybeans (Martineau *et al.*, 1979), wheat (Saadalla *et al.*, 1990), chrysanthemum (Wang and Yeh, 2013; Wang *et al.*, 2008; Yeh and Lin, 2003), leafy radish (Chen *et al.*, 2014), marsh-rosemary (Chang, 2014) and *Phalaenopsis* (Yang and Hu, 2015). The exposure time and midpoint of sigmoidal curve that would able to evaluate heat tolerance of crop (Ingram and Buchanan, 1981; Lester, 1985). Result indicated the position of leaves have increasing the relative injury value after heating at 50°C, the leaves ion leakage showed nearly midpoint among 54.9–55.2°C, therefore cell membrane thermostability should be able to determine at 55°C in *Dendrobium* Lucky Girl (Fig. 1). The position of leaf (L3) showed more sensitively than others leaves.

Some research indicated, silicon improved development and production of crop (Kamenidou *et al.*, 2010; Mattson and Leatherwood, 2010; Sivanesan *et al.*, 2010). In present experiment showed silicon dioxide treatments enhanced root activity and chlorophyll SPAD value (Fig. 2 and Fig. 3). Agarie *et al.* (1998) reported that silicon treatment reduced ion leakage in rice leaves, when it is heating at 42.5°C. Ma (2004) also reported that silicon played a major role against stresses such as drought, freeze, metal ion toxin, salt, pest and disease. Result indicated that leaves (L3–5) ion leakage emerged fluctuation data, it should be discussed the assimilated difference of *Den.* Lucky Girl on ion leakage due to silicon dioxide concentration (Fig. 4), because silicon element absorbed, translocated and accumulated of plant by silicic acid and silica gel form, it should be consider to the ability of the roots to take up this element in monocotyledonous and dicotyledons (Epstein, 1999; Ma and Yamaji, 2006). In addition, silicon dioxide treatments increased the leaves chlorophyll content (Table 1), the result similar like Ahmad *et al.* (1992), Duan *et al.* (2013), Liang (1999) and Ma *et al.* (2002). Agarie *et al.* (1992, 1993) indicated that silicon treatments accelerated development, accumulation of dry weight and photosynthetic rate that attributed silicon maintained the chlorophyll stability. In present experiment also showed that silicon treatments decreased the malondialdehyde concentration of leaves (Table 1). Cell membrane has been received injury in adversity condition that lead to lipid peroxidation and produced the precursor is malondialdehyde (MDA), it would be able to integrate protein and enzyme in order to cause damage of cell membrane structure (Zhang and Yin, 2009; Bailly *et al.*, 1996; Blokhina *et al.*, 2003; Gutteridge and Halliwell, 1990; Liu and Huang, 2000b; Malencic *et al.*, 2000). Lombardi and Wangersky (1991) reported that the marine diatom (*Chaetoceros gracilis*) less silicon supplement lead to break of cell membrane. Agarie *et al.* (1998) indicated silicon treatment prevented cell membrane function and structure damage in rice during stress environment. Wang and Galletta (1998) also reported that silicon treatment caused alternation glycolipids and phospho-

lipids ratio of cell membrane. In other hand, silicon treatment also decreased lipid peroxidation and malondialdehyde content at salt stress on barley (Liang, 1999; Liang *et al.*, 2003, 2006). Even silicon treatments did not enhance the plant height and number of node, however it has been increased number of leaves during vegetative growth stage, 150 mg·L⁻¹ silicon dioxide treatments appeared lower flower abortion rate (Table 2 and Table 3).

In summary, silicon element has been protected the cell membrane to stress that discussed respect to membrane integrity, stability and function of crop. Furthermore, beneficial element silicon on physiological advantage and mechanism of horticultural crop should be more investigated.

AUTHOR CONTRIBUTIONS

Shing-Kuan WU, designed this study and wrote the initial draft of the manuscript. Ikuo MIYAJIMA, offered advices on tropical horticultural crops research for efficiency experiments on heat tolerance, and revised the manuscript and inspected final data. Kuang-Liang HUANG, joined orchid research team work, designed the cultural environment for orchid research under light intensity and temperature control for studies, and offered suggestion for research. Ya-Chin KUO, worked on experimental process of fertilizer supplement and data collection. Ruey-Song LIN, organized the research protocol through physiological and biochemical experiments on heat tolerance of crops and managed lab process. The final version of this manuscript was approved by all authors.

REFERENCES

- Agarie, S., W. Agata, F. Kubota and P. B. Kaufman. 1992 Physiological roles of silicon in photosynthesis and dry matter production in rice plant. I. Effect of silicon and shading treatments. *Jpn. J. Crop Sci.*, **61**: 200–206
- Agarie, S., H. Uchida, W. Agata, F. Kubota and P. B. Kaufman. 1993 Effect of silicon on growth dry matter production and photosynthesis in rice plant. *Crop Prod. Improve. Tech.*, **34**: 225–234
- Agarie, S., H. Uchida, W. Agata, F. Kubota and P. B. Kaufman. 1998 Effect of silicon on transpiration and leaf conductance in rice plants. *Plant Prod. Sci.*, **1**: 89–95
- Ahmad, R., S. H. Zaheer and S. Ismail. 1992 Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Sci.*, **85**: 43–50
- Arnon, D. I., M. B. Allen and F. R. Whatley. 1954 Photosynthesis by isolated chloroplasts. *Nature*, **174**: 394–396
- Bailly, C., A. Benamar, F. Corbineau and D. Côme. 1996 Changes in malondialdehyde content and in superoxide dismutase catalase and glutathione reductase activities in sunflower seeds as related to deterioration during accelerated aging. *Physiol. Plant.*, **97**: 104–110
- Blokhina, O., E. Virolainen and K. V. Fagerstedt. 2003 Antioxidants oxidative damage and oxygen deprivation stress: a review. *Ann. Bot.*, **91**: 179–194
- Chaisompongpan, N., P. H. Li, D. W. David and A. H. Markhart. 1990 Photosynthetic responses to heat stress in common bean genotypes differing in heat acclimation potential. *Crop Sci.*, **30**: 100–104
- Chang, Y. T. 2014 Breeding of low vernalization-requiring culti-

- vars/lines, selection indices, thermal time of flowering, and phylogenetic relationship in *Limonium*. PhD dissertation. Dept. of Hort. and Landscape Architecture. Natl. Taiwan Univ.
- Chen, W. L., W. J. Yang, H. F. Lo and D. M. Yeh. 2014 Physiology anatomy and cell membrane thermostability selection of leafy radish (*Raphanus sativus* var. *oleiformis* Pers.) with different tolerance under heat stress. *Sci. Hortic.*, **179**: 367–375
- Chuang, H. T., K. L. Huang, R. S. Shen, I. Miyajima and S. T. Hsu. 2014. Using cut-column pollination method overcome crossing barriers in *Phalaenopsis* Sunrise Goldmour 'KHM637'. *J. Fac. Agr., Kyushu Univ.*, **59**: 265–271
- Duan, X. Y., M. Tang and W. S. Wang. 2013 Effect of silicon on physiology and biochemistry of *Dendrobium moniliforme* plantlets under cold stress. *Agri. Biotechnol.*, **2**: 18–21
- Epstein, E. 1999 Silicon. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, **50**: 641–664
- Gao, X. P., C. Zou, L. J. Wang and F. S. Zhang. 2006 Silicon decreases transpiration rate and conductance from stomata of maize plants. *J. Plant Nutri.*, **29**: 1637–1647
- Gutteridge, J. M. C. and B. Halliwell. 1990 The measurement and mechanism of lipid peroxidation in biological systems. *Trends. Biochem. Sci.*, **15**: 129–135
- Heath, R. L. and L. Packer. 1968 Photoperoxidation in isolated chloroplasts I. Kinetics and stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.*, **125**: 189–198
- Ingram, D. L. and D. Buchanan. 1981 Measurement of direct heat injury of root of three woody plant. *HortScience*, **16**: 769–771
- Kamamoto, H., T. Amore and A. R. Kuehnle. 1999 *Breeding Dendrobium Orchids in Hawaii*, Hawaii Univ. Press, (Honolulu), pp. 9
- Kamenidou, S., T. J. Cavins and S. Marek. 2010 Silicon supplements affect floricultural quality traits and elemental nutrient concentration of greenhouse produced gerbera. *Sci. Hortic.*, **123**: 390–394
- Ke, Y. 2006 *Plant Physiology*, 1st ed. Yi-Hsien Publishing Co, (Taipei), pp. 682.
- Kuo, W. J. 2011 Effect of environmental factors, substrates and fertilizers on growth and flowering of nobile type *Dendrobium*. Master's thesis. Dept. of Plant Ind. Natl. Pingtung Univ. of Sci. and Tech.
- Lasseigne, F. T., S. L. Warren, F. A. Blazich and T. G. Ranney. 2007 Day/night temperature affects growth and photosynthesis of cultivated *Salvia* taxa. *J. Amer. Soc. Hort. Sci.*, **132**: 492–500
- Lester, G. E. 1985 Leaf cell membrane thermostabilities in *Cucumis melo*. *J. Amer. Soc. Hort. Sci.*, **110**: 506–509
- Liang, Y. C. 1999 Effect of silicon on enzyme activity and sodium potassium and calcium concentration in barley under salt stress. *Plant Soil*, **209**: 217–224
- Liang, Y. C., Q. Chen, Q. Liu, W. H. Zhang and R. X. Ding. 2003 Exogenous silicon increase antioxidant enzyme activity and reduces lipid peroxidation in root of salt stress barley (*Hordeum vulgare* L.). *J. Plant Physiol.*, **160**: 1157–1164
- Liang, Y. C., W. H. Zhang, Q. Chen, Y. L. Liu and R. X. Ding. 2006 Effect of exogenous silicon on H⁺-ATPase activity phospholipids and fluidity of plasma membrane in leaves of salt stress barley (*Hordeum vulgare* L.). *Environ. Exp. Bot.*, **57**: 212–219
- Liao, Y. W., P. L. Huang, K. L. Huang, I. Miyajima and S. T. Hsu. 2015 Factors affecting to somatic embryogenesis and plant regeneration from callus and in vitro ontogeny of *Doritaenopsis* Taisuco Ladylip. *J. Fac. Agr., Kyushu Univ.*, **60**: 13–22
- Liu, X. Z. and B. R. Huang. 2000a Carbohydrate accumulation in relation to heat stress tolerance in two creeping bentgrass cultivar. *J. Amer. Soc. Hort. Sci.*, **125**: 442–447
- Liu, X. Z. and B. R. Huang. 2000b Heat stress injury in relation to membrane lipid peroxidation in creeping bentgrass. *Crop. Sci.*, **40**: 503–510
- Lombardi A. T. and P. J. Wangersky. 1991 Influence of phosphorus and silicon on lipid class production by the marine diatom *Chaetoceros gracilis* growth in turbidostat cage culture. *Mar. Ecol. Prog. Ser.*, **77**: 39–47
- Ma, J. F. 2004 Role of silicon in enhancing the resistance of plants to biotic and abiotic stresses. *Soil Sci. Plant Nutr.*, **50**: 11–18
- Ma, J. F. and N. Yamaji. 2006 Silicon uptake and accumulation in higher plants. *Trends Plant Sci.*, **11**: 392–397
- Ma, C. C., Q. F. Li, L. Z. Shu and J. Y. Zhang. 2002 Preliminary examination of the mechanism about effects of silicon on maize seed germination and seedling growth. *Acta Agron. Sin.*, **28**: 665–669
- Mail, M. and N. C. Aery. 2009 Effect of silicon on growth biochemical constituents and mineral nutrition of cowpea. *Commun. Soil Sci. Plant Anal.*, **40**: 1041–1052
- Malencic, Dj., O. Gasic, M. Popovic and P. Boza. 2000 Screening for antioxidant properties of *Salvia reflexa* Hornem. *Phytotherapy Res.*, **14**: 546–548
- Marcum, K. B. 1998 Cell membrane thermostability and whole-plant heat tolerance of Kentucky bluegrass. *Crop Sci.*, **38**: 1214–1218
- Marker, A. F. H. 1972 The use of acetone and methanol in the estimation of chlorophyll in the presence of pheophytin. *Freshwater Biol.*, **2**: 361–385
- Martineau, J. R., J. E. Specht, J. H. Williams and C. Y. Sullivan. 1979 Temperature tolerance in soybeans. I. Evaluation of a technique for assessing cellular membrane thermostability. *Crop Sci.*, **19**: 75–78
- Mattson, N. S. and W. R. Leatherwood. 2010 Potassium silicate drenches increase leaf silicon content and affect morphological traits of several floriculture crop grown in a peat-based substrate. *HortScience*, **45**: 43–47
- Miao, C., R. Q. Li and J. B. Wang. 1994 Ultrastructural study in leaf cell of *Brassica oleracea* var. *capitata* under heat stress. *Acta Bot. Sin.*, **36**: 730–732
- Prabagar, S., M. J. Hodson and D. E. Evans. 2011 Silicon amelioration of aluminium toxicity and cell death in suspension cultures of Norway spruce (*Picea abies* (L.) Karst.). *Environ. Exp. Bot.*, **70**: 266–276
- Saadalla, M. M., J. F. Shanahan and J. S. Quick. 1990 Heat tolerance in winter wheat: I. Hardening and genetic effects on membrane thermostability. *Crop Sci.*, **30**: 1243–1247
- Sivanesan, I., M. S. Son, J. P. Lee and B. R. Jeong. 2010 Effects of silicon on growth of *Tagetes patula* L. Boy orange and Yellow boy seedlings cultured in an environment controlled chamber. *Propag. Ornament. Plants*, **10**: 136–140
- Steponkus, P. L. and F. O. Lanphear. 1967 Refinement of triphenyltetrazolium chloride method of determining cold injury. *Plant Physiol.*, **42**: 1423–1426
- Wahid, A., S. Gelani, M. Ashraf and M. R. Foolad. 2007 Heat tolerance in plants: An overview. *Environ. Exp. Bot.*, **61**: 199–223
- Wang, C. H. and D. M. Yeh. 2013 Assessment of cell Membrane thermostability of chrysanthemum and its application to screening for heat-tolerant seedlings. *J. Taiwan Soc. Hort. Sci.*, **59**: 153–166
- Wang, S. Y. and G. J. Galletta. 1998 Foliar application of potassium silicate induces metabolic changes in strawberry plants. *J. Plant Nutr.*, **21**: 157–167
- Wang, B. L., M. Xu, Q. H. Shi and J. S. Cao. 2004 Effect of high temperature stress on antioxidant systems, chlorophyll and chlorophyll fluorescence parameters in early cauliflower leaves. *Sci. Agr. Sin.*, **37**: 1245–1250
- Wang, C. H., D. M. Yeh and C. S. Sheu. 2008 Heat tolerance and flowering-heat- delay sensitivity in relation to cell membrane thermostability in chrysanthemum. *J. Amer. Soc. Hort. Sci.*, **133**: 754–759
- Wu, S. K. 2016 Effect of essential nutritional elements, plant growth regulators on growth and flowering, and simulated transportation in Nobile *Dendrobium*. PhD dissertation. Dept. of Hort. Natl. Chung Hsing Univ.
- Wu, G. S., Y. J. Wang, W. H. Cao, Y. W. Jiang and L. R. Zhang. 1995 Studies on mode of high temperature injury and screening method for heat tolerant Chinese cabbage. *Acta Agr. Boreall Sin.*, **10**: 111–115
- Yang, Y. and W. C. Hu. 2015 Study on heat tolerance index of

- Phalaenopsis*. Tainan District Agricultural Research and Extension Station Research Bull., **65**: 1–9
- Yao, Y. G., X. H. Shi, J. G. Yang and S. Y. Wang. 2000 Relation of the permeability of plasma membrane of leaves and the contents of several biochemical matter to heat tolerance of pepper. *J. Hunan Agr. Univ.*, **26**: 97–99
- Yeh, D. M. and H. F. Lin. 2003 Thermostability of cell membranes as a measure of heat tolerance and relationship to flowering delay in chrysanthemum. *J. Amer. Soc. Hort. Sci.*, **128**: 656–660
- Yoshida, S. 1975 The physiology of silicon in rice. *Asian and Pacific Council Food and Fertilizer Technology Center Technical Bull.*, **25**: 1–27
- Zhang, Y. F. and B. Yin. 2009 Influences of salt and alkali mixed stresses on antioxidative activity and MDA content of *Medicago sativa* at seedling stage. *Acta Prataculturae Sin.*, **18**: 46–50