

## Laplacian energy of directed graphs and minimizing maximum outdegree algorithms

Perera, Kissani  
Graduate School of Mathematics, Kyushu University

Mizoguchi, Yoshihiro  
Faculty of Mathematics, Kyushu University

<http://hdl.handle.net/2324/18650>

---

出版情報 : MI Preprint Series. 2010-35, 2010-11-29. Faculty of Mathematics, Kyushu University  
バージョン : draft  
権利関係 :



# **MI Preprint Series**

**Kyushu University  
The Global COE Program  
Math-for-Industry Education & Research Hub**

## **Laplacian energy of directed graphs and minimizing maximum outdegree algorithms**

**Kissani Perera  
& Yoshihiro Mizoguchi**

**MI 2010-35**

( Received November 29, 2010 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# Laplacian Energy of Directed Graphs and Minimizing Maximum Outdegree Algorithms

K.K.K.R. Perera<sup>a</sup>, Y.Mizoguchi<sup>b</sup>

<sup>a</sup>*Graduate School of Mathematics, Kyushu University, Japan*

<sup>b</sup>*Faculty of Mathematics, Kyushu University, Japan*

---

## Abstract

Energy has been studied in mathematical perspective as well as physical perspective for several years ago. In spectral graph theory, the eigenvalues of several kinds of matrices have been studied, of which Laplacian matrix attracted the greatest attention [5, 12]. Recently, in 2009, Adiga considered Laplacian energy of directed graphs using skew Laplacian matrix, in which degree of vertex is considered as total of the out-degree and the in-degree. Since directed graphs play an important role in identifying the structure of web-graphs as well as communication graphs, we consider Laplacian energy of simple directed graphs and find some relations by using the general definition of Laplacian matrix. Unlike in [1], we derived two types of equations for simple directed graphs and symmetric directed graphs with  $n \geq 2$  vertices by considering out-degree of vertex. Further we consider the class  $P(\alpha)$  which consists of non isomorphic graphs with energy less than some  $\alpha$  and find 47 non isomorphic directed graphs for class  $P(10)$ . Our objective extended to enumerate the structure of directed graphs using the Laplacian energy concept. Minimization maximum outdegree(MMO) algorithm defined in [3] can be used to find the directed graphs with minimum Laplacian energy.

*Keywords:* Laplacian energy, directed graph, MMO algorithms

---

---

☆

*Email address:* [kissani@math.kyushu-u.ac.jp](mailto:kissani@math.kyushu-u.ac.jp) (K.K.K.R. Perera)

## 1. Introduction

The relation between Hückel theory and the theory of graph spectra was observed for a long time. The basic problem in Hückel theory is to determine the eigenvalues and eigenvectors of the graph representing carbon atom connectivity of a given conjugated system. An interesting quantity is the sum of the energies of all the electrons in a molecule, called total  $\pi$ -electron energy [5, 6, 9, 4, 8]. Several criteria relate to energy such as energy change due to edge addition, maximal energy, equal energy has been considered in [2, 9, 4]. In spectral graph theory, the eigenvalues of several other matrices have been studied, of which Laplacian matrix attracted the greatest attention [5]. Therefore based on definitions of energy, in 2006, Gutman [8] defined

Laplacian energy for undirected graph  $G(n, m)$  as  $LE_g(G) = \sum_{i=1}^n |\mu_i - 2m/n|$ , and derive some lower bounds and upper bounds. He use axillary eigenvalues  $\gamma_i, i = 1, \dots, n$  defined by  $\gamma_i = \mu_i - \frac{2m}{n}$  which satisfy  $\sum_{i=1}^n \gamma_i = 0$  and  $\sum_{i=1}^n \gamma_i^2 = 2M$  where  $M = m + \frac{1}{2} \sum_{i=1}^n (d(i) - \frac{2m}{n})^2$  and  $d(i)$  is the degree of a vertex. The following bounds were obtained from the definition of  $LE_g(G)$ .

**Theorem 1.1 (Gutman[8]).**

$$LE_g(G) \leq \sqrt{2Mn} \quad (1)$$

**Theorem 1.2 (Gutman[8]).** *If  $G$  is a graph with one component then,*

$$LE_g(G) \leq \frac{2m}{n} + \sqrt{(n-1)(2M - (\frac{2m}{n})^2)} \quad (2)$$

**Theorem 1.3 (Gutman[8]).**

$$2\sqrt{M} \leq LE_g(G) \leq 2M \quad (3)$$

Further in [12], Total  $\pi$ -electron energy and Laplacian energy was compared. A similar problem for the usual Laplacian energy has been considered in [10] for undirected graphs using second spectral moment. According to [10]

Laplacian energy was defined as  $LE_k(G) = \sum_{i=1}^n \mu_i^2$  for eigenvalues  $\mu_i$  of undirected Laplacian matrix  $L = D - A$ . Following results were obtained in [10].

**Theorem 1.4 (Kragujevac[10]).** For any graph  $G$  on  $n$  vertices whose degree are  $d(1), d(2), \dots, d(n)$ ,

$$LE_k(G) = \sum_{i=1}^n d(i)(d(i) + 1) \quad (4)$$

**Theorem 1.5 (Kragujevac[10]).** For any connected graph  $G$  on  $n \geq 2$  vertices,

$$LE_k(G) \geq 6n - 8. \quad (5)$$

Equality holds iff  $G$  is a path  $P_n$  on  $n$  vertices.

**Theorem 1.6 (Kragujevac[10]).** For any  $\alpha > 4$ , the class  $p(\alpha)$  of all non-isomorphic connected graphs with the property  $LE_k(G) \leq \alpha$  is finite.

By using eigenvalues of Laplacian matrix, Laplacian Estrada index was defined in [11] and derived some upper and lower boundaries. Further in 2009, Gutman found various relationships using incident energy [7]. In 2009, Adiga

[1] introduce skew Laplacian energy for directed graphs as  $SLE(G) = \sum_{i=1}^n \mu_i^2$ ,

which is similar to [10]. Eigenvalues  $\mu_i$  are the eigenvalues of skew Laplacian matrix  $SL(G) = D - S(G)$  where  $S(G)$  is the adjacency matrix with  $s_{ij} = 1$  and  $s_{ji} = -1$  whenever there is a arc from  $i \rightarrow j$  and 0 otherwise.  $D$  is a diagonal matrix with  $D(i, i) = d(i) = d^{out}(i) + d^{in}(i)$  where  $d^{out}(i)$  is the outdegree and  $d^{in}(i)$  is the indegree of vertex  $i$ . Later upper and lower bound

for skew Laplacian energy  $SLE(G) = \sum_i |\mu_i - \frac{2m}{n}|$  of simple directed graphs

similar to equation (1), (2) and (3) are also derived as in the equation (7) in Theorem 1.7. Skew Laplacian eigenvalues of  $L$  satisfy the following relations.

$$\begin{aligned} \sum_{i=1}^n \mu_i &= 2m \\ \sum_{i=1}^n \mu_i^2 &= \sum_{i=1}^n d(i)(d(i) - 1) \end{aligned} \quad (6)$$

**Theorem 1.7 (Adiga[1]).**

$$\begin{aligned} SLE(G) &\leq \sqrt{2M_1 n} \\ SLE(G) &\leq k + \sqrt{(n-1)[2M_1 - k^2]} \\ 2\sqrt{M} &\leq SLE(G) \leq 2M_1 \end{aligned} \quad (7)$$

where  $M_1 = M + 2m = m + \frac{1}{2} \sum_{i=1}^n (d(i) - \frac{2m}{n})^2$ .

Most real world networks such as communication networks, web graphs etc. are directed graphs. Even though the energy concept originated in chemistry to find the energy of molecular structure, our objective is to investigate the applicability of the concept to find the structure of the web graphs. We enumerate graph structure in which Laplacian energy is less than some value. As an example, we consider the class  $P(\alpha)$  and find the structure of directed graphs belong to the class  $P(10)$ . This paper is a small attempt to find the Laplacian energy and its behavior due to several criteria. We introduce notations and derive formulas for  $LE(G)$ , the Laplacian energy of a directed graph  $G$  by using Kirchoff matrix as in the Section 2. Then we find some relations between undirected and directed graphs of  $LE$  in Section 3. Finally we analyze the MMO algorithm [3] and discuss how it is useful to find the directed graphs with minimum Laplacian energy in Section 4.

## 2. Laplacian energy of directed graphs

A graph  $G$ , which has directed edge or arc is called a directed graph. Adjacency matrix  $A$  of  $G$  is the  $n \times n$  matrix  $A = (a_{ij})$ , where  $a_{ij} = 1$  whenever  $(v_i, v_j)$  is an directed edge and 0 otherwise. A directed graph having no multiple edges or self loops is called a *simple directed graph*. i.e.,  $a_{ij} \in \{0, 1\}$  and  $a_{ij} = 1 \Rightarrow a_{ji} = 0$ . A graph in which each edge is bidirected is called a *symmetric directed graph*. i.e.,  $a_{ij} = 1 \Rightarrow a_{ji} = 1$ . Let  $D = \text{diag}(d^{\text{out}}(1), d^{\text{out}}(2), d^{\text{out}}(3), \dots, d^{\text{out}}(n))$  be diagonal matrix with outdegree of the vertices  $v_1, v_2, \dots, v_n$ . Then we call  $L(G) = D(G) - A(G)$ , Laplacian matrix and its eigenvalues are denoted by  $\{\mu_1, \mu_2, \dots, \mu_n\}$ . Since  $L(G)$  is asymmetric matrix it does not give real eigenvalues always.

**Definition 2.1.** Let  $A(G)$  be the adjacency matrix of a directed graph  $G$ .

Then Laplacian energy of  $G$  is defined as  $LE(G) = \sum_{i=1}^n \mu_i^2$  where  $n$  is the order of  $G$  and  $\mu_i, (i = 1, \dots, n)$  are the eigenvalues of the Laplacian matrix. Let  $G_1 = (V(G_1), E(G_1))$  and  $G_2 = (V(G_2), E(G_2))$  be two finite, directed graphs with disjoint sets of vertices  $V(G_1)$  and  $V(G_2)$ . Then the direct sum  $G = G_1 \oplus G_2$  of these graphs is defined by  $V(G) = V(G_1) \cup V(G_2)$  and  $E(G) = E(G_1) \cup E(G_2)$ .

**Theorem 2.1.** *If  $G$  is a disconnected directed graph with components  $G_1, G_2, \dots, G_n$ ,*

$$LE(G) = \sum_{i=1}^n LE(G_i). \quad (8)$$

**Theorem 2.2.** *Let  $G$  be a directed graph with vertex degrees  $d^{out}(1), d^{out}(2), \dots, d^{out}(n)$ . Then the following relations are hold.*

*If  $G$  is a simple directed graph then*

$$LE(G) = \sum_{i=1}^n d^{out}(i)^2$$

*If  $G$  is a symmetric directed graph then*

$$LE(G) = \sum_{i=1}^n d^{out}(i)(d^{out}(i) + 1)$$

(Proof) Suppose  $G$  is a simple directed graph. Let  $D$  be a diagonal matrix with  $D(i, i) = d^{out}(i)$  for  $i \in V$ . If  $i \rightarrow j$  is an arc then  $a_{ij} = 1$  and  $a_{ji} = 0$ . From Viéte Rule, it is clear that  $\sum_i \mu_i = \text{Trace}(L) = \sum_{i=1}^n d^{out}(i)$  and sum of the determinant of all the  $2 \times 2$  principal sub matrices are  $\sum_{i < j} \mu_i \mu_j$ . i.e.,

$$\begin{aligned} \sum_{i < j} \mu_i \mu_j &= \sum_{i < j} \det \begin{pmatrix} d^{out}(i) & -a_{ij} \\ 0 & d^{out}(j) \end{pmatrix} \\ &= \sum_{i < j} d^{out}(i) d^{out}(j) \end{aligned}$$

For every  $i < j$ ,

$$\begin{aligned} \sum_{i \neq j} \mu_i \mu_j &= 2 \sum_{i < j} \mu_i \mu_j \\ &= \sum_{i \neq j} d^{out}(i) d^{out}(j) \\ &= \sum_{i \neq j} d^{out}(i) d^{out}(j) \end{aligned}$$

Therefore

$$\begin{aligned}
LE(G) &= \sum_i \mu_i^2 = \left(\sum_i \mu_i\right)^2 - \sum_{i \neq j} \mu_i \mu_j \\
&= \left(\sum_i d^{out}(i)\right)^2 - \left(\sum_{i \neq j} d^{out}(i) d^{out}(j)\right) \\
&= \sum_{i=1}^n (d^{out}(i))^2
\end{aligned} \tag{9}$$

If  $G$  is a symmetric directed graph then  $d^{out}(i) = d^{in}(i) = d(i)$  for each node  $i$ . Hence Laplacian energy of symmetric directed graph is similar to the undirected graph as given by  $LE(G_u) = \sum_{i=1}^n d(i)(d(i) + 1)$  in [10]. We can replace  $d(i)$  with  $d^{out}(i)$  and obtained the result.  $\square$

**Corollary 2.3.** *For any directed graph  $G$ , its Laplacian energy  $LE(G)$  is an integer.*

(Proof) Since degree of vertex is an integer we have integer values for  $LE(G) = \sum_{i=1}^n (d^{out}(i))^2$  or  $LE(G) = \sum_{i=1}^n d^{out}(i)(d^{out}(i) + 1)$  for  $i = 1, 2, \dots, n$ .  $\square$

**Corollary 2.4.** *The Laplacian energy of a simple directed path  $P_n$  with  $n \geq 2$  is  $(n - 1)$ .*

(Proof) Since every directed path  $P_n$  has exactly  $(n - 1)$  vertices with out-degree 1 and one vertex of degree 0, using theorem (2.2), we conclude that  $LE(P_n) = (n - 1)$ .  $\square$

**Corollary 2.5.** *The Laplacian energy of a simple directed cycle  $C_n$  with  $n \geq 3$  is  $n$ .*

(Proof) Since every vertex in  $C_n$  has out degree one, it follows from theorem (2.2) that  $LE(G) = n$ .  $\square$

**Corollary 2.6.** *For any connected directed graph on  $n \geq 2$  vertices, we have*

$$n - 1 \leq LE(G) \leq n^2(n - 1). \tag{10}$$

Moreover  $LE(G) = n^2(n - 1)$  if and only if  $G$  is a complete directed graph  $K_n$  and  $LE(G) = n - 1$  if and only if  $G$  is a directed path  $P_n$  on  $n$  vertices.



(Proof) Let  $G$  be a connected directed graph with  $n(\geq 2)$  vertices. Maximum degree of any vertex is less than or equal to  $(n-1)$ . If  $G$  is a simple connected graph then  $LE(G) = \sum_{i=1}^n \mu_i^2 = \sum_{i=1}^n (d^{out}(i))^2 < \sum_{i=1}^n d^{out}(i)(d^{out}(i) + 1)$ .

If  $G$  is a symmetric directed graph then  $LE(G) = \sum_{i=1}^n d^{out}(i)(d^{out}(i) + 1) \leq n^2(n-1)$ . This implies that for any directed graph  $G$ ,  $LE(G) \leq n^2(n-1)$ . Since each vertex of a complete directed graph has exactly  $n-1$  degrees, it is clear that the maximum Laplacian energy of directed graphs with  $n$  vertices is achieved for the complete directed graph  $K_n$ .

We prove left side of inequality (10) by induction. As we know, to form a directed graph we need at least two nodes. Only connected graph which has two node is a simple or bi directed path. Since eigenvalues of  $L(P_2) = \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$  is 1 and 0 and  $L(K_2) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$  is 2 and 0, the result is true for  $n = 2$ . Suppose the result is true for any connected directed graph with  $n-1$  vertices. i.e.,  $LE(G) \geq n-2$ . Then we need to prove the result for any arbitrary connected directed graph with  $n$  vertices. Let  $G$  be a connected directed graph with  $n$  vertices. Then, there is an induced subgraph  $H \subset G$  on  $n-1$  vertices which is also connected. Let  $V(H) = \{v_1, v_2, \dots, v_{n-1}\}$ ,  $V(G) = V(H) \cup \{v_n\}$  and  $LE(H) \geq n-2$ . It is easy to show that  $LE(G) \geq LE(H)+1$ . So we have  $LE(G) \geq n-1$ . We can also prove that if  $G$  is a simple, connected directed graph with  $n$  vertices such that  $LE(G) = n-1$ , then  $G$  must be a directed path  $P_n$ . Suppose  $LE(G) = n-1$ . Let  $n = 2$ . Then  $LE(G) = 1$ .

Since  $LE(G) = \sum_{i=1}^2 (d^{out}(i))^2$ , we have  $d^{out}(1) + d^{out}(2) = 1$ . This happened when we have a one vertex with outdegree 1. That is there exists one directed edge between two nodes. Therefore  $G$  should be a directed path.  $\square$

### 3. Relations between undirected graphs and directed graphs

Every undirected graph can be converted to a directed graph by assigning directions. If each edge is replaced by two way directions then it is similar to the undirected graph.

**Definition 3.1.** For a given directed graph  $G_d = (V_d, E_d)$ , we define an undirected graph  $U(G_d) = (U(V_d), U(E_d))$  by  $U(V_d) = V_u$  and  $U(E_d) = \{\{v_1, v_2\} \mid (v_1, v_2) \in E_d \text{ or } (v_2, v_1) \in E_d\}$ .

Let  $A(U(G)) = (a_{ij})$  be adjacency matrix of  $U(G)$  and let  $A'(G) = (a'_{ij})$  be adjacency matrix of  $G$ . Then

$$a_{ij} = \begin{cases} 1 & \text{if } a'_{ij} = 1 \text{ or } a'_{ji} = 1 \\ 0 & \text{otherwise} \end{cases}$$

**Example 1.** Figure 1 shows four directed graphs  $G_{d_1}, G_{d_2}, G_{d_3}, G_{d_4}$  with  $U(G_{d_i}) = G_u$ . The Laplacian energies  $LE(G_{d_i})$  are different.




Figure 1: Representation of undirected to directed graphs

**Theorem 3.1.** For any directed graph  $G_d$ ,  $LE(G_d) \leq LE_k(U(G_d))$

(Proof) Let  $G_d = (V_d, E_d)$  and  $U(G_d) = (V_u, E_u)$ . For any node  $v \in V_d$ ,  $d^{out}(v) \leq d(v)$ .

$$\begin{aligned} LE_k(U(G_d)) &= \sum_{i=1}^n d(i)(d(i) + 1) \\ &= \sum_{i=1}^n d(i)^2 + \sum_{i=1}^n d(i) \\ &\geq \sum_{i=1}^n (d^{out}(i))^2 + \sum_{i=1}^n d^{out}(i) \\ &\geq LE(G_d) \end{aligned}$$

Equality occur iff  $G_d$  is a symmetric directed graph. □

**Theorem 3.2.** *Let  $G$  be a directed graph and  $G' = G - e$  be a directed graph obtaining by deleting arc  $e$ . Then  $LE(G') \leq LE(G)$ .*

(Proof) Let  $G = (V_d, E_d)$  be a directed graph with  $|V_d| = n$ . Let  $H = (V_h, E_h)$  be edge induced sub graph with  $|V_h| = n_1$  and  $|E_h| = e$  edges. Define  $G'$  as  $H \oplus (n - n_1)K_1$ . Then Laplacian  $L(G)$  of  $G$  is the  $L(G - E_h) + L(G')$ .  $L(G')$  is a square matrix with maximum eigenvalue 1 and all other 0. Then  $\sum_{i=1}^n \mu_i(G) - \sum_{i=1}^n \mu_i(G') = 1$ . This implies that  $\sum_{i=1}^n \mu_i(G) > \sum_{i=1}^n \mu_i(G')$  and exists at least one  $\mu_i(G') < \mu_i(G)$ . Hence prove the theorem.  $\square$

**Proposition 3.1.** *Let  $G_d$  be non-symmetric directed graph with  $U(G_d) = G_u$ . Then there exists a directed graph  $G'_d$  such that  $U(G'_d) = G_u$  and  $LE(G_d) < LE(G'_d)$ .*

(Proof) Let  $G_d$  be non-symmetric directed graph. Then there exists  $\{v_i, v_j\} \in E(G_u)$  and  $\{v_i, v_j\} \notin E(G_d)$  or  $\{v_j, v_i\} \notin E(G_d)$ . Suppose  $\{v_i, v_j\} \notin E(G_d)$ . Let  $G'_d$  be a connected directed graph with  $V(G'_d) = V(G_d)$ ,  $E(G'_d) = E(G_d) \cup \{v_i, v_j\}$ . Then  $\sigma(G'_d) = G_u$  and by theorem 3.2,  $LE(G'_d) > LE(G_d)$ . By adding arcs for each node we can transform given non-symmetric graph to symmetric graph, which is identical to the undirected graph.  $\square$

**Proposition 3.2.** *Let  $G_d$  be a non-simple directed graph with  $U(G_d) = G_u$ . Then there exists a directed graph  $G'_d$  such that  $U(G'_d) = G_u$  and  $LE(G_d) > LE(G'_d)$ .*

(Proof) Suppose  $G_d$  be a non-simple directed graph. Then there exists  $\{v_i, v_j\} \in E(G_u)$  and  $\{v_i, v_j\} \in E(G_d)$  and  $\{v_j, v_i\} \in E(G_d)$ . Suppose  $\{v_i, v_j\} \in E(G_d)$ . Let  $G'_d$  be a connected directed graph with  $V(G'_d) = V(G_d)$  and  $E(G'_d) = E(G_d) - \{v_i, v_j\}$ . Then  $U(G'_d) = G_u$  and by Theorem 3.2,  $LE(G'_d) < LE(G_d)$ . By deleting arcs from each node we can transform given non-simple graph to simple graph.  $\square$

**Theorem 3.3.** *Let  $P(\alpha) = \{G \mid LE(G) \leq \alpha, G \text{ is a simple connected directed graph}\}$ . For any  $\alpha \geq 1$ , the class  $P(\alpha)$  of all non-isomorphic connected directed graphs with the property  $LE(G) \leq \alpha$  is finite.*

(Proof) Let  $G$  be a directed graph with  $n$  vertices and  $m$  arcs such that  $LE(G) \leq \alpha$ . By Corollary 2.6,  $n - 1 \leq LE(G) \leq \alpha$ . Hence we obtain  $n - 1 \leq \alpha$ . Since  $n$  is finite, class  $P(\alpha)$  is also finite.  $\square$

**Corollary 3.4.** *The class  $P(10)$  contains exactly 47 directed graphs. More exactly 29 directed graphs with  $n \leq 10$ , 8 directed cycles with  $n \leq 10$  and 10 directed paths with  $n \leq 11$ . Some of the graphs are listed in Figure 2.*

(Proof) Let  $\alpha = 10$ . Every simple connected directed graph with  $n$  vertices has at least  $(n-1)$  arcs. Notice that for  $n = 12$ ,  $LE(G) \geq (n-1) = 11 > 10$ . For  $n = 11$ ,  $LE(G) \geq 10$ . Therefore all directed graphs from the class  $P(10)$  have at most 11 vertices. Since  $LE(P_n) = n-1$  it has 10 directed path  $P_n$  with  $n \leq 11$  and since  $LE(C_n) = n$  it has 8 directed cycle with  $n \leq 10$ .  $\square$

**Theorem 3.5.** *Let  $G = (V_d, E_d)$  be a simple connected directed graph with  $|V_d| = n$  and  $|E_d| = m$ . If  $\Delta = \max\{d^{out}(v) \mid v \in V_d\}$  and  $\delta = \min\{d^{out}(v) \mid v \in V_d\}$  then*

$$\frac{m^2}{n} \leq LE(G) \leq m(\Delta + \delta) - n\delta\Delta.$$

(Proof) By Cauchy Schwarz inequality,  $LE(G) = \sum_{i=1}^n \mu_i^2 = \sum_{i=1}^n (d^{out}(i))^2 \geq$

$\frac{1}{n} \left( \sum_{i=1}^n d^{out}(i) \right)^2 = \frac{m^2}{n}$ . Lets consider the  $(d^{out}(i) - \delta)(d^{out}(i) - \Delta)$ . For all  $i$ ,

$(d^{out}(i) - \delta) \geq 0$  and  $(d^{out}(i) - \Delta) \leq 0$ . Therefore  $(d^{out}(i) - \Delta)(d^{out}(i) - \delta) \leq 0, \forall i \in V_d$ . Further  $\sum_i (d^{out}(i) - \Delta)(d^{out}(i) - \delta) = \sum_i (d^{out}(i))^2 - (\Delta +$

$\delta) \sum_i d^{out}(i) + n\delta\Delta \leq 0$ . This shows that  $\sum_i (d^{out}(i))^2 \leq m(\Delta + \delta) - n\delta\Delta \leq 0$ .

Hence  $LE(G) \leq m(\Delta + \delta) - n\delta\Delta$ .  $\square$

Remark: If  $\Delta = \delta$  then  $m = \sum_{i=1}^n d^{out}(i) = n\Delta$  and  $m(\Delta + \delta) - n\delta\Delta$   
 $= 2n\Delta^2 - n\Delta^2 = n\Delta^2 = \frac{m^2}{n}$ .

#### 4. Minimizing maximum outdegree algorithms

In this section we describe the relationship between minimum Laplacian energy of directed graphs and Minimizing maximum out degree algorithms called MMO algorithms [3].




Figure 2: Directed graphs with nodes < 10

**Definition 4.1.** Let  $G_u$  be an undirected graph. The optimal directed Laplacian energy  $LE_{opt}(G_u)$  of  $G_u$  is defined by

$$LE_{opt}(G_u) = \min\{LE(G_d) \mid G_d \text{ is a directed graph and } U(G_d) = G_u\}.$$

**Definition 4.2.** For a directed graph  $G_d = (V_d, E_d)$ , the maximum out degree  $\Delta(G_d)$  of  $G_d$  is defined by

$$\Delta(G_d) = \max\{d^{out}(v) \mid v \in V_d\}.$$

**Definition 4.3.** Let  $G_u$  be an undirected graph. The optimal maximum out-degree  $\Delta_{opt}(G_u)$  of  $G_u$  is defined by

$$\Delta_{opt}(G_u) = \min\{\Delta(G_d) \mid G_d \text{ is a directed graph and } U(G_d) = G_u\}.$$

**Definition 4.4.** Let  $G_d = (V_d, E_d)$  be a directed graph. We denote  $v \rightarrow w$  if  $(v, w) \in E_d$ . We also denote  $v \Rightarrow w$  if there exists  $v_1, v_2, \dots, v_k \in V_d$ , ( $k \geq 1$ ) such that  $v = v_1, v_1 \rightarrow v_2, \dots, v_{k-1} \rightarrow v_k$  and  $v_k = w$ .

**Proposition 4.1.** Let  $G_u = (V_u, E_u)$  be an undirected graph and  $G_d = (V_d, E_d)$  be a directed graph satisfying  $U(G_d) = G_u$ . If  $\Delta(G_d) = p$  and  $\{d^{out}(v) \mid v \in V_d\} = \{p, p-1\}$  then  $LE(G_d) = LE_{opt}(G_u)$ .

(Proof) Let  $k = |\{v \in V_d \mid d^{out}(v) = p\}|$  and  $l = |\{v \in V_d \mid d^{out}(v) = p-1\}|$ . Since  $\sum\{d^{out}(v) \mid v \in V_d\} = |E_u|$ , we have  $pk + (p-1)l = |E_u|$ . Since  $k+l = |V_u|$ ,  $k$  and  $l$  is uniquely determined.  $|E_u| = kp + l(p-1) = |V_u|p + (k - |V_u|)$  and  $|V_u|p = |E_u| + |V_u| - k$ . Hence  $p = \frac{|E_u|}{|V_u|} + 1 - \frac{k}{|V_u|}$ . Since  $1 \leq k \leq |V_u|$  then  $0 \leq 1 - \frac{k}{|V_u|} < 1$ . So we have  $p = \lceil \frac{|E_u|}{|V_u|} \rceil$ .

Let  $h(x_1, x_2, \dots, x_n, \lambda) = \sum_{i=1}^n x_i^2 + 2\lambda(\sum_{i=1}^n x_i - |E_u|)$ . Since  $\frac{\partial h}{\partial x_i} = 2x_i + 2\lambda = 0$  and  $\frac{\partial h}{\partial \lambda} = \sum_{i=1}^n x_i - |E_u| = 0$ , the function  $h$  is minimum at  $x_i = -\lambda$ , ( $i = 1, \dots, n$ ) and  $\sum_{i=1}^n x_i = |E_u|$ . Since  $\sum_{i=1}^n x_i = -|V_u|\lambda = |E_u|$ , we have

$\lambda = -\frac{|E_u|}{|V_u|}$  and  $x_i = \frac{|E_u|}{|V_u|}$ , ( $i = 1, \dots, n$ ). If all  $x_i$ 's are integer, the function  $h$  have minimum value with  $x_i \in \{\lceil \frac{|E_u|}{|V_u|} \rceil, \lceil \frac{|E_u|}{|V_u|} \rceil - 1\}$  for ( $i = 1, \dots, n$ ). We consider  $x_i = d^{out}(v_i)$ . Then  $h(x_1, x_2, \dots, x_n, \lambda) = LE(G_d)$ . So  $LE(G_d) = kp^2 + l(p-1)^2$  gives the optimal solution.  $\square$

Finding the orientation of simple graph by minimizing maximum out degree of a node is studied in literature [13, 3] and defined as MMO (Minimizing maximum out degree) algorithms. In order to minimize the maximum out degree and find a optimal solution to MMO problems, [3] use simple algorithm called reverse algorithm as in Table 4.

Input	An undirected graph $G_u = (V_u, E_u)$
Output	Oriented graph $MMO(G_u) = G_d = (V_d, E_d)$
Step 1:	Set $E_d = \emptyset$
Step 2:	Find arbitrary orientation and update $E_d$
Step 3:	Compute out degree $d^{out}(v)$ for each $v \in V_d$ . Let $u$ be $\max\{d^{out}(v)   v \in V_d\}$
Step 4:	Find a directed path $P = u \rightarrow v_1 \rightarrow \dots \rightarrow v_k$ of length $k(k \geq 1)$ which satisfy $d^{out}(v_i) \leq d^{out}(u), \forall 1 \leq i \leq k-1$ and $d^{out}(v_k) \leq d^{out}(u) - 2$ If such $P$ exists then set $E_d = E_d \setminus \{P \cup \bar{P}\}$ , where $\bar{P} = v_k \rightarrow \dots \rightarrow v_1 \rightarrow u$ and goto Step 2. Otherwise halt.

Table 1: MMO Algorithm

Reverse path cause to reduce the maximum outdegree by one and increase the outdegree of terminal vertex by one. It is proved in [3] that if  $G_d = MMO(G_u)$  then  $\Delta_{opt}(G_u) = \Delta(G_d)$ .

**Example 2.** In Figure 3, we demonstrate two directed graphs  $G_{d_1}$  and  $G_{d_2}$  with  $U(G_{d_1}) = U(G_{d_2}) = G_u$ . The maximum outdegree of  $G_{d_1}$  and  $G_{d_2}$  are same. But  $LE(G_{d_1}) \neq LE(G_{d_2})$ . In Figure 4 we demonstrate four directed graphs  $G_{d_1}, G_{d_2}, G_{d_3}, G_{d_4}$  with  $U(G_{d_i}) = G_u$  and  $LE(G_{d_i}) = LE_{opt}(G_u), (i = 1, 2, 3, 4)$ . We can see  $\{d^{out}(v) | v \in V_{d_i}\} = \{1, 2\}$  and  $|\{v \in V_{d_i} | d^{out}(v) = 1\}| = |\{v \in V_{d_j} | d^{out}(v) = 1\}|$  and  $|\{v \in V_{d_i} | d^{out}(v) = 2\}| = |\{v \in V_{d_j} | d^{out}(v) = 2\}|$  for  $(i, j = 1, 2, 3, 4)$ .

**Proposition 4.2.** Let  $G_u = (V_u, E_u)$  be undirected graph and  $G_d = (V_d, E_d) = MMO(G_u)$ . Let  $v_0 \in V_d$  be a vertex with  $d^{out}(v_0) = \Delta(G_d)$ . Define  $V' = \{w \in V_d | v_0 \Rightarrow w\}$ .  $G'_u = (V', E'_u)$  be the induced undirected graph where  $E'_u = \{\{u, v\} \in E_u | u, v \in V'\}$  and  $G'_d = (V', E'_d)$  be the directed graph where  $E'_d = \{(u, v) \in E_d | u, v \in V'\}$ . Then the following relations are hold.

1.  $d^{out}(v_0) \geq d^{out}(v') \geq d^{out}(v_0) - 1$  for  $v' \in V'$ .
2.  $LE(G'_d) = LE_{opt}(G'_u)$




Figure 3: Equal maximum degree and different Laplacian energy

(Proof)

1. Let  $v' \in V'$ . Then we have a path  $P = v_0 \rightarrow \dots \rightarrow v'$  from  $v_0$  to  $v'$ . Since  $d^{out}(v_0) = \Delta(G_d)$ , we have  $d^{out}(v') \leq d^{out}(v_0)$ . Further  $G_d = MMO(G_u)$  imply that  $d^{out}(v') > d^{out}(v_0) - 2$ . Therefore  $d^{out}(v') \geq d^{out}(v_0) - 1$ .
2. From (1) we have  $\{d^{out}(v) \mid v \in V'\} = \{d^{out}(v_0), d^{out}(v_0) - 1\}$ .  
By Proposition 4.1, we have  $LE(G'_d) = LE_{opt}(G'_u)$ .  $\square$

**Theorem 4.1.** Let  $G_u = (V_u, E_u)$  be an undirected graph and  $G_d = (V_d, E_d) = MMO(G_u)$ . Let  $p = \Delta(G_d)$ ,  $V_p = \{v \in V_d \mid d^{out}(v) = p\}$  and  $V_1 = \{w \in V_d \mid v_p \in V_p, v_p \Rightarrow w\}$ . If  $V_u = V_1$  then  $LE(G_d) = LE_{opt}(G_u)$ .

(Proof) Since  $G_d = MMO(G_u)$  then  $\{d^{out}(v) \mid v \in V_1\} = \{p, p - 1\}$ . If  $V_u = V_1$  then we have  $LE(G_d) = LE_{opt}(G_u)$  by Proposition 4.1.  $\square$

**Example 3.** In figure 5(a)  $MMO(G_u) = (V_d, E_d)$  and  $\{d^{out}(v) \mid v \in V_d\} = \{3, 2, 1\}$ .  $p = \Delta(G_d) = 3$ ,  $V_p = \{9, 11, 12\}$ ,  $V_1 = \{9, 10, 11, 12, 13, 14\}$ . So we cannot apply Theorem 4.1. In Figure 5(b)  $MMO(G_u) = (V_d, E_d)$  and  $\{d^{out}(v) \mid v \in V_d\} = \{2, 1\}$ . We can see  $p = \Delta(G_d) = 2$ ,  $V_p = \{1, 2, 3, 4, 5, 6\}$  and  $V_1 = \{1, 2, 3, 4, 5, 6, 7, 8\}$ . Then we have  $LE(MMO(G_u)) = LE_{opt}(G_u)$  from the Theorem 4.1.






Figure 4: Oriented graphs with minimum Laplacian energy

## 5. Conclusion

We build relations on Laplacian energy of directed graphs. Then we enumerated the structure of the graphs whose Laplacian energy is less than some  $\alpha$  value. Further we considered relationship between MMO algorithms and Laplacian energy. It is remained for the future to further analysis whether MMO algorithms always gives the optimal solution for minimum Laplacian energy.

## 6. Acknowledgment

The authors thank Dr.Tetsuji Taniguchi for his valuable suggestions and discussions. This work has been partially supported by Kyushu University




Figure 5: Graphs consists of optimal Laplacian Energy

Global COE Program "Education-and-Research Hub for Mathematics-for-Industry".

## References

- [1] C. Adiga and M. Smitha. On the skew Laplacian energy of a digraph. *International Math. Forum*, 39:1907–1914, 2009.
- [2] S. Akbari, E. Ghorbani, and M.R. Oboudi. Edge addition, singular values, and energy of graphs and matrices. *Linear Algebra and its Applications*, 430:2192–2199, 2009.
- [3] Y. Asahiro, E. Miyano, H. Ono, and K. Zenmyo. Graph orientation algorithms to minimize the maximum outdegree. *Int. J. Found. Comput. Sci.*, 18:197–215, 2007.
- [4] V. Brankov, D. Stevanovic, and I. Gutman. Equienergetic chemical trees. *J.Seb.Chem.Soc.*, 69:549–553, 2004.
- [5] Cvetković, D.M. and Doob, M. and Sachs, H. *Spectra of Graphs: Theory and Applications*. Academic Press, New York, 1980.

- [6] I. Gutman. Topology and stability of conjugated hydrocarbons. the dependence of total  $\pi$ -electron energy on molecular topology. *J.Seb.chem.Soc.*, 70:441–456, 2005.
- [7] I. Gutman, D. Kiani, M. Mirzakhah, and B. Zhou. On incidence energy of a graph. *Linear Algebra and its Applications*, 431:1223–1233, 2009.
- [8] I. Gutman and B. Zhou. Laplacian energy of a graph. *Linear Algebra and its Applications*, 414:29–37, 2006.
- [9] J.H. Koolen, V. Moulton, and I. Gutman. Improving the McClelland inequality for total  $\pi$ -electron energy. *Chemical Physics Letters*, 320:213–216, 2000.
- [10] M. Lazić. On the Laplacian energy of a graph. *Czechoslovak Math.Journal*, 56:1207–1213, 2006.
- [11] J. Li, W.C. Shiu, and A. Chang. On the Laplacian Estrada index of a graph. *Appl. Anal. Discrete Math.*, 3:147–156, 2009.
- [12] S. Radenković and I. Gutman. Total  $\pi$ -electron energy and Laplacian energy: How far the analogy goes? *Journal of the Serbian Chemical Society*, 73:1343–1350, 2007.
- [13] V. Venkateswaran. Minimizing maximum indegree. *Discrete Applied Mathematics*, 143:374–378, 2004.

# List of MI Preprint Series, Kyushu University

The Global COE Program  
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA  
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO  
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU  
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI  
Torsion points of abelian varieties with values in infinite extensions over a  $p$ -adic field
- MI2008-6 Yoshiyuki TOMIYAMA  
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI  
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA  
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA  
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA  
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA  
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO  
On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA  
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA  
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA  
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO  
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI  
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI  
Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI  
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI  
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA  
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI  
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI  
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA  
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO  
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA  
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA  
Hypergeometric  $q$ -functions of the  $q$ -Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA  
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA  
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI  
Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE  
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO  
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI  
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI  
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO  
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA  
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO  
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI  
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO  
On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection

- MI2009-26 Manabu YOSHIDA  
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI  
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA  
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA  
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA  
On asymptotic behaviors of solutions to parabolic systems modelling chemo-taxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI  
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA  
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA  
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA  
Projective reduction of the discrete Painlevé system of type  $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI  
Finite element computation for scattering problems of micro-hologram using DtN map



- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA  
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA  
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA  
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA  
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI  
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI  
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI  
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA  
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO  
Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA  
An algebraic approach to underdetermined experiments

- MI2010-10 Kei HIROSE & Sadanori KONISHI  
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA  
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDO  
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA  
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI  
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO  
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE  
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU  
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA  
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE  
Lagrangian approach to weakly nonlinear stability of an elliptical flow

- MI2010-21 Hiroki MASUDA  
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU  
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiko FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU  
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra
- MI2010-25 Toshimitsu TAKAESU  
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA  
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA  
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA  
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE  
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI  
Spin-spin correlation functions of the  $q$ -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and spike detection via Gaussian basis expansions

- MI2010-32 Nobutaka NAKAZONO  
Hypergeometric  $\tau$  functions of the  $q$ -Painlevé systems of type  $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI  
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI  
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI  
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms