Frequency of SCEs in Japanese Infants Lactationally Exposed to Organochlorone Pesticides

Nagayama, Junya Laboratory of Environmental Molecular Epidemiology, School of Health Sciences, Faculty of Medicine, Kyushu University

Nagayama, Mayumi Laboratory of Environmental Molecular Epidemiology, School of Health Sciences, Faculty of Medicine, Kyushu University

Nakagawa, Reiko Fukuoka Institute of Health and Environmental Sciences

Hirakawa, Hironori Fukuoka Institute of Health and Environmental Sciences

他

https://doi.org/10.15017/18756

出版情報:福岡醫學雑誌.94(5), pp.166-173,2003-05-25. 福岡医学会 バージョン: 権利関係:

Frequency of SCEs in Japanese Infants Exposed to Dioxins and PCBs through the Breast Milk

Junya NAGAYAMA and Mayumi NAGAYAMA

Laboratory of Environmental Molecular Epidemiology, School of Health Sciences, Faculty of Medicine, Kyushu University, Fukuoka 812-8582

Takao IIDA, Hironori HIRAKAWA and Takahiko MATSUEDA

Fukuoka Institute of Health and Environmental Sciences, Fukuoka 818-0135

Jun'ichiro FUKUSHIGE Fukuoka Children's Hospital, Fukuoka 810-0063

Abstract Frequency of sister chromatid exchanges (SCEs), which has been considered an index to the synthetic and sharp genotoxic and clastogenic potencies, was examined by using the infant lymphocytes postnatal of around ten months, in order to evaluate the genotoxic or clastogenic potency of exposures to PCBs and dioxins through the breast milk. SCE frequency (mean \pm S.D.) as the control culture treated with the solvent, DMSO, alone (SCEcontrol) was 8.3 ± 1.1 /cell and that as the culture treated with 7,8-benzo-flavone (SCEANF) was 11.9 ± 1.5 /cell. In addition, the difference between SCEANF and SCEcontrol, that is, \triangle SCEs was calculated 3.6 ± 1.5 /cell.

The intake of PCBs (mean \pm S.D.) through the breast milk in these infants was 133 ± 85 mg/kg body weight and that of dioxins 24 ± 13 TEQ-ng/kg body weight. PCBs intake in the average was about 5,500 times greater than that of dioxins. We however, could not find any significant effect of PCBs on the frequencies of SCEs. Dioxins, on the contrary, showed a significant negative relationship to the frequency of \triangle SCEs. Therefore, exposure to dioxins through the breast milk seemed to elicit some genotoxic or clastogenic effects on Japanese general infants postnatal of around ten months.

Introduction

Foods in Japan have been polluted with some organochlorine pesticides, polychlorinated biphenyls (PCBs) and dioxins which are polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs) and coplanar PCBs (Co-PCBs)¹⁴⁾²⁷⁾. So, Japanese people have also been contaminated with these compounds⁴⁾⁶⁾⁸⁾. Consequently, PCBs and dioxins have been determined in Japanese breast milk and respective median concentrations of PCBs and dioxins on lipid weight basis were about 110 ppb and 24 ppt as the 2,3,7,8-tetrachlorodibenzo-p-dioxin toxic equivalent (2,3,7,8TCDD TEQ)¹⁵⁾²⁶⁾. These results indicate that the median level of PCB was about 4, 600 times higher than that of dioxins in Japanese breast milk. Therefore, we should give due attention to possible health effects of not only dioxins but also PCBs in Japanese infants.

In particular, prenatal and suckling stages of human life as well as animal life are considered to be the highest sensitivity to these hazardous chemicals, because of their remarkably fast differentiation and growth. We have already done some works from this viewpoint and got several worthwhile findings¹⁶⁾¹⁷⁾¹⁸⁾¹⁹⁾.

In the meantime, the formation mecha-

158

nism of sister chromatid exchanges (SCEs) has not been well clarified yet at present. SCEs, however, seem to be a good index to the synthetic and sharp genotoxic and clastogenic potencies for several chemicals⁹⁾¹⁰⁾²⁹⁾³²⁾. We have also been carried out several studies from this point of view and some important results have been obtained²⁰⁾²¹⁾²²⁾²³⁾²⁴⁾.

In this study, changes of SCE frequency in the lymphocytes of Japanese infants lactationally exposed to PCBs and dioxins were investigated, in order to evaluate their genotoxic and clastogenic potencies.

Materials and Methods

Fifty to 100 ml of breast milk at the postpartum period of 2 to 4 month were collected from 124 healthy mothers in total, mean age : 29 years old and the range : 22 \sim 41 years old, in July to October, 1994 and in June to October, 1995 and 1996. These samples of the breast milk were analyzed for PCBs by ECD gas chromatography method¹¹⁾ and for dioxins by HRGC-HRMS technique using a Finnigan MAT-95 mass spectrometer (Germany) directly interfaced with Varian Model 3400 gas chromatograph⁷⁾¹²⁾¹⁵⁾.

Lymphocytes in the whole blood were stimulated with phytohemagglutinin and cultured for two replicative cycles in the presence of bromodeoxyuridine (100 mM) as detailed elsewhere²⁰⁾²³⁾²⁴⁾. Differential staining of sister chromatids was obtained by a fluorochrome plus Giemsa technique and the frequencies of control, solvent (DMSO) treated SCEs (SCEcontrol), 7,8-benzoflavone (ANF) treated SCEs (SCEANF) and \triangle SCEs (SCEANF - SCEcontrol) were evaluated.

In order to get normal distribution, frequencies of SCE control, SCEANF and \bigtriangleup SCEs and lactational exposures to PCBs and dioxins, namely, intakes of these compounds through the breast milk were transformed by a natural logarithm. Then, relationship between frequencies of SCEcontrol, SCEANF or \triangle SCEs and the intakes of PCBs or dioxins was statistically examined by Spearman rank correlation method.

Toxic equivalent (TEQ) concentrations of the dioxins were calculated by using 1998 WHO TEF values³⁰⁾. TEQ-sum of all congeners of the dioxins determined in every breast milk sample was summarized as the total 2,3,7,8-TCDD TEQ concentration or level. Lactational exposure to PCBs or dioxins was estimated as a product of their respective daily intake, which was calculated with their respective level in the milk times an expected intake of breast milk in Japanese infant, that is, 120g/kg body weight, multiplied by individual duration (days) of breast feeding.

Five to 10 ml of the peripheral blood of 105 infants (60 males and 45 females with the mean age of 10 months old) born of these mothers were individually obtained by venipuncture in January \sim March of 1995, 1996 and 1997. Among them, SCE frequency of the lymphocytes was measured in 66 infants.

Results

1) SCE frequencies of the lymphocytes in the blood of Japanese infants

The frequency distributions of SCEcontrol, SCEANF and \triangle SCEs of lymphocytes in the blood of infants are shown in Fig. 1. The mean value of SCEcontrol was 8.3/cell, and minimum and maximum ones were 6.5 and 12.4/cell, respectively. The average value of SCEANF was 11.9/cell, and the range was 9. $1 \sim 15.6$ /cell. The mean of \triangle SCE was 3.6/

J. Nagayama et al.





cell, and respective values of minimum and maximum were 0.2 and 7.2/cell.

2) Intakes of PCBs and dioxins through the breast milk in Japanese infants

The distributions of total intakes of PCBs and dioxins in Japanese infants during breast-fed periods are indicated in Fig. 2. In case of PCBs, the average intake was 133 mg/kg body weight, and the lowest and highest ones were 28 and 388 mg/kg body weight, respectively. The mean intake of dioxins was 24 TEQ-ng/kg body weight with the range of $3.9 \sim 65$ TEQ-ng/kg body weight. The average intake of PCBs was about 5,500 times greater than that of dioxins in Japanese infants of this study.

 Relationships between lactational exposures to PCBs and dioxins and SCE frequencies of the lymphocytes

We could not find any correlation of the lactational exposures to PCBs with the frequencies of SCE control, SCEANF or \triangle SCEs of the lymphocytes in the blood of Japanese infants.

In case of exposure to dioxins through the breast milk, dioxins did not show any signif-





160





icant relationship with frequencies of SCEcontrol and SCEANF of the lymphocyte, as indicated in Fig. 3. Frequency of \triangle SCEs, however, of the lymphocytes in the blood of infants postnatal of around 10 months was significantly decreased with the increasing lactational exposure to dioxins (r=-0.292, p=0.020), as shown in Fig. 4.

Discussion

Some conflicting results have been reported about the effect of aging on the frequency of SCEs. In some studies, it has been shown that the SCE frequency in human lymphocytes was significantly in-



Fig. 4 Relationship between the lactational exposure to dioxins and the frequency of \triangle SCEs of the lymphocytes in the Japanese infants around postnatal of ten months (r=-0.292, p=0.020)

creased with an $age^{3)28)31}$, and in others not¹⁾⁵⁾¹³⁾. In our studies, frequencies of both SCEcontrol and SCEANF in human lymphocytes at 20 ~ 64-year-old people were clearly enhanced with $age^{21)24}$. In these investigations, average frequencies of SCEcontrol and SCEANF were 10.1 and 13. 6/cell, respectively. In SCE frequency of this study, the former was 8.3/cell, and the latter 11.9/cell, which were apparently lower than the respective values in our previous studies. Based on these findings, it seems the SCE frequency probably increased with age in Japanese people.

In 0 \sim 75-year-old persons, SCE frequency of the lymphocytes in the umbilical cord blood was the highest, that at 1 \sim 5-year-old is the lowest and after these ages it increased with age²). Further, we have to examine SCE frequency of the lymphocytes in umbilical cord blood or in the blood of newborn infant, in order to confirm their findings.

The estimated mean intake of PCBs was 133 mg/kg body weight and that of dioxins 24 TEQ-ng/kg body weight, which was four orders of magnitude lower than that of PCBs. In spite of such relatively large amounts of intake, we have never observed the correlation of PCBs with any frequencies of SCEcontrol, SCEANF and \triangle SCEs in the lymphocyte of Japanese breastfed infants postnatal of around 10 months. Contrastively, lactational exposure to dioxins was significantly and negatively correlated with the \triangle SCE frequency. At present, although we do not known the significance of this finding, the average intake of dioxins through the breast milk in Japanese general infants is about 30 times higher than the tolerable daily intake of dioxins in Japan, that is, 4 TEQ-pg/kg body weight²⁵⁾. Therefore, this level of intake of dioxins may cause some adverse health effects in sucklings, which are one of the most sensitive stages in not only humans but also animals to toxic chemicals. In adults, however, we could not find any significant effect of dioxin like chemicals, mostly PCDFs, on the frequencies of SCEcontrol, SCEANF and \triangle SCEs of the lymphocytes at the levels of more than 300 TEQ-pg/g on lipid weight basis in the blood, which was about 10 times higher than the Japanese normal contamination level of dioxins²³⁾. Accordingly, infants or sucklings are considered more sensitive to dioxins and dioxin like chemicals than adults.

This study, however, was carried out with rather small number of infants, so in order to get more conclusive findings for the effects of dioxins on SCE frequency and for the evaluation of their genotoxic and clastogenic potencies, further large-scale investigations are needed.

References

 Bender MA, Preston RJ, Leonard RC, Pyatt BE and Gooch PC: Chromosomal aberration and sister-chromatid exchange frequencies in peripheral blood lymphocytes of a large human population sample.II. Extension of age range. Mutat. Res. 212: 149-154, 1989.

- Das BC, Rani R, Mitra AB and Luthra UK: Baseline frequency of sisterchromatid exchanges (SCE) in newborn lymphocytes and its relationship to in vivo aging in humans. Mutat. Res. 144: 85-88, 1985.
- Dutkowski RT, Lesh R, Staiano-Coico L, Thaler H, Darlington GJ and Weksler ME: Increased chromosomal instability in lymphocytes from elderly humans. Mutat. Res. 149: 505-512, 1985.
- Hirakawa H, Iida T, Matsueda T and Nagayama J: Concentrations and distribution of PCDDs, PCDFs and Co-PCBs in various tissues of Japanese. Organohal. Comp. 30: 127-130, 1996.
- 5) Hirsh BA, Sentz KK and McGue M: Genetic and environmental influences on baseline SCE. Environ. Mol. Mutagen. 20: 2-11, 1992.
- 6) Iida T, Hirakawa H, Matsueda T, Nakagawa R, Hori T and Nagayama J: Concentrations of PCDDs, PCDFs, Co-PCBs and organochlorine pesticides in the blood and breast milk in Japanese women. Organohal. Comp. 44: 123-127, 1999.
- Iida T, Hirakawa H, Matsueda T, Takenaka S and Nagayama J: Polychlorinated dibenzo-p-dioxins and related compounds in breast milk of Japanese primiparas and multiparas. Chemosphere 38: 2461-2466, 1999.
- Kashimoto T, Takayama K, Mimura M, Miyata H, Murakami Y and Matsumoto H: PCDDs, PCDFs, PCBs, coplanar PCBs and organochlorinated pesticides in human adipose tissue in Japan. Chemosphere 19: 921-926, 1989.
- 9) Latt SA: Sister chromatid exchange formation. Ann. Rev. Genet. 15: 11-55, 1981.
- Latt SA, Allen JW, Shuler L, Loveday KS and Munrow SH: The detection and induction of sister chromatid exchange (INC-UCLA Symposium on Molecular and Cellular Biology). Mol. Hum. Cytogenet. X II. 315-334, 1977.
- 11) Masuda Y, Kagawa R and Kuratsune M : Polychlorinated biphenyls in Yusho patients and ordinary persons. Fukuoka Acta Med. 65 : 17-24, 1974 (in Japanese).

163

- 12) Matsueda T, Iida T, Hirakawa H, Fukamachi K, Tokiwa H and Nagayama J: Toxic evaluation of PCDDs, PCDFs and coplanar PCBs in breast-fed babies of Yusho and healthy mothers. Chemosphere 27: 187-194, 1993.
- 13) Musilova J, Michalova K and Pacovsky V: Induction of sister chromatid exchanges by mitomycin C in lymphocytes of young and old human donors. Gerontology 30: 365-370, 1984.
- 14) Nakagawa R, Hirakawa H and Hori T: Estimation of 1992-1993 dietary intake of organochlorine and organophosphorus pesticides in Fukuoka, Japan. J.AOAC International 78: 921-929, 1995.
- 15) Nakagawa R, Hirakawa H, Iida T, Matsueda T and Nagayama J: Maternal body burden of organochlorine pesticides and dioxins. J. AOAC International 82: 716-724, 1999.
- 16) Nagayama J, Fukushige J, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T and Watanabe T: Developmental condition in 10-month-old Japanese infants perinatally exposed to organochlorine pesticides, PCBs and dioxins. Organohal. Comp. 48: 244-247, 2000.
- 17) Nagayama J, Fukushige J, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T and Watanabe T: Effects of exposure to organochlorine pesticides, PCBs and dioxins through human milk on total development in 10month-old Japanese infants. Organohal. Comp. 48: 240-243, 2000.
- 18) Nagayama J, Tsuji H, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T, Fukushige J and Watanabe T : Condition of helper and suppressor T lymphocyte subpopulations in 10-month-old Japanese infants perinatally exposed to organochlorine pesticides, PCBs and dioxins. Organohal. Comp. 49: 87-90, 2000.
- 19) Nagayama J, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T, Fukushige J and Watanabe T: Condition of thyroid hormone system in 10-month-old Japanese infants perinatally exposed to organochlorine pesticides, PCBs and dioxins. Organohal. Comp. 48:

236-239, 2000.

- 20) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T and Masuda Y: Effects of highly toxic organochlorine compounds retained in human body on induction of sister chromatid exchanges in cultured human lymphocytes. Chemosphere 29: 2349-2345, 1994.
- Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H: Effects of donor age on frequency of sister chromatid exchanges and accumulation of dioxins and related chemicals in healthy Japanese. Organohal. Comp. 44: 197-200, 1999.
- 22) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H: Comparison of frequency of sister chromatid exchanges and contamination level of dioxins and related chemicals in healthy Japanese and "Yusho" patients. Organohal. Comp. 44: 31-35, 1999.
- 23) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H : Comparison between "Yusho" patients and healthy Japanese in contamination level of dioxins and related chemicals and frequency of sister chromatid exchanges. Chemosphere 43: 931-936, 2001.
- 24) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H: Effects of donor age on frequency of sister chromatid exchanges in human lymphocytes cultured in vitro. Chemosphere 43: 845-849, 2001.
- 25) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Yanagawa T and Fukushige J: Effect of dioxins in mother' s milk on sister chromatid exchange frequency in infant lymphocytes. Fukuoka Acta Med. 92: 177-183, 2001.
- 26) Nagayama J, Tsuji H, Okamura K, Iida T, Hirakawa H, Matsueda T, Hasegawa M, Sato K, Tomita A, Yanagawa T, Igarashi H, Fukushige J and Watanaba T: Perinatal exposure to polychlorinated biphenyls on lymphocyte subpopulations and thyroid hormone status in Japanese breast-fed infants. Organohal. Comp. 37: 163-167, 1998.
- 27) Ogaki J, Takayama K, Miyata H and Kashimoto T: Levels of PCDDs and

PCDFs in human tissues and various foodstuffs in Japan. Chemosphere 16: 2047-2056, 1987.

- 28) Sarto F, Faccioli MC, Cominato I and Levis AG: Aging and smoking increase the frequency of sister-chromatid exchanges (SCE) in man. Mutat. Res. 144: 183-187, 1985.
- 29) Stetka DG and Wolff S: Sister chromatid exchanges as assay for genetic damage induced by mutagens-carcinogens II. In vitro test for compounds requiring metabolic activation. Mutat. Res. 41: 343-350, 1976.
- 30) Van den Berg M, Birnbaum LS, Bosveld ATC, Brunstorm B et al.: Toxic

equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ. Health Perspect. 106: 775-792, 1998.

- 31) Wen WN, Liew TL, Wuu SW and Jan KY: The effect of age and cell proliferation on the frequency of sister chromatid exchange in human lymphocytes cultured in vitro. Mech. Ageing Dev. 21: 377-384, 1983.
- 32) Wolff S: Chromosome aberrations, sister chromatid exchanges and the lesions that produce them. In Wolff S (ed): Sister Chromatid Exchange. pp. 41-57, Wiley, New York, 1982.

(Received for publication April 1, 2003)

(和文抄録)

乳児リンパ球細胞の姉妹染色分体交換頻度への授乳による PCBs と ダイオキシン類の影響

九州大学医学部保健学科環境分子疫学研究室

長山淳哉,長山真弓
 福岡県保健環境研究所
 飯田隆雄,平川博仙,松枝隆彦
 福岡市立こども病院
 福重淳一郎

姉妹染色分体交換(Sister chromatid exchanges, SCEs)はこれまで総合的で鋭敏な遺 伝毒性あるいは染色体傷害性の指標として知られ ている。この研究では授乳による PCBs とダイオ キシン類への曝露と遺伝毒性の関係を調べる目的 で,生後10ヶ月前後の一見健康な乳児の末梢血リ ンパ球細胞の SCEs 頻度と PCBs やダイオキシ ン類への曝露との関係を統計学的に解析した。

溶媒のみ処理したコントロール群の SCEs 頻度 (SCEcontrol)(平均値±標準偏差,以下同様)は $8.3\pm1.1/$ 細胞であり,7,8-ベンゾフラボン (ANF)処理した場合のそれ(SCEANF)は11.9± 1.5/細胞であった。また両者の差,つまり、 SCEs(SCEANF - SCEcontrol)は $3.6\pm1.5/$ 細胞 であった。また、これらの乳児の PCBs の母乳か らの摂取量は 133 ± 85 mg/kg 体重であり、ダイオ キシン類では 24 ± 13 TEQ-ng/kg 体重であった。 したがって授乳による PCBs の摂取量はダイオ キシン類よりも約 5,500 倍多かった.

PCB への授乳による乳児の曝露量はダイオキ シン類よりもはるかに多いにもかかわらず, PCBs と SCEs 頻度との関連性は SCEcontrol, SCEANF および \triangle SCEs のいずれに対しても認 められなかった.しかしダイオキシン類の場合に は,統計的に有意ではないけれども母乳からの摂 取量が増えるにしたがって SCEcontrol が上昇し, また逆に SCEANF は減少する傾向を示した.その 結果, \triangle SCEs の有意な低下が観察された.

このような SCEs 頻度への影響がどのような臨 床上の疾病と関連するのか現時点では定かでない が、ダイオキシン類への授乳期の曝露が次世代に 対して何らかの遺伝毒性あるいは染色体傷害性作 用を有する可能性が示唆されたのであり、今後、 さらに詳細な研究を行う必要がある.

Frequency of SCEs in Japanese Infants Lactationally Exposed to Organochlorone Pesticides

Junya NAGAYAMA and Mayumi NAGAYAMA

Laboratory of Environmental Molecular Epidemiology, School of Health Sciences, Faculty of Medicine, Kyushu University, Fukuoka 812-8582

Reiko Nakagawa, Hironori HIRAKAWA, Takahiko Matsueda and Takao IIda

Fukuoka Institute of Health and Environmental Sciences, Fukuoka 818-0135

Jun'ichiro FUKUSHIGE

Fukuoka Children's Hospital, Fukuoka 810-0063

Abstract Changes in the frequency of sister chromatid exchanges (SCEs) in cultured mammalian cells have been considered as indices to the synthetic and sharp genotoxic and clastogenic potencies. Frequency of SCEs in cultured lymphocyte obtained from infants postnatal of around ten months was examined to evaluate the genotoxic or clastogenic potency of lactational exposures to some organochlorine pesticides such as HCHs, DDT and chlordane in this study.

Frequency (median, min. \sim max.) of SCEs as the control culture treated with the solvent, DMSO, alone (SCE control) was 8.0/cell, 6.5 \sim 12.4/cell, that as the culture treated with 7,8-benzoflavone (SCEANF) was 11.8/cell, 9.1 \sim 15.6/cell and the difference between SCEANF and SCEcontrol, that is, \triangle SCEs was 3.9/cell, 0.2: 7.2/cell. Lactational exposures (median, min. \sim max.) to the three organochlorine pesticides were as follows: HCHs; 341 mg/kg body weight, 43 \sim 1449 mg/kg body weight, DDT; 272 mg/kg body weight, 33 \sim 1361 mg/kg body weight and chlordane; 69 mg/kg body weight, 13 \sim 379 mg/kg body weight.

The median exposure level to chlordane through the breast milk was one fourth to one fifth times lower than that to HCHs or DDT and probably due to such small amounts of it's intake, we could not find any correlation of chlordane with frequencies of SCEcontrol, SCEANF and \triangle SCEs in cultured lymphocytes of Japanese infants postnatal of about ten months. In the meantime, frequencies of SCEcontrol and SCEANF showed increasing and decreasing tendencies with the increasing exposure to DDT and also HCHs through the breast milk. In consequence, the frequency of \triangle SCEs was significantly lowered. Therefore, lactational exposure to DDT and HCHs seemed to elicit some genotoxic or clastogenic effects on Japanese general infants at ages of around the months. These compounds were also considered rather potent S-dependent clastogens than non-S-dependent ones. So, further detailed studies are needed not only to get conclusive findings but also from this viewpoint.

Introduction

Our environments including food have been polluted with some organochlorine compounds such as polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), polychlorinated biphenyls (PCBs) and pesticides¹¹⁾²⁷⁾. Accordingly, Japanese people have also been contaminated with these compounds³⁾⁶⁾. Some pesticides such as hexachlorocyclohexans (HCHs), 1,1,1-trichloro-2, 2-bis- (4-chlorophenyl) -ethane (DDT), dieldrin, heptachlor epoxide (HCE) and chlordane have been determined in Japanese breast milk⁴⁾¹²⁾¹³⁾¹⁴⁾¹⁵⁾²⁴⁾. Their levels in the breast milk were more than 100 to 10,000 times higher than those of PCDDs and PCDFs, which were so-called dioxins, in 2,3, 7,8-tetrachlorodibenzo-p- dioxin (2,3,7,8-TCDD) toxic equivalent (TEQ) value, as indicated in the above mentioned studies. Therefore, we should give due attention to possible health consequences of these pesticides as well as PCDDs and PCDFs. In particular, prenatal and suckling stages of not only humans but also animals are considered the highest sensitivity to these compounds, because of their remarkably rapid

differentiation and growth.

In the meantime, cytogenetic changes such as micronucleated cells (MNs) and sister chromatid exchanges (SCEs) have frequently been utilized as indicators of mostly genetic damage due to exposure to different carcinogens or mutagens. These two cytogenetic changes are considered to occur as results of different mechanisms of DNA or chromosome damage. MNs have been considered to be the result of chromosomes lagging behind the genome at cell division. On the other hand, SCEs are formed during the S phase after an initial change in the form of DNA base damage²⁸⁾, when quadriradials as mitotic chiasmas are a consequence of mitotic crossing over³⁰). Changes in frequency of MNs and SCEs have been considered good indices to the synthetic and sharp genotoxic or clastogenic potency for several chemicals⁷⁾⁸⁾²⁶⁾²⁹⁾. We have already carried out several studies from this viewpoint and some valuable results have been obtained¹⁷⁾¹⁸⁾¹⁹⁾²⁰⁾²¹⁾.

In this study, changes of SCEs frequency in cultured lymphocytes of Japanese infants exposed to HCHs, DDT and chlordane through the breast milk were investigated, in order to evaluate their genotoxic and clastogenic potencies.

Materials and Methods

Fifty to 100 ml of breast milk at the postpartum period of 2nd to 4th month were collected from 124 healthy mothers in total, mean age: 29 years old and the range: 22 \sim 41 years old, in July to October, 1994 and in June to October, 1995 and 1996. These samples of the breast milk were analyzed for HCHs, DDT and chlordane by ECD gas chromatography method^{11)12)13)14)15)24).}

Lactational exposure to these pesticides was estimated as a product of their respective daily intake, which was calculated with their respective level in the milk times an expected intake of breast milk in Japanese infants, that is, 120g/kg body weight, multiplied by individual duration (days) of breast feeding.

Five to 10 ml of the peripheral blood of 105 infants (60 males and 45 females with the mean age of 10 months old) born of these mothers were individually obtained by venipuncture in January \sim March of 1995, 1996 and 1997. Among them, SCE frequency of the lymphocytes was measured in 63 infants.

Lymphocytes in the whole blood were stimulated with phytohemagglutinin and cultured for two replicative cycles in the presence of bromodeoxyuridine (100 mM) as detailed elsewhere¹⁷⁾¹⁸⁾¹⁹⁾²³⁾. Differential staining of sister chromatids was obtained by a fluorochrome plus Giemsa technique and the frequencies of control, solvent (DMSO) treated SCEs (SCEcontrol), 7,8-benzoflavone (ANF) treated SCEs (SCEANF) and \triangle SCEs (SCEANF – SCEcontrol) were evaluated.

In order to get normal distribution, fre-

quencies of SCEcontrol, SCEANF and \triangle SCEs and lactational exposures to HCHs, DDT and chlordane, namely, intakes of these compounds through the breast milk were transformed by a natural logarithm. Then, relationship between SCEcontrol, SCEANF or \triangle SCEs and the intakes of the organochlorine pesticides was statistically examined by Spearman rank correlation method.

Results

The distributions in the total intakes of HCHs, DDT and chlordane in Japanese infants during breast-fed periods are shown in Fig. 1. Total intakes (median, min. \sim max.) of HCHs (mostly b-HCH), DDT (sum of p, p'-DDE and p, p'-DDT) and chlordane (sum of oxychlordane, trans-nonachlor and cis-nonachlor) were as follows : HCHs; 341 mg/kg body weight, 43 \sim 1449 mg/kg body weight, DDT ; 272 mg/kg body weight, 33 \sim 1361 mg/kg body weight and chlordane ; 69 mg/kg body weight. Median intake of chlordane were one order of magnitude lower than that of HCHs or DDT.

Frequencies (median, min. \sim max.) of SCEcontrol, SCEANF and \triangle SCEs of cultured lymphocytes in the blood of 63 Japanese breast-fed infants were 8.0/cell, 6. $5 \sim 12.4$ /cell, 11.8/cell, 9.1 \sim 15.6/cell and 3.9/cell, 0.2 \sim 7.2/cell, respectively.

We could not find any significant relationship between the lactational exposure to chlordane and the frequency of SCEcontrol, SCEANF or \triangle SCEs. Significant correlations were not observed either between the frequency of SCEcontrol or SCEANF and the intake of HCHs or DDT. Fig. 2 indicates the correlations of the lactational exposure to DDT with the frequency of SCEcontrol or SCEANF. We, however, could see that the frequency of \triangle SCEs of lymphocytes in the blood of infants postnatal of around ten months was significantly decreased with the increasing intake of DDT through the breast milk (r=-0.247, p=0.050), as shown in Fig. 3. This kind of relationship was also observed between the exposure to HCHs and the frequency of \triangle SCEs (r=-0.274, p=0.065), which was indicated in Fig. 4.



Fig. 1 Distributions in total intakes of HCHs* (left), DDT** (middle) and chlordane*** (right) through the breast milk in Japanese infants postnatal of around ten months
*; mostly β-HCH, **; sum of p, p'-DDE and p, p'-DDT, ***; sum of oxychlordane, *trans*- nonachlor and *cis*-nonachlor



Fig. 2 Relationship between the exposure to DDT through the breast milk and the frequency of SCEcontrol (left, r=0.109, p=0.405) or SCEANF (right, r=-0.029, p=0.824) of the lymphocytes in the blood of the Japanese infants postnatal of around ten months

Discussion

The estimated median intake of chlordane was 69 mg/kg body weight, which was one fourth to one fifth of that of HCHs or DDT. Probably due to such small amounts of intake of chlordane through the breast milk, we could not observe any correlation of chlordane with frequencies of SCEcontrol, SCEANF and \triangle SCEs in cultured lymphocytes of Japanese infants postnatal of around ten months.

Figs. 2 and 3 show relationship between the lactational exposure to DDT and the





Fig. 3 Relationship between the lactational exposure to DDT and the frequency of \triangle SCEs of the lymphocytes in the Japanese infants postnatal of around ten months (r=-0.247, p=0.050)



Fig. 4 Correlation of exposure to HCHs through the breast milk with the frequency of \triangle SCEs of the lymphocytes in the blood of the Japanese infants (r=-0.274, p=0.065)

genotoxic and clastogenic potencies¹⁾⁹⁾¹⁰⁾. Although we do not know the significance of these findings at present, lactational exposures to these organochlorine compounds seem to elicit some genotoxic or clastogenic effects in Japanese general infants postnatal of around ten months.

The types of mutations which could contribute to spontaneous MNs include (a) mutations to kinetochore proteins, centromeres and spindle apparatus that could lead to unequal chromosome distribution or whole chromosome loss at anaphase, and (b) unrepaired DNA-strand breaks induced endogeneously or as a result of exposure to environmental mutagens which may result in acentric chromosome fragments²⁾. Therefore, the assay of MNs can detect both clastogens and spindle poisons and can be preferentially used to estimate the dose of ionizing radiations or truly radiomimetic (non-S-dependent) chemicals to which people have been exposed. On the other hand, SCEs, which are not readily induced by ionizing radiations or non-S-dependent clastogens, have proved to be the most sensitive mammalian genotoxic endpoint for determining exposure to S-dependent chemicals, because SCEs are considered to form during the S-phase of the cell cycle³⁰, probably at DNA replication forks or sites where replication is incomplete⁵⁾²⁵⁾. Therefore, according to the results of this study and our other investigations16)22), DDT, HCHs and dioxins seemed rather potent Sdependent clastogens than non-S-dependent ones.

References

 Colman GW, Lundgren K, Shore D, Thomoson CL and Lucier GW : Effects of a-naphthoflavone on levels of sister chromatid exchanges in lymphocytes from active and passive cigarette smokers : dose-response relationships. Cancer Res. 46 : 6452-6455, 1986.

- Fenech M: The cytokinesis-block micronucleus technique: a detailed description of the method and its application to genotoxicity studies in human populations. Mutation Res. 285: 35-44, 1993.
- Hirakawa H, Iida T, Matsueda T and Nagayama J: Concentrations and distribution of PCDDs, PCDFs and Co-PCBs in various tissues of Japanese. Organohal. Comp. 30: 127-130, 1996.
- 4) Iida T, Hirakawa H, Matsueda T, Nakagawa R, Hori T and Nagayama J: Concentrations of PCDDs, PCDFs, Co-PCBs and organochlorine pesticides in the blood and breast milk in Japanese women. Organohal. Comp. 44: 123-127, 1999.
- 5) Ishii Y and Bender MA : Effects of inhibitors of DNA synthesis on spontaneous and ultraviolet light-induced sisterchromatid exchanges in Chinese hamster cell. Mutation Res. 79 : 19-32, 1980.
- 6) Kashimoto T, Takayama K, Mimura M, Miyata H, Murakami Y and Matsumoto H: PCDDs, PCDFs, PCBs, coplanar PCBs and organochlorinated pesticides in human adipose tissue in Japan. Chemosphere 19: 921-926, 1989.
- 7) Latt SA: Sister chromatid exchange formation. Ann. Rev. Genet. 15: 11-55, 1981.
- 8) Latt SA, Allen JW, Shuler L, Loveday KS and Munrow SH: The detection and induction of sister chromatid exchange (INC-UCLA Symposium on Molecular and Cellular Biology). Mol. Hum. Cytogenet. X II. 315-334, 1977.
- 9) Lundgren K, Andries M, Thompson C and Lucier GW: Dioxin treatment of rats in increased in vitro induction of sister chromatid exchanges by a-naphthoflavone: an animal model for human exposure to halogenated aromatics. Toxicol. Appl. Pharmacol. 85: 198-195, 1986.
- Lundgren K, Collman GW, Wang-Wuu S, Tiernan T, Taylor M, Thompson CL and Lucier GW: Cytogenetic and chemical detection of human exposure to polyhalogenated aromatic hydrocarbons. Environ. Mol. Mutagen. 11: 1-11, 1988.
- 11) Nakagawa R, Hirakawa H and Hori T: Estimation of 1992-1993 dietary intake of

organochlorine and organophosphorus pesticides in Fukuoka, Japan. J.AOAC International 78: 921-929, 1995.

- 12) Nakagawa R, Hirakawa H, Iida T, Matsueda T and Nagayama J: Maternal body burden of organochlorine pesticides and dioxins. J. AOAC International 82: 716-724, 1999.
- 13) Nagayama J, Fukushige J, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T and Watanabe T: Developmental condition in 10-month-old Japanese infants perinatally exposed to organochlorine pesticides, PCBs and dioxins. Organohal. Comp. 48: 244-247, 2000.
- 14) Nagayama J, Fukushige J, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T and Watanabe T: Effects of exposure to organochlorine pesticides, PCBs and dioxins through human milk on total development in 10month-old Japanese infants. Organohal. Comp. 48: 240-243, 2000.
- 15) Nagayama J, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T, Fukushige J and Watanabe T: Condition of thyroid hormone system in 10-month-old Japanese infants perinatally exposed to organochlorine pesticides, PCBs and dioxins. Organohal. Comp. 48: 236-239, 2000.
- 16) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T and Fukushige J: Frequency of SCEs in Japanese infants exposed to dioxins and PCBs through the breast milk. Fukuoka Acta Med. In press.
- 17) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T and Masuda Y: Effects of highly toxic organochlorine compounds retained in human body on induction of sister chromatid exchanges in cultured human lymphocytes. Chemosphere 29: 2349–2345, 1994.
- 18) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H : Comparison between "Yusho" patients and healthy Japanese in contamination level of dioxins and related chemicals and frequency of sister chromatid exchanges. Chemosphere 43: 931-936, 2001.
- 19) Nagayama J, Nagayama M, Iida T, Hira-

kawa H, Matsueda T, Ohki M and Tsuji H: Effects of donor age on frequency of sister chromatid exchanges in human lymphocytes cultured in vitro. Chemosphere 43: 845–849, 2001.

- 20) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H: Effects of donor age on frequency of sister chromatid exchanges and accumulation of dioxins and related chemicals in healthy Japanese. Organohal. Comp. 44: 197-200, 1999.
- 21) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Ohki M and Tsuji H: Comparison of frequency of sister chromatid exchanges and contamination level of dioxins and related chemicals in healthy Japanese and "Yusho" patients. Organohal. Comp. 44: 31-35, 1999.
- 22) Nagayama J, Nagayama M and Masuda Y: Frequency of micronuclei induced in cultured human lymphocytes by highly toxic organochlorine congeners. Fukuoka Acta Med. 84: 189–194, 1993.
- 23) Nagayama J, Nagayama M, Iida T, Hirakawa H, Matsueda T, Yanagawa T and Fukushige J: Effect of dioxins in mother' s milk on sister chromatid exchange frequency in infant lymphocytes. Fukuoka Acta Med. 92: 177-183, 2001.
- 24) Nagayama J, Tsuji H, Iida T, Nakagawa R, Matsueda T, Hirakawa H, Astuti ET, Yanagawa T, Fukushige J and Watanabe T : Condition of helper and suppressor T lymphocyte subpopulations in 10-monthold Japanese infants perinatally exposed to organochlorine pesticides, PCBs and dioxins. Organohal. Comp. 49: 87-90, 2000.
- Painter RB: A replication model for sister-chromatid exchange. Mutation Res. 70: 337-341, 1980.
- 26) Stetka DG and Wolff S: Sister chromatid exchanges as assay for genetic damage induced by mutagens-carcinogens II. In vitro test for compounds requiring metabolic activation. Mutat. Res. 41: 343-350, 1976.
- 27) Swedish Environmental Protection Agency: Persistent Organic Pollutants pp. 9-129, ISBN 91-620-1189-8, ISSN 1100-23X, 1998.

(93)

²⁸⁾ Uggla AH and Natarajan AT: X-ray-

induced SCEs and chromosomal aberrations in CHO cells. Influence of nitrogen and air during irradiation in different stages of the cell cycle. Mutation Res. 122: 193-200, 1983.

- 29) Wolff S: Chromosome aberrations, sister chromatid exchanges and the lesions that produce them. In Wolff S (ed): Sister Chromatid Exchange. pp. 41-57, Wiley, New York, 1982.
- 30) Wolff S, Bodycote J and Painter RB: Sister chromatid exchanges induced in Chinese hamster cells by UV irradiation of different stages of the cell cycle: the necessity for cells to pass through S. Mutation Res. 25: 73-81, 1974.

(Received for publication April 1, 2003)

(和文抄録)

乳児リンパ球細胞の姉妹染色分体交換頻度への授乳による農薬の影響

九州大学医学部保健学科環境分子疫学研究室 長山淳哉,長山真弓 福岡県保健環境研究所 中川礼子,平川博仙,松枝隆彦,飯田隆雄 福岡市立こども病院 福重淳一郎

総合的で鋭敏な遺伝毒性指標として知られてい る姉妹染色分体交換(Sister chromatid exchanges, SCEs)頻度を用いて,授乳による HCHs, DDT およびクロルデンへの曝露と乳児リ ンパ球細胞の SCEs 頻度との関係を数理統計学的 に調べた.

リンパ球細胞の培養中に溶媒 (DMSO) のみ処 理したコントロール群の SCEs 頻度すなわち SCEcontrol (中央値,最小値~最大値,以下同様) は8.0/細胞, 6.5~12.4/細胞, 7,8-ベンゾフラボ ン (ANF) 処理した場合の SCEs 頻度, SCEANF は11.8/細胞, 9.1~15.6/細胞, また両者の差 SCEANF - SCEcontrol, \triangle SCEs は3.9/細胞, 0. 2~7.2/細胞であった.母乳からの有機塩素系農薬 の摂取量は次のようである. HCHs; 341 mg/kg 体重, 43~1449 mg/kg 体重, DDT; 272 mg/kg 体重, 33~1361 mg/kg 体重, クロルデン; 69 mg/ kg 体重, 13~379 mg/kg 体重.

クロルデンの授乳による曝露量は中央値で比較 するとHCHsとDDTの4分の1から5分の1 程度と少なく, SCEs 頻度との関係も認められな かった.一方, HCHs と DDT については母乳から のこれらの農薬の摂取量が増加するにつれて有意 ではないけれども SCEcontrol が上昇し、逆に SCEANF が減少する傾向が観察された。その結 果,いずれの農薬でも△ SCEs 頻度の有意な低下 が認められた. このような SCEs への影響が臨床 上どのような意義があるのか不明であるが, HCHsや DDT への授乳期の曝露が次世代に対し て何らかの遺伝毒性作用を及ぼす可能性が示唆さ れたのであり, 今後, さらに詳細な研究が必要で ある.また同様の結果は母乳からのダイオキシン 類への曝露でも観察されているが、SCEs への影 響は細胞分裂のS期依存型遺伝毒性物質とみなさ れ,この視点からの研究も重要である.

(95)