Introducing Viewpoints of Mechanics into Basic Growth Analysis : (XV) Relationships between Richards Growth Function and Basic Growth Function

Shimojo, Masataka

Nakano, Yutaka

University Farm, Faculty of Agriculture, Kyushu University

Ishimatsu, Satoshi

School of Medicine, Kyushu University

Kakihara, Hidetoshi

Laboratory of Regulation in Metabolism and Behavior, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

他

<https://doi.org/10.5109/18839>

出版情報 : 九州大学大学院農学研究院紀要. 55 (2), pp.259-260, 2010-10-29. Faculty of Agriculture, Kyushu University

バージョン :

権利関係 :

Introducing Viewpoints of Mechanics into Basic Growth Analysis – (XV) Relationships between Richards Growth Function and Basic Growth Function –

Masataka SHIMOJO*, Yutaka NAKANO¹, Satoshi ISHIMATSU²,
Hidetoshi KAKIHARA³, Tao SHAO and Manabu TOBISA⁴

Laboratory of Regulation in Metabolism and Behavior, Division of Animal and Marine Bioresource Sciences,
Department of Bioresource Sciences, Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan
and Department of Grassland and Forage Science, College of Animal Sciences and Technology,
Nanjing Agricultural University, Weigang 1, Nanjing 210095, the People's Republic of China
(Received June 30, 2010 and accepted July 9, 2010)

This study was designed to investigate relationships between Richards growth function and basic growth function by comparing them using growth dynamics. There are inflection points in Richards growth function, but basic growth function expresses exponential curves only. To compare these two functions that are greatly different, the first and second derivatives were incorporated into each of them. The results obtained were as follows. Richards growth function was related to basic growth function when combined with derivatives and $m = 1$, though $m = 1$ was prohibited in Richards growth function itself. The occurrence of this relationship was a mathematical result of growth dynamics based on differential. It was suggested from the present study that Richards growth function was related to basic growth function when combined with the first and second derivatives.

INTRODUCTION

The basic growth analysis is a simple method to analyze weight changes of the plant (Blackman, 1919; Watson, 1952; Radford, 1967; Hunt, 1990) and the animal (Brody, 1945). There is, however, a limitation in the application to overall growth curve. The problem of this analysis is that basic growth function gives only an exponential increase, and does not give inflection points that appear in the curve of actual growth. It is well known that there are various improved growth functions that predict actual growth of plants and animals. Logistic function, which is one of them, was related with basic growth function using a concept of growth dynamics (Shimojo *et al.*, 2009). Osumi and Ishikawa (1983) reported that Richards growth function (Richards, 1959) had strong points in describing, analyzing and predicting actual growth, probably a superior function to others in the flexibility to fit growth curves of various types.

The present study was designed to investigate relationships between Richards growth function and basic growth function by comparing them using growth dynamics.

RICHARDS GROWTH FUNCTION AND BASIC GROWTH FUNCTION AS RELATED USING GROWTH DYNAMICS

Dynamics of Richards growth function

Richards growth function (Richards, 1959) is given by

$$W_R = A \cdot (1 - b \cdot \exp(-k \cdot t))^{1/(1-m)}, \quad (1)$$

where A , b , k and m are constants, $m \neq 1$, t = time. Introducing viewpoints of mechanics into Richards growth function (1) gives

$$\begin{aligned} dW_R/dt &= (A/(1-m)) \cdot (1-b \cdot \exp(-k \cdot t))^{m/(1-m)} \cdot (b \cdot k \cdot \exp(-k \cdot t)), \\ d^2W_R/dt^2 &= (A/(1-m)^2) \cdot (1-b \cdot \exp(-k \cdot t))^{(2m-1)/(1-m)} \\ &\quad \cdot (b \cdot k^2 \cdot \exp(-k \cdot t)) \cdot (m-1+b \cdot \exp(-k \cdot t)), \end{aligned} \quad (2)$$

where dW_R/dt = growth rate, d^2W_R/dt^2 = growth acceleration, $m \neq 1$.

Combining functions (1), (2) and (3) leads to

$$\begin{aligned} (dW_R/dt)^2 &= W_R \cdot (d^2W_R/dt^2) \cdot ((b \cdot \exp(-k \cdot t)) / \\ &\quad (m-1+b \cdot \exp(-k \cdot t))). \end{aligned} \quad (4)$$

Dynamics of basic growth function

Basic growth function is given by

¹ University Farm, Faculty of Agriculture, Kyushu University, Kasuya, Fukuoka 811–2307

² School of Medicine, Kyushu University, Fukuoka 812–8582

³ Laboratory of Regulation in Metabolism and Behavior, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

⁴ Faculty of Agriculture, Miyazaki University, Miyazaki 889–2192, Japan

* Corresponding Author (E-mail: mshimojo@agr.kyushu-u.ac.jp)

$$W_B = W_0 \cdot \exp(r \cdot t), \quad (5)$$

where W = weight, t = time, $W_0 = W$ at $t = 0$, r = relative growth rate.

Introducing viewpoints of mechanics into basic growth function (5) gives

$$dW_B/dt = r \cdot W_0 \cdot \exp(r \cdot t), \quad (6)$$

$$d^2W_B/dt^2 = r^2 \cdot W_0 \cdot \exp(r \cdot t), \quad (7)$$

where dW_B/dt = growth rate, d^2W_B/dt^2 = growth acceleration.

Combining functions (5), (6) and (7) leads to

$$(dW_B/dt)^2 = W_B \cdot (d^2W_B/dt^2). \quad (8)$$

Relating Richards growth function with basic growth function using growth dynamics

It is possible to substitute 1 for m in function (4), though $m \neq 1$ is required in Richards growth function (1) and its first derivative (2) and second derivative (3). Thus,

$$\begin{aligned} & (dW_R/dt)^2 \\ &= W_R \cdot (d^2W_R/dt^2) \cdot ((b \cdot \exp(-k \cdot t)) / \\ & \quad (1 - 1 + b \cdot \exp(-k \cdot t))) \\ &= W_R \cdot (d^2W_R/dt^2). \end{aligned} \quad (9)$$

By $m = 1$, equation (4) results in equation (9). Equation (9) takes the same form as that of equation (8),

$$(dW_R/dt)^2 = W_R \cdot (d^2W_R/dt^2), \quad (9)$$

$$(dW_B/dt)^2 = W_B \cdot (d^2W_B/dt^2). \quad (8)$$

This result suggests a kind of collapse of Richards growth function into basic growth function. However, it was reported that Richards growth function tends to Gompertz function when m tends to 1 (Richards, 1959). This contradiction remains to be investigated.

Conclusions

It is suggested from the present study that Richards growth function is related to basic growth function when combined with the first and second derivatives.

ACKNOWLEDGEMENTS

This study was supported by a Grant-in-Aid for Challenging Exploratory Research from Japan Society for the Promotion of Science (No. 21658089).

REFERENCES

- Blackman, V. H. 1919 The compound interest law and plant growth. *Ann. Bot.*, **33**: 353–360
- Brody, S. 1945 Time relations of growth of individuals and populations. In “Bioenergetics and growth – with special reference to the efficiency complex in domestic animals”, Reinhold Publishing Corporation, New York, pp. 484–574
- Hunt, R. 1990 *Basic Growth Analysis*. Unwin Hyman Ltd., London.
- Osumi, S. and Y. Ishikawa 1983 Applicability of the Richards' growth function to analysis of growth of tree. *The Sci. Rep. Kyoto Pref. Univ. Agric.* **35**: 49–76
- Radford, P. J. 1967 Growth analysis formulae—their use and abuse. *Crop Sci.*, **7**: 171–175
- Richards, F. J. 1959 A flexible growth function for empirical use. *J. Exp. Bot.* **10**: 290–300
- Shimojo, M., T. Shao, S. Ishimatsu, J. Tanoue, H. Kakiyama, C. Sata, H. Fukudome, R. Ishiwaka, Y. Asano, Y. Nakano, M. Tobisa and Y. Masuda 2009 Introducing viewpoints of mechanics into basic growth analysis – (XIII) Comparing growth mechanics between logistic functions and basic growth functions –. *J. Fac. Agr., Kyushu Univ.*, **54**: 361–363
- Watson, D. J. 1952 The physiological basis of variation in yield. *Adv. Agron.*, **4**: 101–145