

## Anderson-Darling test and the Malliavin calculus

Hirotsu, Takehiro  
Nisshin Fire & Marine Insurance Company, Limited

Kondo, Hiroki  
Nisshin Fire & Marine Insurance Company, Limited

Saito, Shingo  
Faculty of Mathematics, Kyushu University

Sato, Takuya  
Faculty of Mathematics, Kyushu University

他

<https://hdl.handle.net/2324/18973>

---

出版情報 : MI Preprint Series. 2011-3, 2011-02-17. 九州大学大学院数理学研究院  
バージョン :  
権利関係 :

# MI Preprint Series

Kyushu University  
The Global COE Program  
Math-for-Industry Education & Research Hub

## Anderson-Darling test and the Malliavin calculus

Takehiro Hirotsu,  
Hiroki Kondo, Shingo Saito,  
Takuya Sato, Tatsushi Tanaka  
& Setsuo Taniguchi

MI 2011-3

( Received February 17, 2011 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# Anderson-Darling test and the Malliavin calculus

TAKEHIRO HIROTSU<sup>†</sup>, HIROKI KONDO<sup>†</sup>, SHINGO SAITO<sup>‡</sup>,  
TAKUYA SATO<sup>†</sup>, TATSUSHI TANAKA<sup>‡</sup>, AND SETSUO TANIGUCHI<sup>‡</sup>

February 14, 2011

## Abstract

The quadratic Wiener functional coming from the Anderson-Darling test statistic is investigated in the framework of the Malliavin calculus. The functional gives a completely new and important example of a quadratic Wiener functional.

*Keywords.* Anderson-Darling test, the Malliavin calculus, quadratic Wiener functional, Legendre polynomial

## 1. INTRODUCTION

In 1952, T. Anderson and D. Darling ([1]) introduced a statistical test, which is now called the Anderson-Darling test, to assert if a sample of data arises from a particular probability distribution. Under the null hypothesis that a sample of data  $\{x_1 < x_2 < \cdots < x_n\}$  comes from a distribution with cumulative distribution function  $F$ , the Anderson-Darling test statistic is defined by

$$A_n^2 = -n - \frac{1}{n} \sum_{j=1}^n (2j-1) \{\log u_j + \log(1 - u_{n-j+1})\}, \quad \text{where } u_i = F(x_i), \quad i = 1, \dots, n.$$

If  $A_n^2$  is large, the null hypothesis is rejected ([1, 2, 9]).

The test statistic  $A_n^2$  admits the integral expression that

$$A_n^2 = n \int_0^1 \left( \frac{\nu_n(t)}{n} - t \right) \psi(t) dt \quad (\nu_n(t) \equiv \#\{k \mid u_k \leq t\}).$$

Then, due to Donsker's invariance principle ([3, 4]), as  $n \rightarrow \infty$ ,  $A_n^2$  converges in law to the random variable

$$\int_0^1 \frac{b_t^2}{t(1-t)} dt,$$

$\{b_t\}_{t \in [0,1]}$  being the pinned Brownian motion ([1]). In [1], Anderson-Darling studied the above random variable and its extension

$$\mathfrak{a}^2(\psi) = \int_0^1 b_t^2 \psi(t) dt,$$

where  $\psi : (0, 1) \rightarrow [0, \infty)$  is Borel measurable, by using the Fourier series decomposition of the Gaussian process  $\{b_t \sqrt{\psi(t)}\}_{t \in [0,1]}$ .

---

<sup>†</sup>Nisshin Fire & Marine Insurance Co., Ltd, 2-3 Kanda Surugadai, Chiyoda-ku, Tokyo, 101-8329, Japan

<sup>‡</sup>Faculty of Mathematics, Kyushu University, Fukuoka 819-0395, Japan

The aim of this paper is to investigate  $\mathfrak{a}^2(\psi)$  from the viewpoint of the Malliavin calculus, which enables one to deal with such Wiener functional as  $\mathfrak{a}^2(\psi)$  in a systematic manner ([6, 7, 10]). A Wiener functional is said to be quadratic if its third derivative in the sense of the Malliavin calculus vanishes. In the study of stochastic oscillatory integrals, which are a probabilistic counterpart of Feynman path integrals, quadratic Wiener functionals play an essential role. Typical examples of quadratic Wiener functionals are the square norm and the variance on the time interval of the standard Brownian motion, and Lévy's stochastic area. See for example [7]. The Wiener functional  $\mathfrak{a}^2(\psi)$  gives a new example of a quadratic Wiener functional.

The Wiener functional  $\mathfrak{a}^2(\psi)$  is a not only new but also very important example. In our knowledge, the concrete examples of quadratic Wiener functionals all relate to the standard Brownian motion. As we shall see later, for  $\psi$  which explodes at the order of  $(1-t)^{-\alpha}$  at  $t=1$  for some  $0 < \alpha < 2$ ,  $\mathfrak{a}^2(\psi)$  is well-defined for the pinned Brownian motion, but not for the standard Brownian motion.

In Section 2, a brief review on the Malliavin calculus, in particular, stochastic oscillatory integrals will be given. Section 3 is devoted to the studies of explicit expressions of the stochastic oscillatory integral associated with  $\mathfrak{a}^2(\psi)$ .

## 2. THE MALLIAVIN CALCULUS

### 2.1. WIENER SPACE OF PINNED BROWNIAN MOTION

In this subsection, we shall give a brief review on the one dimensional classical Wiener space of pinned Brownian motion on  $[0, 1]$ . Let  $\mathcal{W}_0$  be the space of real-valued continuous functions  $w$  on  $[0, 1]$  with  $w(0) = w(1) = 0$ . Being equipped with the uniform convergence norm,  $\mathcal{W}_0$  is a real separable Banach space. Denote by  $\mathcal{H}_0$  the subspace of  $\mathcal{W}_0$  consisting of  $h$  which admits square integrable derivative  $h'$ ;  $h(t) = \int_0^t h'(s)ds$ ,  $t \in [0, 1]$ , and  $\int_0^1 (h'(s))^2 ds < \infty$ .  $\mathcal{H}_0$  is a separable Hilbert space with inner product

$$\langle h, g \rangle = \int_0^1 h'(t)g'(t)dt, \quad h, g \in \mathcal{H}_0.$$

The associated norm of  $\mathcal{H}_0$  is denoted by  $\|\cdot\|$ . It is easily seen that  $\mathcal{H}_0$  is embedded continuously and densely in  $\mathcal{W}_0$ . The dual space  $\mathcal{W}'_0$  of  $\mathcal{W}_0$  is included in  $\mathcal{H}_0$  continuously and densely under the standard identification of  $\mathcal{H}_0$  and its dual space  $\mathcal{H}'_0$ . Let  $\mu$  be a Gaussian measure on  $(\mathcal{W}_0, \mathcal{B}(\mathcal{W}_0))$ ,  $\mathcal{B}(\mathcal{W}_0)$  being the topological  $\sigma$ -field of  $\mathcal{W}_0$ , such that

$$(1) \quad \int_{\mathcal{W}_0} \exp(\sqrt{-1}\ell)d\mu = \exp(-\|\ell\|^2/2) \quad \text{for any } \ell \in \mathcal{W}'_0,$$

where an element of  $\mathcal{W}'_0$  is regarded as a random variable on  $\mathcal{W}_0$  in the natural manner. The triplet  $(\mathcal{W}_0, \mathcal{H}_0, \mu)$  is the one dimensional classical Wiener space of pinned Brownian motion on  $[0, 1]$ .

Let  $b_t$ ,  $t \in [0, 1]$ , be the coordinate mapping on  $\mathcal{W}_0$ ;  $b_t(w) = w(t)$ ,  $w \in \mathcal{W}_0$ . Due to (1),  $\{b_t\}_{t \in [0, 1]}$  is a continuous Gaussian process with mean zero and covariance function

$$\int_{\mathcal{W}_0} b_t b_s d\mu = t \wedge s - ts, \quad t, s \in [0, 1],$$

where  $t \wedge s = \min\{t, s\}$ .

## 2.2. QUADRATIC WIENER FUNCTIONALS

For a separable Hilbert space  $K$ , we say that a  $K$ -valued Wiener functional  $F$  belongs to  $\mathbf{D}^\infty(K)$  if  $F$  is infinitely differentiable in the sense of the Malliavin calculus, and its derivatives of all orders including itself are  $p$ -th integrable with respect to  $\mu$  for any  $p > 1$  (cf. [5, 8]). The  $k$ -th derivative of  $F$  is denoted by  $\nabla^k F$ . Denoting by  $\mathcal{H}_0 \otimes K$  the Hilbert space of Hilbert-Schmidt operators of  $\mathcal{H}_0$  to  $K$  equipped with the Hilbert-Schmidt norm, we define the adjoint operator  $\nabla^* : \mathbf{D}^\infty(\mathcal{H}_0 \otimes K) \rightarrow \mathbf{D}^\infty(K)$  of  $\nabla$  by

$$\int_{\mathcal{W}_0} \langle \nabla^* F, G \rangle d\mu = \int_{\mathcal{W}_0} \langle F, \nabla G \rangle d\mu, \quad F \in \mathbf{D}^\infty(\mathcal{H}_0 \otimes K), G \in \mathbf{D}^\infty(K),$$

where, for the sake of simplicity, we have also used  $\langle \cdot, \cdot \rangle$  to denote the inner products on  $\mathcal{H}_0 \otimes K$  and  $K$ .

Thinking of a symmetric  $U \in \mathcal{H}_0^{\otimes 2} \equiv \mathcal{H}_0 \otimes \mathcal{H}_0$  as a constant function on  $\mathcal{W}_0$  with values in  $\mathcal{H}_0^{\otimes 2}$ , we set

$$Q_U = (\nabla^*)^2 U.$$

If  $U$  is of trace class, we define

$$q_U = Q_U + \text{tr } U.$$

It is easily checked ([5, 8]) that the third derivative of functional  $F \in \mathbf{D}^\infty(\mathbf{R})$  vanishes, i.e.,  $\nabla^3 F = 0$ , if and only if  $F$  admits an expression as

$$F = \frac{1}{2} Q_U + \nabla^* h + c,$$

where

$$U = \nabla^2 F, \quad h = \int_X \nabla F d\nu, \quad \text{and} \quad c = \int_X F d\nu.$$

It is known (cf. [6]) that

**Proposition 1.** *Let  $U \in \mathcal{H}_0^{\otimes 2}$  be symmetric, and  $h \in \mathcal{H}_0$ . For  $\zeta \in \mathbf{C}$  with  $|\zeta| < 1/\|U\|_{\text{op}}$ , where  $\|\cdot\|_{\text{op}}$  denotes the operator norm, it holds that*

$$\int_{\mathcal{W}_0} e^{(\zeta/2)Q_U + \eta \nabla^* h} d\mu = \{\det_2(I - \zeta U)\}^{-1/2} e^{\eta^2 \langle (I - \zeta U)^{-1} h, h \rangle / 2}$$

for every  $\eta \in \mathbf{C}$ , where  $\det_2$  is the Carleman-Fredholm determinant, and  $\langle \cdot, \cdot \rangle$  was extended complex bi-linearly to the complexified Hilbert space  $\mathcal{H}_0 \otimes \mathbf{C}$  of  $\mathcal{H}_0$ . If, in addition,  $U$  is of trace class, then

$$\int_{\mathcal{W}_0} e^{(\zeta/2)q_U + \eta \nabla^* h} d\mu = \{\det(I - \zeta U)\}^{-1/2} e^{\eta^2 \langle (I - \zeta U)^{-1} h, h \rangle / 2},$$

where  $\det$  is the Fredholm determinant.

It is a routine to extend the above identity to  $\zeta$ 's in a much larger domain in  $\mathbf{C}$  by holomorphic continuation.

Such a Wiener integral as in the above proposition is called a stochastic oscillatory integral, and is a probabilistic counterpart to the Feynman path integrals.

### 2.3. SQUARE NORM ON MEASURE SPACE

We review briefly the result in [10].

Let  $E$  be a topological space,  $\mathcal{E}$  its Borel  $\sigma$ -field, and  $\sigma$  a  $\sigma$ -finite measure on  $(E, \mathcal{E})$ . Consider a continuous mapping  $E \ni e \mapsto f_e \in \mathcal{H}_0$ , where the topology of  $\mathcal{H}_0$  is the strong one, i.e., comes from the norm. Assume that

$$(A0) \quad \int_E \|f_e\|^2 \sigma(de) < \infty.$$

Due to Assumption (A0), one can define  $F \in L^2(\mu)$  by the Bochner integral

$$F = \frac{1}{2} \int_E (\nabla^* f_e)^2 \sigma(de),$$

and the non-negative definite, symmetric Hilbert-Schmidt operator  $A : \mathcal{H}_0 \rightarrow \mathcal{H}_0$  of trace class by

$$\langle Ah, g \rangle = \int_E \langle f_e, h \rangle \langle f_e, g \rangle \sigma(de), \quad h, g \in \mathcal{H}_0.$$

Then

$$\text{tr } A = \int_E \|f_e\|^2 \sigma(de) \quad \text{and} \quad F = q_A/2.$$

Denote by  $a_1, a_2, \dots$  the eigenvalues of  $A$ , counted with multiplicity, and  $\{h_n\}_{n=1}^\infty$  the corresponding orthonormal basis (ONB in short) of  $\mathcal{H}_0$ . In conjunction with Proposition 1, we have that

$$(2) \quad \int_{\mathcal{W}_0} e^{\zeta F + \eta \nabla^* h} d\mu = \left\{ \prod_{n=1}^\infty (1 - \zeta a_n) \right\}^{-1/2} \exp\left( \frac{\eta^2}{2} \sum_{n=1}^\infty \frac{\langle h_n, h \rangle_H^2}{1 - \zeta a_n} \right)$$

for  $\zeta, \eta \in \mathbf{C}$  with  $|\zeta| < 1/\max\{a_n : n = 1, 2, \dots\}$ .

## 3. STOCHASTIC OSCILLATORY INTEGRAL ASSOCIATED WITH $\mathfrak{a}^2(\psi)$

In this section, we investigate the stochastic oscillatory integral associated with general  $\mathfrak{a}^2(\psi)$ .

### 3.1. GENERAL $\mathfrak{a}^2(\psi)$

In this subsection, we apply the results in the previous section to general  $\mathfrak{a}^2(\psi)$ .

Let  $\psi : (0, 1) \rightarrow [0, \infty)$  be Borel measurable and assume that

$$(A1) \quad \int_0^1 t(1-t)\psi(t)dt < \infty.$$

For each  $t \in [0, 1]$ , we consider the coordinate mapping  $b_t$  as an element of  $\mathcal{W}'_0 \subset \mathcal{H}_0$ , and, to emphasize this, write  $f_t$  for  $b_t$ ;

$$f_t(w) = w(t), \quad w \in \mathcal{W}_0.$$

For square integrable  $v : [0, 1] \rightarrow \mathbf{R}$ , defining  $h_v \in \mathcal{H}_0$  by

$$h_v(t) = \int_0^t v(s)ds - \left( \int_0^1 v(s)ds \right)t, \quad t \in [0, 1],$$

we have that

$$(3) \quad \langle h_v, g \rangle = \int_0^1 v(s)g'(s)ds, \quad g \in \mathcal{H}_0.$$

Hence  $f_t \in \mathcal{H}_0$  is represented as

$$f_t(s) = t \wedge s - ts, \quad s \in [0, 1].$$

In particular,

$$\|f_t\|^2 = t(1-t).$$

By Assumption (A1),

$$\int_0^1 \|f_t\|^2 \psi(t) dt < \infty,$$

and hence Assumption (A0) in Subsection 2.3 is satisfied with  $E = [0, 1]$  and  $\sigma(dt) = \psi(t)dt$ . Thus, we can now define the quadratic Wiener functional

$$F_\psi = \frac{1}{2} \int_0^1 (\nabla^* f_t)^2 \psi(t) dt.$$

It is easily seen (cf. [5, 8]) that

$$\nabla^* \ell = \ell \quad \text{for every } \ell \in \mathcal{W}'_0.$$

This yields that  $\nabla^* f_t(w) = w(t)$ ,  $w \in \mathcal{W}_0$ , and hence

$$F_\psi = \frac{1}{2} \mathfrak{a}^2(\psi)$$

As was seen in Subsection 2.3,

$$(4) \quad \mathfrak{a}^2(\psi) = 2F_\psi = q_{A_\psi},$$

where the Hilbert-Schmidt operator  $A_\psi : \mathcal{H}_0 \rightarrow \mathcal{H}_0$  is given by

$$\langle A_\psi h, g \rangle = \int_0^1 h(t)g(t)\psi(t)dt, \quad h, g \in \mathcal{H}_0.$$

In what follows, we investigate the eigenvalues of  $A_\psi$ . Since  $A_\psi$  is non-negative definite and symmetric, the eigenvalues of  $A_\psi$  are all non-negative real numbers. Consider the following conditions;

$$(A2) \quad \int_0^1 \{t(1-t)\}^{1/2} \psi(t) dt < \infty.$$

$$(A3) \quad \psi > 0 \text{ a.e. on } (0, 1).$$

$$(A4) \quad \psi \text{ is continuous on } (0, 1).$$

Since  $|h(t)| \leq \sqrt{2t(1-t)} \|h\|$  for  $t \in [0, 1]$ , under (A2),  $h\psi$  is integrable on  $(0, 1)$ .

**Proposition 2.** (i) Assume (A2). Then it holds that

$$(5) \quad (A_\psi h)(t) = - \int_0^t \int_0^s h(u)\psi(u)duds + \left( \int_0^1 \int_0^s h(u)\psi(u)duds \right) t, \quad t \in [0, 1].$$

(ii) Assume (A2) and (A3). Then  $\ker A_\psi = \{0\}$ .

(iii) Assume (A2) and (A4). Let  $h \in \mathcal{H}_0$ . Then  $h$  is an eigenfunction corresponding to an eigenvalue  $\lambda \geq 0$  of  $A_\psi$  if and only if  $h \in C^1([0, 1]) \cap C^2((0, 1))$  and satisfies that

$$(6) \quad \lambda h'' = -\psi h \quad \text{on } (0, 1).$$

Before proceeding to the proof, using the assertion (iii), we give the theorem on the decomposition of  $A_\psi$  and the infinite product expression of the oscillatory integral associated with  $\mathfrak{a}^2(\psi)$ . To do so, we introduce some notation. For  $\varphi \in C([0, 1])$  and  $a \in \mathbf{R}$ , we denote by  $\sigma_{\text{ab}}(\varphi; a)$  the set of all  $g \in C([0, 1]) \cap C^2((0, 1))$  such that  $g \not\equiv 0$ ,  $g(0) = g(1) = 0$  and  $\varphi g'' = ag$  on  $(0, 1)$ , and set

$$\sigma_{\text{ab}}(\varphi) = \{a \in \mathbf{R} \mid \sigma_{\text{ab}}(\varphi; a) \neq \emptyset\}.$$

**Theorem 1.** Assume (A2), (A4), and that

$$(A5) \quad \psi > 0 \text{ on } (0, 1) \text{ and } 1/\psi \in C([0, 1]),$$

where “ $1/\psi \in C([0, 1])$ ” means that  $1/\psi : (0, 1) \rightarrow (0, \infty)$  admits a continuous extension to  $[0, 1]$ .

(i) If  $-\lambda < 0, \in \sigma_{\text{ab}}(1/\psi)$  and there is a  $g \in \sigma_{\text{ab}}(1/\psi; -\lambda)$  such that  $g\psi$  is integrable on  $[0, 1]$ , then  $1/\lambda$  is an eigenvalue of  $A_\psi$ , and  $g$  is an eigenfunction associated with  $1/\lambda$ .

(ii) Suppose that there exist  $-\lambda_n < 0, \in \sigma_{\text{ab}}(1/\psi)$  and  $g_n \in \sigma_{\text{ab}}(1/\psi; -\lambda_n)$ ,  $n = 1, 2, \dots$ , such that each  $g_n\psi$  is integrable on  $[0, 1]$  and  $\{g_n\}_{n=1}^\infty$  is an ONB of  $\mathcal{H}_0$ . Then

$$A_\psi = \sum_{n=1}^\infty \frac{1}{\lambda_n} g_n \otimes g_n,$$

where  $h \otimes h \in \mathcal{H}_0 \otimes \mathcal{H}_0$  is defined by  $(h \otimes h)(g) = \langle h, g \rangle h$ ,  $g \in \mathcal{H}_0$ . Moreover, it holds that

$$(7) \quad \int_{\mathcal{W}_0} e^{\zeta \mathfrak{a}^2(\psi) + \eta \nabla^* h} d\mu = \left\{ \prod_{n=1}^\infty \left( 1 - \frac{2\zeta}{\lambda_n} \right) \right\}^{-1/2} \exp \left( \frac{\eta^2}{2} \sum_{n=1}^\infty \frac{\langle h, g_n \rangle^2}{1 - (2\zeta/\lambda_n)} \right)$$

for  $\zeta, \eta \in \mathbf{C}$  with  $|\zeta| < 1/(2 \max\{1/\lambda_n : n = 1, 2, \dots\})$  and  $h \in \mathcal{H}_0$ .

*Proof.* (i) By the assumption, it holds that

$$g'' = -\frac{1}{\lambda} g\psi \quad \text{on } (0, 1).$$

Since  $g\psi$  is integrable on  $[0, 1]$ , this implies that  $g'$  extends to a continuous function on  $[0, 1]$ . Hence  $g \in C^1([0, 1])$ , and hence  $g \in \mathcal{H}_0$ . By Proposition 2 (iii),  $\lambda$  is an eigenvalue of  $A_\psi$ , and  $g$  is a corresponding eigenfunction.

(ii) In conjunction with (2) and (4), the assertion (i) yields the second assertion.  $\square$



*Proof of Proposition 2.* (i) By an elementary change of variables, we see that

$$\int_0^1 h(t)g(t)\psi(t)dt = \int_0^1 g'(t) \left( - \int_0^t h(u)\psi(u)du \right) dt.$$

By (3), this implies (5).

(ii) Suppose that  $A_\psi h = 0$ . By Assumption (A2) and (5),  $h\psi = 0$  a.e. Then Assumption (A3) and the continuity of  $h$  implies that  $h = 0$ .

(iii) The necessity is an immediate consequence of (i) and the continuity of  $h\psi$  on  $(0, 1)$ .

Conversely, suppose that  $h \in \mathcal{H}_0$  belongs to  $C^1([0, 1]) \cap C^2((0, 1))$  and satisfies (6). Due to the continuity of  $h'$  on  $[0, 1]$  and the integrability of  $h\psi$  on  $[0, 1]$ , integrating (6), we obtain that

$$\lambda h'(s) = \lambda h'(0) - \int_0^s h(u)\psi(u)du, \quad 0 \leq s \leq 1.$$

Since  $h(0) = 0$ , integrating again, we see that

$$\lambda h(t) = \lambda h'(0)t - \int_0^t \int_0^s h(u)\psi(u)duds, \quad 0 \leq t \leq 1.$$

This yields that

$$(8) \quad \lambda h(t) = - \int_0^t \int_0^s h(u)\psi(u)duds + \left( \int_0^1 \int_0^s h(u)\psi(u)duds \right) t, \quad 0 \leq t \leq 1,$$

which, by the assertion (i), means that  $A_\psi h = \lambda h$ .  $\square$

It is an easy exercise of Calculus to see that the eigenvalues of  $A_\psi$  is also computable via the following integral kernel which was used in [1] to obtain the Fourier decomposition of the Gaussian process  $\{b_t \sqrt{\psi(t)}\}_{t \in [0, 1]}$ .

**Proposition 3.** *Assume (A2). Then it holds that*

$$(9) \quad A_\psi h(t) = \int_0^1 (t \wedge s - ts)\psi(s)h(s)ds, \quad t \in [0, 1].$$

*In particular, if  $\lambda > 0$  is an eigenvalue of  $A_\psi$  and  $h \in \mathcal{H}_0$  is a corresponding eigenfunction, then  $g(t) = \sqrt{\psi(t)} h(t)$  satisfies the integral equation with symmetric kernel  $(t \wedge s - ts)\sqrt{\psi(t)}\sqrt{\psi(s)}$ ;*

$$\lambda g(t) = \int_0^1 (t \wedge s - ts)\sqrt{\psi(t)}\sqrt{\psi(s)}g(s)ds, \quad t \in [0, 1].$$

### 3.2. ANDERSON-DARLING TEST STATISTIC

We now restrict our attention to the case where

$$\psi(t) = \psi_{AD}(t) = \frac{1}{t(1-t)},$$

i.e. the case corresponding to the Anderson-Darling test statistic.  $\psi_{\text{AD}}$  satisfies Assumptions (A1)–(A5). Our aim of this subsection is to calculate eigenvalues and eigenfunctions of  $A_{\psi_{\text{AD}}}$ . To state the main result of this subsection, we need some notation. We denote by  $P_n(x)$   $n = 0, 1, \dots$ , the Legendre polynomials defined as the coefficients of the Taylor expansion of  $(1 - 2x\eta + \eta^2)^{-1/2}$  in  $\eta$  ([11]);

$$(1 - 2x\eta + \eta^2)^{-1/2} = \sum_{n=0}^{\infty} P_n(x)\eta^n.$$

It admits the explicit expression as

$$P_n(x) = \sum_{j=0}^{\lfloor n/2 \rfloor} (-1)^j \frac{(2n - 2j)!}{2^n j!(n - j)!(n - 2j)!} x^{n-2j},$$

where  $\lfloor a \rfloor$  stands for the largest integer less than or equal to  $a$ . Our first goal of this subsection is that

**Theorem 2.** *Let*

$$h_n(t) = 4t(1 - t)P'_n(2t - 1), \quad t \in [0, 1]$$

and  $\tilde{h}_n = h_n / \|h_n\|$ ,  $n = 1, 2, \dots$ . *It then holds that*

$$(10) \quad A_{\psi_{\text{AD}}} = \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \tilde{h}_n \otimes \tilde{h}_n,$$

Moreover, it holds that

$$(11) \quad \int_{\mathcal{W}_0} e^{\zeta \mathbf{a}^2(\psi_{\text{AD}}) + \eta \nabla^* h} d\mu = \left\{ \prod_{n=1}^{\infty} \left( 1 - \frac{2\zeta}{n(n+1)} \right) \right\}^{-1/2} \\ \times \exp \left( \frac{\eta}{2} \sum_{n=1}^{\infty} \frac{\langle h, h_n \rangle^2}{1 - (2\zeta / \{n(n+1)\})} \right)$$

for  $\zeta, \eta \in \mathbf{C}$  with  $|\zeta| < 1$  and  $h \in \mathcal{H}_0$ .

*Proof.* The proof is carried out by adapting the observation in [1] to our framework. By Theorem 1, it suffices to study  $\sigma_{\text{ab}}(1/\psi_{\text{AD}}; a)$ . Thus we need to investigate the ordinary differential equation (ODE in short);

$$t(1 - t)h''(t) = \lambda h(t), \quad t \in (0, 1), \quad h(0) = h(1) = 0.$$

Through the change of variables  $t = (x + 1)/2$ , we shall investigate the ODE

$$(12) \quad \begin{cases} (1 - x^2)g''(x) = \lambda g(x), & x \in (-1, 1), \\ g(-1) = g(1) = 0. \end{cases}$$

The Ferrers' associated Legendre function of degree  $n$  and order 1 given by

$$P_{n,1}(x) = (1 - x^2)^{1/2} P'_n(x)$$

is an eigenfunction associated with the eigenvalue  $-n(n+1)$  of the differential operator

$$\mathcal{L}_1 = (1-x^2)\frac{d^2}{dx^2} - 2x\frac{d}{dx} - \frac{1}{1-x^2} \quad \text{on } (-1, 1).$$

For details, see [11]. Moreover, notice that, for any  $\phi \in C^2((-1, 1))$ ,  $g \equiv (1-x^2)^{1/2}\phi$  satisfies that

$$(1-x^2)g'' = (1-x^2)^{1/2}\mathcal{L}_1\phi \quad \text{on } (-1, 1).$$

Thus

$$g_n = (1-x^2)^{1/2}P_{n,1} = (1-x^2)P'_n$$

solves the ODE (12) with  $\lambda = -n(n+1)$ .

Define

$$h_n(t) = 4t(1-t)P'_n(2t-1) = g_n(2t-1), \quad t \in [0, 1].$$

Then

$$h_n \in \sigma_{\text{ab}}(1/\psi_{\text{AD}}; -n(n+1)).$$

Since  $h_n \in C^\infty([0, 1])$  and  $h_n(0) = h_n(1) = 0$ ,  $h_n\psi_{\text{AD}}$  is integrable. By Theorem 1,  $h_n \in \mathcal{H}_0$  and

$$(13) \quad A_{\psi_{\text{AD}}}h_n = \frac{1}{n(n+1)}h_n.$$

We shall now show that  $\{h_n\}_{n=1}^\infty$  is dense in  $\mathcal{H}_0$ . To do so, let

$$k_n(t) = t^{n+1} - t, \quad t \in [0, 1], n = 1, 2, \dots$$

Suppose that  $h \in \mathcal{H}_0$  is perpendicular to all  $k_n$ ,  $n = 1, 2, \dots$ . Then

$$0 = \langle h, k_n \rangle = (n+1) \int_0^1 h'(t)t^n dt, \quad n = 1, 2, \dots$$

This implies that  $h'$  is constant, and hence  $h = 0$ . Thus  $\{k_n\}_{n=1}^\infty$  spans  $\mathcal{H}_0$ . Since  $P'_n$  and  $k_n/\{t(1-t)\}$  are both polynomials of order  $n-1$ ,  $\{h_n\}_{n=1}^\infty$  also spans  $\mathcal{H}_0$ .

Thus the expression (10) holds. The identity (11) is an immediate consequence of (10) and Theorem 1.  $\square$

The infinite product in Theorem 2 can be replaced by an elementary function;

**Corollary 1.** *It holds that*

$$(14) \quad \int_{\mathcal{W}_0} e^{\zeta \mathfrak{a}^2(\psi_{\text{AD}}) + \eta \nabla^* h} d\mu = \left( \frac{-2\pi\zeta}{\cos(\frac{\pi}{2}\sqrt{1+8\zeta})} \right)^{1/2} \\ \times \exp\left( \frac{\eta}{2} \sum_{n=1}^{\infty} \frac{\langle h, h_n \rangle^2}{1 - (2\zeta/\{n(n+1)\})} \right)$$

for  $\zeta, \eta \in \mathbf{C}$  with  $|\zeta| < 1$  and  $\text{Re } \zeta > -1/8$ , and  $h \in \mathcal{H}_0$ .

*Proof.* Recall the well known identity that

$$\cos\left(\frac{\pi}{2}z\right) = \prod_{n=1}^{\infty}\left(1 - \frac{z^2}{(2n-1)^2}\right).$$

Substitute  $z = \sqrt{1+4x}$  to have

$$\cos\left(\frac{\pi}{2}\sqrt{1+4x}\right) = (-4x) \prod_{n=2}^{\infty}\left(1 - \frac{4x+1}{(2n-1)^2}\right).$$

Since

$$1 - \frac{4x+1}{(2n-1)^2} = \left(1 - \frac{x}{n(n-1)}\right)\left(1 - \frac{1}{(2n-1)^2}\right),$$

we obtain that

$$\begin{aligned} \cos\left(\frac{\pi}{2}\sqrt{1+4x}\right) &= (-4x) \prod_{n=1}^{\infty}\left(1 - \frac{x}{n(n+1)}\right) \lim_{x \rightarrow 1} \frac{\cos((\pi/2)x)}{1-x^2} \\ &= -\pi x \prod_{n=1}^{\infty}\left(1 - \frac{x}{n(n+1)}\right). \end{aligned}$$

Thus, by Theorem 2, we obtain the desired identity.  $\square$

## ACKNOWLEDGMENTS

The paper is one of the products achieved in the industry-university joint research “Stochastic models for non-life insurance” by Nisshin Fire & Marine Insurance Co., Ltd, and the Faculty of Mathematics, Kyushu University. The authors thank all those who have engaged in the joint research activity, especially Mr. Sugiyama and Mr. Sugeta. The authors also acknowledge the support in part by the Global COE Program “Education and Research Hub for Mathematics-for-Industry” at Kyushu University.

## REFERENCES

- [1] T. Anderson and D. Darling, Asymptotic theory of certain “goodness of fit” criteria based on stochastic processes, *Annals of Mathematical Statistics* **23** (1952), 193–212.
- [2] T. Anderson and D. Darling, A test of goodness of fit, *Journal of the American Statistical Association* **49** (1954), 765-769.
- [3] P. Billingsley, *Convergence of Probability Measures*, John Wiley & Sons, 1968.
- [4] M. Donsker, Justification and extension of Doob’s heuristic approach to the Kolmogorov-Smirnov theorems, *Annals of Mathematical Statistics* **23** (1952), 277–281.
- [5] N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*, Second Edition, North-Holland/Kodansha, 1989.

- [6] P. Malliavin and S. Taniguchi, Analytic functions, Cauchy formula and stationary phase on a real abstract Wiener space, *Jour. Funct. Anal.*, **143** (1997) 470–528.
- [7] H. Matsumoto and S. Taniguchi, Wiener functionals of second order and their Lévy measures, *Elect. Jour. Probab.*, **7** (2002), Paper No.14, 1–30.
- [8] I. Shigekawa, *Stochastic Analysis*, Trans. Math. Monographs **224**, AMS, 2004.
- [9] M. Stephens, EDF statistics for goodness fit and some comparisons, *Journal of the American Statistical Association* **69** (1974), 730-737.
- [10] S. Taniguchi, Quadratic Wiener functionals of square norms on measure spaces, *Commun. Stoch. Anal.*, **2** (2008), 11-26.
- [11] E.T. Whittaker and G.N. Watson, *A course of modern analysis, an introduction to the general theory of infinite processes and of analytic functions; with an account of the principal transcendental functions*, 4th ed., Cambridge Univ. Press, 1927.

# List of MI Preprint Series, Kyushu University

The Global COE Program  
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA  
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO  
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU  
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI  
Torsion points of abelian varieties with values in infinite extensions over a  $p$ -adic field
- MI2008-6 Yoshiyuki TOMIYAMA  
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI  
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA  
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA  
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA  
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA  
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO  
On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA  
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA  
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA  
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO  
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI  
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI  
Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI  
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI  
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA  
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI  
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI  
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA  
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO  
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA  
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA  
Hypergeometric  $q$ -functions of the  $q$ -Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA  
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA  
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI  
Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain



- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE  
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO  
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI  
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI  
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO  
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA  
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO  
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI  
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO  
On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection

- MI2009-26 Manabu YOSHIDA  
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI  
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA  
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA  
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA  
On asymptotic behaviors of solutions to parabolic systems modelling chemo-taxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI  
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA  
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA  
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA  
Projective reduction of the discrete Painlevé system of type  $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI  
Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA  
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA  
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA  
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA  
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI  
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI  
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI  
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA  
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO  
Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA  
An algebraic approach to underdetermined experiments

- MI2010-10 Kei HIROSE & Sadanori KONISHI  
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA  
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDO  
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA  
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI  
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO  
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE  
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU  
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA  
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE  
Lagrangian approach to weakly nonlinear stability of an elliptical flow

- MI2010-21 Hiroki MASUDA  
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU  
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiko FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU  
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra
- MI2010-25 Toshimitsu TAKAESU  
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA  
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA  
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA  
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE  
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI  
Spin-spin correlation functions of the  $q$ -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and spike detection via Gaussian basis expansions

- MI2010-32 Nobutaka NAKAZONO  
Hypergeometric  $\tau$  functions of the  $q$ -Painlevé systems of type  $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI  
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI  
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI  
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA  
CAP representations of inner forms of  $Sp(4)$  with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER  
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide Fukumoto & Alexander B. Samokhin  
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki Kondo, Shingo Saito & Setsuo Taniguchi  
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro Hirotsu, Hiroki Kondo, Shingo Saito, Takuya Sato, Tatsushi Tanaka & Setsuo Taniguchi  
Anderson-Darling test and the Malliavin calculus