SUCCESSIVE COMPARISONS BETWEEN ORDERED NORMAL MEANS BASED ON CLOSED TESTING PROCEDURE

Imada, Tsunehisa
Liberal Arts Education Center, Kumamoto Campus, Tokai University

<https://doi.org/10.5109/1906488>

出版情報 : Bulletin of informatics and cybernetics. 47, pp.25-36, 2015-12. Research Association
of Statistical Sciences

バージョン :

権利関係 :

SUCCESSIVE COMPARISONS BETWEEN ORDERED NORMAL MEANS BASED ON CLOSED TESTING PROCEDURE

By

Tsune-hisa IMADA*

Abstract

In this study we discuss multiple tests for checking differences among a sequence of normal means with ordered restriction. Specifically, we propose a stepwise multiple comparison procedure based on closed testing procedure using the statistics used by Williams (1971). We give some numerical examples regarding critical values and power of the test intended to compare our procedure and other procedures.

Key Words and Phrases: Multivariate t -distribution, Power of the test, Stepwise test.

1. Introduction

There are independent normal random variables $X_1, X_2, \dots, X_K$. Assume

$$X_k \sim N(\mu_k, \sigma^2)$$

for $k = 1, 2, \dots, K$. Assuming

$$\mu_1 \leq \mu_2 \leq \dots \leq \mu_K, \quad (1)$$

we consider multiple tests to find differences among $\mu_1, \mu_2, \dots, \mu_K$. Lee and Spurrier (1995) set up the null hypothesis and its alternative hypothesis as

$$H_{k,k+1} : \mu_k = \mu_{k+1} \quad \text{vs.} \quad H_{k,k+1}^A : \mu_k < \mu_{k+1}$$

for $k = 1, 2, \dots, K-1$ and proposed a single step multiple comparison procedure for $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$. Shiraishi (2014) developed Lee and Spurrier (1995)'s procedure using closed testing procedure intended to obtain a more powerful procedure. Specifically, Shiraishi (2014) constructed the closed set of hypotheses F based on $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ and proposed a stepwise multiple comparison procedure for F based on closed testing procedure. Furthermore, Douke *et al.* (2006) proposed a sequential rejective step down procedure for $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ and Imada (2015) proposed a step up procedure for them.

On the other hand, Hayter (1990) set up the null hypothesis and its alternative hypothesis as

$$H_{k,l} : \mu_k = \mu_l \quad \text{vs.} \quad H_{k,l}^A : \mu_k < \mu_l$$

* Liberal Arts Education Center, Kumamoto Campus, Tokai University 9-1-1, Toroku Higashi-ku Kumamoto, Japan 862-8652 Japan. tel +81-96-386-2731 timada@ktmail.tokai-u.jp

for $1 \leq k < l \leq K$ and proposed a single step multiple comparison procedure for all $H_{k,l}$ s. Furthermore, Shiraishi (2014) developed Hayter (1990)'s procedure using closed testing procedure.

However, the assumption (1) was not adopted in constructing the procedures described above. The isotonic regression estimators of $\mu_1, \mu_2, \dots, \mu_K$ are the maximum likelihood estimators under (1). Nashimoto and Wright (2005) proposed two types of multiple comparison procedures for all $H_{k,l}$ s ($1 \leq k < l \leq K$) using the isotonic regression estimators. Furthermore, they modified the procedures to avoid the contradiction that $H_{k,l}$ is retained and $H_{k',l'}$ is rejected for $1 \leq k \leq k' < l' \leq l \leq K$.

On the other hand, Williams (1971) proposed a step down multiple comparison procedure for finding minimum k satisfying $\mu_k < \mu_{k+1}$ based on the statistics using the isotonic regression estimators.

In this study we focus on the simultaneous test for $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ and propose another type of stepwise multiple comparison procedure based on closed testing procedure using Williams (1971)'s statistics.

In Section 2, we discuss Lee and Spurrier (1995)'s single step multiple comparison procedure. Specifically, we discuss how to determine the critical value for a specified significance level according to Lee and Spurrier (1995). Furthermore, we formulate the power of the test. In Section 3, we discuss Shiraishi (2014)'s stepwise multiple comparison based on closed testing procedure. In Section 4, we discuss a stepwise multiple comparison based on closed testing procedure using Williams (1971)'s statistics. In Section 5, we give some numerical examples regarding critical values and power of the test intended to compare the procedures. In Section 6, we give some concluding remark.

2. Lee and Spurrier (1995)'s single step procedure

2.1. Determination of the critical value for a specified significance level

Let $x_{k1}, x_{k2}, \dots, x_{kn_k}$ be a sample from $N(\mu_k, \sigma^2)$ for $k = 1, 2, \dots, K$. Lee and Spurrier (1995) used the statistic

$$t_{k,k+1} = \frac{\bar{x}_{k+1} - \bar{x}_k}{\sqrt{\frac{1}{n_{k+1}} + \frac{1}{n_k}}s}$$

for testing $H_{k,k+1}$. Here

$$\bar{x}_k = \frac{1}{n_k} \sum_{i=1}^{n_k} x_{ki}, \quad s = \sqrt{\frac{1}{\phi} \sum_{k=1}^K \sum_{i=1}^{n_k} (x_{ki} - \bar{x}_k)^2}$$

where $\phi = \sum_{k=1}^K n_k - K$. If $t_{k,k+1} > c$ for a specified critical value c , $H_{k,k+1}$ is rejected. Otherwise, it is retained. The critical value c is determined so that

$$P\left(\max_{1 \leq k \leq K-1} t_{k,k+1} > c\right) = \alpha \quad (2)$$

for a specified significance level α under the assumption that $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ are true. The left hand of (2) is the probability that at least one hypothesis among $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ is rejected. (2) is equivalent to

$$P(t_{1,2} \leq c, t_{2,3} \leq c, \dots, t_{K-1,K} \leq c) = 1 - \alpha.$$

We discuss how to calculate $P(t_{1,2} \leq c, t_{2,3} \leq c, \dots, t_{K-1,K} \leq c)$ when $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ are true. $(t_{1,2}, t_{2,3}, \dots, t_{K-1,K})$ is distributed according to the multivariate t -distribution with covariance matrix

$$\mathbf{\Lambda} = \begin{pmatrix} 1 & \rho_{1,2,3} & 0 & 0 & \cdots & 0 & 0 \\ \rho_{1,2,3} & 1 & \rho_{2,3,4} & 0 & \cdots & 0 & 0 \\ 0 & \rho_{2,3,4} & 1 & \rho_{3,4,5} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & \cdots & 1 & \rho_{K-2,K-1,K} \\ 0 & 0 & \cdots & 0 & \cdots & \rho_{K-2,K-1,K} & 1 \end{pmatrix}$$

and ϕ degrees of freedom. Here

$$\rho_{k,k+1,k+2} = -\sqrt{\frac{n_k n_{k+2}}{(n_k + n_{k+1})(n_{k+1} + n_{k+2})}}.$$

Its provability density function is given by

$$f(t_{1,2}, t_{2,3}, \dots, t_{K-1,K}; \mathbf{\Lambda}, \phi) = \frac{\Gamma((K-1+\phi)/2)}{(\phi\pi)^{(K-1)/2} \Gamma(\phi/2) |\mathbf{\Lambda}|^{1/2}} \left(1 + \frac{1}{\phi} \mathbf{t}' \mathbf{\Lambda}^{-1} \mathbf{t}\right)^{-(K-1+\phi)/2}$$

where $\mathbf{t} = (t_{1,2}, t_{2,3}, \dots, t_{K-1,K})'$. Then

$$\begin{aligned} & P(t_{1,2} \leq c, t_{2,3} \leq c, \dots, t_{K-1,K} \leq c) \\ &= \int_{-\infty}^c \int_{-\infty}^c \cdots \int_{-\infty}^c f(t_{1,2}, t_{2,3}, \dots, t_{K-1,K}; \mathbf{\Lambda}, \phi) dt_{1,2} dt_{2,3} \cdots dt_{K-1,K}. \end{aligned}$$

2.2. Power of the test

Next, we consider the power of the test. Assume

$$1 \leq i_1 < i_2 < \cdots < i_k \leq K-1$$

and

$$\mu_{i_j} < \mu_{i_{j+1}} \quad (j = 1, 2, \dots, k), \quad \mu_l = \mu_{l+1} \quad (l \neq i_1, i_2, \dots, i_k). \quad (3)$$

We focus on the all pairs power defined by Ramsey (1978). It is the probability that $H_{i_1, i_1+1}, H_{i_2, i_2+1}, \dots, H_{i_k, i_k+1}$ are rejected. The power is given by

$$P(t_{i_1, i_1+1} > c, t_{i_2, i_2+1} > c, \dots, t_{i_k, i_k+1} > c). \quad (4)$$

To calculate (4) we should specify the difference $\mu_{i_j+1} - \mu_{i_j}$ for $j = 1, 2, \dots, k$ in (3). Specifically, let

$$\mu_{i_j+1} - \mu_{i_j} = \delta_{i_j} (> 0) \quad (j = 1, 2, \dots, k), \quad \mu_l = \mu_{l+1} \quad (l \neq i_1, i_2, \dots, i_k). \quad (5)$$

We discuss how to calculate (4) under (5). Letting

$$t_{0, i_j, i_j+1} = \frac{\bar{x}_{i_j+1} - \bar{x}_{i_j}}{\sqrt{\frac{1}{n_{i_j+1}} + \frac{1}{n_{i_j}}}} \sigma \quad (j = 1, 2, \dots, k), \quad s_0 = \frac{s}{\sigma},$$

we obtain

$$t_{i_j, i_j+1} = \frac{t_{0, i_j, i_j+1}}{s_0} \quad (j = 1, 2, \dots, k)$$

and

$$\begin{aligned} & P(t_{i_1, i_1+1} > c, t_{i_2, i_2+1} > c, \dots, t_{i_k, i_k+1} > c) \\ &= P(t_{0, i_1, i_1+1} > cs_0, t_{0, i_2, i_2+1} > cs_0, \dots, t_{0, i_k, i_k+1} > cs_0). \end{aligned}$$

$(t_{0, i_1, i_1+1}, t_{0, i_2, i_2+1}, \dots, t_{0, i_k, i_k+1})'$ is distributed according to the multivariate normal $N_k(\boldsymbol{\delta}, \boldsymbol{\Lambda}_1)$ where

$$\boldsymbol{\delta} = \left(\frac{\delta_{i_1}}{\sqrt{\frac{1}{n_{i_1+1}} + \frac{1}{n_{i_1}}}\sigma}, \frac{\delta_{i_2}}{\sqrt{\frac{1}{n_{i_2+1}} + \frac{1}{n_{i_2}}}\sigma}, \dots, \frac{\delta_{i_k}}{\sqrt{\frac{1}{n_{i_k+1}} + \frac{1}{n_{i_k}}}\sigma} \right)'$$

and $\boldsymbol{\Lambda}_1$ is the submatrix of $\boldsymbol{\Lambda}$. When the value of s_0 is given,

$$\begin{aligned} & P(t_{0, i_1, i_1+1} > cs_0, t_{0, i_2, i_2+1} > cs_0, \dots, t_{0, i_k, i_k+1} > cs_0 | s_0) \\ &= \int_{cs_0}^{\infty} \int_{cs_0}^{\infty} \cdots \int_{cs_0}^{\infty} f_1(t_{0, i_1, i_1+1}, t_{0, i_2, i_2+1}, \dots, t_{0, i_k, i_k+1}; \boldsymbol{\delta}, \boldsymbol{\Lambda}_1) \\ & \quad \times dt_{0, i_1, i_1+1} dt_{0, i_2, i_2+1} \cdots dt_{0, i_k, i_k+1} \end{aligned}$$

where $f_1(t_{0, i_1, i_1+1}, t_{0, i_2, i_2+1}, \dots, t_{0, i_k, i_k+1}; \boldsymbol{\delta}, \boldsymbol{\Lambda}_1)$ is the probability density function of $N_k(\boldsymbol{\delta}, \boldsymbol{\Lambda}_1)$. On the other hand, since ϕs_0^2 is distributed according to χ^2 -distribution with ϕ degrees of freedom, the probability density function of s_0 is

$$g(s_0) = \frac{\phi^{\phi/2}}{2^{(\phi-2)/2} \Gamma[\phi/2]} s_0^{\phi-1} \exp \left[-\frac{\phi s_0^2}{2} \right].$$

The power is given by

$$\begin{aligned} & P(t_{i_1, i_1+1} > c, t_{i_2, i_2+1} > c, \dots, t_{i_k, i_k+1} > c) \\ &= \int_0^{\infty} g(s_0) \left\{ \int_{cs_0}^{\infty} \int_{cs_0}^{\infty} \cdots \int_{cs_0}^{\infty} f_1(t_{0, i_1, i_1+1}, t_{0, i_2, i_2+1}, \dots, t_{0, i_k, i_k+1}; \boldsymbol{\delta}, \boldsymbol{\Lambda}_1) \right. \\ & \quad \times dt_{0, i_1, i_1+1} dt_{0, i_2, i_2+1} \cdots dt_{0, i_k, i_k+1} \Big\} ds_0. \end{aligned}$$

Since $\boldsymbol{\delta}$ depends on the unknown σ , we should specify the value of σ to calculate the power.

3. Shiraishi (2014)'s stepwise procedure based on closed testing procedure

Let F be the family of hypotheses consisting of $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$ and all sorts of intersections of plural hypotheses chosen from $H_{1,2}, H_{2,3}, \dots, H_{K-1,K}$. F is closed. Specifically, for two hypotheses chosen from F arbitrarily, their intersection is also included in F . Shiraishi (2014) proposed a stepwise multiple comparison procedure for F based on closed testing procedure.

Let I be a subset of $\{1, 2, \dots, K-1\}$ consisting of successive plural integers. $\sharp(I)$ denotes the number of elements of I . Therefore, if $I = \{i, i+1, \dots, j\}$, $\sharp(I) = j - i + 1$. Defining the hypothesis H_I as

$$H_I : \mu_i = \mu_{i+1} = \dots = \mu_j,$$

we obtain

$$H_I = H_{i,i+1} \cap H_{i+1,i+2} \cap \dots \cap H_{j-1,j} \in F.$$

Occasionally, H_I is denoted by $H_{i,j}$. Each hypothesis in F is equal to H_I or $H_{I_1} \cap H_{I_2} \cap \dots \cap H_{I_k}$ where $I_1, I_2, \dots, I_k$ are disjoint. Intended to construct a stepwise procedure for F based on closed testing procedure, we specify a way to test each hypothesis in F . We use the statistic $t_I = \max\{t_{i,i+1}, t_{i+1,i+2}, \dots, t_{j-1,j}\}$ for testing $H_I : \mu_i = \mu_{i+1} = \dots = \mu_j$ and determine the critical value c_I so that $P(t_I > c_I) = \alpha$ under the assumption that H_I is true. If $t_I > c_I$, H_I is rejected. Otherwise, it is retained.

Next, we discuss how to test $H_{I_1} \cap H_{I_2} \cap \dots \cap H_{I_k}$ where $I_1, I_2, \dots, I_k$ are disjoint. Let $M = \sharp(I_1) + \sharp(I_2) + \dots + \sharp(I_k)$. For $l = 1, 2, \dots, k$ we determine $c_{I_l, M}$ so that

$$P(t_{I_l} > c_{I_l, M}) = 1 - (1 - \alpha)^{\sharp(I_l)/M}$$

under the assumption that H_{I_l} is true. Specifically, intended to test $H_{I_1} \cap H_{I_2} \cap \dots \cap H_{I_k}$ we set up the critical value $c_{I_l, M}$ for testing H_{I_l} for $l = 1, 2, \dots, k$. If $t_{I_l} > c_{I_l, M}$ for at least one l , $H_{I_1} \cap H_{I_2} \cap \dots \cap H_{I_k}$ is rejected. Otherwise, it is retained. Shiraishi (2014) indicated the probability that $H_{I_1} \cap H_{I_2} \cap \dots \cap H_{I_k}$ is rejected when it is true is not greater than α . Therefore, we specified the way to test each hypothesis in F satisfying the specified significance level α . We test the hypotheses in F hierarchically. Specifically, if a hypothesis and all hypotheses deriving it are rejected, we reject the hypothesis. Otherwise we retain it. By the principle of closed testing procedure maximum type I FWE (familywise error rate) of this stepwise procedure is not greater than α . It is indicated by Nagata and Yoshida (1997).

Example.

Letting $K = 4$,

$$F = \{H_{1,4}, H_{1,3}, H_{2,4}, H_{1,2} \cap H_{3,4}, H_{1,2}, H_{2,3}, H_{3,4}\}.$$

We test the hypotheses in F as follows.

Step 1.

We test $H_{1,4}$.

Case 1. If $H_{1,4}$ is retained, we retain all other hypotheses in F and stop the test.

Case 2. If $H_{1,4}$ is rejected, go to Step 2.

Step 2.

We test $H_{1,3}, H_{2,4}, H_{1,2} \cap H_{3,4}$.

If $H_{1,3}$ is retained, we retain $H_{1,2}, H_{2,3}$.

If $H_{2,4}$ is retained, we retain $H_{2,3}, H_{3,4}$.

If $H_{1,2} \cap H_{3,4}$ is retained, we retain $H_{1,2}, H_{3,4}$.

Step 3.

We test the hypotheses which are not retained at Step 2 among $H_{1,2}, H_{2,3}, H_{3,4}$.

The power under (5) is the probability that $H_{i_1, i_1+1}, H_{i_2, i_2+1}, \dots, H_{i_k, i_k+1}$ are rejected at the final step in the stepwise test. Since it is difficult to formulate the probability, we calculate it by Monte Carlo simulation.

4. Stepwise procedure based on closed testing procedure using Williams (1971)'s statistic

In Section 3 we discussed the stepwise procedure for F based on closed testing procedure proposed by Shiraishi (2014). In this section we propose a stepwise procedure for F using Williams (1971)'s statistic. First, we consider testing $H_I : \mu_i = \mu_{i+1} = \dots = \mu_j$. Let

$$y_l = \frac{n_l \bar{x}_l + n_{l+1} \bar{x}_{l+1} + \dots + n_j \bar{x}_j}{n_{(l,j)}} \quad (l = i+1, i+2, \dots, j)$$

where $n_{(l,j)} = n_l + n_{l+1} + \dots + n_j$. $M_I = \max\{y_{i+1}, y_{i+2}, \dots, y_j\}$ is the isotonic regression estimator of μ_j under $\mu_i \leq \mu_{i+1} \leq \dots \leq \mu_j$. We use

$$t_I^* = \frac{M_I - \bar{x}_i}{\sqrt{\frac{1}{n_j} + \frac{1}{n_i} s}}$$

for testing H_I . If $t_I^* > c_I^*$, H_I is rejected. Otherwise, it is retained. We determine c_I^* so that

$$P(t_I^* > c_I^*) = \alpha \quad (6)$$

under the assumption that H_I is true. (6) is equivalent to

$$P(t_I^* \leq c_I^*) = 1 - \alpha.$$

To determine c_I^* we consider how to calculate $P(t_I^* \leq c_I^*)$. Letting

$$z_l = \frac{y_l - \bar{x}_i}{\sqrt{\frac{1}{n_j} + \frac{1}{n_i} s}}$$

for $l = i+1, i+2, \dots, j$, we obtain

$$P(t_I^* \leq c_I^*) = P(z_{i+1} \leq c_I^*, z_{i+2} \leq c_I^*, \dots, z_j \leq c_I^*).$$

$(z_{i+1}, z_{i+2}, \dots, z_j)'$ is distributed according to the multivariate t -distribution with covariance matrix

$$\mathbf{\Lambda}_I = \frac{1}{\frac{1}{n_j} + \frac{1}{n_i}} \begin{pmatrix} \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \dots & \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} \\ \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+2,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+2,j)}} & \dots & \frac{1}{n_i} + \frac{1}{n_{(i+2,j)}} \\ \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+2,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+3,j)}} & \dots & \frac{1}{n_i} + \frac{1}{n_{(i+3,j)}} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+2,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+3,j)}} & \dots & \frac{1}{n_{(j-1,j)}} + \frac{1}{n_i} \\ \frac{1}{n_i} + \frac{1}{n_{(i+1,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+2,j)}} & \frac{1}{n_i} + \frac{1}{n_{(i+3,j)}} & \dots & \frac{1}{n_i} + \frac{1}{n_j} \end{pmatrix}$$

and ϕ degrees of freedom. Letting $f(z_{i+1}, z_{i+2}, \dots, z_j; \mathbf{\Lambda}_I, \phi)$ be its provability density function, we obtain

$$\begin{aligned} P(t_I^* \leq c_I^*) &= P(z_{i+1} \leq c_I^*, z_{i+2} \leq c_I^*, \dots, z_j \leq c_I^*) \\ &= \int_{-\infty}^{c_I^*} \int_{-\infty}^{c_I^*} \dots \int_{-\infty}^{c_I^*} f(z_{i+1}, z_{i+2}, \dots, z_j; \mathbf{\Lambda}_I, \phi) dz_{i+1} dz_{i+2} \dots dz_j. \end{aligned}$$

Next, we discuss how to test $H_{I_1} \cap H_{I_2} \cap \cdots \cap H_{I_k}$ where $I_1, I_2, \dots, I_k$ are disjoint. Let $M = \sharp(I_1) + \sharp(I_2) + \cdots + \sharp(I_k)$. For $l = 1, 2, \dots, k$ we determine $c_{I_l, M}^*$ so that

$$P(t_{I_l}^* > c_{I_l, M}^*) = 1 - (1 - \alpha)^{\sharp(I_l)/M}$$

under the assumption that H_{I_l} is true. If $t_{I_l}^* > c_{I_l, M}^*$ for at least one l , $H_{I_1} \cap H_{I_2} \cap \cdots \cap H_{I_k}$ is rejected. Otherwise, it is retained. Therefore, the probability that $H_{I_1} \cap H_{I_2} \cap \cdots \cap H_{I_k}$ is rejected is

$$1 - P(t_{I_1}^* \leq c_{I_1, M}^*, t_{I_2}^* \leq c_{I_2, M}^*, \dots, t_{I_k}^* \leq c_{I_k, M}^*).$$

When $H_{I_1}, H_{I_2}, \dots, H_{I_k}$ are true,

$$1 - P(t_{I_1}^* \leq c_{I_1, M}^*, t_{I_2}^* \leq c_{I_2, M}^*, \dots, t_{I_k}^* \leq c_{I_k, M}^*) \leq \alpha,$$

because

$$P(t_{I_1}^* \leq c_{I_1, M}^*, t_{I_2}^* \leq c_{I_2, M}^*, \dots, t_{I_k}^* \leq c_{I_k, M}^*) \geq \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l, M}^*) \quad (7)$$

and

$$\prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l, M}^*) = 1 - \alpha.$$

The proof of (7) is given in Appendix. Since we specified the way to test each hypothesis in F satisfying the specified significance level α , the stepwise procedure based on closed testing procedure is constructed.

It is also difficult to formulate the power of the test under (5). The power is calculated by Monte Carlo simulation.

5. Simulation results

We discussed three types of multiple comparison procedures intended to find differences among a sequence of normal means with ordered restriction. First, we discussed Lee and Spurrier (1995)'s single step procedure. Next, we discussed Shiraishi (2014)'s stepwise procedure developing Lee and Spurrier (1995)'s procedure based on closed testing procedure. Furthermore, we proposed the stepwise procedure based on closed testing procedure using Williams (1971)'s statistics. In this section we give some simulation results intended to compare five procedures including Douke *et al.* (2006)'s sequential rejective step down procedure and Imada (2015)'s step up procedure. SS, CT, CT-W, SD and SU denote these procedures, respectively.

Let $K = 3, 4, 5$ and $\alpha = 0.05$. We set up a balanced sample size n for each population. Specifically, let $n = 15, 30$. Table 1 gives critical values of SS.

Table 1 : Critical values of SS

K	3	4	5
$n = 15$	2.018	2.179	2.286
$n = 30$	1.988	2.152	2.261

Tables 2,3,4 give critical values of CT for $K = 3, 4, 5$. When we test $H_{1,3} \cap H_{4,5}, H_{1,2} \cap H_{3,5}$ by Table 4, we use the critical values 2.211, 2.188 for testing $H_{1,3}$ and $H_{3,5}$ and use 2.087, 2.066 for testing $H_{4,5}$ and $H_{1,2}$ for $n = 15, 30$, respectively. Tables 5,6,7 give critical values of CT-W for $K = 3, 4, 5$. When we test $H_{1,3} \cap H_{4,5}, H_{1,2} \cap H_{3,5}$ in Table 7, we use the critical values 1.970, 1.953 for testing $H_{1,3}$ and $H_{3,5}$ and use 2.087, 2.066 for testing $H_{4,5}$ and $H_{1,2}$ for $n = 15, 30$, respectively.

Table 2 : Critical values of CT ($K = 3$)

	$n = 15$	$n = 30$
$H_{1,3}$	2.018	1.988
$H_{1,2}, H_{2,3}$	1.682	1.663

Table 3 : Critical values of CT ($K = 4$)

	$n = 15$	$n = 30$
$H_{1,4}$	2.179	2.152
$H_{1,3}, H_{2,4}$	2.004	1.981
$H_{1,2} \cap H_{3,4}$	1.998	1.976
$H_{1,2}, H_{2,3}, H_{3,4}$	1.673	1.659

Table 4 : Critical values of CT ($K = 5$)

	$n = 15$	$n = 30$
$H_{1,5}$	2.286	2.261
$H_{1,4}, H_{2,5}$	2.168	2.146
$H_{1,3} \cap H_{4,5}, H_{1,2} \cap H_{3,5}$	2.211, 2.087	2.188, 2.066
$H_{1,3}, H_{2,4}, H_{3,5}$	1.995	1.977
$H_{1,2} \cap H_{3,4}, H_{1,2} \cap H_{4,5}, H_{2,3} \cap H_{4,5}$	1.989	1.971
$H_{1,2}, H_{2,3}, H_{3,4}, H_{4,5}$	1.667	1.656

Table 5 : Critical values of CT-W ($K = 3$)

	$n = 15$	$n = 30$
$H_{1,3}$	1.758	1.737
$H_{1,2}, H_{2,3}$	1.682	1.663

Table 6 : Critical values of CT-W ($K = 4$)

	$n = 15$	$n = 30$
$H_{1,4}$	1.772	1.755
$H_{1,3}, H_{2,4}$	1.748	1.732
$H_{1,2} \cap H_{3,4}$	1.998	1.976
$H_{1,2}, H_{2,3}, H_{3,4}$	1.673	1.659

Table 7 : Critical values of CT-W ($K = 5$)

	$n = 15$	$n = 30$
$H_{1,5}$	1.777	1.763
$H_{1,4}, H_{2,5}$	1.766	1.752
$H_{1,3} \cap H_{4,5}, H_{1,2} \cap H_{3,5}$	1.970, 2.087	1.953, 2.066
$H_{1,3}, H_{2,4}, H_{3,5}$	1.742	1.729
$H_{1,2} \cap H_{3,4}, H_{1,2} \cap H_{4,5}, H_{2,3} \cap H_{4,5}$	1.989	1.971
$H_{1,2}, H_{2,3}, H_{3,4}, H_{4,5}$	1.667	1.656

Next, we give numerical examples of power of the test. Let $\sigma = 1$. We set up two types of arrangements of means $(\mu_1, \mu_2, \mu_3) = (0, 1, 2), (0, 1, 1)$ for $K = 3$. We set up three types of arrangements of means $(\mu_1, \mu_2, \mu_3, \mu_4) = (0, 1, 2, 3), (0, 1, 2, 2), (0, 1, 1, 1)$ for $K = 4$. We set up four types of arrangements of means $(\mu_1, \mu_2, \mu_3, \mu_4, \mu_5) = (0, 1, 2, 3, 4), (0, 1, 2, 3, 3), (0, 1, 2, 2, 2), (0, 1, 1, 1, 1)$ for $K = 5$. Specifically, for each K we set up $K-1$ types of arrangements of means varying the number of differences among successive normal means from 1 to $K-1$. Although the formulations of the power of SS, SD and SU are obtained, we should obtain the power of CT and CT-W by Monte Carlo simulation. Here we obtain the power of CT and CT-W with 1,000,000 times of experiments. Tables 8 to 13 give the power of the test for five procedures. CT-W is most powerful among five procedures except for the case when the number of differences among successive normal means is equal to $K-1$. CT, SD and SU are more powerful compared to SS except for the case when the number of differences among successive normal means is equal to 1. In each case the differences of power among CT, SD and SU are not remarkably large.

Table 8 : Power of the test ($K = 3, n = 15$)

(μ_1, μ_2, μ_3)	SS	CT	CT-W	SD	SU
(0, 1, 2)	0.546	0.704	0.710	0.706	0.711
(0, 1, 1)	0.763	0.766	0.816	0.766	0.770

Table 9 : Power of the test ($K = 3, n = 30$)

(μ_1, μ_2, μ_3)	SS	CT	CT-W	SD	SU
(0, 1, 2)	0.938	0.972	0.972	0.972	0.972
(0, 1, 1)	0.969	0.971	0.983	0.971	0.973

Table 10 : Power of the test ($K = 4, n = 15$)

$(\mu_1, \mu_2, \mu_3, \mu_4)$	SS	CT	CT-W	SD	SU
(0, 1, 2, 3)	0.290	0.579	0.588	0.578	0.596
(0, 1, 2, 2)	0.458	0.555	0.634	0.553	0.561
(0, 1, 1, 1)	0.712	0.713	0.743	0.714	0.714

Table 11 : Power of the test ($K = 4, n = 30$)

$(\mu_1, \mu_2, \mu_3, \mu_4)$	SS	CT	CT-W	SD	SU
(0, 1, 2, 3)	0.871	0.959	0.959	0.959	0.959
(0, 1, 2, 2)	0.913	0.942	0.957	0.942	0.944
(0, 1, 1, 1)	0.956	0.958	0.968	0.957	0.958

Table 12 : Power of the test ($K = 5, n = 15$)

$(\mu_1, \mu_2, \mu_3, \mu_4, \mu_5)$	SS	CT	CT-W	SD	SU
(0, 1, 2, 3, 4)	0.133	0.468	0.481	0.467	0.501
(0, 1, 2, 3, 3)	0.229	0.396	0.433	0.393	0.408
(0, 1, 2, 2, 2)	0.399	0.485	0.509	0.463	0.466
(0, 1, 1, 1, 1)	0.675	0.677	0.702	0.676	0.675

Table 13 : Power of the test ($K = 5, n = 30$)

$(\mu_1, \mu_2, \mu_3, \mu_4, \mu_5)$	SS	CT	CT-W	SD	SU
(0, 1, 2, 3, 4)	0.791	0.945	0.946	0.945	0.946
(0, 1, 2, 3, 3)	0.839	0.915	0.927	0.915	0.916
(0, 1, 2, 2, 2)	0.891	0.923	0.933	0.915	0.916
(0, 1, 1, 1, 1)	0.945	0.947	0.960	0.946	0.947

6. Conclusions

In this study, we discussed the multiple comparison procedures for finding differences among a sequence of normal means with ordered restriction. Specifically, we proposed the stepwise procedure based on closed testing procedure using Williams (1971)'s statistics and compared our procedure with Lee and Spurrier (1995)'s single step procedure, Shiraishi (2014)'s stepwise procedure based on closed testing procedure, Douke *et al.* (2006)'s sequential rejective step down procedure and Imada (2015)'s step up procedure through the simulation regarding the power of the test. We confirmed that our procedure is most powerful in almost all cases. Specifically, by adopting the assumption that a sequence of normal means is ordered we can construct a powerful procedure.

We gave numerical examples when the number of the populations is not greater than 5. Although the determination of the power of closed testing procedure is accompanied with computational complications when the number of populations is large, we should give numerical examples regarding critical value and power of the test when the number of normal means is large in the future.

Appendix

In this Appendix, we derive

$$P(t_{I_1}^* \leq c_{I_1, M}^*, t_{I_2}^* \leq c_{I_2, M}^*, \dots, t_{I_k}^* \leq c_{I_k, M}^*) \geq \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l, M}^*)$$

given in (7).

Let U be a nonnegative random variable and $f(u)$ be its probability density function. Let $h_1(U)$, $h_2(U)$, $\dots$, $h_l(U)$ be functions which are nonnegative, bounded, continuous and monotone in the same direction. Furthermore, we assume there exists finite expectation $E[h_i(U)]$ for each i . By Corollary A.1.1 in Hsu (1996)

$$E\left[\prod_{i=1}^l h_i(U)\right] \geq \prod_{i=1}^l E[h_i(U)]. \quad (8)$$

(8) is rewritten as

$$\int_0^\infty \prod_{i=1}^l h_i(u) f(u) du \geq \prod_{i=1}^l \int_0^\infty h_i(u) f(u) du. \quad (9)$$

We derive the inequality (7) using (9).

First, let $I = \{i, i+1, \dots, j\}$. We consider the conditional probability $P(t_I^* \leq c_I^* | s_0)$ given s_0 . Letting

$$z_{0,l} = \frac{y_l - \bar{x}_i}{\sqrt{\frac{1}{n_j} + \frac{1}{n_i}} \sigma} \quad (l = i+1, i+2, \dots, j), \quad s_0 = \frac{s}{\sigma},$$

we obtain

$$z_l = \frac{y_l - \bar{x}_i}{\sqrt{\frac{1}{n_j} + \frac{1}{n_i}} s} = \frac{z_{0,l}}{s_0}$$

and

$$P(t_I^* \leq c_I^* | s_0) = P(z_{0,i+1} \leq c_I^* s_0, z_{0,i+2} \leq c_I^* s_0, \dots, z_{0,j} \leq c_I^* s_0 | s_0). \quad (10)$$

Here $(z_{0,i+1}, z_{0,i+2}, \dots, z_{0,j})'$ is distributed according to the multivariate normal $N_{j-i}(\mathbf{0}, \mathbf{\Lambda}_I)$. Letting $f_0(z_{0,i+1}, z_{0,i+2}, \dots, z_{0,j}; \mathbf{\Lambda}_I)$ be its probability density function, we obtain

$$P(t_I^* \leq c_I^* | s_0) = \int_{-\infty}^{c_I^* s_0} \int_{-\infty}^{c_I^* s_0} \cdots \int_{-\infty}^{c_I^* s_0} f_0(z_{0,i+1}, z_{0,i+2}, \dots, z_{0,j}; \mathbf{\Lambda}_I) dz_{0,i+1} dz_{0,i+2} \cdots dz_{0,j} \quad (11)$$

by (10). (11) shows that $P(t_I^* \leq c_I^* | s_0)$ is monotonically increasing for $0 \leq s_0 < \infty$. By (9)

$$\int_0^\infty \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l,M}^* | s_0) g(s_0) ds_0 \geq \prod_{l=1}^k \int_0^\infty P(t_{I_l}^* \leq c_{I_l,M}^* | s_0) g(s_0) ds_0.$$

Since

$$\int_0^\infty P(t_{I_l}^* \leq c_{I_l,M}^* | s_0) g(s_0) ds_0 = P(t_{I_l}^* \leq c_{I_l,M}^*)$$

for $l = 1, 2, \dots, k$, we obtain

$$\int_0^\infty \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l,M}^* | s_0) g(s_0) ds_0 \geq \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l,M}^*). \quad (12)$$

On the other hand, since $t_{I_1}^*, t_{I_2}^*, \dots, t_{I_k}^*$ are independent when the value of s_0 is given, we obtain

$$P(t_{I_1}^* \leq c_{I_1,M}^*, t_{I_2}^* \leq c_{I_2,M}^*, \dots, t_{I_k}^* \leq c_{I_k,M}^*) = \int_0^\infty \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l,M}^* | s_0) g(s_0) ds_0. \quad (13)$$

By (12) and (13) we obtain

$$P(t_{I_1}^* \leq c_{I_1,M}^*, t_{I_2}^* \leq c_{I_2,M}^*, \dots, t_{I_k}^* \leq c_{I_k,M}^*) \geq \prod_{l=1}^k P(t_{I_l}^* \leq c_{I_l,M}^*).$$

Acknowledgement

The author is deeply grateful to the referee and the editors for their valuable comments and suggestions.

References

- Bartholomew, D. J. (1959). A test of homogeneity for ordered alternatives, *Biometrika* **46**, 36–48.
- Douke, H., Imada, T. and Nakamura, T. (2006). Development of sequential step-wise procedures for dose response test. *Journal of the Japan Statistical Society* **36** (1), 45–64. (in Japanese)

- Hayter, A. J. (1990). A one-sided studentized range test for testing against a simple ordered alternative, *Journal of the American Statistical Association* **85**, No. 411, 778-785.
- Hsu, J. C. (1996). *Multiple Comparisons*. Boca Raton : Chapman & Hall.
- Imada, T. (2015). Multiple comparison procedures for checking differences among sequence of normal means with ordered restriction. *Journal of the Japan Statistical Society* **44** (2), 251-270. (in Japanese)
- Lee, R. E. and Spurrier, J. D. (1995). Successive comparisons between ordered treatments. *Journal of Statistical Planning and Inference*, **43**, 323-330.
- Nagata, Y. and Yoshida, M. (1997). *Basic Theory of Statistical Multiple Comparison Procedures*. Tokyo : Scientist Inc. (in Japanese)
- Nashimoto, K. and Wright, F.T. (2005). Multiple comparison procedures for detecting differences in simply ordered means, *Computational Statistics and Data Analysis*, **48**, 291-306.
- Ramsey, P. H. (1978). Power differences between pairwise multiple comparisons. *Journal of the American Statistical Association*, **73**, 479-485.
- Shiraishi, T. (2014). Closed testing procedures in multi-sample models under a Simple Ordered Restriction. *Journal of the Japan Statistical Society* **43** (2), 215-245. (in Japanese)
- Williams, D. A. (1971). A test for differences between treatment means when several dose levels are compared with a zero dose control, *Biometrics* **27**, 103-117.

Received March 20, 2015

Revised September 4, 2015