Occurrences of radiolarian biostratigraphic markers *Lychnocanoma nipponica sakaii* and *Amphimelissa setosa* in Core YK07-12 PC3B from the Okhotsk Sea

Ikenoue, Takahito

Department of Earth & Planetary Sciences, Graduate School of Sciences, Kyushu University

Takahashi, Kozo

Department of Earth & Planetary Sciences, Graduate School of Sciences, Kyushu University

Sakamoto, Tatsuhiko

Earth and Life History Research Program, Institute of Biogeosciences, Japan Agency for Marine-Earth Science and Technology

Sakai, Saburo

Earth and Life History Research Program, Institute of Biogeosciences, Japan Agency for Marine-Earth Science and Technology

他

<https://doi.org/10.5109/19193>

出版情報：九州大学大学院理学研究院紀要：Series D, Earth and planetary sciences. 32 (3), pp.1-10, 2011-03-10. Faculty of Science, Kyushu University

バージョン：

権利関係：

Occurrences of radiolarian biostratigraphic markers *Lychnocanoma nipponica sakaii* and *Amphimelissa setosa* in Core YK07-12 PC3B from the Okhotsk Sea

Takahito Ikenoue*, Kozo Takahashi*, Tatsuhiko Sakamoto**,
Saburo Sakai** and Koichi Iijima**

Abstract

Core YK07-12 PC03B obtained from the central Okhotsk plain (52°36'N, 150°08'E ; water depth: 1049 m) was analyzed every 5 cm, focusing on two radiolarian biostratigraphic markers in the late Pleistocene: *Lychnocanoma nipponica sakaii* and *Amphimelissa setosa*. The last occurrence (LO) datum of *A. setosa* at 67 ka conformed with those of the LO of the previously published studies. However, the LO of *L. nipponica sakaii* at 25 ka was off set and significantly younger than those published in the previous studies in the North Pacific and the marginal seas. We introduce herein the conceptual use of the last common occurrence (LCO) of *L. nipponica sakaii* as the LCO conforms with the LO of this taxon published elsewhere and hence more reliable, especially out side of the northern part of the central Okhotsk plain. The LCO of *L. nipponica sakaii* is 46 ka, 21 kyrs older than the LO, but is approximately the same as the LOs published elsewhere. We postulate the LCO as a practical datum which can be compared with the data from out side of the studied region. As alternative explanations, the obtained significantly younger LO than the LOs of other studies from the out side of the study region may stem from the following two reasons. Because that the present study is specifically focused on the two biostratigraphic marker radiolarian taxa with details our microscopic counts are significantly greater than those performed in the previous studies. This may have caused the documentation of the LO with the rare microscopic counts on multiple microslides, which could have been missed had we counted only one microslide. It is also possible that the effect of bioturbation due to upward transport in the slow sedimentation rate regime caused the appearance of the rare specimens of *L. nipponica sakaii* well above the LCO.

Keywords: Radiolarian biostratigraphy, last occurrence (LO), last common occurrence (LCO), *Lychnocanoma nipponica sakaii*, *Amphimelissa setosa*, Okhotsk Sea, late Pleistocene

1. Introduction

The Okhotsk Sea represents as one of the lowest latitude regions with extensive sea-ice cover in the world. Environmental conditions of the Okhotsk Sea with such sea-ice formation can alter global water circulations because that the formation results in the generation of dense Okhotsk intermediate water which exits into the North Pacific (e.g., Talley, 1991). The Okhotsk Sea is also characterized by a high biological productivity and thus it serves as an efficient biological pump to absorb atmospheric CO₂. Therefore, for further understanding the global carbon cycle it is important to study the region including the past environmental conditions (Takahashi, 1998).

Radiolarians represent as one of the common microzooplankton groups in the pelagic and hemipelagic realms. Their siliceous skeletons are often well preserved in the sediments and hence they can be used in biostratigraphy

Manuscript received on 1 November 2010; accepted on 7 January 2011

* Department of Earth & Planetary Sciences, Graduate School of Sciences, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, JAPAN; Corresponding author's e-mail: ikenoue@geo.kyushu-u.ac.jp

** Earth and Life History Research Program, Institute of Biogeosciences, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Natsushima-cho 2-15, Yokosuka, 237-0061, JAPAN

of deep-sea sediments and reconstruction of paleoceanographic conditions. Pertinent previous works on radiolarian biostratigraphy in the Okhotsk Sea are, for example, published by Takahashi et al. (2000), Matul et al. (2002, 2009), and Okazaki et al. (2003, 2005). The purpose of this study is to examine radiolarian biostratigraphic markers of the late Pleistocene in detail employing Core YK07-12 PC03B obtained in the northern part of the central Okhotsk plain.

2. Materials and Methods

During Cruise YK07-12 of R/V Yokosuka belonging to Japan Agency for Marine-Earth Science and Technology (JAMSTEC) in summer 2007, Core YK07-12 PC03B was obtained from the northern part of the central Okhotsk plain (52°36'N, 150°08'E ; water depth: 1,049 m; Fig. 1). The recovered core length is 14.09 m. Sediment samples were continuously sliced every 1 cm in thickness throughout the core. In this study, samples were analyzed at every 5 cm interval.

Fig. 1. Location map of Core YK07-12 PC03B (solid circle) together with other relevant cores (open circles) in the Okhotsk Sea.

An age model of Core YK07-12 PC3B was established based on $\delta^{18}\text{O}$ stratigraphy with subordinate magnetic susceptibility (MS). Graphic correlations of $\delta^{18}\text{O}$ records were established between the $\delta^{18}\text{O}$ variations of benthic foraminifer shells (*Uvigerina* spp.) in Core PC3B and the standard curve of LR04 Benthic $\delta^{18}\text{O}$ global stack constructed by Lisiecki and Raymo (2005) (Fig. 2). As results of the analyses, Marine Isotope Stages (MIS) 1-12.3 were identified, and it was determined that Core PC3B recorded at least 450 kyrs of the environmental conditions of the northern part of the central Okhotsk plain.

Fig. 2. An age model of Core YK07-12 PC3B based on $\delta^{18}\text{O}$ stratigraphy and subordinated by MS. Changes in (a) MS of Core YK07-12-PC3B; (b) benthic foraminiferal (*Uvigerina* spp.) $\delta^{18}\text{O}$ of Core YK07-12-PC3B; and (c) stacked benthic foraminiferal oxygen isotope curve LR04 (Lisiecki and Raymo, 2005), Lines MR2 and MR3 are tephra horizons recovered in Core YK07-12 PC3B.

Radiolarian counts were conducted on 53 selected samples whose ages ranged from 133 ka at 264.5 cm depth to the present at the core top. Wet samples of 100-1,100 mg were weighed after freeze-drying for 24 hrs. They were treated with 5 ml of 1N hydrochloric acid, 10 ml of 30% hydrogen peroxide, and heated on a hot plate at 90°C for 1 hr in order to remove organic matter and calcium carbonate. Then, Calgon® (hexametaphosphate, surfactant) solution was added to the mixture to further disaggregate the sediments. Our samples contained a large amount of lithogenic grains and thus we employed the elutriation method for extracting opal particles from the residues according to Itaki (2003). After the elutriation, samples were sieved through a stainless screen with 45 µm mesh and filtered through Gellman® membrane filters with a nominal pore size of 0.45 µm. The filtered samples were washed with distilled water to remove salts, dried in an oven at 50°C, and then permanently mounted with Canada Balsam® on microslides. We counted the number of all radiolarian specimens on the slides under a compound light-microscope at x100-200 magnifications. Normally, counts were made on an entire area of

Table 1. Radiolarian counts in Core YK07-12 PC3B.

Section	Total depth at cube center (cm)	Age (ka)	Dry bulk density (g cm ⁻³)	SR (cm ky ⁻¹)	Dry sediment (g)	Number of slides counted	Total Radiolaria				<i>Lychnocanoma nipponika sakaii</i>				<i>Amphimelissa setosa</i>			
							Count	AR (No. rads cm ⁻² ky ⁻¹)	Count	AR (No. rads cm ⁻² ky ⁻¹)	%	Specimens (g dry sed. ⁻¹)	Count	AR (No. rads cm ⁻² ky ⁻¹)	%	Specimens (g dry sed. ⁻¹)		
1	9.3	0.00	0.324	2.49	0.042	1	434	8381	0	0	0	0	0	0	0	0	0	
1	16.0	2.79	0.459	2.49	0.040	1	630	17896	0	0	0	0	0	0	0	0	0	
1	20.5	5.59	0.564	2.49	0.043	1	563	18559	0	0	0	0	0	0	0	0	0	
1	25.0	8.38	0.617	2.49	0.050	1	323	9901	0	0	0	0	0	0	0	0	0	
1	29.5	11.18	0.526	2.49	0.051	1	130	3374	0	0	0	0	0	0	0	0	0	
1	34.0	13.97	0.742	2.49	0.057	1	168	5418	0	0	0	0	0	0	0	0	0	
1	40.9	16.76	0.976	2.49	0.096	1	34	860	0	0	0	0	0	0	0	0	0	
1	45.4	19.69	1.010	1.43	0.118	1	79	970	0	0	0	0	0	0	0	0	0	
1	50.0	23.16	1.005	1.43	0.413	4	282	981	0	0	0	0	0	0	0	0	0	
1	54.5	26.63	0.968	1.43	0.420	4	426	1404	1	3	0	2	0	0	0	0	0	
1	59.1	30.10	1.057	1.43	0.982	8	721	1110	4	6	0.55	4	0	0	0	0	0	
1	66.0	33.57	1.045	1.43	0.408	3	561	2055	12	44	2.14	29	0	0	0	0	0	
1	70.6	37.03	1.002	1.43	0.419	2	1382	4727	2	7	0	5	0	0	0	0	0	
1	75.2	40.50	0.957	1.43	0.427	2	1600	5134	2	6	0	5	0	0	0	0	0	
1	79.3	43.97	1.058	1.43	0.095	1	239	3819	3	48	1.26	32	0	0	0	0	0	
2	79.3	44.66	1.002	1.43	0.096	1	192	2861	0	0	0	0	0	0	0	0	0	
2	84.8	48.13	1.020	1.43	0.441	2	972	3218	89	295	9.16	202	0	0	0	0	0	
2	89.4	51.60	0.938	1.43	0.491	5	326	891	222	606	68.10	452	0	0	0	0	0	
2	94.0	55.07	0.967	1.43	0.527	5	252	661	122	320	48.41	231	0	0	0	0	0	
2	100.9	58.54	0.966	1.43	0.452	8	309	945	195	596	63.11	432	0	0	0	0	0	
2	105.5	62.01	0.914	1.43	0.407	3	590	1894	335	1075	56.78	823	0	0	0	0	0	
2	110.1	65.47	0.946	1.43	0.071	1	122	2331	35	669	28.69	494	0	0	0	0	0	
2	114.7	68.94	1.151	1.43	0.139	1	189	2235	32	378	16.93	230	1	12	0.53	7	7	
2	119.2	72.41	0.730	1.43	0.066	1	232	3670	38	601	16.38	576	0	0	0	0	0	
2	126.1	75.88	0.700	1.43	0.058	1	305	5304	56	974	18.36	972	2	35	0.66	35	35	
2	130.6	79.20	0.706	1.43	0.057	1	279	4979	60	1071	21.51	1060	2	36	0.72	35	35	
2	135.2	81.15	0.493	2.5	0.027	1	223	10294	41	1893	18.39	1536	4	185	1.79	150	150	
2	139.8	83.10	0.566	2.5	0.049	1	357	10352	63	1827	17.65	1291	3	87	0.84	61	61	
2	144.4	85.06	0.619	2.5	0.046	1	359	12211	44	1497	12.26	967	1	34	0.28	22	22	
2	149.0	87.01	0.584	2.5	0.043	1	194	6522	25	840	12.89	576	4	134	2.06	92	92	
2	155.9	88.97	0.652	2.5	0.060	1	495	13505	48	1310	9.70	804	2	55	0.40	34	34	
2	160.5	90.92	0.502	2.5	0.062	1	631	12741	56	1131	8.87	900	19	384	3.01	305	305	
2	165.1	92.88	0.578	2.5	0.056	1	476	12328	71	1839	14.92	1272	7	181	1.47	125	125	
2	169.7	94.83	0.622	2.5	0.056	1	322	8926	71	1968	22.05	1266	7	194	2.17	125	125	
2	174.3	96.79	0.665	2.5	0.064	1	309	8085	33	863	10.68	520	9	235	2.91	142	142	
3	178.9	98.94	0.724	2.5	0.051	1	366	12984	38	1348	10.38	745	9	319	2.46	176	176	
3	184.6	100.89	0.659	2.5	0.067	1	331	8206	6	149	1.81	90	11	273	3.32	165	165	
3	189.0	102.85	0.595	2.5	0.052	1	790	22408	24	681	3.04	458	42	1191	5.32	802	802	
3	195.8	104.80	0.740	2.5	0.058	1	804	25809	22	706	2.74	382	50	1605	6.22	868	868	
3	200.2	106.75	0.626	2.5	0.063	1	2166	53544	23	569	1.06	363	149	3683	6.88	2354	2354	
3	204.8	108.71	0.588	2.5	0.061	1	2522	60421	18	431	0.71	293	199	4768	7.89	3241	3241	
3	209.4	110.66	0.575	2.5	0.064	1	4001	89649	26	583	0.65	405	642	14385	16.05	10000	10000	
3	214.1	112.62	0.518	2.5	0.037	1	3002	104436	36	1252	1.20	968	578	20108	19.25	15538	15538	
3	221.1	114.57	0.510	2.5	0.046	1	2810	78002	30	833	1.07	654	506	14046	18.01	11024	11024	
3	225.7	116.53	0.585	2.5	0.049	1	2925	87846	26	781	0.89	534	470	14115	16.07	9651	9651	
3	230.3	118.48	0.561	2.5	0.033	1	1939	81672	27	1137	1.39	811	203	8551	10.47	6096	6096	
3	234.9	120.44	0.601	2.5	0.043	1	1651	58251	22	776	1.33	516	86	3034	5.21	2019	2019	
3	239.5	122.41	0.626	2.5	0.056	1	1367	37905	14	388	1.02	248	111	3078	8.12	1968	1968	
3	244.1	124.44	0.638	1.85	0.059	1	557	11132	5	100	0.90	85	48	959	8.62	812	812	
3	250.9	126.47	0.687	1.85	0.059	1	762	16461	9	194	1.18	153	45	972	5.91	765	765	
3	255.5	128.50	0.917	1.85	0.081	1	421	8875	8	169	1.90	99	46	970	10.93	571	571	
3	260.0	130.53	0.973	1.85	0.101	1	278	4934	7	124	2.52	69	21	373	7.55	207	207	
3	264.5	132.56	1.021	1.85	0.076	1	344	8582	3	75	0.87	40	22	549	6.40	291	291	

one microslide with a goal of counting >300 specimens. The counts of <200 specimens were resulted in eight cases out of the total of 55 cases whereas the counts exceeding 500 specimens resulted in 22 cases (11 cases >1,000 specimens) (Table 1). In two extremely cases of rare radiolarian occurrences (Samples 59.1 cm and 100.9 cm) eight microslides were prepared and all the slides were counted in order to fulfill the initial goal of counting sufficient number of specimens. In order to achieve the goal as close as possible, multiple microslides were counted in eleven cases in total (Table 1). The classification of *Lychnocanoma nipponica* (Nakaseko) *sakaii* (Morley and Nigrini) and *Amphimelissa setosa* (Cleve) in the samples follow that of Morley and Nigrini (1995) for the former and Bjørklund et al. for the latter (1998), respectively.

A radiolarian accumulation rate (RAR) was calculated using the following equation: $\text{RAR (No. radiolarians cm}^{-2} \text{ kyr}^{-1}) = [\text{No. radiolarians g}^{-1} \text{ dry sediment}] (\text{No. g}^{-1}) \times [\text{dry bulk density}] (\text{g cm}^{-3}) \times [\text{sedimentation rate}] (\text{cm kyr}^{-1})$. Dry bulk density was determined from weights of each of 10 cc sediment samples taken with a polypropylene cube. It was calculated from the wet bulk density multiplied by the water contents (%). The water contents of each sample were determined from weights of the wet and dry samples. The samples were weighed before and after drying in an oven at 50°C for 24 hrs. Sedimentation rates (SR) were assumed between the age control points.

3. Results and Discussion

The temporal changes in total RAR in Core YK07-12 PC3B are shown in Fig. 3. Its pattern tends to follow the temporal patterns displayed by the global $\delta^{18}\text{O}$ changes during the glacial-interglacial cycles (e.g., Lisiecki and Raymo, 2005); total RAR increased during the interglacial periods and decreased during the glacial periods at least for the time span examined. The values during MIS 1 and MIS 5 were relatively high compared to those of the other MIS stages.

Fig. 3. Changes in total RAR in Core YK07-12 PC3B during the last 130 kyrs. Gray areas indicate cold MIS.

In this study we focused on two important radiolarian taxa as biostratigraphic markers in the late Pleistocene: *Lychnocanoma nipponica* (Nakaseko) *sakaii* (Morley and Nigrini) and *Amphimelissa setosa* (Cleve) (Fig. 4). They are quite useful markers in the North Pacific and marginal seas (Bjørklund and Swanberg, 1987; Morley et al., 1995; Matul et al., 2002). In Core YK07-12 PC3B, the LO of *L. nipponica sakaii* occurred in the upper MIS 3 at 25 ka, and the *L. nipponica sakaii* acme event occurred at 81 ka (Table 1, Fig. 5). Our data displayed several differences compared to the results shown by the previous studies. The LO of *L. nipponica sakaii* in the previous studies are listed in Table 2. Specifically, the LO of *L. nipponica sakaii* was 49 ka (Morley et al., 1982) and 50 ka (Morley et al., 1995) in the North Pacific. Tanaka and Takahashi (2005) and Itaki et al. (2009) dated the LO of *L. nipponica sakaii* at 46-52 ka using sediment samples from the Bering Sea. Itaki et al. (2007) also estimated the LO to be 54 ka using sediment samples from the Japan Sea. The LO in four piston cores XP98-PC1, XP98-PC2, XP98-PC4, MD01-2412 from the Okhotsk Sea were determined also as ca. 50 ka (Takahashi et al., 2000; Okazaki et al., 2005). The locations of these four core sites are further south of the core site of the present study.

Fig. 4. Radiolarian taxa from Core YK07-12 PC03B discussed in the text. All scale bars equal to 50 μ m. a. *Lychnocanoma nipponica* (Nakaseko) *sakaii* (Morley and Nigrini), Sample 230.3 cm. b. *Lychnocanoma nipponica* (Nakaseko) *sakaii* (Morley and Nigrini), Sample 230.3 cm. c. *Amphimelissa setosa* (Cleve), sample 230.3 cm. d. *Amphimelissa setosa* (Cleve), Sample 230.3 cm.

On the contrary to the majority of the LOs of ca. 50 ka mentioned above, Matul et al. (2002) and Matul et al. (2009) proposed the LO to have occurred at 28 ka and 34 ka in the central Okhotsk Sea, which is fundamentally the same region as that in our study. As mentioned earlier, the results of our study also show that the LO to be at 25 ka level, approximately the same age as those in Matul et al. (2002, 2009). These LOs from the three piston cores (LV28-42-4, MD01-2412, and YK07-12 PC03B) from the same general region collectively displayed significantly younger ages than those from other previous studies. While there is a discrepancy in the timing of the LOs between the older (ca. 50 ka) and younger (34-25 ka) groups, the overall patterns of the population change with time appear to be conformable between the two groups. Namely, the AR, abundance and relative abundance of *L. nipponica sakaii* increased steadily from ca. 120 ka towards 50 ka and rapidly declined after 50

Fig. 5. Changes in AR, relative abundance and abundance of *L. nipponica sakaii* in Core YK07-12 PC03B. The LO, LCO and Acme datums are shown with arrows.

Fig. 6. Changes in AR, relative abundance and abundance of *A. setosa* in Core YK07-12 PC03B. The LO and LCO are shown with arrows.

ka. Thus, the trend of the changes in AR, abundance and relative abundance in our study are roughly conformable with those in the previous studies.

Furthermore, the LO of *A. setosa* is placed at 67 ka within MIS 4, slightly above the MIS 4/5 boundary (Fig. 6). Thus, the LO of *L. nipponica sakaii* and *A. setosa* obtained from Core YK07-12 PC3B correspond to the LOs of Core LV28-42-4 (Matul et al., 2002) fairly well. It is quite possible that the LO of *L. nipponica sakaii* in the northern part of the central Okhotsk plain is diachronous with respect to the LO of this taxon further south in the Okhotsk plain and in other basins, considering the consistency of the LOs from the same region. It is plausible that the environmental conditions between ca. 50 ka and ca. 34–25 ka in the northern part of the central Okhotsk plain were still favorable for the survival of *L. nipponica sakaii* whereas those of the further south were severe enough to cause an extinction of this taxon at ca. 50 ka. This could well be due to the distribution of the sea-ice cover during the interval in question in the region. It is noteworthy to mention here that the sea-ice cover was more extensive at Site XP98-PC1 than Site XP98-PC2 (Fig. 1), according to Okazaki et al. (2005). Shiga and Koizumi (2000) also reported more open water conditions in the eastern Okhotsk Sea than in the western area during the time period around 22 ka and the last glacial maximum. Based on these pieces of information we interpret that the survival of *L. nipponica sakaii* was prolonged at our study site than the western area and perhaps further south in the Okhotsk Sea. Thus, one of the possible three reasons (see below for the remainders) for the significantly young 25 ka as the LO is the diachronous extinction of this taxon within the entire Okhotsk Sea.

As a part of the second explanation of the discrepancy, we herein propose the introduction of “the last common occurrence (LCO)” concept as a practical biostratigraphic datum. This datum should be far more reliable than the LO for the following details which we have encountered in the present piston core. The tail end of the occurrence of *L. nipponica sakaii* prior to ca. 25 ka (Fig. 5) may well be due to the detailed radiolarian counts performed in our study, which is specifically devoted to illustrate the exact occurrences. Specifically, the total radiolarian counts of 426 (4 microslides) at 27 ka and 721 (8 microslides) at 30 ka were performed (Table 1). These values are apparently far more greater than those proceeded in other published data whereas their exact counts were not published nor available in their results. Our microscopic counts for *L. nipponica sakaii* were one specimen at 27 ka and four specimens at 30 ka, respectively (Table 1). These relatively small count values could well have been missed and recorded as zeros, had the significantly smaller counts (1 microslide or portion of it) than the present study were made. In that case our LO would have been 46 ka in stead. Thus, it makes a sense to use the 46 ka level as the LCO, which is graphically illustrated as the rapid decrease interval in AR, abundance and %*L. nipponica sakaii*.

Table 2.

(a) List of the LO / LCO of *L. nipponica sakaii* and SR for cores used in the recent studies.

Core/ Site	Area	Latitude	Longitude	Age (ka)	SR (cm kyr ⁻¹)	Reference
YK07-12 PC03B	Okhotsk Sea	52°36.01' N	150°08.25' E	25 / 46	2.86	This paper
LV28-42-4	Okhotsk Sea	51°42.89' N	150°59.13' E	28	2-4	Matul et al. (2002)
MD01-2415	Okhotsk Sea	53°57.09' N	149°57.52' E	34	2-4	Matul et al. (2009)
MD01-2412	Okhotsk Sea	44°31' N	145°00' E	47-48	46	Okazaki et al. (2005)
XP98-PC1	Okhotsk Sea	51°00.9' N	152°00.5' E	43-44	8.1	Takahashi et al. (2000)
XP98-PC2	Okhotsk Sea	50°23.7' N	148°19.4' E	53-55	10.5	Takahashi et al. (2000)
XP98-PC4	Okhotsk Sea	49°29.3' N	146°07.7' E	55-56	12.6	Takahashi et al. (2000)
PC-23A	Bering Sea			48.6		Itaki et al. (2009)
ES	Bering Sea			45.9		Tanaka and Takahashi (2005)
BOW-8A	Bering Sea			49.5		Tanaka and Takahashi (2005)
BOW-9A	Bering Sea			49.7		Tanaka and Takahashi (2005)
BOW-12A	Bering Sea			51.7		Tanaka and Takahashi (2005)
MD01-2407 & others	Japan Sea			54		Itaki et al. (2007)
V20-120, RC14-103 & others	North Pacific			49		Morley et al. (1982)
ODP Sites 881, 883 & others	North Pacific			50		Morley et al. (1995)

(b) List of the LO / LCO of *A. setosa* and SR for cores used in the recent studies.

Core/ Site	Area	Latitude	Longitude	Age (ka)	SR (cm kyr ⁻¹)	Reference
YK07-12 PC03B	Okhotsk Sea	52°36.01' N	150°08.25' E	67 / 74	2.86	This paper
LV28-42-4	Okhotsk Sea	51°42.89' N	150°59.13' E	72	2-4	Matul et al. (2002)
MD01-2415	Okhotsk Sea	53°57.09' N	149°57.52' E	64	2-4	Matul et al. (2009)
ES	Bering Sea			85.1		Tanaka and Takahashi (2005)
BOW-8A	Bering Sea			111.5		Tanaka and Takahashi (2005)
BOW-9A	Bering Sea			68		Tanaka and Takahashi (2005)
GAT-3A	Bering Sea			82.6		Tanaka and Takahashi (2005)
	North Pacific			70		Björklund and Swanberg (1987)

As a second alternative, the slow SR observed in the central Okhotsk Sea may have caused the discrepancy in the LO levels. Itaki et al. (2009) speculated that the delay of the LO of *L. nipponica sakaii* by Matul et al. (2002) might be due to the effect of bioturbation in the slow SR environment. When sedimentation rates are slow fossil radiolarian specimens initially located at several cm deep (or even deeper sometimes) in sediments, for example, could have been subsequently transported upward to the sediment surface or nearby by organisms living near the surface or within sediments, causing a bioturbation. Such an activity can alter the sequentially deposited initial signature in a drastic manner. On the other hand, the chances of the effect in high SR regime can minimize such an action because faster burial than the artifact of the bioturbation. The sedimentation rates of the previously studied cores from the Okhotsk Sea are shown in Table 2. The sediment cores with higher SRs than the three piston cores (LV28-42-4, MD01-2412, and PC03B) may have led the LO of *L. nipponica sakaii* to be ca. 50 ka. Thus, the time lag of the LO of *L. nipponica sakaii* with respect to the LCO in YK07-12 PC03B could possibly be due to the effect of bioturbation in the slow SR regime. Note that the first disappearances of *L. nipponica sakaii* and *A. setosa* after the major occurrence maxima are 46 ka and 74 ka, respectively. These ages are nearly equal to the LO of other previously published studies. One drawback of such an explanation with the slow SR is that the LO of *A. setosa* at 68 ka did not cause any significant time lag compared to the data from the fast SR regimes. Because that the occurrence of total radiolarians is about the same at both at 25 ka and 67 ka (Table 1, Fig. 3) it is difficult to explain why we did not encounter a time lag problem in the LO of *A. setosa* in the present piston core.

4. Conclusions

A focused detailed study on two important radiolarian biostratigraphic markers of the late Pleistocene was performed in the northern part of the central Okhotsk plain: *Lychnocanoma nipponica* (Nakaseko) *sakaii* (Morley and Nigrini) and *Amphimelissa setosa* (Cleve). We have employed Core YK07-12 PC3B from the northern part of the central Okhotsk plain with the aid of benthic foraminiferal $\delta^{18}\text{O}$ stratigraphy. The introduction of the concept of “the last common occurrence (LCO)” makes a sense where the LO and the LCO differ significantly in age, especially in the case of the present study. The LCO of the 46 ka in the present study serves as a comparable datum with respect to the LO at ca. 50 ka of *L. nipponica sakaii* found further south in the Okhotsk Sea and elsewhere in the marginal seas of the North Pacific and subarctic Pacific. It is possible that the extinction datum of *L. nipponica sakaii* in the northern part of the central Okhotsk plain (25 ka) specifically lagged behind of that of elsewhere by ca. 25 kyrs. This may well have caused the diachronous LO compared to those of elsewhere. We interpret that the sea-ice conditions were favorable to allow the survival of *L. nipponica sakaii* for additional ca. 16-25 kyrs in the study region compared to elsewhere. As alternative explanations for the discrepancy in the LOs, we also offer (1) the detailed counts of the radiolarians in this devoted study; and (2) the slow SR where bioturbation may have caused the apparent time lag of the LO at 25 ka compared to the LO of ca. 50 ka in the studies performed other than in the northern part of the central Okhotsk plain. Furthermore, the LO of *Amphimelissa setosa* at 67 ka obtained from the present study was conformable with that of Core LV28-42-4.

5. Acknowledgments

We thank scientists, Captain and crew participated in Cruise YK07-12 of R/V Yokosuka who assisted us on board for this work. We thank Dr. Yusuke Okazaki of JAMSTEC for his constructive review of an early draft of this paper as well as further discussion. The senior author received and benefitted by the Professor Tatsuro Matsumoto Scholarship Fund.

6. References

- Björklund, K.R., and Swanberg, N.R. (1987) The distribution of two morphotypes of the radiolarian *Amphimelissa setosa* Cleve (Nassellarida): a result of environmental variability. *Sarsia*, **72**, 245-254.
- Björklund, K.R., Cortese, G., Swanberg, N., and Schrader, H.J. (1998) Radiolarian faunal provinces in surface sediments of the Greenland, Iceland and Norwegian (GIN) Seas. *Marine Micropaleontology*, **35**, 105-140.
- Itaki, T. (2003) Elutriation technique for the extraction of radiolarian skeletons from sandy sediment. *Fossils*, **73**,

38-41.

- Itaki, T., Komatsu, N., and Motoyama, I. (2007) Orbital- and millennial-scale changes of radiolarian assemblages during the last 220 kyrs in the Japan Sea. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **247**, 115-130.
- Itaki, T., Uchida, M., Kim, S., Shin, H-S., and Tada, R. (2009) Late Pleistocene stratigraphy and palaeoceanographic implications in northern Bering Sea slope sediments: evidence from the radiolarian species *Cycladophora davisiana*. *Journal of Quaternary Science*, **24**, 856-865.
- Lisiecki, L.E., and Raymo, M.E., 2005. A Plio-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography*, **20**, 1003.
- Morley, J.J., and Nigrini, J.D. (1995) Miocene to Pleistocene radiolarian biostratigraphy of North Pacific sites 881, 884, 885, 886, and 887. In: Rea, D.K., Basov, I.A., Scholl, D.W., Allan, J.F. (Eds.), *Proc. ODP, Sci. Results*, **145**, College Station, TX (Ocean Drilling Program), 55-91.
- Morley, J.J., Hays, J.D., and Robertson, J.H. (1982) Stratigraphic framework for the late Pleistocene in the northwest Pacific Ocean. *Deep-Sea Research*, **29**, 1485-1499.
- Morley, J.J., Tiase, V.L., Ashby, M.M., and Kashgarian, M. (1995) A high-resolution stratigraphy for Pleistocene sediments from North Pacific Sites 881, 883, and 887 based on abundance variations of the radiolarian *Cycladophora davisiana*. In: Rea, D.K., Basov, I.A., Scholl D.W., Allen, J.F. (Eds.), *Proc. ODP, Sci. Results*, **145**, College Station, TX (Ocean Drilling Program), 133-140.
- Matul, A., Abelmann, A., Tiedemann, R., Kaiser, A., and Nürnberg, D. (2002) Late Quaternary polycystine radiolarian datum events in the Sea of Okhotsk. *Geo-Marine Letters*, **22**(1), 25-32.
- Matul, A., Abelmann, A., Nürnberg, D., and Tiedemann, R. (2009) Stratigraphy and major paleoenvironmental changes in the Sea of Okhotsk during the last million years inferred from radiolarian data. *Oceanology*, **49**(1), 93-100.
- Okazaki, Y., Takahashi, K., Yoshitani, H., Nakatsuka, T., Ikehara, M., and Wakatsuchi, M. (2003) Radiolarians under the seasonally sea-ice covered conditions in the Okhotsk Sea: flux and their implications for paleoceanography. *Marine Micropaleontology*, **49**(3), 195-230.
- Okazaki, Y., Takahashi, K., Katsuki, K., Ono, A., Hori, J., Sakamoto, T., Uchida, M., Shibata, Y., Ikehara, M., and Aoki, K. (2005) Late Quaternary paleoceanographic changes in the southwestern Okhotsk Sea: evidence from geochemical, radiolarian, and diatom records. *Deep-Sea Research II*, **52**, 2332-2350.
- Shiga, K., and Koizumi, I., (2000) Latest Quaternary oceanographic changes in the Okhotsk Sea based on diatom records. *Mar. Micropaleontol.* **38**, 91-117.
- Takahashi, K. (1998) The Bering and Okhotsk Seas: modern and past paleoceanographic changes and gateway impact. *Journal Asian Earth Science*, **16**, 49-58.
- Takahashi, K., Okazaki, Y., and Yoshitani, H. (2000) Radiolarian fossils and paleoceanography: accumulation changes in the Okhotsk Sea. *Chikyu Monthly*. **22**, 623-630. [in Japanese]
- Talley, L. D. (1991) An Okhotsk Sea water anomaly: implications for ventilation in the North Pacific. *Deep-Sea Research*, **38**, S171-S190.
- Tanaka, S. and Takahashi, K. (2005) Late Quaternary paleoceanographic changes in the Bering Sea and the western subarctic Pacific based on radiolarian assemblage. *Deep-Sea Research II*, **52**, 2131-2149.