

Regulation of expression and trafficking of perforin-2 by LPS and TNF- α

熊, 鵬

<https://hdl.handle.net/2324/1928626>

出版情報：九州大学, 2017, 博士（医学）, 課程博士
バージョン：
権利関係：やむを得ない事由により本文ファイル非公開（2）



氏 名：熊 鵬

論 文 名：Regulation of expression and trafficking of perforin-2 by LPS and TNF- α

(パーフォリン 2 の LPS と TNF- α による発現と輸送の調節)

区 分：甲

論 文 内 容 の 要 旨

Perforin-2 is constitutively expressed in macrophages that are required for bacterial control. In this study, we found that perforin-2 is expressed in human macrophages with two isoforms: full-length perforin-2a and a splice variant, perforin-2b. Two isoforms show different subcellular distributions. Perforin-2a was predominantly localized to the membrane of endosome-like vesicles by a C-terminal transmembrane domain. In contrast, the short isoform perforin-2b lacking the transmembrane domain failed to localize to the membrane of vesicles. Furthermore, we determined that the pro-inflammatory stimuli LPS and TNF- α induced perforin-2a expression via the NF- κ B pathway and triggered perforin-2a vesicles fusion with lysosomes. On the other hand, we detected the secretion of perforin-2b in response to LPS stimulation. Taken together, our data provide the evidence that membrane-bound and secretory isoforms of perforin-2 are present in human macrophages and may play important roles in immune defense.