

First Report of Soft Rot Disease of Papaya Caused by *Klebsiella variicola* in Bangladesh

HOSSAIN, Shahadat

Department of Plant Pathology, Bangabandhu Sheikh Mujibur Rahman Agricultural University |
Laboratory of Plant Pathology, Division of Agrobiological Science, Department of Bioresource
Sciences, Faculty of Agriculture, Kyushu University

KHAN, Abu Ashraf

Department of Plant Pathology, Bangabandhu Sheikh Mujibur Rahman Agricultural University |
Laboratory of Plant Pathology, Division of Agrobiological Science, Department of Bioresource
Sciences, Faculty of Agriculture, Kyushu University

RAHMAN, Md. Mahbubur

Department of Biotechnology, Bangabandhu Sheikh Mujibur Rahman Agricultural University |
Laboratory of Plant Pathology, Division of Agrobiological Science, Department of Bioresource
Sciences, Faculty of Agriculture, Kyushu University

IYAMA, Kazuhiro

Laboratory of Plant Pathology, Division of Agrobiological Science, Department of Bioresource
Sciences, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/1955381>

出版情報：九州大学大学院農学研究院紀要. 63 (2), pp.201-205, 2018-09-01. Faculty of
Agriculture, Kyushu University

バージョン：

権利関係：

First Report of Soft Rot Disease of Papaya Caused by *Klebsiella variicola* in Bangladesh

Shahadat HOSSAIN¹, Abu Ashraf KHAN^{1*}, Md. Mahbubur RAHMAN²,
Kazuhiro IYAMA³ and Naruto FURUYA³

Laboratory of Plant Pathology, Division of Agrobiological Science, Department of Bioresource Sciences,
Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan
(Received April 5, 2018 and accepted May 8, 2018)

Twelve bacteria were isolated from soft rotten papaya collected from markets of different locations of Gazipur district, Bangladesh. Among these, five isolates (CP01–CP05) caused soft rot symptoms on potato slices. The isolates also produced dark brown to blackish characteristics soft rot symptoms on papaya fruit by artificial inoculation. The isolates were Gram-negative and also negative in oxidase, methyl red, arginine dihydrolase, gelatin liquefaction and indole tests. All of the isolates were positive in catalase, oxidative fermentative, nitrate reduction, acetoin and urease production tests. The isolates grew well at 41°C and 5% of salt concentration. They utilized lactose, glucose, rhamnose, sucrose, melibiose, arabinose, mannitol, inositol, sorbitol and citrate but not adonitol as sole sources of carbon. Phylogenetic analysis based on 16 S rRNA gene sequence indicated that an isolate CP03 was closely related with *K. variicola*. These results suggested that the isolates from the diseased papaya were *K. variicola*. This study reports for the first time *K. variicola* causing soft rot of papaya in Bangladesh.

Key words: Soft rot, papaya, *Klebsiella variicola*, Bangladesh

INTRODUCTION

Papaya (*Carica papaya*) is very nutritious and commonly used as fruits and vegetables. It is infected by various pathogens including fungi, bacteria and virus. Among these infectious diseases, soft rot caused by bacterial pathogen is the one of the most common and destructive disease causing severe economic loss.

Several bacterial species are known to macerate tissues of a wide range fruits and vegetables enzymatically. Species of *Erwinia* belonging to the “*carotovora*” group including *Erwinia carotovora* (synonym: *Pectobacterium carotovorum*), *E. chrysanthemi* (synonym: *Dickeya dadantii*), are usually referred as the soft rot bacteria. Moreover, pseudomonads such as *Pseudomonas viridiflava* and *P. marginalis* were reported to cause soft rot of different fruits and vegetables (Perombelon and Kelman, 1980; Kim *et al.*, 2002). Although *Klebsiella* species are isolated from clinical samples, the bacteria are associated with several plants such as banana, rice, sugarcane and maize (Rosenblueth *et al.*, 2004). *K. variicola* was reported to cause soft rot of banana in China (Fan *et al.*, 2016) and *K. pneumoniae* induces soft rot of onion (Liu *et al.*, 2015).

In Bangladesh, *P. carotovorum*, *D. dadantii* and *Pseudomonas* species were reported in various fruits and vegetables as soft rot bacteria (Meah and Khan, 1987; Hossain, 2016).

In this study, several bacteria were isolated from soft rotted papaya in Bangladesh. Although *P. carotovorum* was reported to cause soft rot in papaya (Himel *et al.*, 2016), the isolates were different from *P. carotovorum* in the preliminary tests. There is no report showing other bacterial pathogens cause soft rot in papaya in the country. Therefore, identification of the bacterial pathogen was carried out in this study.

MATERIALS AND METHODS

Isolation of bacterial isolates from soft rotted papaya

Bacteria were isolated from the diseased papaya fruits collected from markets of different locations of Gazipur district, Bangladesh by the previously described method (Mortensen, 1997). In brief, small part from the margin of rotted tissues was cut and surface sterilized with 1% sodium hypochlorite (NaOCl) for 2–3 min. Sterilized samples were washed several times with sterilized distilled water to remove the residual sodium hypochlorite. The samples were placed in Petri dishes containing sterilized distilled water and were crushed with a sterile scalpel. After crushing, the Petri dishes were kept undisturbed for 10–15 min to release the bacteria associated with rotted tissues. One loopful of the resulting suspension was streaked on YPDA (yeast extract peptone dextrose agar medium: yeast extract 3 g, peptone 0.6 g, dextrose 3 g, agar 15 g, distilled water 1 liter, pH 7.2) plate. The plates were incubated at 30°C for 48 h. Preferential individual bacterial colonies that appeared on plate were picked and re-streaked on another fresh plate to obtain pure culture. The isolates were preserved in 30% glycerol and stored at –20°C for further study.

¹ Department of Plant Pathology, Bangabandhu Sheikh Mujibur Rahman Agricultural University

² Department of Biotechnology, Bangabandhu Sheikh Mujibur Rahman Agricultural University

³ Laboratory of Plant Pathology, Division of Agrobiological Science, Department of Bioresource Sciences, Faculty of Agriculture, Kyushu University

* Corresponding author (E-mail: ashraf@bsmrau.edu.bd)

Potato tissue maceration test

All of the bacterial isolates originated from single colonies were tested for their ability to macerate tissue using potato (De Boer and Kelman, 2001), since ability of tissue maceration correlates with the pectolytic characteristic and pathogenicity. Potato tubers were sterilized with 70% ethyl alcohol, rinsed in sterilized distilled water and aseptically cut into slices. The potato slices were put in Petri dishes containing sterilized filter paper impregnated with 2 ml of sterilized distilled water. The potato slices were inoculated with the isolates by the needle pricking method. The inoculated slices were maintained in moistened Petri dishes and incubated at 30°C for 2 days. The bacterial isolates caused soft rot on potato slices were selected for further studies.

Pathogenicity test of bacterial isolates on papaya

Healthy papaya fruit was used for pathogenicity test. Papaya fruit was sterilized with 70% ethyl alcohol, rinsed in sterilized distilled water and aseptically punctured with multiple needles. Fresh culture of bacteria was inoculated at the injured site. Then, the inoculated papaya fruit was covered with a polyethylene bag to maintain moisture and incubated at 30°C for 3 days.

Physiological and biochemical tests to characterize pathogenic bacterial isolates

A series of physiological and biochemical tests were performed to characterize the bacterial isolates. The tests were Gram reaction (Suslow *et al.*, 1982), catalase production (Hayward, 1992), oxidative fermentative test (Hugh and Leifson, 1953), oxidase test (Kovacs, 1956), gelatin liquefaction test (Schaad, 1988), urease production (Schaad, 1988), nitrate reduction test (Lelliot and Dickey, 1984), indole test (Lelliot and Dickey, 1984), acetoin production (Dye, 1968), methyl red test (Schaad, 1988), arginine dihydrolase (Thornley, 1960), growth at 41°C temperature (Schaad, 1988), growth in 5% NaCl (Schaad, 1988), and utilization of carbon sources (Ayers *et al.*, 1919).

16S rRNA gene sequencing

A representative bacterial isolate CP03 was selected for 16S rRNA gene sequencing. The bacterial isolate was cultured in nutrient broth at 28°C in a shaker incubator. Approximately 24 h-old culture was subjected for genomic DNA isolation. Genomic DNA was extracted by using Gene JET Genomic DNA Purification Kit (Thermo Scientific Ltd.) following the manufacturer's protocol. Amplification of the targeted 16S rRNA gene was performed with the universal primer sets 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGATACCTTGTACGACTT-3'). Individual PCR mixture contained 6 µl of 25 mM MgCl₂, 10 µl of 10 × PCR buffer, 2.0 µl of 10 mM dNTP mix, 5.0 µl of 20 µM each primer, 5.0 µl (150 ng) of DNA template, 1 µl (5 U/µl) of *Taq* DNA polymerase and 66 µl of sterile double-distilled water thus the total volume of 100 µl. A negative control (without template DNA) was also included in the PCR run. The PCR amplification was performed in a PCR

Thermocycler (Eppendorf Ltd.). The condition for PCR was set as follows: an initial denaturation step at 94°C for 5 min; 35 cycles of a denaturation step at 94°C for 1 min, an annealing at 55°C for 40 sec and an extension at 72°C for 1 min and a final extension step at 72°C for 5 min. The PCR product was purified by using a commercial Gene JET PCR Purification Kit (Thermo Scientific Ltd.) following the manufacturer's protocol. The purified PCR product was sequenced from the Centre for Advanced Research in Sciences, University of Dhaka, Bangladesh.

Phylogenetic analysis

Nucleotide sequences were aligned by the Clustal X (Thompson *et al.*, 1997), and the root phylogenetic tree was drawn using the software njplot (Saitou and Nei, 1987). The statistical confidence of the nodes was estimated by bootstrapping using 1,000 resample.

RESULTS AND DISCUSSION

Isolation of bacterial pathogens from soft rotted papaya fruits

A total of 12 bacteria were isolated from soft rotted papaya fruits collected from markets of different locations of Gazipur district, Bangladesh. Among 12 isolates, five isolates designated as CP01, CP02, CP03, CP04 and CP05 induced characteristic soft rot symptoms on potato slices. The rotted tissues were yellow in the center surrounded by dark brown edge (Fig. 1A). The results suggested the papaya isolates had pectolytic ability. These five isolates were tested for pathogenicity on their original host papaya by artificial inoculation. All isolates caused soft rot on papaya (Table 1). The dark brown to blackish rotted symptoms developed on papaya fruits within 2–3 days (Fig. 1B), which were similar to those observed in the natural infected fruits. The same bacterium was re-isolated from the inoculated fruits (data not shown). Colony morphology of the isolates on YPDA was white to grayish white, smooth, round, glistening, convex raised (Fig. 2).



Fig. 1. Rotted symptoms developed on potato slice (A) and papaya fruit (B) by a papaya isolate, CP03.



Fig. 2. Colony morphology of a papaya isolate, CP03 on YPDA medium.

Physiological and biochemical characteristics of the isolates

The isolates were Gram-negative and also negative in oxidase, gelatin liquefaction, indole, methyl red and arginine dihydrolase tests. All of the isolates were positive in catalase, oxidative fermentative, nitrate reduction, acetoin and urease production tests. The isolates were grown well at 41°C and 5% of salt concentration. They utilized lactose, glucose, melibiose, sucrose, rhamnose, arabinose, mannitol, inositol, sorbitol and citrate but not adonitol as sole sources of carbon (Table. 1).

The results of physiological and biochemical tests of the papaya isolates were consistent with descriptions of Holt *et al.* (1994), Alves *et al.* (2006) and Zheng *et al.* (2014) for *K. variicola*. Bacteriological characteristics

Table 1. Physiological and biochemical characteristics of papaya isolates

Characteristics	Present isolates ¹⁾ (n=5)	Reference species ²⁾			
		<i>Klebsiella variicola</i>	<i>Klebsiella pneumoniae</i>	<i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>	<i>Dickeya dadantii</i>
Potato soft rot test	+	nt	nt	+	+
Pathogenicity to papaya	+	nt	nt	nt	nt
Gram reaction	-	-	-	-	-
Catalase	+	+	+	+	+
Oxidative fermentative	+	+	+	+	+
Oxidase	-	-	-	-	-
Gelatin liquefaction	-	-	-	+	+
Nitrate reduction	+	+	+	+	+
Indole	-	-	-	-	+
Methyl red	-	-	-	+	-
Acetoin	+	+	+	+	+
Urease	+	+	+	-	-
Arginine dihydrolase	-	-	-	-	-
Growth at 41°C	+	+	+	+	+
Growth in 5% NaCl	+	+	+	+	+
Utilization of:					
lactose	+	+	+	+	-
glucose	+	+	+	+	+
melibiose	+	+	+	+	+
sucrose	+	+	+	+	+
rhamnose	+	+	+	+	+
arabinose	+	+	+	+	+
adonitol	-	-	+	-	-
mannitol	+	+	+	+	+
inositol	+	+	+	+	-
sorbitol	+	+	+	-	-
citrate	+	+	+	+	+

+, positive; -, negativ; nt, not tested

1) CP01-CP05

2) Results from Holt *et al.* (1994), Khan *et al.* (2000), Alves *et al.* (2006) and Zheng *et al.* (2014)




Fig. 3. Phylogenetic tree based on 16S rRNA gene sequences. Percentage bootstrap values only higher than 80% of 1,000 replicates are indicated at branching nodes.

of the papaya isolates were different from the characteristics of common bacterial soft rot pathogens *P. carotovorum* and *D. dadantii*; but very close to those of *K. variicola* and *K. pneumoniae*. Papaya isolates did not use adonitol like *K. variicola* but *K. pneumoniae* utilized adonitol as sole source of carbon. Rosenblueth *et al.* (2004) demonstrated that *K. variicola* is a very close species of *K. pneumoniae* but isolated genetically.

Identification and phylogenetic analysis of bacterial isolate based on 16S rRNA gene sequences

Analysis of the 16S rRNA gene sequence of the papaya isolate CP03 was performed to identify the bacterium. The partial sequence was deposited in the GenBank with accession number KX227610. The partial sequence of present isolate CP03 showed 99% similarity with 16S rRNA gene sequences of *Klebsiella* species containing *K. pneumoniae*, *K. variicola*.

The related sequences were retrieved from the NCBI and used for phylogenetic analysis.

The papaya isolate CP03 formed a cluster with *Klebsiella variicola* F2R9 and At-22 in the phylogenetic tree based on 16S rRNA gene sequence, but not supported by high bootstrap value (Fig. 3). Based on the results of the physiological and biochemical tests and phylogenetic analysis suggested that papaya soft rot causing isolates were *K. variicola*. For further detailed identification, multi locus sequence typing will be useful (Maatallah *et al.*, 2014)

K. variicola and *K. pneumoniae* are reported to cause soft rot in onion and banana (Liu *et al.*, 2015; Fan *et al.*, 2016). Although *Klebsiella* spp. are well recognized as clinical pathogens, previously they were isolated from different plant hosts such as rice, maize, sugarcane, banana, onion etc. (Rosenblueth *et al.*, 2004; Lin *et al.*, 2015; Liu *et al.*, 2015; Fan *et al.*, 2016). The results of this study also revealed the plant pathogenic nature

of *K. variicola*. To our knowledge, bacterial soft rot of papaya caused by *K. variicola* is a new disease in Bangladesh. Management of the disease at postharvest level will be the subject for future study.

AUTHOR CONTRIBUTIONS

S. Hossain conducted experiments and drafted the manuscript. A. A. Khan planned and initiated the research works and edited the manuscript. M. M. Rahman contributed the materials and provide support during molecular works. K. Iiyama performed bioinformatics analysis and edited the manuscript. N. Furuya provided valuable suggestions and edited the manuscript.

REFERENCES

- Alves, M. S., R. C. da Silva Dias, A. C. Dias de Castro, L. W. Riley and B. M. Moreira 2006 Identification of clinical isolates of indole positive and indole negative *Klebsiella* spp. *J. Clin. Microbiol.*, **44**: 3640–3646
- Ayers, S. H., P. Rupp and W. T. Johnson 1919 *A study of the alkali forming bacteria in milk*. U. S. Dept. Agric. Bull. No. 782
- De Boer, S. H. and A. Kelman 2001 Gram-negative bacteria: *Erwinia* soft rot group. In "Laboratory guide for identification of plant pathogenic bacteria, 3rd ed.", ed. by N. W. Schaad, J. B. Jones and W. Chun., pp. 56–72
- Dye, D. N. 1968 A taxonomic study of the genus *Erwinia*. I. The amylovora group. *New Zealand J. Sci.* **11**: 590–607
- Fan, H. C., L. Zeng, P. W. Yang, Z. X. Guo and T. T. Bai 2016 First report of banana soft rot caused by *Klebsiella variicola* in China. *Plant Dis.*, **100**: 517
- Hayward, A. C. 1992 Identification of *Pseudomonas solanacearum*. In "SAVERNET Bacterial wilt training course held on October 5 to November 16 AVRDC" pp. 101–102
- Himel, R. M., A. A. Khan, A. M. Akanda and M. Karim 2016 Characterization and identification of soft rot bacterial pathogens of different fruits in Bangladesh. *Int. J. Biosci.*, **9**: 1–9
- Holt, J. G., N. R. Krieg, P. H. Sneath, J. T. Staley and S. T. Williams 1994 *Bergey's manual of determinative bacteriology*. Wil-

- liams and Wilkins Co., Baltimore
- Hossain, M. S. 2016 Characterization of soft rot causing bacteria from fruits and vegetables and their control. An MS Thesis, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh
- Hugh, R. and E. Leifson 1953 The taxonomic significance of fermentative versus oxidative metabolism of carbohydrates by various gram-negative bacteria. *J. Bacteriol.*, **66**: 24–26
- Khan, A. A., N. Furuya, H. Ura and N. Matsuyama 2000 Rapid identification of *Erwinia chrysanthemi* isolated from soft rotted eggplant and *Phalaenopsis* sp. by lipid and fatty acid profiling. *J. Fac. Agr. Kyushu Univ.*, **44**: 257–263
- Kim, Y. K., S. D. Lee and C. S. Choi 2002 Soft rot of onion bulbs caused by *Pseudomonas marginalis* under low temperature storage. *Plant Pathol. J.*, **18**: 199–203
- Kovacs, N. 1956 Identification of *Pseudomonas solanacearum* by the oxidase reaction. *Nature*, **178**: 703
- Lelliot, R. A. and R. S. Dickey 1984 Genus VII. *Erwinia*. In “Bergey’s Manual of Systematic Bacteriology” vol. 1, ed. by N. R. Krieg and J. G. Holt, The Williams & Wilkins Co., Baltimore, Md. pp. 469–476
- Lin, Li., C. Wei, M. Chen, H. Wang, Y. Li, Y. Li, L. Yang and Q. An 2015 Complete genome sequence of endophytic nitrogen-fixing *Klebsiella variicola* strain DX120E. *Stand. Genomic Sci.*, **10**: 22
- Liu, S., M. Lv, Y. Gu, and J. Zhou 2015 First report of bulb disease of onion caused by *Klebsiella pneumoniae* in China. *Plant Dis.*, **99**: 1853
- Maatallah, M., M. Vading, M. H. Kabir, A. Bakhrouf, M. Kalin, P. Nauc ler, S. Brisse and C. G. Giske 2014 *Klebsiella variicola* is a frequent cause of bloodstream infection in the stockholm area, and associated with higher mortality compared to *K. pneumoniae*. *PLoS One*, **9**: e113539
- Meah, M. B. and A. A. Khan 1987 *Check list of fruit and vegetable diseases in Bangladesh. Survey of diseases of some important fruits and vegetables in Bangladesh*. Bangladesh Agricultural University, Mymensingh, Bangladesh
- Mortensen, C. N. 1997 *Seed bacteriology laboratory guide. Danish Government Institute of Seed Pathology (DGISP) for developing countries*. Copenhagen, Denmark. pp. 1–2
- Perombelon, M. C. M. and A. Kelman 1980 Ecology of soft rot *Erwinias*. *Ann. Rev. Phytopathol.*, **18**: 361–387
- Rosenblueth, M., L. Martinez, J. Silva and M. Romero 2004 *Klebsiella variicola*, a novel species with clinical and plant associated isolates. *Syst. Appl. Microbiol.*, **27**: 27–35
- Saitou, N. and M. Nei 1987 The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.*, **4**: 406–425
- Schaad, N. W. 1988 *Laboratory guide for identification of plant pathogenic bacteria*. 2nd ed. American Phytopathological Society Press
- Suslow, T. V., M. N. Schroth and M. Isaka 1982 Application of a rapid method for Gram differentiation of plant pathogenic and saprophytic bacteria without staining. *Phytopathology*, **72**: 917–918
- Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin and D. G. Higgins 1997 The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.*, **25**: 4876–4882
- Thornley, M. J. 1960 The differentiation of *Pseudomonas solanacearum* from other Gram-negative bacteria on the basis of arginine metabolism. *J. App. Bacteriol.*, **23**: 37–52
- Zheng, P., L. Zhang, L. Titan, L. Zhang, F. Chen, B. Z. Li and Z. Cui 2014 Isolation and characterization of novel bacteria containing deaminase from the rhizosphere resource on dry-farming lands. *Pak. J. Bot.*, **46**: 1905–1910