

Freezing Tolerance of *Lactuca sativa* and Induction of CBF and GolS Genes during Cold Treatment

HONJOH, Ken-ichi

Laboratory of Food Hygienic Chemistry, Division of Food Science and Biotechnology, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University

OKANO, Hitomi

Laboratory of Food Hygienic Chemistry, Division of Food Science and Biotechnology, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University

KAWABATA, Aya

Laboratory of Food Hygienic Chemistry, Division of Food Science and Biotechnology, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University

KUROKAWA, Masaru

Laboratory of Food Hygienic Chemistry, Division of Food Science and Biotechnology, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/1955390>

出版情報：九州大学大学院農学研究院紀要. 63 (2), pp.249-257, 2018-09-01. 九州大学大学院農学研究院

バージョン：

権利関係：



Freezing Tolerance of *Lactuca sativa* and Induction of *CBF* and *GolS* Genes during Cold Treatment

Ken-ichi HONJOH*, Hitomi OKANO¹, Aya KAWABATA¹, Masaru KUROKAWA¹, Taiki KIMURA¹, Takeshi MACHIDA², Yoshimitsu MASUDA and Takahisa MIYAMOTO

Laboratory of Food Hygienic Chemistry, Division of Food Science and Biotechnology,
Department of Bioscience and Biotechnology, Faculty of Agriculture,
Kyushu University, Fukuoka 812–8581, Japan

(Received May 1, 2018 and accepted May 8, 2018)

Plants respond to several environmental changes such as low- and high-temperatures, drought, flood, and so on, leading to development of stress tolerance. Acquisition abilities of freezing tolerance of three lettuce cultivars were investigated. All the investigated cultivars showed improvement of freezing tolerance at –3°C after exposure to non-freezing low-temperature at 2°C for 7 days. Out of them, papa lettuce cultivar was used to investigate expression levels of cold-responsive genes. The cultivar showed induction of *CBF* and *GolS* genes, which are well known as cold-responsive genes in other plants, during cold treatment. These results showed that lettuce has at least one cold-responsive signal pathway, suggesting a possibility that genetical modification might enhance the freezing tolerance.

Key words: C-repeat/DRE binding factor, freezing tolerance, galactinol synthase, lettuce

INTRODUCTION

Extreme weather events such as drought, flooding, typhoon, cold summer, and so on have threatened the stable production and the supply of agricultural produce such as crops and vegetables. Although lettuce is one of important produces and is affected by extreme weather events, there are not many reports concerning its response to environmental changes such as abiotic stress. Several researchers reported responses of lettuce to drought or salt stress (Garrido *et al.*, 2013; Leyva *et al.*, 2011; Porcel *et al.*, 2005; Ruiz-Lozano *et al.*, 1996), however there are few reports concerning responses of lettuce to cold stress.

From a model plant, *Arabidopsis thaliana*, one cDNA clone corresponding to a gene encoding cold-inducible transcriptional activator was isolated (Stockinger *et al.*, 1997). Now, the activator is well known as a protein binding to a cis-acting DNA regulatory element, C-repeat/dehydration responsive element (DRE), in plants and the protein stimulates expression of genes responsible for enhancement of drought tolerance or freezing tolerance of the plants. The activator protein was called as the C-repeat/DRE binding factor (CBF) and the homologous genes have been cloned from several plants. Introduction of a *CBF* gene from *Arabidopsis* into *Brassica napus* led to expression of the genes induced by CBF proteins and enhancement of freezing tolerance of the plant, showing the existence of

cold-responsive pathways in *B. napus* (Jaglo *et al.*, 2001). Although there are many reports concerning freezing tolerance or cold response of several plants such as *Arabidopsis*, there are few reports concerning cold response of plants such as lettuce which is not strong. If freezing tolerance of lettuce is improved, lettuce can be kept at lower temperatures than normal refrigerated temperature for storage. Thus, not only lettuce plants can acclimate to a rapid decline in temperature during cultivation, but also postharvest-storage period of lettuce is expected to be prolonged. In order to improve the freezing tolerance of lettuce plant, it is important to know the molecular response of the plant exposed to cold.

Arabidopsis overexpressing CBF3/DREB1A transcriptional factor accumulated raffinose and enhancement of freezing tolerance (Gilmour *et al.*, 2000). Raffinose family oligosaccharides (RFO) including galactinol and raffinose are well studied as protectants against abiotic stress such as drought, salinity, cold stress in plant cells (Taji *et al.*, 2002). For accumulation of galactinol and raffinose in plants, galactinol synthase catalyzes a first reaction for biosynthesis of RFO (Taji *et al.*, 2002). Introduction of cold-inducible galactinol synthase gene from *Medicago sativa* into tobacco plant enhanced multiple tolerance of the plant to abiotic stresses (Zhuo *et al.*, 2013).

In the present paper, we investigated freezing tolerance of three lettuce cultivars. From one of three cultivars, partial cDNA clones corresponding to *CBF* and *GolS* genes were isolated. Then, expression patterns of the genes during cold treatment of lettuce plants were investigated.

MATERIALS AND METHODS

Plant materials and growth conditions

Seeds of three kinds of *Lactuca sativa* L. (Papa let-

¹ Laboratory of Food Hygienic Chemistry, Division of Food Science and Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka 812–8581, Japan

² Department of Immunology, School of Medicine, Fukushima Medical University, Fukushima 960–1295, Japan

* Corresponding author (E-mail: honjoh@agr.kyushu-u.ac.jp)

tuce, Mama lettuce, and Berkley) were obtained from Nakahara Seeds Co. Ltd. (Fukuoka, Japan). The seeds were disinfected with 70% ethanol for 10–15 s, and then dipped in 0.2% sodium hypochlorite solution for 10 min. The seeds were then rinsed three times with sterile water. The seeds were sown in pots (7.5 cm diameter) with soil and lettuce plants were cultivated in a growth chamber (model MLR-350; SANYO, Co. Ltd., Tokyo, Japan) at 22°C under a photosynthetic photon flux density of about 40 $\mu\text{mol}/\text{m}^2\text{s}$ with a 16 h photoperiod. Sterilized tap water was poured to the soil 2–3 day intervals.

Cold treatment

Lettuce plants grown for 2 weeks were incubated in a refrigerator (model SPR-T1281; SANYO) at 2°C for indicated period. After this treatment, lettuce plants were used for following freezing treatment or harvested for preparation of total RNA. Harvested leaves were immediately frozen in liquid nitrogen and stored at –80°C until preparation of RNA.

Freezing test

Low-temperature-treated or non-treated lettuce plants were moved into incubator and kept at –3°C for 30 min. Then, crashed ice, which had been pre-incubated at –3°C, covered soil of the plant pot for ice nucleation. The lettuce plants were kept at –3°C for 24 h. For thawing, the frozen plants were moved to refrigerator and kept at 2°C for further 24 h. And then, the plants were moved to normal growth conditions at 22°C and further cultivated for a week to evaluate freezing tolerance. For evaluation of freezing tolerance, 5–7 lettuce plants were cultivated in a pot. Three pots were used for each experimental data point and four independent experiments were done. Viability was determined by expressing the number of plants, which survived after one week of freezing-thawing, as a percentage of number of plants subjected to freezing test.

Preparation of total RNA

For preparation of total RNA, a several leaves were

harvested. The harvested leaves were grounded in liquid nitrogen using mortar and pestle until the leaves became power. Total RNA of lettuce leaves was prepared from this grounded powder using RNeasy Plant Mini Kit (QIAGEN, Tokyo, Japan) according to the manufacturer's instructions.

Preparation of poly(A)⁺RNA

Poly (A)⁺RNA was purified from total RNA using Oligotex[™]-dT30<Super> mRNA Purification Kit (Takara, Kyoto, Japan), according to the manufacturer's instruction. The purified poly(A)⁺RNA was dissolved in RNase-free water and the concentration of RNA solution was adjusted to 0.5 $\mu\text{g}/\mu\text{l}$.

Synthesis of first strand cDNA

SMART Scribe[™] Reverse Transcriptase (Clontech, Mountain View, CA, USA) was used for cDNA synthesis. At first, a mixture of 2 μl of total RNA or poly(A)⁺RNA, 0.5 μl of oligo (dT) primer, and 2.5 μl water was heated at 72°C for 4 min in order to denature the RNA conformation. After cooling on ice, 2 μl of 5 × First-Strand buffer, 1 μl of 10 mM dNTP, 1 μl of 20 mM DTT, and 1 μl of 100 U/ μl of SMART Transcriptase RT (Clontech) were added to the mixture. The mixture was incubated at 42°C for 2 h. The reaction was terminated by heating at 72°C for 15 min. Obtained reaction mixture was used as first strand cDNA solution.

Determination of nucleotide sequences of partial cDNA fragments for *LsCBF*, *LsGolS1*, and *LsGolS2*

To determine sequence of cDNA of *LsCBF*, its partial fragments were obtained by using PCR with several primers (Table 1) and RACE fragments of each cDNA were obtained by using SMARTer RACE 5'/3' kit (Clontech) according to instructions' manual. Partial cDNA fragments of *LsGolS1* and *LsGolS2* were also obtained by using PCR with primers listed in Table 2. Then, sequencing of the obtained fragments was directly performed by a contractor (Macrogen Japan, Kyoto, Japan). Determined sequences were assembled for fur-

Table 1. Nucleotide sequences of the primers for amplification of partial fragments and determination of sequence of *LsCBF*

Primer name	Sequence
LsCBF No.1 (F)	5'-TCCGG(C/T)AAGTGGGT(C/T)TGTGA-3'
LsCBF No.2 (R)	5'-GCCGC(C/T)TT(C/T)TG(A/G)ATGTCCTT-3'
LsCBF No.3 (F)	5'-CCGGTAAGTGGGTTTGTGAG-3'
LsCBF No.4 (F)	5'-TCAGCGTGTTTGAATTTTGC-3'
LsCBF No.5 (R)	5'-GCCACACCCTCGTTTTCTTA-3'
LsCBF No.6 (F)	5'-TGCCGGTCCCGGAATCTAGT-3'
LsCBF No.7 (R)	5'-TTGCGTATCCATCAATCTGG-3'
LsCBF No.8 (F)	5'-GGAAGAAGTTCAAGGAAACACG-3'
LsCBF No.9 (R)	5'-CAATGGCACATTTGCGTATC-3'
3' CDS III primer (–T ₃₀)	5'-ATTCTAGAGGCCGAGGCGGCCGACATG-3'

ther sequence analysis.

Sequence analysis

Sequence alignment and phylogenetic analysis were performed using the Clustal W program (<http://clustalw.ddbj.nig.ac.jp/index.php?lang=ja>).

PCR and semi-quantitative RT-PCR

KOD Fx Neo DNA polymerase (Toyobo, Osaka, Japan) was used for PCR. Standard reaction was carried out according to the manufacturer's instruction. The PCR reaction was carried out with a 20 μ l mixture containing 0.2 U of KOD Fx Neo DNA polymerase, 1 \times KOD Fx Neo buffer, 400 μ M dNTP, 20 pmol of each appropriate primer (Tables 1 and 2), and 1 μ l of the reverse transcription (RT) reaction mixture. The mixtures were subjected to PCR amplification consisting of template DNA denaturation at 98°C for 10 s, primer-template annealing at 55°C for 30 s, and primer extension at 68°C for 30 s. Total RNA or poly(A)⁺RNA solutions were used in place of the RT reaction mixture in order to confirm the absence of genomic DNA in the RNA solution.

The relative levels of transcription were calculated by normalization of expression level of *actin* gene from lettuce (Accession number as mRNA, AY260165.1).

qPCR

qPCR was performed on Mx3000P™ Real-Time PCR System (Stratagene, La Jolla, CA, USA) with the SYBR Premix Ex Taq™ (Perfect Real Time) according to the manufacturer's instruction. Each reaction was carried out at least three times, with threshold cycle (C_T) determined in duplicate. Used primers were shown in Table 2.

Southern blot analysis

Genomic DNA from lettuce was prepared using a DNeasy Plant Mini Kit (Qiagen Co. Ltd., Hilden, Germany). Ten micrograms of the genomic DNA was digested with *Eco*RI, fractionated on an agarose gel, and blotted onto a Hybond N⁺ nylon membrane (GE healthcare, Heidenberg, Germany). The membrane was subjected to Southern blot analysis. For preparation of a DIG-labeled probe, a PCR DIG Probe Synthesis Kit (Roche, Basel, Switzerland) was used and PCR was car-

ried out using LsCBF No. 3 and No. 7 primers and a partial fragment of LsCBF as a template. Then, the blotted membrane was incubated at 55°C in a DIG Easy Hyb solution (Roche), and the re-incubated in the same solution supplemented the DIG-labeled probe overnight. The membrane was washed with 2 \times SSC and 0.1% SDS at room temperature, followed by 0.1 \times SSC and 0.1% SDS at 68°C. After washing, detection was carried out by using a DIG Nucleic Acid Detection Kit (Roche).

Statistical analysis

Statistical analysis was performed using Excel 2011 (Microsoft, USA) with the add-in software Satcel 2 (Yanai 2004).

RESULTS AND DISCUSSION

Freezing tolerance of lettuce plants

In order to study freezing tolerance of lettuce, three cultivars (Papa lettuce, Mama lettuce, and Berkley) were grown at 22°C for two weeks. Then, they were cold-treated at 2°C for indicated periods and moved into a freezer at -3°C. Freezing tolerance was evaluated with viability of lettuce plants which had been frozen at -3°C for 24 h and then grown at 22°C for one week. Survival rates of lettuce plants, which had been cold-treated at 2°C for 4–7 days, were significantly enhanced compared to those of untreated plants (Figs. 1 and 2). Furthermore, viabilities of lettuce plants after freezing and thawing were increased with the length of cold treatment period. Berkley showed the highest viability among three cultivars cold-treated for 4 days, however, viabilities of Papa lettuce and Mama lettuce cultivars treated for 7 days were enhanced as same as that of Berkley cultivar treated for 7 days. So, any remarkable differences in freezing tolerance among three cultivars were not observed after 7 days. In further experiments, Papa lettuce cultivar was used for further experiments.

To our knowledge, there are no research papers regarding to acquisition of freezing tolerance of lettuce plants. In Japan, lettuce is cultivated at cool areas located in high altitude in summer and at areas located in low altitude in winter. Thus, lettuce is estimated to relatively be strong against non-freezing low tempera-

Table 2. Nucleotide sequences of the primers of the target genes for semi-quantitative RT-PCR or qPCR.

Gene (Accession number)	Primer	Nucleotide sequence	Ampricon (bp)
<i>LsCBF</i> (LC380915)	LsCBF NO.7(F)	5'-GGAAGAAGTTCAAGGAAACACG-3'	530
	LsCBF No. 8(R)	5'-CAATGGCACATTTGCGTATC-3'	
<i>LsGolS1</i> (DY975888)	LsGolS1C(F)	5'-TTT ACC CAC CGG AAA GTC AG-3'	185
	LsGolS1C(R)	5'-GTC CAT CAC CGC GTA AAA GT-3'	
<i>LsGolS2</i> (DY971098)	LsGolS2C(F)	5'-CCC AAT CTC TCC ACC TAC CA-3'	201
	LsGolS2C(R)	5'-GCA GTA GTG CAC CAC CTT GA-3'	
<i>LsActin</i> (AY260165.1)	LsActin (F)	5'-AAG CCC AAT CGA AAA GAG GT-3'	199
	LsActin (R)	5'-GCC TCT GTA AGG AGC ACA GG-3'	




Fig. 5. Photos of lettuce plants after freezing–thawing.

Lettuce plants were incubated at 2°C for indicated periods, frozen at –3°C for 24 h, thawed at 2°C for 24 h, and then grown under normal cultivation conditions at 22°C. Samples of 0 day were not incubated at 2°C. The photos were taken one week after freezing–thawing. Representative pictures of the samples are shown here.




Fig. 2. Freezing tolerance of lettuce evaluated by viability. (A) Papa lettuce, (B) Mama lettuce and (C) Berkley. Values are means \pm S.D. Significance is indicated as: * $p < 0.05$ or ** $p < 0.01$.

ture. Our results showed that cold treatment at 2°C improved freezing tolerance of lettuce at –3°C. This indicates a possibility that freezing tolerance of lettuce might be further improved by using any means such as transgenic approaches. Although lettuce is major produce, there is one report regarding improvement of freezing tolerance of lettuce by using transgenic technique. Vanjildorj *et al.* (2005) reported that transgenic lettuce with transcriptional factor *ABF3* gene enhanced cold tolerance of the lettuce plant at –4°C. Although detail mechanisms of the improvement of freezing tolerance were not mentioned in the paper, overexpression of the *ABF3* gene would lead to expression of *ABF3*–inducible genes in the lettuce plant. Thus, transgenic approach has a possibility that freezing tolerance of lettuce plants might be dramatically improved dependently on introduced genes.

Regarding to response of lettuce to abiotic stress such as salt stress, there are a few research papers of stress response of *L. sativa* using techniques of molecular biology except for a study on drought–induced expression of *DREB2A* gene, designated *LsDREB2A*, from *L. sativa* by Kudo *et al.* (2014). Park *et al.* (2005) also improved salt and drought tolerance of lettuce by constitutive expression of *LEA* with transgenic technique. *DREB2A* is a transcriptional factor regulating stress–inducible genes and *LEA* probably work as a protectant against several stresses. Thus, utilization of transcriptional factors or stress tolerance plants would be helpful for enhancement of freezing tolerance.

Cloning of a partial fragment of cDNA for *LsCBF* gene

CBF(*DREB1*) and *DREB2* are well known as ones of major stress responsive transcriptional factors (Mizoi *et al.*, 2012). Although cDNA corresponding to *LsDREB2A*




Fig. 3. Comparison of deduced amino acid sequence of partial fragment of *LsCBF* with those of other CBF proteins.

Identical amino acid residues and relatively similar amino acid residues are shaded in black and grey colors, respectively. Dashes are no amino acids.

AP2 domains and other two conserved regions of CBF proteins, PKKP/RAGR xKF xETRHP and DSAWR regions, are also shown. Numbers in parenthesis are NCBI reference or GenBank accession numbers. Plants explained in abbreviations are as follows. At: *Arabidopsis thaliana*, Cs: *Cichorium endivia*, Ls: *Lactuca sativa*, Rc: *Ricinus communis*, Vv: *Vitis vinifera*.

was isolated from lettuce and studied by Kudo *et al.* (2014), another type such as *CBF(DREB1)* had not been identified in lettuce so far. This means that expression pattern of *CBF* gene during cold treatment of lettuce has not been investigated. At first, we designed degenerate primers for amplification of a partial fragment of cDNA corresponding to *CBF* gene (*LsCBF*) from lettuce (Table 1). Then, we carried out PCR by using cDNA derived from poly(A)⁺RNA of papa lettuce. Nucleotide sequence of the obtained PCR fragment was determined. After obtaining a part of cDNA of *LsCBF*, RACE method and several degenerated primers were used for determination of sequence of cDNA of *LsCBF*. Although we could not obtain full-length sequence of cDNA for *LsCBF*, a partial fragment composing of 672 bp (DDBJ accession number, LC380915) was obtained and the fragment encoded 191 amino acids. The deduced amino acid sequence showed similarity to those of CBF proteins from other plant sources (Fig. 3). CBF transcriptional factor belongs to transcriptional factors of AP2 family (Jaglo *et al.*, 2001). AP2 domain was found in the deduced amino acid sequence of the obtained fragment. Furthermore, PKKP/RAGR xKF xETRHP and DSAWR regions, which are specific to common of CBF proteins, were found in the sequence. Thus, the obtained fragment was confirmed to be a part of cDNA for *LsCBF* gene. This result shows a possibility that lettuce has a




Fig. 4. Southern blot analyses of genomic DNA of lettuce. Lanes: 1, Positive control; 2, Intact lettuce genomic DNA; 3, *EcoRI* digested lettuce genomic DNA. PCR product of *LsCBF* was used as a probe.




Fig. 5. Phylogenetic analysis using the Neighbor-Joining method and based on predicted amino acid sequences of confirmed or putative CBF/DREB1 and DREB2 proteins from different plant species. Comparison sequences were Aa, *Ageratina adenophora*; At, *Arabidopsis thaliana*; Cs, *Cucumis sativa*; Dv, *Dendranthema vestitum*; Fv, *Fragaria vesca*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Sl, *Solanum lycopersicum*; Vv, *Vitis vinifera*; Zm, *Zea mays*. Numbers in parenthesis are NCBI reference or GenBank accession numbers.

cold responsive pathway as same as other plants such as *Arabidopsis thaliana* which acquires freezing tolerance by exposure to cold.

Because there are over 10 homologs for *CBF* in *Arabidopsis*, it was expected that there are several homologs for *LsCBF* genes in the present study. Thus, by using a partial fragment of *LsCBF* as a probe, we tried to perform genomic Southern blot hybridization (Fig. 4). At least one strongly-hybridized band and several weakly-hybridized bands were detected, suggesting existence of other *LsCBF-like* genes. At present, a *CBF* homolog gene, *LsDREB2A* (accession number AB641390.1) was cloned from lettuce by another researcher (Kudo *et al.*, 2014). We compared deduced amino acid sequence of the cloned *LsCBF* to those of other *CBF* homologs and made a phylogenetic tree of *CBF* proteins (Fig. 5). The results showed that *LsCBF* protein was categorized to DREB1 group and *LsDREB2A* protein was categorized to DREB2 group. DREB1 is major regulator of cold-response and DREB2 is regulator of heat and drought response (Mizoi *et al.*, 2012). Thus, the corresponding gene to isolated cDNA was considered to be newly identified for cold regulation factor in the present paper.




Fig. 6. Differential expression analysis of *CBF* genes in lettuce under cold treatment.

Expression levels of *LsCBF* were estimated by (A) semi-quantitative PCR and (B) quantitative RT-PCR. Total RNAs were extracted from leaves of plants treated with 2°C for 0, 1, 6, and 24h, and subjected to respective measurement methods. Quantification of transcriptional levels for *LsCBF* gene, compared with a control, was calculated after normalization to the actin gene of *L. sativa*.

Response of *LsCBF* to cold treatment

The expression pattern of *CBF* gene was investigated by using semi-quantitative and real-time RT-PCR (Fig. 6), *CBF* gene was remarkably induced at 1 and 6 h of cold treatment. However, at 24 h of cold treatment, its transcriptional level returned to the almost same level in non-cold treated lettuce plant. This suggests that lettuce has a signal pathway responding to cold as well as other plants which acquire freezing tolerance. Although expression of a *CBF* gene in tomato plants is induced by cold treatment, the expression level of the gene, at 24 h of cold treatment, returned to the level in warm-grown plants (Jaglo *et al.*, 2001). As well as freezing-sensitive plants such as tomato, even if *CBF* gene from lettuce might be induced by cold once, the expression level of the gene appeared to be lowered soon. Furthermore, Jaglo *et al.* (2001) suggested that there were few cold-regulated genes responsive to CBF/DRE elements in tomato. Similarly, there might be fewer cold-regulated genes in lettuce than those in other freezing-tolerant plants.

Zhang *et al.* (2004) reported that tomato (*Lycopersicon esculentum*) has three homologues for *CBF* gene (*LeCBF1*, *LeCBF2*, *LeCBF3*) and only *LeCBF1* gene was cold-inducible. They also confirmed that *LeCBF1* functions in transgenic *Arabidopsis* as well as *AtCBF3*, leading to enhancement of freezing tolerance. However, overexpression of *LeCBF1* or *AtCBF3*




Fig. 7. Phylogenetic analysis using the Neighbor-Joining method and based on predicted amino acid sequences of confirmed or putative GolS proteins from different plant species. Comparison sequences were Al: *Arabidopsis lyrata*, At: *Arabidopsis thaliana*, Cs: *Cichorium endivia*, Gm: *Glycine max*, Mt: *Medicago truncatula*, Ls: *Lactuca sativa*, Pt: *Populus trichocarpa*, Rc: *Ricinus communis*, Vv: *Vitis vinifera*, Zm: *Zea mays*. Numbers in parentheses are NCBI reference or GenBank accession numbers.




Fig. 8. Differential expression analysis of *LsGolS1* and *LsGolS2* genes in lettuce under cold treatment. Expression levels of *LsGolS1* and *LsGolS2* were estimated by (A) semi-quantitative PCR and (B) quantitative RT-PCR. Total RNAs were extracted from leaves of plants treated with 2°C for 0, 1, 6, and 24h, and subjected to respective measurement methods. Quantification of transcriptional levels for *LsGolS1* and *LsGolS2* genes, compared with a control, was calculated after normalization to the actin gene of *L. sativa*.

gene in transgenic tomato plants did not enhance freezing tolerance of the plants. Furthermore, by using microarray analysis, there were only four genes which were up-regulated by LeCBF1 or AtCBF3 (Zhang *et al.*, 2004).

As described above, tomato is sensitive to freezing stress (Jaglo *et al.*, 2001). In the present paper, it was confirmed that at least one *CBF1* gene homologue (*LsCBF*) exists in lettuce (Fig. 4) and was up-regulated by cold

(Fig. 3). However, similar case to that in tomato (Zhang *et al.*, 2004) was anticipated. That is, even if CBF proteins are expressed in lettuce, freezing tolerance of the lettuce plant expressing the proteins might not be improved. The fact suggests that there might be only few genes which are up-regulated by *LsCBF* in lettuce. Furthermore, it also suggests that even if expression level of *LsCBF* in lettuce is enhanced, improvement of freezing tolerance of lettuce is not expected. Thus, we targeted another gene with functions leading to improvement of freezing tolerance.

Cloning of partial fragments of cDNA for *LsGolS* genes and cold response of the genes

GolS is a key enzyme for raffinose synthesis and catalyzes synthesis of galactinol using UDP-galactose and myo-inositol as substrates (Taji *et al.*, 2002). One of the corresponding genes is known as cold responsive genes (Zhuo *et al.*, 2013). Thus, we started to analyze GolS genes in lettuce. By searching lettuce EST database, two candidates of EST for GolS genes were found and GenBank accession numbers of the candidate ESTs were DY975888 and DY971098. Deduced amino acid sequences were determined from nucleotide sequences of respective EST clones. The two EST clones were estimated to be derived from genes encoding GolS proteins and the corresponding genes were designated *LsGolS1* and *LsGolS2*. The deduced amino acid sequences of *LsGolS1* and *LsGolS2* showed high similarity to those of *GolS* gene from other higher plants and phylogenetic tree was constructed (Fig. 7).

By using Primer3 (<http://primer3.ut.ee>), primers for *GolS* genes were designed for analysis with semi-quantitative and qPCR. As expected based on the sequences, about 180-bp and 200-bp PCR products were amplified (data not shown). Nucleotide sequences of the products were determined and the amplified products were confirmed as parts of *LsGolS1* and *LsGolS2* genes, respectively. Then, semi-quantitative RT-PCR and qPCR were done for analysis of expression levels of respective *LsGolS* genes during cold treatment (Fig. 8). The results showed that the expression of both the genes were induced by cold. As well as *GolS* genes from other plants (Liu *et al.*, 1998; Taji *et al.*, 2002; Phan *et al.*, 2009), *LsGolS* genes were confirmed to be induced by cold. *LsGolS1* was induced at 24 h and *LsGolS2* was induced at 6-h cold treatment. Expression level of *LsGolS2* seemed higher than that of *LsGolS1*.

As shown in Fig. 2, freezing tolerance of lettuce at -3°C was enhanced. The level of the freezing tolerance might be enough to overcome frost damage at around -3°C during cultivation period, however, it is not so high compared to those of other frost hardy plants such as spinach and *Arabidopsis* that acquire freezing tolerance at around -10°C (Guy *et al.*, 1985; Gilmour *et al.*, 1991). Overexpression of galactinol synthase in transgenic plants improved tolerance of oxidative stress, salinity and osmotic stress, chilling stress, or freezing stress (Nishizawa *et al.*, 2008; Sun *et al.*, 2013; Shimosaka and Ozawa, 2015; Zhuo *et al.*, 2012). Thus, by using trans-

genic approach, constitutive expression of *GolS* has a possibility to further improve freezing tolerance of lettuce at lower temperatures.

In the present paper, we showed that lettuce has a cold response pathway, suggesting a possibility to improve freezing tolerance by using gene manipulation.

ACKNOWLEDGEMENTS

This study was financially supported by The SKYLARK Food Science Institute.

REFERENCES

- Garrido, Y., J. A. Tudela, A. Marin, T. Mestre, V. Martinez and M. Gil 2013 Physiological, phytochemical and structural changes of multi-leaf lettuce caused by salt stress. *J. Sci. Food. Agric.*, **94**: 1592–1599
- Gilmour, S. J., A. M. Sebolt, M. P. Salazar, J. D. Everard and M. F. Thomashow 2000 Overexpression of the Arabidopsis *CBF3* transcriptional activator mimics multiple biochemical changes associated with cold acclimation. *Plant Physiol.*, **124**: 1854–1865
- Gilmour, S. J. and M. F. Thomashow 1991 Cold acclimation and cold-regulated gene expression in ABA mutants of *Arabidopsis thaliana*. *Plant Mol. Biol.*, **17**: 1233–1240
- Guy, C. L., K. J. Niemi and R. Brambl 1985 Altered gene expression during cold acclimation of spinach. *Proc. Natl. Acad. Sci. USA*, **82**: 3673–3677
- Jaglo, K. R., S. Kleff, K. L. Amundsen, X. Zhang, V. Haake, J. Z. Zhang, T. Deits and M. F. Thomashow 2001 Components of the Arabidopsis C-repeat/dehydration responsive element binding factor cold-response pathway are conserved in *Brassica napus* and other plant species. *Plant Physiol.*, **127**: 910–917
- Kudo, K., T. Oi, and Y. Uno 2014 Functional characterization and expression profiling of a *DREB2*-type gene from lettuce (*Lactuca sativa* L.). *Plant Cell Tissue Org. Cult.*, **116**: 97–109
- Leyva, R., E. Sánchez-Rodríguez, J. J. Ríos, M. M. Rubio-Wilhelmi, L. Romero, J. M. Ruiz and B. Blasco 2011 Beneficial effects of exogenous iodine in lettuce plants subjected to salinity stress. *Plant Sci.*, **181**: 195–202
- Liu, J.-J., D. C. Krenz, A. F. Galvez, and B. O. de Lumen 1998 Galactinol synthase (GS): increased enzyme activity and levels of mRNA due to cold and desiccation. *Plant Sci.*, **134**: 11–20
- Mizoi, J., K. Shinozaki and K. Yamaguchi-Shinozaki 2012 AP2/ERF family transcription factors in plant abiotic stress responses. *Biochim. Biophys. Acta–Gene Regul. Mech.*, **1819**: 86–96
- Nicolle, C., N. Cardinault, E. Gueux, L. Jaffrelo, E. Rock, A. Mazur, P. Amouroux and C. Révész 2004 Health effect of vegetable-based diet: lettuce consumption improves cholesterol metabolism and antioxidant status in the rat. *Clin. Nutr.*, **23**: 605–614
- Phan, T., Y. Ishibashi, T. Yuasa and M. Iwaya-Inoue 2009 Chilling stress induces galactinol synthase (*OsGolS1*) in rice seedling. *Cryobiol. Cryotechnol.*, **55**: 139–146
- Porcel, R., R. Azcón and J. M. Ruiz-Lozano 2005 Evaluation of the role of genes encoding for dehydrin proteins (LEA D-11) during drought stress in arbuscular mycorrhizal *Glycine max* and *Lactuca sativa* plants. *J. Exp. Bot.*, **56**: 1933–1942.
- Ruiz-Lozano, J. M., R. Azcón and J. M. Palma 1996 Superoxide dismutase activity in arbuscular mycorrhizal *Lactuca sativa* plants subjected to drought stress. *New Phytol.*, **134**: 327–333
- Shimosaka, E. and K. Ozawa 2015 Overexpression of cold-inducible wheat galactinol synthase confers tolerance to chilling stress in transgenic rice. *Breed. Sci.*, **65**: 363–371
- Stockinger, E. J., S. J. Gilmour and M. F. Thomashow 1997 *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl*

- Acad Sci USA*, **94**: 1035–1040
- Sun, Z., Q. W. Qi, Z. Wang, P. Li, C. Wu, H. Zhang and Y. Zhao 2013 Overexpression of *TsGOLS2*, a galactinol synthase, in *Arabidopsis thaliana* enhances tolerance to high salinity and osmotic stresses. *Plant Physiol. Biochem.*, **69**: 82–89
- Vanjildorj, E., T.-W. Bae, K.-Z. Riu, S.-Y. Kim and H.-Y. Lee 2005 Overexpression of *Arabidopsis ABF3* gene enhances tolerance to drought and cold in transgenic lettuce (*Lactuca sativa*). *Plant Cell Tissue Org. Cult.*, **83**: 41–50
- Yanai, H. 2004 Statcel—the useful add-in software forms on Excel, 2nd edn. OMS, Tokyo
- Zdravković, J. M., G. S. Aćamović–Djoković, J. D. Mladenović, R. M. Pavlović and M. S. Zdravković 2014 Antioxidant capacity and contents of phenols, ascorbic acid, β -carotene and lycopene in lettuce. *Hem. Ind.*, **68**: 193–198
- Zhang, X., S. G. Fowler, H. Cheng, Y. Lou, S. Y. Rhee, E. J. Stockinger and M. F. Thomashow 2004 Freezing-sensitive tomato has a functional CBF cold response pathway, but a CBF regulon that differs from that of freezing-tolerant *Arabidopsis*, *Plant J.*, **39**: 905–919
- Zhuo, C., T. Wang, S. Lu, Y. Zhao, X. Li and Z. Guo 2013 A cold responsive galactinol synthase gene from *Medicago falcate* (*MfGolS1*) is induced by myo-inositol and confers multiple tolerances to abiotic stresses. *Physiol. Plant.*, **149**: 67–78
- Zuther, E., K. Büchel, M. Hundertmark, M. Stitt, D.K. Hinch and A. G. Heyer 2004 The role of raffinose in the cold acclimation response of *Arabidopsis thaliana*. *FEBS Lett.*, **576**: 169–173