Biochemical Changes in Chinese Pond Mussel *Anodonta woodiana* Following Exposure to Copper

LIU, Hongbo

Key Laboratory of Fishery Ecological Environment Assessment and Resource Conservation in Middle and Lower Reaches of the Yangtze River, Freshwater Fisheries Research Center, Chinese Academy of Fishery Sciences | Laboratory of Marine Environmental Science, Faculty of Agriculture, Kyushu University

CHEN, Xiubao

Key Laboratory of Fishery Ecological Environment Assessment and Resource Conservation in Middle and Lower Reaches of the Yangtze River, Freshwater Fisheries Research Center, Chinese Academy of Fishery Sciences | Laboratory of Marine Environmental Science, Faculty of Agriculture, Kyushu University

OSHIMA, Yuji

Aquatic Biomonitoring and Environmental Laboratory, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University | Laboratory of Marine Environmental Science, Faculty of Agriculture, Kyushu University

SHIMASAKI, Yohei

Aquatic Biomonitoring and Environmental Laboratory, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University | Laboratory of Marine Environmental Science, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/1955400>

出版情報：九州大学大学院農学研究院紀要. 63 (2), pp.311-318, 2018-09-01. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Biochemical Changes in Chinese Pond Mussel *Anodonta woodiana* Following Exposure to Copper

Hongbo LIU¹, Xiubao CHEN¹, Yuji OSHIMA², Yohei SHIMASAKI²,
Tao JIANG¹ and Jian YANG^{1,*}

Laboratory of Marine Environmental Science, Faculty of Agriculture,
Kyushu University, Fukuoka 812–8581, Japan

(Received May 1, 2018 and accepted May 8, 2018)

We evaluated the potential utility of biochemical parameters as indicators of exposure to water-borne copper (Cu) in the freshwater mussel *Anodonta woodiana*. Semi-static renewal tests showed that the 96 h median lethal concentration of Cu was 3.4 mg/L (95% confidence interval [CI]: 2.1–6.5 mg/L) for juveniles and 22.1 mg/L (CI: 13.8–28.6 mg/L) for adult *A. woodiana*. In vivo bioassays were performed to determine the activity of the antioxidant enzymes superoxide dismutase (SOD) and catalase (CAT), and the levels of glutathione (GSH) in the adductor muscle, foot, gill, mantle, and visceral mass of *A. woodiana* after 48 and 96 h of exposure to Cu concentrations of 0.125 and 0.25 mg/L for juveniles, and 0.75, 1.5, 3 mg/L for adults. The levels of the lipid peroxidation product, malondialdehyde (MDA), were evaluated in five adult tissues. Our results indicated that SOD and CAT activity, and GSH levels in juvenile mussel foot and adductor muscle of adults might be sensitive to Cu exposure. This antioxidant defense system (CAT, GSH) was significantly enhanced in the mantle of juvenile mussel exposed to 0.25 mg Cu/L. The effect of Cu on the foot and mantle of juveniles, and the adductor muscle of adults seemed to be directly correlated with the levels of reactive oxygen species and antioxidant defense system components. Our data suggest that the biochemical responses (SOD and/or CAT, GSH) in the foot and mantle of juveniles, and the adductor muscle of adult *A. woodiana* might be good biomarkers of Cu contamination.

Key words: *Anodonta woodiana*; copper; tissue; biochemical changes; biomarker

INTRODUCTION

Copper (Cu) is an important micronutrient and essential element for bivalve development; however, it is also a widespread pollutant found in aquatic environments (Lorenzo *et al.*, 2005; Rivas *et al.*, 2007), ranking as the metal of highest concern for aquatic organisms (Donnachie *et al.*, 2014). At low levels of exposure (<0.1 mg/L) mussels can bioaccumulate Cu to various extents in different tissues (Amiard-Triquet *et al.*, 1986). Excess Cu is frequently found following chemical treatments, such as fungicide application in aquaculture ponds to eradicate filamentous algae; application of copper sulfate as high as 2 mg/L has been reported (Li *et al.*, 2007). Such high concentration may result in serious injury to aquatic organisms through the production of reactive oxygen species (ROS) and oxidative damage.

Antioxidant defenses are present in all aerobic organisms, and involve enzymatic and non-enzymatic constituents (Szaleczky *et al.*, 1999). Antioxidant enzymes (superoxide dismutase [SOD] and catalase [CAT]) and free radical scavengers (reduction of superoxide anion), which remove ROS and maintain cell homeostasis, rep-

resent the first line of defense against the toxic effects of xenobiotics that may occur following exposure to pollutants (Doyotte *et al.*, 1997). Nevertheless, when organisms are exposed to high levels of pollutants, antioxidant systems may be inhibited and oxidative stress may lead to enzymatic inactivation and lipid peroxidation (LPO) (Al-Subiai *et al.*, 2011; Doyotte *et al.*, 1997).

The development of biochemical biomarkers of non-lethal exposure may provide sensitive and rapid indicators that can be used to assess the cellular effects of contaminants. A recent study that measured oxidative stress and the antioxidant response of mussel to environmental chemicals was able to demonstrate that mussel condition can be rapidly documented (Farris and van Hassel, 2007). SOD, CAT, glutathione (GSH), and malondialdehyde (MDA) are the most common markers of antioxidant defense or oxidative damage that are affected by metal exposure (Eertman *et al.*, 1995; Goswami *et al.*, 2014; Radak *et al.*, 2017). The antioxidant defense system involved in the detoxification of xenobiotics and their metabolites (biotransformation and antioxidant enzymes) has been extensively studied (Cossu *et al.*, 1997, 2000; Doyotte *et al.*, 1997), and has well-defined impacts on mussels.

The nature of antioxidant defense may differ for euryoxic and stenoxic organisms (Livingstone *et al.*, 1992). However, few investigators have successfully used these reliable, sensitive, and definitive measures for freshwater bivalves. In fact, in the freshwater mussel, *Unio tumidus*, reduced glutathione was shown to be the most sensitive parameter indicating toxicant exposure (Doyotte *et al.*, 1997). Therefore, freshwater mus-

¹ Key Laboratory of Fishery Ecological Environment Assessment and Resource Conservation in Middle and Lower Reaches of the Yangtze River, Freshwater Fisheries Research Center, Chinese Academy of Fishery Sciences, Wuxi 214081, China

² Aquatic Biomonitoring and Environmental Laboratory, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan

* Corresponding author: E-mail: jiany@ffrc.cn (J. Yang)

sels are potentially suitable species for assessing aquatic contaminants and biochemical responses, allowing freshwater environments to be monitored.

The swan mussel *Anodonta woodiana* Lea, 1834 is typical among unionid bivalves; it has largely expanded its worldwide distribution and is one of the most successful invasive mussel species (especially in Europe) (Douda *et al.*, 2012; Guarneri *et al.*, 2014). This species easily meets the requirements for environmental monitoring due to its wide distribution, sessile lifestyle, and high tolerance to chemical contaminants (Chen and Xie, 2005; Liu *et al.*, 2010; Yokoyama and Park, 2002). *A. woodiana* has been successfully bred and cultured under artificial conditions in our laboratory (Chen *et al.*, 2015). Since artificially-propagated *A. woodiana* displayed low background levels of heavy metals (Chen *et al.*, 2012), it is suitable for standardized monitoring or toxicity testing of waterborne contaminants. Thus far, the toxicity and bioaccumulation of organotins (Yang *et al.*, 2008), organochlorines (Bian *et al.*, 2009), and heavy metals (Liu *et al.*, 2010) have been studied using *A. woodiana* from freshwater fisheries and aquaculture. However, less is known about the biochemical response of these contaminants or the potential of target tissues in the mussel at different life history stages.

Since we hypothesize that Cu exposure will lead to oxidative stress and induce LPO (Regoli, 1998), and that such disruption may have age-, tissue-, and dose-specific tendencies in *A. woodiana*, the aim of this study was to investigate the biochemical responses of five tissues of *A. woodiana* (i.e., adductor muscle, gill, foot, mantle, and visceral mass) following exposure to nonlethal Cu concentrations. The impaired response was assessed by measuring the levels of biomarkers of oxidative damage, namely SOD, CAT, GSH, and LPO. In this study, LPO was determined by measuring MDA. MDA is a common end-product of LPO, and is highly reactive and can form simple and complex adducts with proteins (Que *et al.*, 2013). This study was performed to elucidate the biochemical response of bivalves to Cu exposure and to develop biomarkers for the assessment of Cu ecotoxicity in this freshwater bivalve mollusk.

MATERIALS AND METHODS

Chemicals

American Standard Testing Method (ASTM)-reconstituted water (ASTM, 2006) was prepared by adding reagent-grade salts (NaHCO_3 , MgSO_4 , $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$, and KCl; analytical reagent grade quality) to deionized water, and was used in all reference toxicant tests. The nominal concentrations of Cu, sodium (Na), magnesium (Mg), potassium (K), and calcium (Ca) were 0, 30, 15, 2, and 10 mg/L, respectively. Copper sulfate ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, analytical reagent grade quality) was chosen as the test chemical and was prepared by adding and dissolving adequate chemical stock to the ASTM-reconstituted water. The chemicals were purchased from Sinopharm Chemical Reagent Co., Ltd., Shanghai, China.

Sampling

Six-month-old (mean $\pm$ standard deviation: shell length 25.0 ± 1.5 mm, shell height 15.2 ± 1.5 mm, shell width 6.6 ± 0.9 mm) and 4-year-old adults (shell length 85.3 ± 3.6 mm, shell height 53.4 ± 2.8 mm, shell width 33.7 ± 1.9 mm) of *A. woodiana* were collected from our culture pond at Nanquan Aquatic Base, Freshwater Fisheries Research Center, Chinese Academy of Fishery Sciences. The culture contains an artificially-propagated, standardized, healthy *A. woodiana* population with successful recruitment. Live mussels were transported directly to the laboratory and kept in sediment-free container ($1 \times 1 \times 1$ m) supplied with 30 L of dechlorinated tap water. Aeration was provided continuously by plastic tubes with a gentle air flow. The mussels were acclimatized to laboratory conditions for 5 days prior to the experimental period. Mussels were fed with 1 g/L of *Chlorella* sp. (Chlorella Industry Co., Tokyo, Japan) once every 2 days and maintained under a 16:8 h light/dark photoperiod. Mussel survival exceeded 95%. The criteria required for a death record were prolonged valve opening and no response to tactile stimulation (APHA, 1999).

Toxicity experiment

Juvenile mussel treatment

The test treatments (acute toxicity tests) were performed according to the methods set forth in the Standard Guide for Conducting Laboratory Toxicity Tests with Freshwater Mussels of the American Society for Testing and Materials (ASTM, 2006). Five juvenile mussels were placed in a 200 mL glass crystallizing dish (122 mm diameter, 22 mm deep) containing 150 mL ASTM-reconstituted water with a Cu concentration of 0, 0.625, 1.25, 2.5, 5, or 10 mg/L (nominal concentration hereinafter). Each concentration was tested in three dishes (five mussels per dish) and the 96 h EC_{50} value (Wang *et al.*, 2009, 2011) was calculated via a probit analysis.

Based on the results of the acute toxicity tests, the effects of nonlethal Cu concentrations on antioxidant enzyme activity and GSH in the tissues of juvenile mussels were investigated. After a 48 h acclimation period, mussels were placed into glass aquaria ($20 \times 20 \times 27.5$ cm) with 2 L of ASTM-reconstituted water and exposed to different concentrations of Cu (control, 0.125, and 0.25 mg/L), with three replicates per treatment for 48 and 96 h. Approximately 75% of the test solution was changed after 48 h. Mussels were not fed during the test.

Adult mussel treatment

The experimental procedures followed the ASTM E2455–06R13 guideline for testing chemicals with mussels (ASTM, 2006). Before the experiments, 90 adult mussels were randomly selected and introduced into 18 glass aquaria ($20 \times 20 \times 27.5$ cm) containing 3 L of ASTM-reconstituted water and acclimated for 48 h. No mortality was observed during the acclimation period. Test mussels were not fed for 24 h prior to Cu exposure. Copper sulfate was prepared at concentrations of 0, 18.75, 37.5, 75, 150, and 300 mg/L (nominal concentrations) in ASTM-reconstituted water, and each set-up

consisted of three replicate containers. Photoperiod, duration of toxicity tests, test water changes, pH range, and water temperature were the same as those used in the juvenile toxicity tests. Dead mussels were removed daily whenever detected.

The effects of nonlethal concentrations of Cu on biochemical responses were bioassayed in a similar way. Mussels were exposed to Cu at concentrations of 0 (control), 0.75, 1.5, and 3 mg/L for 96 h. Renewal of test solutions, and water quality and photoperiod conditions were the same as those described for the juvenile mussel toxicity tests.

Water chemical analysis

Water samples (20 mL) were collected from each test concentration after 48 h of exposure. Copper analyses were performed with an inductively-coupled plasma mass spectrometer (ICP-MS, Agilent 7500ce, Agilent Technologies, USA) as described by Chen *et al.* (2012). Quality assurance and quality control was performed by spike recoveries; ranged from 99.7% to 101.5%.

The measured Cu, Na, Mg, K, and Ca concentrations were 0.00, 28.16, 13.25, 2.31, and 8.39 mg/L in ASTM-reconstituted waters, respectively. Water Cu concentrations in the toxicity test were 0.00, 0.126, and 0.23 mg/L for juveniles, and 0.00, 0.59, 1.42, and 2.84 mg/L for adults. The observed results will be presented based on the nominal concentration of Cu.

Tissue collection and determination of biochemical parameters

The adductor muscle, foot, gill, mantle, and visceral mass from each individual were obtained and rinsed with cold saline solution for further analysis. Tissues were weighed and homogenized in 9-fold (w/v) physiological saline. After centrifugation at $2500 \times g$ for 10 min at 4°C, the supernatant was collected and SOD and GSH were measured using a microplate reader (Multiscan MK3; Thermo Fisher Scientific, Waltham, MA, USA), and CAT and MDA were measured using a spectrophotometer (722, Shanghai Jingke Instrument Plant, China) with commercially-available kits (SOD, A001-3; CAT, A007-1; GSH, A006-2; MDA, A003-1; Nanjing Jiancheng Institute of Biotechnology, Nanjing, China), according to the manufacturer's instructions. Briefly, SOD activity was assayed at 450 nm using xanthine oxidase as a superoxide generator and expressed as U/mg protein (Moustafa *et al.*, 2010). CAT activity was measured following the method of Shimizu (Shimizu *et al.*, 1984), and changes in absorbance were recorded at 405 nm. Results were expressed as U/mg protein. The GSH content was estimated using the method described by Jollow *et al.* (1974). The developed yellow color was read immediately at 405 nm and the content was expressed as $\mu\text{mol/mg}$ protein. The amount of malonaldehyde was assayed in the form of thiobarbituric acid reacting substances (TBARS) using the method described by Esterbauer and Cheesman (1990), and the MDA formed in each sample was measured at 532 nm. The final concentration of MDA was expressed as nmol/mg protein.

Statistical analysis

The Cu EC_{50} at 96 h was calculated by probit analysis based on the nominal concentrations. EC_{50} was presented with 95% confidence intervals (CI) as EC_{50} (95% CI). The no-observable effect concentration (NOEC) and the lowest-observable effect concentration (LOEC) were determined using the method described by Wright and Welbourn (2002). The maximum allowable toxicant concentration (MATC) was calculated by taking the geometric means of the NOEC and LOEC (Wright and Welbourn, 2002).

The activities of Cu/Zn-SOD and CAT and the levels of GSH and MDA in treated swan mussel were compared with those of the control group at different sampling times (48 and 96 h) and data were expressed as means $\pm$ standard deviation ($n = 3$). Dunnett's test of one-way analysis of variance (ANOVA) was used to compare differences between the control and treatment groups. Linear regression analysis was also used to describe the relationship between biochemical parameters and Cu concentrations. All statistical analyses were performed using SPSS Statistics v.16.0 software (IBM Corp., Armonk, NY, U.S.A.) and only $P < 0.05$ was accepted as significant.

RESULTS

Copper exposure test

The results of acute Cu^{2+} toxicity tests are listed in Tables 1 and 2. The 96 h acute Cu EC_{50} was 3.4 mg/L for juvenile *A. woodiana* and 22.1 mg/L for adult *A. woodi-*

Table 1. Number of juvenile *Anodonta woodiana* deaths (mean $\pm$ standard error [SE]) following exposure to Cu for different durations (mean $\pm$ SE), and 48 and 96 h EC_{50} values of Cu for juvenile *A. woodiana* (mg/L)

Cu concentration	48 h	96 h
0	0	0
0.625	0.25 ± 0.50	1.00 ± 0.82
1.25	0.50 ± 0.58	1.25 ± 0.96
2.5	1.25 ± 0.96	2.25 ± 0.96
5	1.25 ± 0.96	2.75 ± 0.50
10	1.75 ± 0.96	3.75 ± 0.96

Table 2. Number of adult *Anodonta woodiana* deaths (mean $\pm$ SE) following exposure to Cu for different durations (mean $\pm$ SE), and 48- and 96- h EC_{50} values of Cu for adult *A. woodiana* (mg/L)

Cu concentration	48 h	96 h
0	0	0
18.75	1.33 ± 0.58	2.00 ± 0.00
37.5	1.67 ± 0.58	4.00 ± 0.00
75	4.67 ± 0.58	5.00 ± 0.00
150	5.00 ± 0.00	5.00 ± 0.00
300	5.00 ± 0.00	5.00 ± 0.00

ana. The 96 h LOEC and MATC were 0.625 and 0.25 mg/L for juveniles, and 18.75 and 5.63 mg/L for adults, respectively. Based on these data, the Cu gradient of sublethal exposure was selected as 0.125 and 0.25 mg/L for juvenile *A. woodiana*, and 0.75, 1.5, and 3 mg/L for adult *A. woodiana*. No mortality was observed in mussels exposed to a Cu gradient during the experiment.

Biochemical responses

Different response patterns for SOD and CAT activity, and GSH levels were observed in the adductor muscle (Fig. 1A, B, C), foot (Fig. 1D, E, F), gill (Fig. 1G, H, I), mantle (Fig. 1J, K, L), and visceral mass (Fig. 1M, N, O) of juvenile mussels after 48 and 96 h of Cu exposure.

After 96 h, SOD activity was decreased in each tissue, CAT activity was increased only in the mantle, and GSH levels were increased in the foot and mantle. Furthermore, the mantle exhibited significantly higher CAT activity (569% of control, $P = 0.005$) and GSH level (3081.3% of control, $P = 0.005$) compared with the control after 96 h of exposure to 0.25 mg/L Cu.

Additionally, adverse changes were found in antioxidant enzyme activities and GSH levels in the foot between mussels exposed to Cu for 48 and 96 h. SOD and CAT

activities, and GSH levels were maximum (SOD_{48h}: 3275.8 ± 636.1 U/mg protein; CAT_{48h}: 537.2 ± 39.8 U/mg protein; GSH_{48h}: 556.2 ± 156.4 μ mol/L protein) after 48 hours of exposure to 0.25 mg/L Cu, followed by a rapid decrease (SOD_{96h}: 79.9 ± 20.6 U/mg protein; CAT_{96h}: 15.5 ± 3.7 U/mg protein; GSH_{96h}: 36.5 ± 13.3 μ mol/L protein) after 96 h of exposure.

Conversely, moderate variations in SOD and CAT activity, and GSH and MDA contents were observed in adult mussels collected from the groups exposed to 0.75 and 1.5 mg/L Cu. Although no significant changes in biochemical parameters were found in adult mussel tissue after 96 h of exposure to 3 mg Cu/L, considerable augmentation of SOD and CAT activity, and GSH levels (SOD_{48h}: 297.9% of control, $P = 0.012$, Fig. 2A; CAT_{48h}: 241.3% of control, $P = 0.034$, Fig. 2B; GSH_{48h}: 660.1% of control, $P = 0.005$, Fig. 2C) was recorded at 48 h in the adult adductor muscle compared with the control.

Overall, maximum variation in antioxidant enzyme activity and GSH levels was observed in juvenile mussel foot. CAT activity and GSH levels were significantly enhanced in the mantle of juvenile mussels in the group exposed to 0.25 mg Cu/L.

Fig. 1. Superoxide dismutase (SOD) and catalase (CAT) activities, and glutathione (GSH) levels in the tissues of juvenile *Anodonta woodiana* exposed to nonlethal concentrations of Cu (mg/L). ** Indicates a significant difference between the experimental and control groups according to Dunnett's *t*-tests, $P < 0.01$; * indicates a significant difference between the experimental and control groups according to Dunnett's *t*-tests, $P < 0.05$. Adductor muscle: Figs. 1A–C; foot: Figs. 1D–F; gill: Figs. 1G–I; mantle: Figs. 1J–L; visceral mass: Figs. 1M–O.

Fig. 2. Superoxide dismutase (SOD) and catalase (CAT) activities, and glutathione (GSH) and malondialdehyde (MDA) levels in the tissues of adult *Anodonta woodiana* exposed to nonlethal concentrations of Cu (mg/L). ** Indicates a significant difference between the experimental and control groups according to Dunnett's *t*-tests, $P < 0.01$; * indicates a significant difference between the experimental and control groups according to Dunnett's *t*-tests, $P < 0.05$. Adductor muscle: Figs. 2A–D; foot: Figs. 2E–H; gill: Figs. 2I–L; mantle: Figs. 2M–P; visceral mass: Figs. 2Q–T.

DISCUSSION

The results of the toxicity test demonstrated that the EC_{50} following 96 h of Cu exposure was much lower for the juvenile *A. woodiana* (3.4 mg/L) than for the adults (22.1 mg/L).

Acute toxicity studies have demonstrated that chemical toxicity can vary by an order of magnitude among life stages and species (Cherry *et al.*, 2002; Augspurger *et al.*, 2003). The early life stages of freshwater mussels are more vulnerable to some pollutants (e.g., Cu) (Keller and Zam *et al.*, 1991; Wang *et al.*, 2011) than adult mussels, because adult mussels can close their valves for a long time to reduce harmful exposure (Harrison *et al.*, 1984). Conversely, the EC_{50} for the adults in the present study was substantially higher than that for other mussel species, such as *Lamellidens corrianus* Lea, 1834, *Mytilus coruscus* Gould, 1861, and *Mytilus galloprovincialis* Lamarck, 1819 (Bat *et al.*, 2013; Harrison *et al.*, 1984; Patil and Mahajan, 2012). Notably, the 96 h EC_{50} value for Cu in the adult marine gastropod, *Onchidium struma* was 74.8 mg/L (Li *et al.*, 2009), which was comparable to that reported here for adult *A. woodiana*. Species-specific differences in tolerance to pollutants may contribute to their diversity under different environmental conditions (e.g., temperature, test medium)

(Harrison *et al.*, 1984), the type of exposure in the field or laboratory (Goswami *et al.*, 2014), and the metabolic capacity for pollutants (Salánki and V-Balogh, 1989). Our observations indicate that the response of swan mussels to Cu is dependent on life stage, with the adult *A. woodiana* being insensitive to Cu compared with adults of other marine bivalves.

Antioxidant defense systems and LPO have been used to monitor oxidative damage induced by heavy metals in mollusk (Company *et al.*, 2006; Farris *et al.*, 1994). However, knowledge of such damage in freshwater mussels remains scarce. The present study is the first to report data on the antioxidant enzymes SOD and CAT, the free radical scavenger GSH, and the peroxidation of the cell constituent MDA in cultured juvenile and adult *A. woodiana*. Exposure of bivalves to Cu resulted in increased CAT and/or SOD activity accompanied by increased GSH content in the foot, mantle of juveniles, or GSH content and MDA product in the adductor muscle of adults. Variations in the antioxidant defense systems illustrate a hormesis effect in the aforementioned tissues.

According to the hormesis theory, there is a non-monotonic response of biological systems to chemicals and toxins exposures (Radak *et al.*, 2017). At low-concentrations (0.125 mg/L for juveniles; 0.75 and 1.5 mg/L for adults, respectively), relatively low levels of ROS may

be rapidly eliminated by an array of anti-oxidant enzymes in order to alleviate oxidative stress and maintain antioxidant enzyme activity at near-normal levels. At a higher concentration (0.25 mg/L for juvenile; 3 mg/L for adult), Cu clearly stimulated the generation of ROS (MDA) in the adult adductor muscle, and significantly increased the activity of antioxidant defense systems (SOD and/or CAT, GSH) in the foot of juveniles at 48 h, in the mantle of juveniles after 96 h, and in the adult adductor muscle throughout the period of exposure. In a comparable study (Eertman *et al.*, 1995), contaminated sediment caused a similar non-dose-dependent induction of SOD and CAT activities in blue mussel *Mytilus edulis*.

Our data suggest that Cu levels in the foot, mantle of juveniles, and the adductor muscle of adults are good indicators of Cu pollution, although enzyme activity levels in the gill have been widely used as biomarkers of metal pollution, including Cu (Rajalakshmi and Mohandas, 2005; Xia *et al.*, 2016). Differential expression of antioxidant enzymes in various tissues of *A. woodiana* is likely associated with the different physiological functions of those tissues. The adductor muscle is the main muscular system in bivalve mollusks and it enables the animal to close its valves tightly when necessary. The foot serves as a fleshy anchor when the animal is stationary or moving forward. The gill and mantle are the main interface between the organism and its environment, and may therefore concentrate contaminants in their tissues (Li *et al.*, 2015). The visceral mass generally holds the bulk of the digestive, reproductive, and excretory systems, including the hepatopancreas, which is involved in digestion, neutralizes large amounts of toxicants that are ingested by the organism (Xia *et al.*, 2016). The lysosomal system in hepatopancreatic cells plays an important role in the detoxification of toxic metals (Rajalakshmi and Mohandas, 2005). Therefore, it is not surprising that the activities of SOD and CAT were markedly low in the visceral mass of juveniles exposed to Cu. Similar variations in antioxidant enzymes have been observed in the hepatopancreas of a freshwater mussel *Lamellidens corrianus* (Lea) exposed to different concentrations of Cu (Rajalakshmi and Mohandas, 2005). More pronounced changes in enzymatic (SOD and/or CAT) activity and GSH levels in juveniles, and in the adult adductor muscle can be attributed to a more sensitive response to Cu during the early-life stage, or to increased levels of oxidative stress in the adult muscular system in response to the metal. Conversely, the absence of any significant change in the foot, gill, mantle, visceral mass of adult *A. woodiana* may be due to valve closure elicited by the high ambient metal concentration, or to metallothionein (MT) synthesis and thus the sequestration of accumulated metal, maintaining the near-normal activity of the antioxidant defense system (Rajalakshmi and Mohandas, 2005).

The response of antioxidant defense systems (SOD and CAT) to Cu might result in the formation of complexes (especially MDA) (Company *et al.*, 2006; Doyotte *et al.*, 1997) or the oxidation of GSH (Livingstone *et al.*,

1992; Regoli, 1998). Differences in response patterns might be related to different pathways of ROS production in tissues (Company *et al.*, 2006), the physiologic function of tissues, and their susceptibility to oxidative stress or properties of the enzymes present in the analyzed tissues (Goswami *et al.*, 2014). Our results demonstrated that antioxidant responses of *A. woodiana* to Cu stress varied amongst tissues and life stages. The foot, mantle of juveniles, and the adductor muscle of adults might be more susceptible to Cu toxicity because marked fluctuations in biochemical indices and strong antioxidant responses were observed in those tissues.

Contaminants can influence unionid bivalves on many different levels (Farris and van Hassel, 2007). An ideal approach would include the use of multiple biomarkers to assess the condition of unionids. A number of related studies have demonstrated the combined effects of environmental stressors on the immune response of mussels (Al-Subiai *et al.*, 2011; Banni *et al.*, 2010; Goswami *et al.*, 2014; Netpae *et al.*, 2012). However, we detected that Cu exerted its effects on a cellular level. To better overcome this shortcoming, other biomarkers, such as molecular, cytological, physiological, and autecological markers, should be taken into consideration, and investigated in further studies (Bourgeault *et al.*, 2010; Farris and van Hassel, 2007). In addition to metal, other biotic and abiotic factors, including sex, salinity, temperature, and season of sampling, can contribute to variations in the antioxidant defense system in mussels (Baraj, 2011). The benefits of using a multi-assay approach to monitor environmental pollution using "standardized" *A. woodiana* are highlighted.

AUTHOR CONTRIBUTIONS

H. Liu designed the study, performed the toxicity experiment, analyzed the data and wrote the paper. X. Chen assisted the toxicity experiment, analyzed the data and wrote the paper. Y. Oshima designed the study, supervised the work and wrote the paper. Y. Shimasaki supervised the work. T. Jiang assisted the water chemical analysis. J. Yang designed the study, supervised the work, analyzed the data and wrote the paper. All authors assisted in editing of the manuscript and approved the final version.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (Grant Nos. 31072214, 31502166) and JSPS RONPAKU (CSC11315) Program. The authors would like to thank Mr. Davison D. Khumbanyiwa for critically reading and providing comments that improved the manuscript.

REFERENCES

- Al-Subiai, S. N., A. J. Moody, S. A. Mustafa and A. N. Jha 2011 A multiple biomarker approach to investigate the effects of copper on the marine bivalve mollusc, *Mytilus edulis*. *Ecotoxicol.*

- Environ. Safety*, **74**: 1913–1920
- American Standard Testing Methods 2013 Standard guide for conducting laboratory toxicity tests with freshwater mussels (E2455–06R13), ASTM International, West Conshohocken, PA.
- American Public Health Association (APHA) 1999 American Water Works Association, Water Environment Federation, Standard methods for the examination of water and wastewater. American Public Health Association, Washington, DC.
- Amiard-Triquet, C., B. Berthet, C. Metayer and J. C. Amiard 1986 Contribution to the ecotoxicological study of cadmium, copper and zinc in the mussel *Mytilus edulis*. *Mar. Biol.*, **92**: 7–13
- Augsburger, T., A. E. Keller, M. C. Black, W. G. Cope and F. J. Dwyer 2003 Water quality guidance for protection of freshwater mussels (Unionidae) from ammonia exposure. *Environ. Toxicol. Chem.*, **22**: 2569–2575
- Banni, M., A. Negri, A. Dagnino, J. Jebali, S. Ameer and H. Boussetta 2010 Acute effects of benzo[a]pyrene on digestive gland enzymatic biomarkers and DNA damage on mussel *Mytilus galloprovincialis*. *Ecotoxicol. Environ. Saf.*, **73**: 842–848
- Baraj, B., F. Niencheski, G. Fillmann and C. D. M. G. Martins 2011 Assessing the effects of Cu, Cd, and exposure period on metallothionein production in gills of the Brazilian brown mussel *Perna perna* by using factorial design. *Environ. Monit. Assess.*, **179**: 155–162
- Bat, L., F. Üstün, O. G. Baki and F. Şahin 2013 Effects of some heavy metals on the sizes of the Mediterranean mussel *Mytilus galloprovincialis* Lamarck, 1819. *Fresenius Environ. Bull.*, **22**: 1933–1938
- Bian, X. S., H. B. Liu, J. L. Gan, R. Li and J. Yang 2009 HCH and DDT residues in bivalves *Anodonta woodiana* from the Taihu lake, China. *Arch. Environ. Contam. Toxicol.*, **56**: 67–76
- Bourgeault, A., C. Gourlay-Francé, F. Vincent-Hubert, F. Palais, A. Geffard, S. Biagianti-Risbourg, S. Pain-Devin and M. H. Tusseau-Vuillemin 2010 Lessons from a transplantation of zebra mussels into a small urban river: an integrated ecotoxicological assessment. *Environ. Toxicol.*, **25**: 468–478
- Chen, J. and P. Xie 2005 Seasonal dynamics of the hepatotoxic microcystins in various organs of four freshwater bivalves from the large eutrophic lake Taihu of subtropical China and the risk to human consumption. *Environ. Toxicol.*, **20**: 572–584
- Chen, X. B., J. Yang, H. B. Liu, Y. P. Su, L. Sun and Y. Oshima 2012 Element concentrations in a unionid mussel *Anodonta woodiana* at different stages. *J. Fac. Agric. Kyushu Univ.*, **57**: 139–144
- Chen, X. B., H. B. Liu, Y. P. Su and J. Yang 2015 Morphological development and growth of the freshwater mussel *Anodonta woodiana* from early juvenile to adult. *Invert. Reproduct. Develop.*, **59**: 131–140
- Cherry, D. S., J. H. Van Hassel, J. L. Farris, D. J. Soucek and R. J. Neves 2002 Site-specific derivation of the acute copper criteria for the Clinch River, Virginia. *Human Ecol. Risk Assessm.*, **8**: 591–601
- Company, R., A. Serafim, R. Cosson, A. Fiala-Médioni, D. Dixon and M. J. Bebianno 2006 Temporal variation in the antioxidant defence system and lipid peroxidation in the gills and mantle of hydrothermal vent mussel *Bathymodiolus azoricus*. *Deep-Sea Res. Part I – Oceanogr. Res. Papers*, **53**: 1101–1116
- Cossu, C., A. Doyotte, M. C. Jacquin, M. Babut, A. Exinger and P. Vasseur 1997 Glutathione reductase, selenium-dependent glutathione peroxidase, glutathione levels, and lipid peroxidation in freshwater bivalves, *Unio tumidus*, as biomarkers of aquatic contamination in field studies. *Ecotoxicol. Environ. Safety*, **38**: 122–131
- Cossu, C., A. Doyotte, M. Babut, A. Exinger and P. Vasseur 2000 Antioxidant biomarkers in freshwater bivalves, *Unio tumidus*, in response to different contamination profiles of aquatic sediments. *Ecotoxicol. Environ. Safety*, **45**: 106–121
- Donnachie, R. L., A. C. Johnson, C. Moeckel, M. G. Pereira and J. P. Sumpter 2014 Using risk-ranking of metals to identify which poses the greatest threat to freshwater organisms in the UK. *Environ. Pollut.*, **194**: 17–23
- Douda, K., M. Vrtilek, O. Slavík and M. Reichard 2012 The role of host specificity in explaining the invasion success of the freshwater mussel *Anodonta woodiana* in Europe. *Biol. Invas.*, **14**: 127–137
- Doyotte, A., C. Cossu, M. C. Jacquin, M. Babut and P. Vasseur 1997 Antioxidant enzymes, glutathione and lipid peroxidation as relevant biomarkers of experimental or field exposure in the gills and the digestive gland of the freshwater bivalve *Unio tumidus*. *Aquat. Toxicol.*, **39**: 93–110
- Eertman, R. H. M., C. L. F. M. G. Groenink, B. Sandee and H. Hummel 1995 Response of the blue mussel *Mytilus edulis* L. following exposure to PAHs or contaminated sediment. *Mar. Environ. Res.*, **39**: 169–173
- Esterbauer, H. and K. H. Cheesman 1990 Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymol.*, **186**: 407–421
- Farris, J. L., J. L. Grudzien, S. E. Belanger, D. S. Cherry and J. Cairns 1994 Molluscan cellulolytic activity responses to zinc exposure in laboratory and field stream comparisons. *Hydrobiologia*, **287**: 161–178
- Farris, J. L. and J. H. van Hassel 2007 Freshwater Bivalve Ecotoxicology, CRC Press, Boca Raton, FL and SETAC Press, Pensacola
- Goswami, P., G. Hariharan, N. Godhantaraman and N. Munuswamy 2014 An integrated use of multiple biomarkers to investigate the individual and combined effect of copper and cadmium on the marine green mussel (*Perna viridis*). *J. Environ. Sci. Health Part A Tox. Hazard Subst. Environ. Eng.*, **49**: 1564–1577
- Guarneri, I., O. P. Popa, L. Gola, L. Kamburska, R. Lauceri, M. Lopes-Lima, L. O. Popa and N. Riccardi 2014 A morphometric and genetic comparison of *Sinanodonta woodiana* (Lea, 1834) populations: does shape really matter? *Aquat. Invas.*, **9**: 183–194
- Harrison, F. L., J. P. Knezovich and D. W. R. Jr 1984 The toxicity of copper to the adult and early life stages of the freshwater clam, *Corbicula manilensis*. *Arch. Environ. Contam. Toxicol.*, **13**: 85–92
- Jollow, D. J., J. R. Mitchell, N. Zampaglione and J. R. Gillete 1974 Bromobenzene induced liver necrosis. Protective role of glutathione and evidence for 3,4-bromobenzene oxide as a hepatotoxic metabolite. *Pharmacol.*, **11**: 151–169
- Keller, A. E. and S. G. Zam 1991 The acute toxicity of selected metals to the freshwater mussel, *Anodonta imbecillis*. *Environ. Toxicol. Chem.*, **10**: 539–546
- Li, N., Y. L. Zhao and J. Yang 2007 Impact of waterborne copper on the structure of gills and hepatopancreas and its impact on the content of metallothionein in juvenile giant freshwater prawn *Macrobrachium rosenbergii* (Crustacea: Decapoda). *Arch. Environ. Contam. Toxicol.*, **52**: 73–79
- Li, X. B., X. L. Hou, Q. Mao, Y. L. Zhao, Y. X. Cheng and Q. Wang 2009 Toxic effects of copper on antioxidative and metabolic enzymes of the marine gastropod, *Onchidium struma*. *Arch. Environ. Contam. Toxicol.*, **56**: 776–784
- Li, Y. Q., H. Z. Yang, N. Liu, J. X. Luo, Q. Wang and L. Wang 2015 Cadmium Accumulation and Metallothionein Biosynthesis in Cadmium-Treated Freshwater Mussel *Anodonta woodiana*. *PLoS One*, **10**: e0117037
- Liu, H. B., J. Yang and J. L. Gan 2010 Trace element accumulation in bivalve mussels *Anodonta woodiana* from Taihu Lake, China. *Arch. Environ. Contam. Toxicol.*, **59**: 593–601
- Livingstone, D. R., F. Lips, P. G. Martinez and R. K. Pipe 1992 Antioxidant enzymes in the digestive gland of the common mussel *Mytilus edulis*. *Mar. Biol.*, **112**: 265–276
- Lorenzo, J. I., R. Beiras, V. K. Mubiana and R. Blust 2005 Copper uptake by *Mytilus edulis* in the presence of humic acids. *Environ. Toxicol. Chem.*, **24**: 973–980
- Moustafa, D. A., N. Jain, N. Sriranganathan and R. Vemulapalli 2010 Identification of a single-nucleotide insertion in the promoter region affecting the *sodC* promoter activity in *Brucella neotomae*. *PLoS One*, **5**: e14112
- Netpae, T., C. Phalaraksh and W. Wongkham 2012 Antioxidant Enzyme Activities and DNA Damage as Biomarker of Copper

- Effect on *Corbicula fluminea*. *eJBio*, **8**: 19–23
- Patil, P. B. and A. Y. Mahajan 2012 Evaluation of acute toxicity of heavy metal copper chloride and copper sulphate to a freshwater Bivalve *lamellidens corrianus* (lea). *Int. J. Innova Biosciences*, **2**: 85–88
- Que X., G. F. Widhopf II, S. Amir, K. Hartvigsen, L. F. Hansen, D. Woelkers, S. Tsimikas, C. J. Binder, T. J. Kipps and J. L. Witztum 2013 IGHV1–69–encoded antibodies expressed in chronic lymphocytic leukemia react with malondialdehyde–acetaldehyde adduct, an immunodominant oxidation–specific epitope. *PLoS One*, **8**: e65203
- Radak, Z., K. Ishihara, E. Tekus, C. Varga, A. Posa, L. Balogh, I. Boldogh and E. Koltai 2017 Exercise, oxidants, and antioxidants change the shape of the bell-shaped hormesis curve. *Redox Biology*, **12**: 285–290
- Rajalakshmi, S. and A. Mohandas 2005 Copper-induced changes in tissue enzyme activity in a freshwater mussel. *Ecotoxicol. Environ. Saf.*, **62**: 140
- Regoli F. 1998 Trace metals and antioxidant enzymes in gills and digestive gland of the Mediterranean mussel *Mytilus galloprovincialis*. *Arch. Environ. Contam. Toxicol.*, **34**: 48–63
- Rivas, B. L., S. Villegas, B. Ruf and I. Peric 2007 Removal of metal ions with impact on the environment by water–insoluble functional copolymers: synthesis and metal–ion uptake properties. *J. Chil. Chem. Soc.*, **52**: 1164–1168
- Salánki, J. and K. V–Balogh 1989 Physiological background for using freshwater mussels in monitoring copper and lead pollution. *Hydrobiologia*, **188/189**: 445–454
- Shimizu, N., K. Kobayashi and K. Hayashi 1984 The reaction of superoxide radical with catalase. Mechanism of the inhibition of catalase by superoxide radical. *J. Biol. Chem.*, **259**: 4414–4418
- Szalecrazy, E., J. Prechl, J. Fehér and A. Somogyi 1999 Alterations in enzymatic antioxidant defence in diabetes mellitus – a rational approach. *Postgrad. Med. J.*, **75**: 13–17
- Wang, N., C. A. Mebane, J. L. Kunz, C. G. Ingersoll, T. W. May, W. R. Arnold, R. C. Santore, T. Augspurger, F. J. Dwyer and M. C. Barnhart 2009 Evaluation of acute copper toxicity to juvenile freshwater mussels (fatmucket, *Lampsilis siliquoidea*) in natural and reconstituted waters. *Environ. Toxicol. Chem.*, **28**: 2367–2377
- Wang, N., C. A. Mebane, J. L. Kunz, C. G. Ingersoll, W. G. Brumbaugh, R. C. Santore, J. W. Gorsuch and W. R. Arnold 2011 Influence of dissolved organic carbon on toxicity of copper to an unionid mussel (*Villosa iris*) and a cladoceran (*Ceriodaphnia dubia*) in acute and chronic water exposures. *Environ. Toxicol. Chem.*, **30**: 2115–2125
- Wright, D. A. and P. Welbourn 2002 Environmental Toxicology. Cambridge University Press, Cambridge, pp. 630
- Xia, X. C., C. X. Hua, S. P. Xue, B. Q. Shi, G. X. Gui, D. X. Zhang, X. Y. Wang and L. H. Guo 2016 Response of selenium–dependent glutathione peroxidase in the freshwater bivalve *Anodonta woodiana* exposed to 2,4–dichlorophenol, 2,4,6–trichlorophenol and pentachlorophenol. *Fish Shellfish Immunol.*, **55**: 499–509
- Yang, J., H. Harino, H. B. Liu and N. Miyazaki 2008 Monitoring the organotin contamination in the Taihu Lake of China by bivalve mussel *Anodonta woodiana*. *Bull. Environ. Contam. Toxicol.*, **81**: 164–168
- Yokoyama, A. and H. D. Park 2002 Mechanism and prediction for contamination of freshwater bivalves (Unionidae) with the cyanobacterial toxin microcystin in hypereutrophic Lake Suwa, Japan. *Environ. Toxicol.*, **17**: 424–433