A spirooxazine-based chemical sensor for ferric-ion detection with the application in drinking water

Pattaweepaiboon, Supak

Department of Chemistry and Centre of Excellence for Innovation in Chemistry, Faculty of Science, Kasetsart University

Phiromphu, Natchayapak Department of Chemistry and Centre of Excellence for Innovation in Chemistry, Faculty of Science, Kasetsart University

Kaewchangwat, Narongpol Functional Nanomaterials and Interfaces Laboratory, National Nanotechnology Center (NANOTEC), National Science and Technology Development Agency (NSTDA)

Suttisintong, Khomson

Functional Nanomaterials and Interfaces Laboratory, National Nanotechnology Center (NANOTEC), National Science and Technology Development Agency (NSTDA)

他

https://doi.org/10.15017/1960665

出版情報: Proceedings of International Exchange and Innovation Conference on Engineering & Sciences (IEICES). 4, pp.50-53, 2018-10-18. 九州大学大学院総合理工学府 バージョン: 権利関係: Supak Pattaweepaiboon¹, Natchayapak Phiromphu¹, Narongpol Kaewchangwat², Khomson Suttisintong², Weekit Sirisaksoontorn^{1,*}

¹Department of Chemistry and Centre of Excellence for Innovation in Chemistry, Faculty of Science, Kasetsart University, Chatuchak, Bangkok, 10900 Thailand

²Functional Nanomaterials and Interfaces Laboratory, National Nanotechnology Center (NANOTEC), National Science and Technology Development Agency (NSTDA), Pathum Thani, 12120 Thailand.

Corresponding author email: fsciwks@ku.ac.th

Abstract: A solution of the spirooxazine $(1.0 \times 10^{-4} \text{ M})$ was freshly prepared in 1,4-dioxane with the subsequent addition of one equivalent solution of each relevant cation such as Na⁺, K⁺, Mg²⁺, Ca²⁺, Sr²⁺, Ba²⁺, Sn²⁺, Pb²⁺, Mn²⁺, Fe²⁺, Co²⁺, Ni²⁺, Cu²⁺, Zn²⁺, Cd²⁺, Hg²⁺, Cr³⁺ and Fe³⁺ (1.0 $\times 10^{-4} \text{ M})$. The metal-ligand complexes were mainly investigated by UV-Vis spectroscopy. In the case of ferric ions, spirooxazine provides the new absorption response at 654 nm, indicating the complex formation of Fe³⁺ with spirooxazine in the open form. Meanwhile, other metal ions did not strongly bind to spirooxazine, giving rise to no absorption peak at 654 nm observed. The stoichiometry of complex was investigated via Job's method as the 1:1 ratio of spirooxazine with Fe³⁺. The detection limit of ferric ions is 2.61 μ M. Moreover, the spirooxazine-based molecular sensor was further developed for determination of spiked ferric ions in drinking water.

Keywords: spirooxazine; chemical sensor; ferric ion

1. INTRODUCTION

Nowadays, one of the most important problems facing many countries is the environmental pollution, especially contaminated toxic metals. Several activities, including geogenic, industrial, agricultural human activities, can cause the release of those toxic metals into the environmental water resource [1,2]. For the positive perspective, some relevant metals, for example Co, Cu, Cr, Fe, Mg, Mn, Mo, Ni, as well as Zn play an important role as the micronutrient for many biological systems [3]. Among other metals, iron has considered to be part of many biological mechanisms, namely oxygen metabolism, catalysis and electron transfer [4]. Importantly, the ferrous $(Fe^{2+})/ferric$ (Fe^{3+}) electron transfer in the human body involves the production of the reactive oxygen-based compounds which can cause the damage to targeted cells and tissues [5]. And also, the excess accumulation of ferric, especially, can lead to the neurodegenerative diseases; i.e. Alzheimer and Parkinson [6-8]. As a result, several techniques have been developed for the need of more effective and selective methods for the Fe^{3+} detection [9].

Spirooxazine is a photochromic molecule that can reversibly isomerize from a closed-ring form to an open-ring form upon the light activation. It has been known that the isomerization occurs through the spiro C-O bond breaking, giving rise to the formation of a zwitterionic open form of spirooxazine, also called merocyanine [10-12]. Interestingly, when a colorless closed form of spirooxazine is activated by light, the solution turns to be colored with the presence of an open-form merocyanine. Besides light activation, the effect of metal coordination on the specific binding site also manages to facilitate the ring-opening process of spiroozaxine in a proper solvent system [13-14]. This phenomena is defined as ionochromism. From the previous reports, a wide variety of metal ions such as Zn^{2+} , Co^{2+} , Cd^{2+} , Mg^{2+} , Ag^+ , Pb^{2+} , Cu^{2+} , Tb^{3+} and Hg^{2+} , can selectively form chelating complexes with an openform merocyanine [14-17]. As a consequence, assynthesized spriooxazine derivatives have been employed as a potential chemical sensor to detect the specific metal ion.

In this research, we aim to synthesize a spirooxazinebased compound for being used as a chemical sensor to detect ferric ions. The selective and optical behaviors of as-synthesized spirooxazine were investigated by the color change of complexation. The selective interaction of spirooxazine molecule with ferric ion was also studied with the structural isomerization from a closed-form spirooxazine to an open-form merocyanine. The resulting colored complex originated from the formation of a merocyanine-Fe³⁺ complex. For the application, the assynthesized spirooxazine was further used to monitor the amount of spiked ferric ion in drinking water in comparison with the standard spectroscopic technique.

2. EXPERIMENT

2.1 Chemicals and instrumentation

All chemicals and reagents are used as received without any purification. All inorganic salts were obtained as chloride salts. Drinking water was obtained from a water dispenser located in the Kasetsart University campus, Bangkok, Thailand. The ¹H and ¹³C NMR spectra were recorded on a Bruker Avance 500 for ¹H (500 MHz) and ¹³C (125 MHz) in CDCl₃ or (CD₃)₂SO solution. Mass spectrometry (MS) was performed on a Micromass Quattro micro TM API. UV-Vis spectra were recorded on a UV-visible spectrophotometer (Shimadzu, UV-1800). The amount of ferric ions was also determined by an atomic absorption spectrometer (Perkin-Elmer analyst 800).

2.2 Methods

The selectivity and sensitivity of a spirooxazine sensor for binding with metal ions were investigated by UV-Vis spectroscopy. Firstly, a spirooxazine solution (1.0 x 10⁻⁴ M) was prepared in 1,4-dioxane at room temperature. The mixtures of spirooxazine with each metal ion (1.0 x 10⁻⁴ M) as follows: Na⁺, K⁺, Mg²⁺, Ca²⁺, Sr^{2+} , Ba^{2+} , Sn^{2+} , Pb^{2+} , Mn^{2+} , Fe^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} , Zn^{2+} , Cd^{2+} , Hg^{2+} , Cr^{3+} , and Fe^{3+} were freshly prepared. Job's plot was demonstrated by using the equimolar solutions of spirooxazine and metal ions with the variation of volumetric ratios between Fe3+ ions and spirooxazine solution. Regarding the sensitivity study of Fe³⁺ ions, the various concentrations of metal ions were systematically added to the spirooxazine solution. To investigate the effect of competing ions, the mixture was prepared by adding 5.0 equivalent of Fe³⁺ into spirooxazine solution with the presence of 5.0 equivalent of other competing metal ions.

The as-synthesized spirooxazine was applied for the quantitative analysis of ferric ions in sample obtained from the water dispenser. Initially, drinking water was spiked by the standard solution of Fe³⁺ with 3 and 5 μ M, respectively. Then, the amount of spiked ferric ions in the water sample was monitored by the spirooxazine sensor in comparison with the flame atomic absorption spectroscopy.

3. RESULTS AND DISCUSSION

3.1 Spirooxazine synthesis

A spirooxazine-based derivative was successfully prepared by the synthetic route in Scheme 1. The asobtained spirooxazine was structurally confirmed by ¹H NMR, ¹³C NMR and mass spectrometry. In brief, 12.0 mmol of ZnCl₂ was added to mixed solution of tetrahydrofuran and water (1:1 v/v) containing 28.0 mmol of 4-(3,4-dihydroisoquinoline-2(1H)-yl)-1-nitrosonaphthalen-2-ol (1). The reaction went completion after 30 min at room temperature, the raw product of 4-(3,4-dihydroisoquinoline-2(1H)-yl)-1-nitrosonaphthalen-2-ol zinc salt (2) was filtrated, washed with a copious amount of water, and then dried for 2 h in vacuo. The resulting product is brown solid, which was collected and used for next step without purification. 9.0 mmol of compound (2) was dissolved in ethanol and stirred under argon at refluxing temperature. Then, the solution of 7.6 mmol 3,3-dimethyl-2-methylene-1-phenethyl indoline (3) in ethanol was subsequently added and the mixture was stirred for 8 h. After that, the remaining solvent was removed under the reduced pressure. Finally, the crude product was further purified by column chromatography with 10% ethyl acetate in hexane to obtain black solid of the resulting spirooxazine (4) with 37% yield.



Scheme 1. Synthesis of a spirooxazine-based molecular sensor.

3.2 Optical behavior of as-synthesized spirooxazine

With the addition of metal ions into the spirooxazine solution in 1,4-dioxane, the result found that the color of spirooxazine solution was changed from colorless to blue with the presence of ferric ion (Fig. 1). No color change was observed by the naked eye in the case of other relevant metal ions in the spirooxazine solution, indicating no complex formation.

In Fig. 2, the absorption spectra of the solution containing spirooxazine and metal ions show similar absorption peak at 300-400 nm due to a closed form spirooxazine. Since ferric ion was added, the appearance of the absorption response of complexes exhibits a new band at 654 nm. This new absorption peak indicates the formation of merocyanine-Fe³⁺ complex. In the case of other metals, they could not induce the ring-opening process of spirooxazine. Hence, no new absorption peak was observed.



Fig. 1. Color change on the presence of one equivalent of metal ions in spirooxazine solution in 1,4-dioxane.



Fig. 2. Absorption spectra of spirooxazine $(1 \times 10^{-4} \text{ M})$ with the presence of metal ions $(1 \times 10^{-4} \text{ M})$ in 1,4-dioxane.

The Job's plot of spirooxazine and Fe^{3+} suggests that the absorbance at 654 nm reached the maximum at 0.5 mole fraction (Fig. 3). It confirms that the stoichiometric ratio of spirooxazine and ferric ion was found to be 1:1 for the complexation.



Fig. 3. Job's plot of spirooxazine and ferric ion in 1,4-dioxane.

In terms of complex formation, spirooxazine forms a complex by breaking the spiro C-O bond in the presence of ferric ions. The open-form merocyanine chelates with

ferric ions by using both nitrogen and oxygen atoms for complexation as proposed in Fig. 4.



Fig. 4. The formation of merocyanine-Fe³⁺ complex.

To investigate the binding ability between spirooxazine and ferric ions, the titration method of spirooxazine with various ferric concentrations was carried out. The result suggests that the absorbance intensity has a linear response toward ferric ions with the concentrations ranging from 1 to 30 μ M (Fig. 5). Based on the linear correlation, the detection limit of ferric ion in 1,4-dioxane was determined to be 2.61 μ M, which is lower than 5.36 μ M, the standard maximum permissible level of ferric ion in drinking water approved by Environmental Protection Agency (EPA) [18].



Fig. 5. Absorption titration at 654 nm of a spirooxazine solution (2.0 x 10^{-5} M) with increasing the amount of ferric ions in 1,4-dioxane.

The absorption spectra in Fig. 6 show the effect of increasing concentrations of ferric ions in the spirooxazine solution in 1,4-dioxane. With the ferric concentrations from 0 to 10 equivalents, the absorbance at 654 nm was increased with a slightly bathochromic shift. This can be ascribed to the formation of J-type aggregates of merocyanine with head-to-head arrangement at the high concentration of ferric ions [11].



Fig. 6. Absorption spectra of solution containing spirooxazine $(1 \times 10^{-4} \text{ M})$ with concentrations of ferric ions from 0 to 10 equivalents.

To investigate the competitive selectivity of spirooxazine toward ferric ion compared with other relevant metal ions, we added 5.0 equivalent of ferric ion into spirooxazine (5 x 10^{-5} M) with the subsequent addition of 5.0 equivalent of competing metal ions. This result suggests that the absorbance enhancement caused by the solution of spirooxazine and ferric ion was not significantly affected by those competing metal ions (Fig. 7). The as-synthesized spirooxazine has a highly selective behavior toward ferric ions.



Fig. 7. Competitive selectivity of spirooxazine toward ferric ions with the subsequent addition of competing metal ions in 1,4-dioxane. (SP = spirooxazine)

In this research, we aim to use the sensing spirooxazine molecule to monitor ferric ion in drinking water by the naked-eye detection. Hence, we need to optimize the solvent system and found the suitable condition by mixing water to 1,4-dioxane (6:4 v/v). This condition was further utilized for monitoring the amount of ferric ion in the water sample.

3.3 Application

To test the sensing performance of spirooxazine toward the determination of ferric ions in drinking water, the results from Table 1 demonstrate that the %recoveries for spiked concentration levels of ferric ions in drinking water detected by a spirooxazine sensor were obtained in the range of 93.75–103.75 % which is relatively comparable to the amount of spiked ferric ions monitored by the standard AAS method. This indicates the excellent performance of the as-synthesized spirooxazine as a chemical sensor.

Table 1. Recoveries of ferric ions in drinking water.

Fe ³⁺	Fe ³⁺ determined (µM)			
spiked (µM)	Sensor	%Recovery	AAS	%Recovery
0	0	-	0	-
3	2.81	93.75	3.04	101.40
5	5.19	103.75	5.15	102.99

4. CONCLUSION

In summary, the spirooxazine was successfully synthesized with the structural confirmation. The high selectivity of spirooxazine for binding with ferric ions was clearly observed. The detection limit of ferric ions is 2.61 μ M. We further demonstrate the development of spirooxazine for the sensitive detection of ferric ions in drinking water. The %recoveries of ferric ions in drinking water was obtained in a range of 93.75–103.75 % with a comparable extent of the standard spectroscopic method.

5. ACKNOWLEDGEMENTS

This work was financially supported by the Graduate School, Department of Chemistry, Faculty of Science, Kasetsart University, the Centre of Excellence for Innovation in Chemistry (PERCH-CIC) and the Young Scientist and Technologist Programme (YSTP) of National Science and Technology Development Agency (NSTDA).

6. REFERENCES

- I.M. Ocheri, Environmental health hazards and national survival and stability: a need for education, Benue State University Journal of Education (BSUJE). 4 (2014) 67-174.
- [2] Z.L. He, X.E. Yang, P.J. Stoffella, Trace elements in agroecosystems and impacts on the environment, J. Trace Elem. Med. Biol. 19 (2005) 125–140.
- [3] A.B. Morrison, J.A. Campbell, Trace elements in human nutrition, Canad. Med. Ass. J. 88 (1963) 523-527.
- [4] E.I. Solomon, et al., Geometric and electronic structure/function correlations in nonheme iron enzymes, Chem. Rev. 100 (2000) 235-349.
- [5] B. Silva, P. Faustino, An overview of molecular basis of iron metabolism regulation and the associated pathologies, Biochim. Biophys. Acta. 1852 (2015) 1347-1359.
- [6] E. Madsen, J.D. Gitlin, Copper and iron disorders of the brain, Annu. Rev. Neurosci. 30 (2007) 317-337.
- [7] Z. Tian, et al., Spirooxazine to merooxazine interconversion in the presence and absence of zinc: approach to a bistable photochemical switch, J. Phys. Chem. A. 114 (2010) 11900-11909.
- [8] W.J. Roberta, et al., The role of iron in brain ageing and neurodegenerative disorders, Lancet Neurol. 13 (2014) 1045-1060.
- [9] D.T. Quang, J.S. Kim, Fluoro- and chromogenic chemodosimeters for heavy metal ion detection in solution and biospecimens, Chem. Rev. 110 (2010) 6280-6301.
- [10] H. Bouas-Laurent, H. Durr, Organic photochromism, Pure Appl. Chem. 73 (2001) 639-665.
- [11] V.I. Minkin, Photo-, thermos-, solvato-, and electrochromic spiroheterocyclic compounds, Chem. Rev. 104 (2004) 2751-2776.
- [12] J.C. Crano, R.J. Guglielmetti, Organic photochromic and thermochromic compounds, first ed., Kluwer Academic, New York, 2002.
- [13] K. Nakatani, et al., Photochromic materials: properties and applications, first ed., Wiley-VCH Verlag GmbH & Co. KGaA., Chinese, 2016.
- [14] S.V. Paramonov, V. Lokshin, O.A. Fedorova, Spiropyran, chromene or spirooxazine ligands: Insights into mutual relations between complexing and photochromic properties, J. Photochem.

Photobiol. C: Photochem. Rev. 12 (2011) 209-236.

- [15] H.S. Kumbhar, et al., A highly selective quinaldine–indole based spiropyran with intramolecular H-bonding for visual detection of Cu (II) ions, Sens. Actuator B-Chem. 222 (2016) 35-42.
- [16] J. Ji, et al., Promoted colorimetric response of spirooxazine derivative: A simple assay for sensitive mercury (II) detection. Res. Chem. Intermed. 42 (2016) 5597.
- [17] L. Popov, et al., Studies of complex formation of photochromic hybrid spirooxazine with metal ions. Russ. J. Phys. Chem. A. 83 (2013) 2309.
- [18] United States Environmental Protection Agency, National Primary Drinking Water Regulations, EPA 816-F-09-0004; EPA, Washington, DC, 2009 (http://www.epa. gov/safewater/)