A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space

Kawakami, Yu
Faculty of Science, Yamaguchi University

<http://hdl.handle.net/2324/20095>

出版情報 : Geometriae Dedicata, 2013-08. Springer Netherlands
バージョン :
権利関係 : (C) Springer Science+Business Media Dordrecht 2013

MI Preprint Series

**Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub**

A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space

Yu Kawakami

MI 2011-16

(Received October 14, 2011)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

A RAMIFICATION THEOREM FOR THE RATIO OF CANONICAL FORMS OF FLAT SURFACES IN HYPERBOLIC THREE-SPACE

YU KAWAKAMI

ABSTRACT. We provide an effective ramification theorem for the ratio of canonical forms of a weakly complete flat front in the hyperbolic three-space. Moreover we give the two applications of this theorem, the first one is to show an analogue of the Ahlfors islands theorem for it and the second one is to give a simple proof of the classification of complete nonsingular flat surfaces in the hyperbolic three-space.

INTRODUCTION

It is well-known that any complete nonsingular flat surface in the hyperbolic 3-space $\mathbf{H}^3$ must be a horosphere or a hyperbolic cylinder, that is, a surface equidistance from a geodesic ([20], [21]). However if we consider flat fronts (namely, projections of Legendrian immersions) and define the notion of weakly completeness, there exist many examples and interesting global properties. For instance, more recently, Martin, Umehara and Yamada [14] showed that there exists a weakly complete bounded flat front in $\mathbf{H}^3$.

The ratio ρ of canonical forms plays important roles in investigating the global properties of weakly complete flat fronts in $\mathbf{H}^3$. Indeed, Kokubu, Rossman, Saji, Umehara and Yamada [8] showed that a point p is a singular point of a flat front in $\mathbf{H}^3$ if and only if $|\rho(p)| = 1$. Moreover the author and Nakajo [6] obtained the best possible upper bound for the number of exceptional values of ρ of a weakly complete flat front in $\mathbf{H}^3$.

The purpose of the present paper is to study the value distribution properties of the ratio of canonical forms of weakly complete flat fronts in $\mathbf{H}^3$. The paper is organized as follows: In Section 1, we recall some definitions and fundamental properties of flat fronts in $\mathbf{H}^3$, which are used throughout this paper. In Section 2, we provide a ramification theorem for the ratio of canonical forms of a weakly complete flat front in $\mathbf{H}^3$ (Theorem 2.2). The theorem is effective in the sense that it is sharp (see Corollary 3.4 and the comment below) and has some applications. We note that it corresponds to the defect relation in Nevanlinna theory ([7], [15], [16] and [18]). In Section 3, we give the two applications of this theorem. The first one is to show an analogue of a special case of the

2010 *Mathematics Subject Classification.* Primary 30D35, 53A35; Secondary 53C42.

Date: 14 October, 2011.

Key words and phrases. flat front, weakly complete, ramification theorem, the Ahlfors islands theorem, Bernstein type theorem.

The author is partially supported by the Grant-in-Aid for Young Scientists (B) No. 21740053, The Ministry of Education, Culture, Sports, Science and Technology, Japan.

Ahlfors Islands Theorem [2, Theorem B.2] for the ratio of canonical forms of a weakly complete flat front in $\mathbf{H}^3$ (Corollary 3.3). We remark that Klotz and Sario [11] investigated the number of islands for the Gauss map of minimal surfaces in the Euclidean 3-space $\mathbf{R}^3$. The second one is to give a simple proof of the classification of complete nonsingular flat surfaces in $\mathbf{H}^3$ (Corollary 3.5).

Finally, the author would like to particular thank to Masatoshi Kokubu, Junjiro Noguchi, Yusuke Okuyama, Wayne Rossman, Masaaki Umehara and Kotaro Yamada for their useful advice.

1. PRELIMINARIES

We briefly summarize here definitions and basic facts on flat fronts in $\mathbf{H}^3$ which we shall need. For more details, we refer the reader to [4], [5], [8], [9], [10], [12], [13] and [19].

Let $\mathbf{L}^4$ be the Lorentz-Minkowski 4-space with inner product of signature $(-, +, +, +)$. Then the hyperbolic 3-space is given by

$$(1.1) \quad \mathbf{H}^3 = \{(x_0, x_1, x_2, x_3) \in \mathbf{L}^4 \mid -(x_0)^2 + (x_1)^2 + (x_2)^2 + (x_3)^2 = -1, x_0 > 0\}$$

with the induced metric from $\mathbf{L}^4$, which is a simply connected Riemannian 3-manifold with constant sectional curvature -1 . Identifying $\mathbf{L}^4$ with the set of 2×2 Hermitian matrices $\text{Herm}(2) = \{X^* = X\}$ ($X^* := {}^t\overline{X}$) by

$$(1.2) \quad (x_0, x_1, x_2, x_3) \longleftrightarrow \begin{pmatrix} x_0 + x_3 & x_1 + ix_2 \\ x_1 - ix_2 & x_0 - x_3 \end{pmatrix}$$

where $i = \sqrt{-1}$, we can write

$$(1.3) \quad \begin{aligned} \mathbf{H}^3 &= \{X \in \text{Herm}(2) ; \det X = 1, \text{trace } X > 0\} \\ &= \{aa^* ; a \in SL(2, \mathbf{C})\} \end{aligned}$$

with the metric

$$\langle X, Y \rangle = -\frac{1}{2} \text{trace}(X\tilde{Y}), \quad \langle X, X \rangle = -\det(X),$$

where $\tilde{Y}$ is the cofactor matrix of Y . The complex Lie group $PSL(2, \mathbf{C}) := SL(2, \mathbf{C})/\{\pm \text{id}\}$ acts isometrically on $\mathbf{H}^3$ by

$$(1.4) \quad \mathbf{H}^3 \ni X \longmapsto aXa^*,$$

where $a \in PSL(2, \mathbf{C})$.

Let Σ be an oriented 2-manifold. A smooth map $f: \Sigma \rightarrow \mathbf{H}^3$ is called a *front* if there exists a Legendrian immersion

$$L_f: \Sigma \rightarrow T_1^*\mathbf{H}^3$$

into the unit cotangent bundle of $\mathbf{H}^3$ whose projection is f . Identifying $T_1^*\mathbf{H}^3$ with the unit tangent bundle $T_1\mathbf{H}^3$, we can write $L_f = (f, n)$, where $n(p)$ is a unit vector in $T_{f(p)}\mathbf{H}^3$ such that $\langle df(p), n(p) \rangle = 0$ for each $p \in M$. We call n a *unit normal vector field* of the

front f . A point $p \in \Sigma$ where $\text{rank}(df)_p < 2$ is called a *singularity* or *singular point*. A point which is not singular is called *regular point*, where the first fundamental form is positive definite.

The *parallel front* f_t of a front f at distance t is given by $f_t(p) = \text{Exp}_{f(p)}(tn(p))$, where “Exp” denotes the exponential map of $\mathbf{H}^3$. In the model for $\mathbf{H}^3$ as in (1.1), we can write

$$(1.5) \quad f_t = (\cosh t)f + (\sinh t)n, \quad n_t = (\cosh t)n + (\sinh t)f,$$

where n_t is the unit normal vector field of f_t .

Based on the fact that any parallel surface of a flat surface is also flat at regular points, we define flat fronts as follows: A front $f: \Sigma \rightarrow \mathbf{H}^3$ is said to be *flat* if, for each $p \in M$, there exists a real number $t \in \mathbf{R}$ such that the parallel front f_t is a flat immersion at p . By definition, $\{f_t\}$ forms a family of flat fronts. We note that an equivalent definition of flat fronts is that the Gaussian curvature of f vanishes at all regular points. However, there exists a case where this definition is not suitable. For details, see [13, Remark 2.2].

We assume that f is flat. Then there exists a (unique) complex structure on Σ and a holomorphic Legendrian immersion

$$(1.6) \quad \mathcal{E}_f: \tilde{\Sigma} \rightarrow SL(2, \mathbf{C})$$

such that f and L_f are projections of $\mathcal{E}_f$, where $\tilde{\Sigma}$ is the universal covering surface of Σ . Here $\mathcal{E}_f$ being a holomorphic Legendrian map means that $\mathcal{E}_f^{-1}d\mathcal{E}_f$ is off-diagonal (see [4], [12], [13]). We call $\mathcal{E}_f$ the *holomorphic Legendrian lift* of f . The map f and its unit normal vector field n are

$$(1.7) \quad f = \mathcal{E}_f \mathcal{E}_f^*, \quad n = \mathcal{E}_f e_3 \mathcal{E}_f^*, \quad e_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

If we set

$$(1.8) \quad \mathcal{E}_f^{-1}d\mathcal{E}_f = \begin{pmatrix} 0 & \theta \\ \omega & 0 \end{pmatrix},$$

the first and second fundamental forms $ds^2 = \langle df, df \rangle$ and $dh^2 = -\langle df, dn \rangle$ are given by

$$(1.9) \quad \begin{aligned} ds^2 &= |\omega + \bar{\theta}|^2 = Q + \bar{Q} + (|\omega|^2 + |\theta|^2), \quad Q = \omega\theta \\ dh^2 &= |\theta|^2 - |\omega|^2 \end{aligned}$$

for holomorphic 1-forms ω and θ defined on $\tilde{\Sigma}$, with $|\omega|^2$ and $|\theta|^2$ well-defined on Σ itself. We call ω and θ the *canonical forms* of f . The holomorphic 2-differential Q appearing in the $(2, 0)$ -part of ds^2 is defined on Σ , and is called the *Hopf differential* of f . By definition, the umbilic points of f coincide with the zeros of Q . Defining a meromorphic function on $\tilde{\Sigma}$ by the ratio of canonical forms

$$(1.10) \quad \rho = \frac{\theta}{\omega},$$

then $|\rho|: \Sigma \rightarrow [0, +\infty]$ is well-defined on Σ , and $p \in \Sigma$ is a singular point if and only if $|\rho(p)| = 1$ ([8]).

Note that the $(1, 1)$ -part of the first fundamental form

$$(1.11) \quad ds_{1,1}^2 = |\omega|^2 + |\theta|^2 = (1 + |\rho|^2)|\omega|^2$$

is positive definite on Σ because it is the pull-back of the canonical Hermitian metric of $SL(2, \mathbf{C})$. Moreover $2ds_{1,1}^2$ coincides with the pull-back of the Sasakian metric on $T_1^*\mathbf{H}^3$ by the Legendrian lift L_f of f (which is the sum of the first and third fundamental forms in this case, see [13, Section 2] for details). The complex structure on Σ is compatible with the conformal metric $ds_{1,1}^2$. Note that any flat front is orientable ([9, Theorem B]). In the present paper, for each flat front $f: \Sigma \rightarrow \mathbf{H}^3$, we always regard Σ as a Riemann surface with this complex structure. A flat front $f: \Sigma \rightarrow \mathbf{H}^3$ is said to be *weakly complete* if the metric $ds_{1,1}^2$ as in (1.11) is complete. We note that the universal cover of a weakly complete flat front is also weakly complete.

Finally, we give examples which play important roles in the following sections.

EXAMPLE 1.1 ([13, Example 4.1], flat fronts of revolution). Let $\bar{\Sigma} = \mathbf{C} \cup \{\infty\}$ and set

$$\omega = -\frac{1}{c^2}z^{-2/(1-\alpha)}dz, \quad \theta = \frac{c^2\alpha}{(1-\alpha)^2}z^{2\alpha/(1-\alpha)}dz,$$

for some constants $\alpha \in \mathbf{R} \setminus \{1\}$ and $c \in \mathbf{R}$. We define Σ by $\Sigma = \bar{\Sigma} \setminus \{0\}$ for the case where $\alpha = 0$ and $\Sigma = \bar{\Sigma} \setminus \{0, \infty\}$ for the case where $\alpha \neq 0$, respectively. Then we can construct a flat front $f: \Sigma \rightarrow \mathbf{H}^3$ whose canonical forms are ω and θ . Indeed, these data give a Legendrian immersion

$$\mathcal{E} = \begin{pmatrix} \frac{z^{-\alpha/(1-\alpha)}}{c} & \frac{c\alpha z^{1/(1-\alpha)}}{1-\alpha} \\ \frac{z^{-1/(1-\alpha)}}{c} & \frac{cz^{\alpha/(1-\alpha)}}{1-\alpha} \end{pmatrix},$$

and the corresponding flat front $f = \mathcal{E}\mathcal{E}^*$ is well-defined on Σ . Moreover f is weakly complete because, for each end $p \in \bar{\Sigma} \setminus \Sigma$ of f , it holds that

$$\text{ord}_p ds_{1,1}^2 = \min\{\text{ord}_p |\omega|^2, \text{ord}_p |\theta|^2\} \leq -1.$$

The ratio of canonical forms of f is given by

$$\rho = \frac{\theta}{\omega} = -\frac{c^4\alpha}{1-\alpha}z^{2(1+\alpha)/(1-\alpha)}.$$

Thus if $\alpha = 0$ or -1 , then ρ is constant. We note that f is a horosphere if $\alpha = 0$ or a hyperbolic cylinder if $\alpha = -1$.

Moreover we can obtain weakly complete flat fronts in $\mathbf{H}^3$ of Voss type ([17, Theorem 8.3], [22]).

PROPOSITION 1.2. *Let E be an arbitrary q points on the Riemann sphere, where $q \leq 3$. Then there exists a weakly complete flat front in $\mathbf{H}^3$ whose image of the ratio of canonical forms omits precisely the set E .*

PROOF. We set $E = \{\alpha_1, \dots, \alpha_{q-1}, \alpha_q\} \subset \mathbf{C} \cup \{\infty\}$ and $\Sigma := \mathbf{C} \cup \{\infty\} \setminus E$. Then we may assume without loss of generality that $\alpha_q = \infty$. We take a holomorphic universal covering map $\xi: \tilde{\Sigma} \rightarrow \Sigma$, where $\tilde{\Sigma}$ is either the complex plane $\mathbf{C}$ or the unit disk. If we set

$$\omega = \frac{d\xi}{\prod_{i=1}^{q-1} (\xi - \alpha_i)}, \quad \rho = \xi$$

and use the representation (1.6), (1.7) and (1.8) on $\tilde{\Sigma}$, we obtain a flat front in $\mathbf{H}^3$ whose the ratio of canonical forms omits precisely the points of E . Moreover it is weakly complete. Indeed, a divergent curve Γ in $\tilde{\Sigma}$ must tend to one of the point w_i ($1 \leq i \leq q$), and we have

$$\int_{\Gamma} ds_{1,1} = \int_{\Gamma} \sqrt{1 + |\rho|^2} |\omega| = \int_{\Gamma} \frac{\sqrt{1 + |\xi|^2}}{\prod_{i=1}^{q-1} |\xi - \alpha_i|} |d\xi| = \infty,$$

when $q \leq 3$. □

2. MAIN THEOREM

In this section, we give an effective ramification theorem for the ratio of canonical forms of a weakly complete flat front in $\mathbf{H}^3$. We first recall the case where the ratio is constant.

FACT 2.1. [6, Propotion 4.4] *Let $f: \Sigma \rightarrow \mathbf{H}^3$ be a weakly complete flat front. If the ratio of canonical forms of f defined by (1.10) is constant, then f is congruent to a horosphere or a hyperbolic cylinder.*

The following is the main result of the present paper.

THEOREM 2.2. *Let $f: \Sigma \rightarrow \mathbf{H}^3$ be a weakly complete flat front. Let $q \in \mathbf{N}$, $\alpha_1, \dots, \alpha_q \in \mathbf{C} \cup \{\infty\}$ be distinct and $m_1, \dots, m_q \in \mathbf{N}$. Suppose that*

$$(2.1) \quad \gamma = \sum_{j=1}^q \left(1 - \frac{1}{m_j} \right) > 3.$$

If the ratio of canonical forms $\rho: \tilde{\Sigma} \rightarrow \mathbf{C} \cup \{\infty\}$ of f satisfies the property that all α_j -points of ρ have multiplicity at least m_j , then f must be congruent to a horosphere or a hyperbolic cylinder.

Before proceeding to the proof of Theorem 2.2, we recall two function-theoretical lemmas. For two distinct values $\alpha, \beta \in \mathbf{C} \cup \{\infty\}$, we set

$$|\alpha, \beta| := \frac{|\alpha - \beta|}{\sqrt{1 + |\alpha|^2} \sqrt{1 + |\beta|^2}}$$

if $\alpha \neq \infty$ and $\beta \neq 0$, and $|\alpha, \infty| = |\infty, \alpha| := 1/\sqrt{1+|\alpha|^2}$. Note that, if we take $v_1, v_2 \in \mathbf{S}^2$ with $\alpha = \varpi(v_1)$ and $\beta = \varpi(v_2)$, we have that $|\alpha, \beta|$ is a half of the chordal distance between v_1 and v_2 , where ϖ denotes the stereographic projection of the 2-sphere $\mathbf{S}^2$ onto $\mathbf{C} \cup \{\infty\}$.

LEMMA 2.3 ([3, Corollary 1.4.15]). *Let ρ be a nonconstant meromorphic function on $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ ($0 < R \leq \infty$). Let $q \in \mathbf{N}$, $\alpha_1, \dots, \alpha_q \in \mathbf{C} \cup \{\infty\}$ be distinct and $m_1, \dots, m_q \in \mathbf{N}$. Suppose that*

$$\gamma = \sum_{j=1}^q \left(1 - \frac{1}{m_j}\right) > 2.$$

If ρ satisfies the property that all α_j -points of ρ have multiplicity at least m_j , then, for arbitrary constants $\eta \geq 0$ and $\delta > 0$ with $\gamma - 2 > \gamma(\eta + \delta)$, it holds that

$$(2.2) \quad \frac{|\rho'|}{1 + |\rho|^2} \frac{1}{\left(\prod_{j=1}^q |\rho, \alpha_j|^{1-1/m_j}\right)^{1-\eta-\delta}} \leq C \frac{R}{R^2 - |z|^2},$$

where C is some constant depending only on γ, η, δ , and $L := \min_{i < j} |\alpha_i, \alpha_j|$.

LEMMA 2.4 ([3, Corollary 1.6.7]). *Let $d\sigma^2$ be a conformal flat metric on an open Riemann surface Σ . Then, for each point $p \in \Sigma$, there exists a local diffeomorphism Φ of a disk $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ ($0 < R \leq +\infty$) onto an open neighborhood of p with $\Phi(0) = p$ such that Φ is a local isometry, that is, the pull-back $\Phi^*(d\sigma^2)$ is equal to the standard Euclidean metric ds_{Euc}^2 on Δ_R and, for a point a_0 with $|a_0| = 1$, the Φ -image Γ_{a_0} of the curve $L_{a_0} = \{w := a_0 s; 0 < s < R\}$ is divergent in Σ .*

Proof of Theorem 2.2. This is proved by contradiction. Suppose that ρ is nonconstant. For our purpose, we may assume $\alpha_q = \infty$ and that $\tilde{\Sigma}$ is biholomorphic to the unit disk because Theorem 2.2 is obvious in the case where $\tilde{\Sigma} = \mathbf{C}$ by Nevanlinna theory [15, Section 3 in Chapter X]. We choose some δ such that $\gamma - 3 > 2\gamma\delta > 0$ and set

$$(2.3) \quad \eta := \frac{\gamma - 3 - 2\gamma\delta}{\gamma}, \quad \lambda = \frac{1}{1 + \gamma\delta}.$$

Then if we choose a sufficiently small positive number δ depending only on γ , for the constant $\varepsilon_0 = (\gamma - 3)/2\gamma$ we have

$$(2.4) \quad 0 < \lambda < 1, \quad \frac{\varepsilon_0 \lambda}{1 - \lambda} \left(= \frac{\gamma - 3}{2\gamma^2 \delta} \right) > 1.$$

Now we define a new metric

$$(2.5) \quad d\sigma^2 = |h_z|^{2/(1-\lambda)} \left(\frac{1}{|\rho'_z|} \prod_{j=1}^{q-1} \left(\frac{|\rho - \alpha_j|}{\sqrt{1+|\alpha_j|}} \right)^{\eta_j(1-\eta-\delta)} \right)^{2\lambda/(1-\lambda)} |dz|^2$$

on the set $\tilde{\Sigma}' = \{z \in \tilde{\Sigma}; \rho'_z(z) \neq 0 \text{ and } \rho(z) = \alpha_j \text{ for all } j\}$ where $\omega = h_z dz$, $\rho'_z = d\rho/dz$ and $\eta_j = 1 - 1/m_j$. Take a point $p \in \tilde{\Sigma}'$. Since the metric $d\sigma^2$ is flat on $\tilde{\Sigma}'$, by Lemma 2.4,

there exists a local isometry Φ satisfying $\Phi(0) = p$ from a disk $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ ($0 < R \leq +\infty$) with the standard metric ds_{Euc}^2 onto an open neighborhood of p in $\tilde{\Sigma}'$ with the metric $d\sigma^2$ such that, for a point a_0 with $|a_0| = 1$, the Φ -image Γ_{a_0} of the curve $L_{a_0} = \{w := a_0 s; 0 < s < R\}$ is divergent in $\tilde{\Sigma}'$. For brevity, we denote the function $\rho \circ \Phi$ on Δ_R by ρ in the followings. By Lemma 2.3, we get

$$(2.6) \quad R \leq C \frac{1 + |\rho(0)|^2}{|\rho'_z(0)|} \prod_{j=1}^q |\rho(0), \alpha_j|^{\eta_j(1-\eta-\delta)} < +\infty.$$

Hence

$$L_{d\sigma}(\Gamma_{a_0}) = \int_{\Gamma_{a_0}} d\sigma = R < +\infty,$$

where $L_{d\sigma}(\Gamma_{a_0})$ denotes the length of Γ_{a_0} with respect to the metric $d\sigma^2$.

Now we prove that Γ_{a_0} is divergent in $\tilde{\Sigma}$. If not, then Γ_{a_0} must tend to a point $p_0 \in \tilde{\Sigma} \setminus \tilde{\Sigma}'$ where $\rho'_z(p_0) = 0$ or $\rho(p_0) = \alpha_j$ for some j because Γ_{a_0} is divergent in $\tilde{\Sigma}'$ and $L_{d\sigma}(\Gamma_{a_0}) < +\infty$. Taking a local complex coordinate ζ in a neighborhood of p_0 with $\zeta(p_0) = 0$, we can write the metric $d\sigma^2$ as

$$d\sigma^2 = |\zeta|^{2k\lambda/(1-\lambda)} v |dz|^2$$

with some positive smooth function v and some real number k . If $\rho - \alpha_j$ has a zero of order $m (\geq m_j \geq 2)$ at p_0 for some $j \leq q-1$, then ρ'_z has a zero of order $m-1$ at p_0 and $h_z(p_0) \neq 0$. Then we have

$$\begin{aligned} k &= -(m-1) + m \left(1 - \frac{1}{m_j}\right) (1 - \eta - \delta) \\ &= 1 - \frac{m}{m_j} - \frac{m}{m_j} (m_j - 1) (\eta + \delta) \\ &\leq -(\eta + \delta) \leq -\varepsilon_0. \end{aligned}$$

For the case where ρ has a pole of order $m (\geq m_q)$, ρ'_z has a pole of order $m+1$, h_z has a zero of order m at p_0 and each component $\rho - \alpha_j$ in the right side of (2.5) has a pole of order m at p_0 . Using the identity $\eta_1 + \cdots + \eta_{q-1} = \gamma - \eta_q$ and (2.4), we get

$$\begin{aligned} k &= \frac{m}{\lambda} + (m+1) - m(\gamma - \eta_q)(1 - \eta - \delta) \\ &= m\eta_q(1 - \eta - \delta) - (m-1) \leq -\varepsilon_0. \end{aligned}$$

Moreover, for the case where $\rho'_z(p_0) = 0$ and $\rho(p_0) \neq \alpha_j$ for all j , we see $k \leq -1$. Thus, in any case, $k\lambda/(1-\lambda) \leq -1$ by (2.4) and there exists a positive constant C' such that

$$d\sigma \geq C' \frac{|d\zeta|}{|\zeta|}$$

in a neighborhood of p_0 . Hence we have

$$R = \int_{\Gamma_{a_0}} d\sigma \geq C' \int_{\Gamma_{a_0}} \frac{|d\zeta|}{|\zeta|} = +\infty,$$

which contradicts (2.6).

On the other hand, since $d\sigma^2 = |dz|^2$, we obtain by (2.5)

$$(2.7) \quad |h_z| = \left(|\rho'_z| \prod_{j=1}^{q-1} \left(\frac{\sqrt{1+|\alpha_j|^2}}{|\rho - \alpha_j|} \right)^{\eta_j(1-\eta-\delta)} \right)^\lambda.$$

By Lemma 2.3, we have

$$\begin{aligned} \Phi^* ds_{1,1} &= |h_z| \sqrt{1+|\rho|^2} |dz| \\ &= \left(|\rho_z| (1+|\rho|^2)^{1/2\lambda} \prod_{j=1}^{q-1} \left(\frac{\sqrt{1+|\alpha_j|^2}}{|\rho - \alpha_j|} \right)^{\eta_j(1-\eta-\delta)} \right)^\lambda |dz| \\ &= \left(\frac{|\rho'_z|}{1+|\rho|^2} \frac{1}{\prod_{j=1}^q |\rho, \alpha_j|^{\eta_j(1-\eta-\delta)}} \right)^\lambda |dz| \\ &\leq C^\lambda \left(\frac{R}{R^2 - |z|^2} \right)^\lambda |dz|. \end{aligned}$$

Thus if we denote the distance $d(p)$ from a point $p \in \tilde{\Sigma}$ to the boundary of $\tilde{\Sigma}$ as the greatest lower bound of the lengths with respect to the metric $ds_{1,1}^2$ of all divergent paths in $\tilde{\Sigma}$, then we have

$$d(p) \leq \int_{\Gamma_{a_0}} ds_{1,1} = \int_{L_{a_0}} \Phi^* ds_{1,1} = C^\lambda \int_{L_{a_0}} \left(\frac{R}{R^2 - |z|^2} \right)^\lambda |dz| \leq C^\lambda \frac{R^{1-\lambda}}{1-\lambda} < +\infty$$

because $0 < \lambda < 1$. However it contradicts the assumption that $ds_{1,1}^2$ is complete. $\square$

3. APPLICATIONS

This section is devoted to prove two applications of the main theorem.

3.1. The Ahlfors Islands Theorem. We recall the definition of an island of a meromorphic function on a Riemann surface.

DEFINITION 3.1. Let Σ be a Riemann surface and $f: \Sigma \rightarrow \mathbf{C} \cup \{\infty\}$ a meromorphic function. Let $V \subset \mathbf{C} \cup \{\infty\}$ be a Jordan domain. A simply-connected component U of $f^{-1}(V)$ with $\bar{U} \subset \Sigma$ is called an *island* of f over V . Note that $f|_U: U \rightarrow V$ is a proper map. The degree of this map is called the *multiplicity* of the island U . An island of multiplicity one is called a *simple island*.

Since the ratio ρ of canonical forms of a flat front $f: \Sigma \rightarrow \mathbf{H}^3$ is a meromorphic function on $\tilde{\Sigma}$, we can consider an island of ρ . When all islands of ρ are small disks, we get the following result by applying Theorem 2.2.

COROLLARY 3.2. *Let $f: \Sigma \rightarrow \mathbf{H}^3$ be a weakly complete flat front. Let $q \in \mathbf{N}$, $\alpha_1, \dots, \alpha_q \in \mathbf{C}$ be distinct, $D(\alpha_j, \varepsilon) := \{z \in \mathbf{C} : |z - \alpha_j| < \varepsilon\}$ ($1 \leq j \leq q$) be pairwise disjoint and*

$m_1, \dots, m_q \in \mathbf{N}$. Suppose that

$$(3.1) \quad \sum_{j=1}^q \left(1 - \frac{1}{m_j}\right) > 3.$$

Then there exists $\varepsilon > 0$ such that if the ratio of canonical forms of f has no island of multiplicity less than m_j over $D(\alpha_j, \varepsilon)$ for all $j \in \{1, \dots, q\}$ then f must be congruent to a horosphere or a hyperbolic cylinder.

PROOF. If such an ε does not exist, for any ε , we can find a weakly complete flat front whose the ratio of canonical forms ρ is nonconstant and has no island of multiplicity less than m_j over $D(\alpha_j, \varepsilon)$. However this implies that all α_j -points of ρ have multiplicity at least m_j , contradicting Theorem 2.2. $\square$

The important special case of Corollary 3.2 is the case where $q = 7$ and $m_j = 2$ for all j . This corresponds to a weak version of the so-called Five Islands Theorem in the Ahlfors theory of covering surfaces ([1], [15, Chapter XIII]).

COROLLARY 3.3. *Let $f: \Sigma \rightarrow \mathbf{H}^3$ be a weakly complete flat front. Let $\alpha_1, \dots, \alpha_7 \in \mathbf{C}$ be distinct and $D(\alpha_j, \varepsilon) := \{z \in \mathbf{C} : |z - \alpha_j| < \varepsilon\}$ ($1 \leq j \leq 7$). Then there exists $\varepsilon > 0$ such that if the ratio of canonical forms ρ of f has no simple island of over any of the small disks $D(\alpha_j, \varepsilon)$ then f must be congruent to a horosphere or a hyperbolic cylinder.*

3.2. The classification of complete flat surfaces in $\mathbf{H}^3$. As another application of Theorem 2.2, we obtain the best possible upper bound for the number of exceptional values of the ratio of canonical forms of a weakly complete flat front in $\mathbf{H}^3$.

COROLLARY 3.4 ([6, Theorem 4.5]). *Let f be a weakly complete flat front in $\mathbf{H}^3$. If the ratio of canonical forms ρ of f omits more than three values, then f must be congruent to a horosphere or a hyperbolic cylinder.*

PROOF. In Theorem 2.2, if ρ does not take a value α_j , then we may set $m_j = \infty$ in (2.1), and if ρ omits all values α_j ($1 \leq j \leq q$), (2.1) means $q > 3$, which is the case of this result. $\square$

The number “three” is sharp because there exist examples in Proposition 1.2. As an application of this corollary, we give a simple proof of the classification of complete nonsingular flat surfaces in $\mathbf{H}^3$.

COROLLARY 3.5 ([20], [21]). *Any complete flat surface in $\mathbf{H}^3$ must be congruent to a horosphere or a hyperbolic cylinder.*

PROOF. Because a flat surface has no singularities, the complement of the image of ρ contains at least the set $\{|\rho| = 1\} \subset \mathbf{C} \cup \{\infty\}$. On the other hand, we have

$$ds^2 = |\omega + \bar{\theta}|^2 \leq |\omega|^2 + |\theta|^2.$$

Thus if a flat surface in $\mathbf{H}^3$ is complete, then it is also weakly complete. Therefore, by Corollary 3.4, it must be congruent to a horosphere or a hyperbolic cylinder. $\square$

REFERENCES

- [1] L. V. Ahlfors, Zur Theorie der Überlagerungsflächen, *Acta Math.* **65** (1935), 157–194, and *Collected Papers Vol. I*, pp. 163–173.
- [2] W. Bergweiler, The role of the Ahlfors five islands theorem in complex dynamics, *Conform. Geom. Dyn.* **4** (2000), 22–34.
- [3] H. Fujimoto, Value distribution theory of the Gauss map of minimal surfaces in $\mathbf{R}^m$, Vieweg, Braunschweig/Wiesbaden, 1993.
- [4] J. A. Gálvez, A. Martínez and F. Milán, Flat surfaces in hyperbolic 3-space, *Math. Ann.* **316** (2000), 419–435.
- [5] Y. Kawakami, Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space, to appear in *Houston Journal of Mathematics*, arXiv:0908.1307.
- [6] Y. Kawakami and D. Nakajo, Value distribution of the Gauss map of improper affine spheres, to appear in *Journal of the Mathematical Society of Japan*, arXiv:1004.1484.
- [7] R. Kobayashi, Toward Nevanlinna theory as a geometric model of Diophantine approximation, *Sugaku Expositions* **16** (2003), 39–79.
- [8] M. Kokubu, W. Rossman, K. Saji, M. Umehara and K. Yamada, Singularities of flat fronts in hyperbolic space, *Pacific J. Math.* **221** (2005), 303–351.
- [9] M. Kokubu, W. Rossman, M. Umehara and K. Yamada, Flat fronts in hyperbolic 3-space and their caustics, *J. Math. Soc. Japan* **59** (2007), 265–299.
- [10] M. Kokubu, W. Rossman, M. Umehara and K. Yamada, Asymptotic behavior of flat surfaces in hyperbolic 3-space, *J. Math. Soc. Japan* **61** (2009), 799–852.
- [11] T. Klotz and L. Sario, Gaussian mapping of arbitrary minimal surfaces, *J. Analyse Math.* **17** (1966), 209–217.
- [12] M. Kokubu, M. Umehara and K. Yamada, An elementary proof of Small’s formula for null curves in $\mathrm{PSL}(2, \mathbf{C})$ and an analogue for Legendrian curves in $\mathrm{PSL}(2, \mathbf{C})$, *Osaka J. Math.* **40** (2003), 697–715.
- [13] M. Kokubu, M. Umehara and K. Yamada, Flat fronts in hyperbolic 3-space, *Pacific J. Math.* **216** (2004), 149–175.
- [14] F. Martín, M. Umehara and K. Yamada, Calabi-Yau problem for Legendrian curves in $\mathbf{C}^3$ and applications, preprint, arXiv:1107.0106.
- [15] R. Nevanlinna, *Analytic Function*, Translated from the second German edition by Phillip Emig. *Die Grundlehren der mathematischen Wissenschaften*, Springer, New York, 1970.
- [16] J. Noguchi and T. Ochiai, *Geometric Function Theory in Several Complex Variables*, Transl. Math. Monog. **80**, Amer. Math. Soc., Providence, RI, 1990.
- [17] R. Osserman, *A survey of minimal surfaces*, Second edition. Dover Publications, Inc., New York, 1986.
- [18] M. Ru, *Nevanlinna theory and its relation to Diophantine approximation*, World Sci. Publ. Co. Inc., River Edge, NJ, 2001.
- [19] K. Saji, M. Umehara and K. Yamada, The geometry of fronts, *Ann. of Math.* **169** (2009), 491–529.
- [20] S. Sasaki, On complete flat surfaces in hyperbolic 3-space, *Kōdai Math Sem. Rep.* **25** (1973), 449–457.
- [21] Y. A. Volkov and S. M. Vladimirova, Isometric immersions of the Euclidean plane in Lobachevskii space (Russian), *Mat. Zametki* **10** (1971), 327–332.
- [22] K. Voss, Über vollständige minimalflächen, *L’Enseignement Math.*, **10** (1964), 316–317.

GRADUATE SCHOOL OF SCIENCE AND ENGINEERING,
YAMAGUCHI UNIVERSITY,
YAMAGUCHI, 753-8512, JAPAN
E-mail address: y-kwkami@yamaguchi-u.ac.jp

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p -adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection

- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments

- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow

- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra
- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions

- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing
- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine

- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space