

Asymptotic behavior of blow-up solutions to a degenerate parabolic equation

Anada, Koichi
Waseda University Senior High School

Ishiwata, Tetsuya
Department of Mathematical Sciences, Shibaura Institute of Technology

<https://hdl.handle.net/2324/20606>

出版情報 : Journal of Math-for-Industry (JMI). 3 (C), pp.1-8, 2012-01-27. Faculty of Mathematics, Kyushu University

バージョン :

権利関係 :



Asymptotic behavior of blow-up solutions to a degenerate parabolic equation

Koichi Anada and Tetsuya Ishiwata

Revised on October 25, 2011

Abstract. We consider the Dirichlet problems for a degenerate parabolic equation, $u_t = u^\delta(\Delta u + \lambda u)$ in a bounded domain in $\mathbb{R}^n$ with a smooth boundary. It has been known that if $\delta \geq 2$ then there exists u which blows up faster than the rate of $(T - t)^{-1/\delta}$, where T is the blow-up time of u . The solutions are called “Type 2”. In this paper we investigate features for asymptotic behavior of “Type 2” solutions for the case of $\delta = 2$ and $\delta > 2$.

Keywords. degenerate parabolic equations, blow-up, asymptotic behavior, type 2, eventual monotonicity

1. INTRODUCTION

Let Ω be a bounded domain in $\mathbb{R}^n$ with a smooth boundary $\partial\Omega$. Suppose that a function $u_0 \in C^\infty(\Omega) \cap C(\bar{\Omega})$ satisfies

$$u_0 > 0 \quad \text{in } \Omega \quad \text{and} \quad u_0(x) = 0 \quad \text{for } x \in \partial\Omega. \quad (1.1)$$

Then, we consider the following initial-boundary value problems.

$$\begin{cases} \frac{\partial u}{\partial t} = u^\delta(\Delta u + \lambda u) & x \in \Omega, t > 0, \\ u(x, 0) = u_0(x) & x \in \bar{\Omega}, \\ u(x, t) = 0 & x \in \partial\Omega, t \geq 0, \end{cases} \quad (1.2)$$

where $\delta > 0$ and λ is greater than the first eigenvalue, $\lambda_1(\Omega)$ of $-\Delta$ in Ω , that is,

$$\lambda > \lambda_1(\Omega).$$

In this paper we discuss classical solutions which are approximated by functions u_ε solving the following problems:

$$\begin{cases} \frac{\partial u_\varepsilon}{\partial t} = u_\varepsilon^\delta(\Delta u_\varepsilon + \lambda u_\varepsilon) & x \in \Omega, t > 0, \\ u_\varepsilon(x, 0) = u_0(x) + \varepsilon & x \in \bar{\Omega}, \\ u_\varepsilon(x, t) = \varepsilon & x \in \partial\Omega, t \geq 0. \end{cases} \quad (1.3)$$

It has been proved that each of them blows up at a finite time, that is, for any solutions approximated by u_ε , there exists $T > 0$ such that

$$\limsup_{t \nearrow T} \|u(\cdot, t)\|_\infty = \infty,$$

where $\|u(\cdot, t)\|_\infty = \sup_{x \in \Omega} u(x, t)$. (For instance, see [2].)

Here the constant T is called “the blow-up time” of u . Moreover it has been known that they are classified into two types by their blow-up rates as follows.

Definition 1. Let u and T be a solution of (1.2) and the blow-up time of u , respectively. Then u is called “Type 1” if it satisfies

$$\limsup_{t \nearrow T} (T - t)^{\frac{1}{\delta}} \|u(\cdot, t)\|_\infty < +\infty \quad (1.4)$$

and u is called “Type 2” if it satisfies

$$\limsup_{t \nearrow T} (T - t)^{\frac{1}{\delta}} \|u(\cdot, t)\|_\infty = +\infty. \quad (1.5)$$

Precisely, It has been proved that if $\delta \geq 2$ then there exists solutions of (1.2) which are “Type 2” and if $0 < \delta < 2$ then u is “Type 1”. (See [1], [3], [4], [6], [9], [10] and so force.) Our purpose of this paper is to investigate asymptotic behavior of “Type 2” solutions for $\delta \geq 2$.

First, we consider the case of $\delta = 2$. S. Angenent gave important asymptotic profiles in [3] for curve shortening problems as follows.

$$v_t = v^2(v_{xx} + v) \quad \text{with periodic boundary conditions.}$$

The first profile is that there exists $t_0 > 0$ such that $v_t(x, t) = v(x, t)^2(v_{xx}(x, t) + v(x, t)) > 0$ for any x and $t > t_0$. This property is called “eventual monotonicity”.

Secondly, it was proved that $V(x, t) := \frac{v(x, t)}{\|v(\cdot, t)\|_\infty}$ satisfies

$$V(x, t) \rightarrow \cos(x - x_0) \quad \text{if } |x - x_0| \leq \frac{\pi}{2}, \quad (1.6)$$

as $t \nearrow T$ for some x_0 . Besides, an upper bound for blow-up rates of solutions was given. In order to prove its existence, Angenent investigated relations between $\int_\Omega \log v(x, t) dx$ and $\log(T - t)$.

Their results hold under a special situation in curve shortening problems but we have not known whether they can be proved without any special assumptions.

In Section 3 we will discuss asymptotic behavior of solution to (1.2) without any special assumptions in the case of $\delta = 2$. Our method is based on behavior of $\int_{\Omega} u^{1-\delta} u_t dx$ as t tends to the blow-up time of u . Precisely, we will prove in Section 3 that if $\delta = 2$ then $2(T-t) \int_{\Omega} u^{-1} u_t dx$ becomes positive and has an upper bound after a finite time. In particular, the positivity implies that there exists $t_0 \in (0, T)$ such that $\int_{\Omega} u^{-1} u_t dx = \int_{\Omega} u(\Delta u + \lambda u) dx > 0$ for $t \in [t_0, T)$ and we call this property “weak eventual monotonicity”.

The positivity and the upper boundedness of $2(T-t) \int_{\Omega} u^{-1} u_t dx$ on $\delta = 2$ lead to two profiles for asymptotic behavior of solutions to (1.2). One is that there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that $t_k \nearrow T$ and

$$\frac{\int_{\Omega} |\nabla U(x, t_k)|^2 dx}{\int_{\Omega} U(x, t_k)^2 dx} \rightarrow \lambda \quad \text{as } k \rightarrow \infty, \quad (1.7)$$

where $U(x, t) := \frac{u(x, t)}{\|u(\cdot, t)\|_{\infty}}$. We can regard (1.7) as a generalization of (1.6). Another is that if $\delta = 2$ and $\log u_0 \in L^1(\Omega)$ then $\int_{\Omega} \log \left[(T-t)^{\frac{1}{2}} u(x, t) \right] dx$ has an upper bound, that is, there exists a positive constant $C > 0$ such that

$$\int_{\Omega} \log \left[(T-t)^{\frac{1}{2}} u(x, t) \right] dx \leq C \quad (1.8)$$

for $t \in [t_0, T)$, where C depends on T and u_0 only. It may give a important profile for blow-up rates of “Type 2” that (1.5) holds for $\delta = 2$ but satisfies (1.8).

In Section 4 the case of $\delta > 2$ will be discussed. We expect that behavior of solutions for $\delta > 2$ may be a little different from $\delta = 2$. In the situation, we investigate features of $\int_{\Omega} u^{1-\delta} u_t dx$, $\int_{\Omega} u^{2-\delta} dx$ and others by numerical computing. And then we will present two conjectures for asymptotic behavior of solutions of (1.2) by numerical observations. Moreover, we will prove under their conjectures that $\delta(T-t)^{\frac{\delta}{2}} \int_{\Omega} u^{1-\delta} u_t dx$ has an upper bound after a finite time and (1.7) holds for some sequences.

2. SOME LEMMAS

Let u be a solution of (1.2) approximated by u_{ε} solving (1.3). The following two lemmas have been well-known in previous studies for asymptotic behavior of u .

Lemma 1. *It holds that*

$$u_t(x, t) \geq -\frac{1}{\delta t} u(x, t) \quad \text{for } x \in \Omega \text{ and } t \in (0, T).$$

Lemma 2. *If $\{t_k\}_{k \in \mathbb{N}}$ is a sequence such that $t \nearrow T$ as $k \rightarrow \infty$, then there is $\mu \in (0, 1)$ and a set $S_{\mu} \subset \Omega$ with positive Lebesgue measure such that*

$$u(x, t) \geq \mu \|u(\cdot, t)\|_{\infty} \quad \text{for } x \in S_{\mu} \text{ and } t \in (0, T)$$

for all k from an appropriate subsequence. In particular, the Lebesgue measure of $\left\{x \in \Omega \mid \limsup_{t \nearrow T} u(x, t) = \infty\right\}$ is positive.

They are very important properties for our purpose and can be proved by the maximum principle for parabolic equations. (For instance, see [9] and [10] for their proofs.)

Moreover, we give another important property for u .

Lemma 3. *If $u_0 \in C^{\infty}(\Omega) \cap C(\bar{\Omega})$ satisfies (1.1) and $u_0^{2-\delta} \in L^1(\Omega)$ then*

$$\left(\int_{\Omega} u^{1-\delta} u_t dx \right)^2 < \int_{\Omega} u^{2-\delta} dx \int_{\Omega} u^{-\delta} (u_t)^2 dx \quad (2.1)$$

for any $t \in (0, T)$.

Remark 1. In Lemma 3, $u^{1-\delta} u_t$, $u^{-\delta} (u_t)^2$ and $u^{2-\delta}$ is integrable in Ω for any $t \in (0, T)$ because u is a classical solution of $u_t = u^{\delta} (\Delta u + \lambda u)$.

Remark 2. When $\delta = 2$, the assumption $u_0^{2-\delta} \in L^1(\Omega)$ is not required and then we can prove

$$\left(\int_{\Omega} u^{-1} u_t dx \right)^2 < |\Omega| \int_{\Omega} u^{-2} (u_t)^2 dx,$$

where $|\Omega|$ is the Lebesgue measure of Ω .

Proof. By Schwartz's inequality,

$$\left(\int_{\Omega} u^{1-\delta} u_t dx \right)^2 \leq \int_{\Omega} u^{2-\delta} dx \int_{\Omega} u^{-\delta} (u_t)^2 dx. \quad (2.2)$$

Note that $\int_{\Omega} u^{2-\delta} dx = |\Omega|$ if $\delta = 2$. Now, $u^{2-\delta} \not\equiv u^{-\delta} (u_t)^2$ because $u^{-1} u_t$ is not constant in Ω . Hence, the equality in (2.2) does not hold and we get this lemma. $\square$

Remark 3. (2.1) implies that When $\delta > 2$, if u_0 satisfies (1.1) and $u_0^{2-\delta} \in L^1(\Omega)$ then

$$E(t) := \int_{\Omega} u^{2-\delta} dx \int_{\Omega} u^{-\delta} (u_t)^2 dx - \left(\int_{\Omega} u^{1-\delta} u_t dx \right)^2 > 0. \quad (2.3)$$

We will discuss the ratio of $E(t)$ to $\left(\int_{\Omega} u^{1-\delta} u_t dx\right)^2$ near the blow-up time in Section 4.

Let $\{u_{0,n}\}_{n \in \mathbb{N}} \in C^\infty(\Omega)$ be such that $\|u_{0,n} - u_0\|_{L^\infty(\Omega)} < \frac{1}{k}$ and $\alpha_n \Theta \leq u_{0,n} \leq \beta_n \Theta$ holds with constants $0 < \alpha_n < \beta_n$, where Θ denotes the principal eigenfunction of $-\Delta$ in Ω with $\max_{\Omega} \Theta = 1$. Then it has been shown in [9] that if $\delta > 1$ then the problems

$$\begin{cases} \frac{\partial u_n}{\partial t} = u_n^\delta (\Delta u_n + \lambda u_n) & x \in \Omega, t \in (0, T(u_n)), \\ u_n(x, 0) = u_{0,n}(x) & x \in \bar{\Omega}, \\ u_n(x, t) = 0 & x \in \partial\Omega, t \geq 0, \end{cases}$$

have unique solutions $u_n \in C(\bar{\Omega} \times [0, T(u_n))) \cap C^\infty(\Omega \times (0, T(u_n)))$, $T(u_n) \rightarrow T$, which may be approximated by function $u_{n,\varepsilon}$ solving (1.3) with $u_{n,\varepsilon} = u_n + \varepsilon$ on the parabolic boundary. As $n \rightarrow \infty$, $u_n \rightarrow u$ in the topology of $C_{loc}(\bar{\Omega} \times [0, T)) \cap C_{loc}^\infty(\Omega \times (0, T))$.

Furthermore, u_n have the property that for each $n \in \mathbb{N}$ and each $\tau > 0$ there is a constant $c(n, \tau) > 0$ such that

$$|\text{dist}(x, \Omega)^{\alpha-1+j(2-\delta)} \partial_x^\alpha \partial_t^j u_n(x, t)| \leq c(n, \tau)$$

for $j = 0, 1$, $\alpha = 0, 1, 2$ and $(x, t) \in \Omega \times (0, T - \tau)$. In particular, $|\nabla(u_n)_t| \leq \text{dist}(x, \partial\Omega)^{\delta-2} c(n, \tau)$ for $(x, t) \in \Omega \times (0, T - \tau)$ and u_n satisfies

$$\begin{aligned} & \int_{\Omega} u_n^{-\delta} \left((u_n)_t \right)^2 dx \\ &= \int_{\Omega} (\Delta u_n + \lambda u_n) (u_n)_t dx \\ &= \int_{\Omega} \left(-\nabla u_n \cdot \nabla (u_n)_t + \lambda u_n (u_n)_t \right) dx \\ &= \frac{1}{2} \left(\int_{\Omega} (-|\nabla u_n|^2 + \lambda u_n^2) dx \right)_t \\ &= \frac{1}{2} \left(\int_{\Omega} u_n^{1-\delta} (u_n)_t dx \right)_t. \end{aligned} \quad (2.4)$$

Therefore, by Remark 2, we obtain the following lemma.

Lemma 4. *If $\delta = 2$ then*

$$2 \left(\int_{\Omega} u_n^{-1} (u_n)_t dx \right)^2 < |\Omega| \left(\int_{\Omega} u_n^{-1} (u_n)_t dx \right)_t$$

for any $t \in (0, T(u_n))$.

And Lemma 4 leads to the following corollary.

Corollary 1. *If $\delta = 2$ then $\int_{\Omega} u_n^{-1} (u_n)_t dx$ is strictly increasing with respect to t .*

Proof. By Lemma 4, if $\delta = 2$ then

$$\left(\int_{\Omega} u_n^{-1} (u_n)_t dx \right)_t > \frac{2}{|\Omega|} \left(\int_{\Omega} u_n^{-1} (u_n)_t dx \right)^2 > 0$$

for any $t \in (0, T(u_n))$. $\square$

3. THE CASE OF $\delta = 2$

3.1. THE MAIN THEOREM FOR $\delta = 2$

We consider the case of $\delta = 2$, that is, the following problem.

$$\begin{cases} \frac{\partial u}{\partial t} = u^2 (\Delta u + \lambda u) & (x, t) \in \Omega \times (0, T), \\ u(x, 0) = u_0(x) & x \in \bar{\Omega}, \\ u(x, t) = 0 & (x, t) \in \partial\Omega \times (0, T), \end{cases} \quad (3.1)$$

where T is the blow-up time of u . Then we can prove the following theorem.

Theorem 1. *Let $\delta = 2$ and u be a solution approximated by u_ε solving (1.3). Suppose that $u_0 \in C^\infty(\Omega) \cap C(\bar{\Omega})$ satisfies (1.1) and $\log u_0 \in L^1(\Omega)$. Then there exists $t_0 > 0$ such that*

$$0 < 2(T - t) \int_{\Omega} u^{-1} u_t dx \leq |\Omega|$$

for any $t \in [t_0, T)$, where $T > 0$ is the blow-up time of u .

Remark 4. The left side inequality in Theorem 1 implies that $\int_{\Omega} u^{-1} u_t dx = \int_{\Omega} u (\Delta u + \lambda u) dx > 0$ for $t \in [t_0, T)$. In this paper we call this property “weak eventual monotonicity”.

Proof of Theorem 1. Let $\tau > 0$ be small enough and $\{t_k\}_{k=1,2,\dots}$ be a sequence with $t_k > \tau$, $t_k \nearrow T$ and $\|u(\cdot, t_k)\|_\infty \nearrow \infty$ as $k \rightarrow \infty$. Since Lemma 1 implies $(\log u)_t \geq -\frac{1}{2t}$, it holds that

$$\log u(x, t_k) - \log u(x, \tau) \geq -\frac{1}{2}(\log t_k - \log \tau) > -\frac{1}{2} \log \frac{T}{\tau}.$$

By Lemma 2, for an appropriate subsequence, there exists $\mu \in (0, 1)$ and a set S_μ such that $|S_\mu| > 0$ and

$$\begin{aligned} & \int_{\Omega} \log u(x, t_k) dx \\ &= \int_{S_\mu} \log u(x, t_k) dx + \int_{\Omega \setminus S_\mu} \log u(x, t_k) dx \\ &\geq |S_\mu| \log(\mu \|u(\cdot, t_k)\|_\infty) \\ &\quad - \frac{1}{2} |\Omega \setminus S_\mu| \log \frac{T}{\tau} + \int_{\Omega \setminus S_\mu} \log u(x, \tau) dx. \end{aligned}$$

Since the last term is bounded for a fixed positive number $\tau \in (0, T)$ because of $\log u_0 \in L^1(\Omega)$, we can get

$$\int_{\Omega} \log u(x, t_k) dx \rightarrow \infty \quad \text{as } k \rightarrow \infty \text{ for any } n \in \mathbb{N}$$

which implies that there exists t_0 such that

$$\int_{\Omega} u(x, t_0)^{-1} u_t(x, t_0) dx = \frac{d}{dt} \int_{\Omega} \log u(x, t_0) dx > 0.$$

Hence, by Corollary 1,

$$\int_{\Omega} u^{-1} u_t dx > 0 \quad \text{for any } t \in [t_0, T).$$

On the other hands, when $\delta = 2$, Lemma 4 leads to

$$\frac{2}{|\Omega|} + \left[\left(\int_{\Omega} u_n^{-1} (u_n)_t dx \right)^{-1} \right]_t < 0 \quad \text{for } t \in (0, T(u_n))$$

for u_n defined in Section 2. Integrating from t to s , we have

$$\begin{aligned} \frac{2(s-t)}{|\Omega|} + \left(\int_{\Omega} u_n(x, s)^{-1} (u_n)_t(x, s) dx \right)^{-1} \\ - \left(\int_{\Omega} u_n(x, t)^{-1} (u_n)_t(x, t) dx \right)^{-1} < 0. \end{aligned}$$

Since

$$\begin{aligned} \int_{\Omega} u_n(x, s)^{-1} (u_n)_t(x, s) dx \\ > \int_{\Omega} u_n(x, t)^{-1} (u_n)_t(x, t) dx > 0 \end{aligned}$$

if $t_0 \leq t < s < T(u_n)$, it holds that

$$2(s-t) \int_{\Omega} u_n(x, t)^{-1} (u_n)_t(x, t) dx < |\Omega|,$$

where $t_0 \leq t < s < T(u_n)$. Then it is verified by letting $s \nearrow T(u_n)$ that

$$\begin{aligned} 2(T(u_n) - t) \int_{\Omega} u_n(\Delta u_n + \lambda u_n) dx \\ = 2(T(u_n) - t) \int_{\Omega} u_n^{-1} (u_n)_t dx \leq |\Omega| \end{aligned}$$

for any $t \in [t_0, T(u_n))$.

Now, we mentioned in Section 2 that $T(u_n) \rightarrow T$ and $u_n \rightarrow u$ in the topology of $C_{loc}^0(\bar{\Omega} \times [0, T)) \cap C_{loc}^\infty(\Omega \times (0, T))$ as $n \rightarrow \infty$. This implies that $u_n(\cdot, t) \rightarrow u(\cdot, t)$ uniformly in $\bar{\Omega}$ and $\Delta u_n(\cdot, t) \rightharpoonup \Delta u(\cdot, t)$ weakly in $C^*(\Omega)$ as $n \rightarrow \infty$ for any $t \in [t_0, T)$. Then we have

$$\begin{aligned} 2(T-t) \int_{\Omega} u^{-1} u_t dx \\ = 2(T-t) \int_{\Omega} u(\Delta u + \lambda u) dx \leq |\Omega| \end{aligned}$$

which completes this proof. $\square$

3.2. BEHAVIOR NEAR THE BLOW-UP TIME

We discuss some important profiles for asymptotic behavior of $U(x, t) := \frac{u(x, t)}{\|u(\cdot, t)\|_\infty}$, where u is a ‘‘Type 2’’ blow-up solution to (3.1).

First, suppose that u_0 satisfies (1.1) and $\Delta u_0 + \lambda u_0 \geq 0$ in Ω . Then it is easily verified by the maximum principle

that $u > 0$, $\Delta u + \lambda u \geq 0$ and $U(\Delta U + \lambda U) \geq 0$ in $\Omega \times (0, T)$. Besides, Theorem 1 leads to

$$\begin{aligned} 0 < \int_{\Omega} U(\Delta U + \lambda U) dx &= \frac{1}{\|u(\cdot, t)\|_\infty^2} \int_{\Omega} u(\Delta u + \lambda u) dx \\ &= \frac{1}{\|u(\cdot, t)\|_\infty^2} \int_{\Omega} u^{-1} u_t dx \\ &\leq \frac{|\Omega|}{2(T-t)\|u(\cdot, t)\|_\infty^2} \end{aligned}$$

for $t \in [t_0, T)$. In addition, ‘‘Type 2’’ means

$$\limsup_{t \nearrow T} (T-t)^{\frac{1}{2}} u(x, t) = +\infty.$$

Therefore, if u is ‘‘Type 2’’ and u_0 satisfies $\Delta u_0 + \lambda u_0 \geq 0$ in Ω , then we obtain that there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that $t_k \nearrow T$ as $k \rightarrow \infty$ and U satisfies

$$\Delta U(x, t_k) + \lambda U(x, t_k) \rightarrow 0 \text{ or } U(x, t_k) \rightarrow 0$$

for almost all of $x \in \Omega$ as $k \rightarrow \infty$.

On the other hands, we assume (1.1) and $\log u_0 \in L^1(\Omega)$ only. Then Theorem 1 can lead to the following corollary such that we can regard as a generalization of the profile for asymptotic behavior of $U(x, t)$ near the blow-up time.

Corollary 2. *Let $\delta = 2$ and u be a ‘‘Type 2’’ solution of (3.1) approximated by u_ε solving (1.3). Suppose that $u_0 \in C^\infty(\Omega) \cap C(\bar{\Omega})$ satisfies (1.1) and $\log u_0 \in L^1(\Omega)$. Then there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that $t_k \nearrow T$ as $k \rightarrow \infty$ and $U(x, t) := \frac{u(x, t)}{\|u(\cdot, t)\|_\infty}$ satisfies*

$$\frac{\int_{\Omega} |\nabla U(x, t_k)|^2 dx}{\int_{\Omega} U(x, t_k)^2 dx} \rightarrow \lambda \quad \text{as } k \rightarrow \infty.$$

Proof. If $\delta = 2$, then

$$\begin{aligned} \int_{\Omega} u^{-1} u_t dx &= \int_{\Omega} u(\Delta u + \lambda u) dx \\ &= \int_{\Omega} (-|\nabla u|^2 + \lambda u^2) dx. \end{aligned}$$

By Theorem 1, we have

$$0 < \int_{\Omega} (-|\nabla u(x, t)|^2 + \lambda u(x, t)^2) dx \leq \frac{|\Omega|}{2(T-t)}$$

for $t \in [t_0, T)$. Hence, $U(x, t) = \frac{u(x, t)}{\|u(\cdot, t)\|_\infty}$ satisfies

$$\int_{\Omega} |\nabla U(x, t)|^2 dx < \lambda \int_{\Omega} U(x, t)^2 dx \quad (3.2)$$

and

$$\lambda \int_{\Omega} U(x, t)^2 dx - \frac{|\Omega|}{2(T-t)\|u(\cdot, t)\|_\infty^2} \leq \int_{\Omega} |\nabla U(x, t)|^2 dx. \quad (3.3)$$

Since if $\delta = 2$ and u is Type 2, then

$$\limsup_{t \nearrow T} (T - t)^{\frac{1}{2}} \|u(\cdot, t)\|_{\infty} = \infty,$$

there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that

$$t_k \nearrow T \text{ and } (T - t_k) \|u(\cdot, t_k)\|_{\infty}^2 \rightarrow \infty \text{ as } k \rightarrow \infty.$$

Furthermore, Lemma 2 implies that

$$0 < \mu^2 |S_{\mu}| \leq \int_{\Omega} U(x, t_k)^2 dx \leq |\Omega| \text{ for any } k. \quad (3.4)$$

Therefore, we have

$$\frac{\int_{\Omega} |\nabla U(x, t_k)|^2 dx}{\int_{\Omega} U(x, t_k)^2 dx} \rightarrow \lambda \text{ as } k \rightarrow \infty. \quad \square$$

3.3. A PROFILE FOR BLOW-UP RATES

We mentioned in Section 1 that there exists “Type 2” solutions of (3.1) for $\delta = 2$ which satisfies

$$\limsup_{t \nearrow T} (T - t)^{\frac{1}{2}} \|u(\cdot, t)\|_{\infty} = \infty \quad (3.5)$$

The following corollary gives another feature for asymptotic behavior of $(T - t)^{\frac{1}{2}} u(x, t)$ as $t \nearrow T$.

Corollary 3. *Let $\delta = 2$ and u be a solution of (3.1) approximated by u_{ε} solving (1.3). Suppose that $u_0 \in C^{\infty}(\Omega) \cap C(\bar{\Omega})$ satisfies (1.1) and $\log u_0 \in L^1(\Omega)$. Then there exists a constant $C > 0$ such that*

$$\int_{\Omega} \log \left[(T - t)^{\frac{1}{2}} u(x, t) \right] dx \leq C$$

for $t \in [t_0, T)$, where C depends on T and u_0 only.

Proof. Since Theorem 1 implies

$$0 < \frac{d}{dt} \int_{\Omega} \log u dx \leq \frac{|\Omega|}{2(T - t)} \text{ for any } t \in [t_0, T),$$

we have

$$\frac{d}{dt} \int_{\Omega} \log \left[(T - t)^{\frac{1}{2}} u(x, t) \right] dx \leq 0 \text{ for any } t \in [t_0, T).$$

Therefore,

$$\int_{\Omega} \log \left[(T - t)^{\frac{1}{2}} u(x, t) \right] dx \leq \int_{\Omega} \log \left[T^{\frac{1}{2}} u_0(x) \right] dx$$

which completes this proof. $\square$

4. THE CASE OF $\delta > 2$

4.1. CONJECTURES FOR $\delta > 2$

For the case of $\delta > 2$, we expect that behavior of blow-up solutions may be a little different from $\delta = 2$. In this section we try to establish some profiles for $\delta > 2$ by numerical computing.

Precisely, we investigated features for asymptotic behavior of blow-up solutions to (1.2) by numerical computing with many kinds of initial data. Note that all of our computations are based on the scheme provided in [5].


For example, we consider the case of $\lambda = 1.0$, $\Omega = (0, 20) \in \mathbb{R}^1$ and

$$u_0(x) = \frac{1}{100} \left(\sin \frac{\pi}{20} x \right) \left(\sin^2 \frac{3\pi}{20} x \right). \quad (4.1)$$


Figure 1 (a) and (b) are behavior of

$$I(t) := \int_{\Omega} u^{2-\delta} dx$$

for $\delta = 2.5$ and $\delta = 4.0$ near the blow-up time, respectively.



(a) Behavior of $I(t)$ for $\delta = 2.5$



(b) Behavior of $I(t)$ for $\delta = 4.0$

Figure 1: Examples of behavior of $I(t) = \int_{\Omega} u^{2-\delta} dx$

They show that $I(t)$ is decreasing near the blow-up time. Since $\frac{d}{dt}I(t) = (2 - \delta) \int_{\Omega} u^{1-\delta} u_t dx$, this observation implies that if $\delta > 2$ then $\int_{\Omega} u^{1-\delta} u_t dx$ should be positive near the blow-up time, that is, u should satisfy the same property as “weak eventual monotonicity” in Remark 4 for the case of $\delta = 2$. We are computing numerically with so many initial data but we have not gotten what does not satisfy “weak eventual monotonicity”.

Remark 5. The property of “weak eventual monotonicity” means that $\int_{\Omega} u^{1-\delta} u_t dx$ should change positive even if it is negative near the initial time $t = 0$, that is, $I(t) = \int_{\Omega} u^{2-\delta} dx$ should be strictly decreasing after a finite time. For instance, Figure 2 is an example of behavior near the initial time of $I(t) = \int_{\Omega} u^{2-\delta} dx$ with the initial data $u_0(x) = K^2 - x^2$, where $\Omega = (-K, K) \in \mathbb{R}^1$, $\delta = 2.5$ and $K = 1.575$.




Figure 2: An example of behavior of $I(t)$ near $t = 0$

Next, we consider the ratio of $E(t)$ to $\left(\int_{\Omega} u^{1-\delta} u_t dx\right)^2$ to discuss another profile for asymptotic behavior near the blow-up time, where $E(t)$ is defined in (2.3) as follows:

$$E(t) = \int_{\Omega} u^{2-\delta} dx \int_{\Omega} u^{-\delta} (u_t)^2 dx - \left(\int_{\Omega} u^{1-\delta} u_t dx\right)^2 > 0.$$


We set

$$\begin{aligned} F(t) &:= E(t) \left(\int_{\Omega} u^{1-\delta} u_t dx\right)^{-2} \\ &= \frac{\int_{\Omega} u^{2-\delta} dx \int_{\Omega} u^{-\delta} (u_t)^2 dx}{\left(\int_{\Omega} u^{1-\delta} u_t dx\right)^2} - 1 \end{aligned}$$


and

$$G(t) := F(t)^{-\frac{\delta}{\delta-2}}.$$

Then we numerically compute $G(t)$ with many kinds of initial data, too. Figure 3 (a) and (b) show the behavior of $G(t)$ near the blow-up time for the case of $\lambda = 1.0$, $\Omega = (0, 20)$ and the initial function (4.1).



(a) Behavior of $G(t)$ for $\delta = 2.5$



(b) Behavior of $G(t)$ for $\delta = 4.0$

Figure 3: Examples of behavior of $G(t) = F(t)^{-\frac{\delta}{\delta-2}}$

They indicate that $G(t)$ should decay as t closed to the blow-up time and the rate is faster than or equal to $O(T-t)$ as $t \nearrow T$. We have not found what does not satisfy this property, too.

From the above numerical observations, we would like to present a conjectures for behavior of $\int_{\Omega} u^{1-\delta} u_t dx$ and $F(t)$ as follows.

Conjectures. Let u be a solution of (1.2) approximated by u_{ε} solving (1.3) and T the blow-up time of u . When $\delta > 2$, there exists $t_0 \in (0, T)$, $C_1 > 0$ and $C_2 > 0$ such that

$$(C1) \quad \int_{\Omega} u^{1-\delta} u_t dx \geq C_1 \text{ for } t \in [t_0, T).$$

$$(C2) \quad F(t) := E(t) \left(\int_{\Omega} u^{1-\delta} u_t dx\right)^{-2} \text{ satisfies}$$

$$F(t) \geq C_2(T-t)^{-\frac{\delta-2}{\delta}} \text{ for } t \in [t_0, T).$$

4.2. THE MAIN THEOREM FOR $\delta > 2$

If two conjectures, (C1) and (C2), are true then we can prove the following theorem which are similar to Theorem 1.

Theorem 2. *Let $\delta > 2$. u be a solution of (1.2) approximated by u_ε solving (1.3). suppose that $u_0 \in C^\infty(\Omega) \cap C(\bar{\Omega})$ satisfies (1.1) and $u_0^{2-\delta} \in L^1(\Omega)$. If (C1) and (C2) hold then*

$$\delta(T-t)^{\frac{2}{\delta}} \int_{\Omega} u^{1-\delta} u_t dx \leq \frac{1}{C_2} \int_{\Omega} u^{2-\delta} dx.$$

for $t \in [t_0, T)$, where $t_0 \in (0, T)$ and $C_2 > 0$ are given in Conjectures, (C1) and (C2).

Remark 6. If $u_0^{2-\delta} \in L^1(\Omega)$ then $u(\cdot, t)^{2-\delta} \in L^1(\Omega)$ for any $t \in (0, T)$.

Proof. Let $\{u_n\}_{n \in \mathbb{N}}$ be defined in Section 2. Then it is verified by Fatou's Lemma and (C2) that

$$\begin{aligned} & \int_{\Omega} u^{2-\delta} dx \cdot \liminf_{n \rightarrow \infty} \int_{\Omega} u_n^{-\delta} \left((u_n)_t \right)^2 dx \\ &= \int_{\Omega} u^{2-\delta} dx \cdot \liminf_{n \rightarrow \infty} \int_{\Omega} (\Delta u_n + \lambda u_n) (u_n)_t dx \\ &\geq \int_{\Omega} u^{2-\delta} dx \int_{\Omega} (\Delta u + \lambda u) u_t dx \\ &\geq \int_{\Omega} u^{2-\delta} dx \int_{\Omega} u^{-\delta} (u_t)^2 dx \\ &\geq \left(C_2(T-t)^{-\frac{\delta-2}{\delta}} + 1 \right) \left(\int_{\Omega} u^{1-\delta} u_t dx \right)^2 \end{aligned}$$

for $t \in [t_0, T)$. Besides,

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\Omega} u_n^{1-\delta} (u_n)_t dx &= \lim_{n \rightarrow \infty} \int_{\Omega} u_n (\Delta u_n + \lambda u_n) dx \\ &= \int_{\Omega} u (\Delta u + \lambda u) dx \\ &= \int_{\Omega} u^{1-\delta} u_t dx \end{aligned}$$

because $u_n(\cdot, t) \rightarrow u(\cdot, t)$ uniformly in $\bar{\Omega}$ and $\Delta u_n \rightharpoonup \Delta u$ weakly in $C^*(\Omega)$ as $n \rightarrow \infty$ for any $t \in (0, T)$. Hence, for $\eta \in (0, C_1^2)$, there exists $N \in \mathbb{N}$ such that

$$\begin{aligned} & \int_{\Omega} u^{2-\delta} dx \left(\int_{\Omega} u_n^{-\delta} \left((u_n)_t \right)^2 dx + \eta \right) \\ &\geq \left(C_2(T-t)^{-\frac{\delta-2}{\delta}} + 1 \right) \left[\left(\int_{\Omega} u_n^{1-\delta} (u_n)_t dx \right)^2 - \eta \right] > 0 \end{aligned}$$

and $\left(\int_{\Omega} u_n^{1-\delta} (u_n)_t dx \right)^2 > \left(\int_{\Omega} u^{1-\delta} u_t dx \right)^2 - \eta > C_1^2 - \eta$ for $n \geq N$ and $t \in [t_0, T)$. Furthermore, by (2.4) and (C1),

we have

$$\begin{aligned} & \frac{2\eta}{C_1^2 - \eta} \left(C_2(T-t)^{-\frac{\delta-2}{\delta}} + 1 + \int_{\Omega} u(x, t_0)^{2-\delta} dx \right) \\ &\geq 2\eta \left(C_2(T-t)^{-\frac{\delta-2}{\delta}} + 1 + \int_{\Omega} u(x, t)^{2-\delta} dx \right) \\ &\quad \times \left(\int_{\Omega} u_n^{1-\delta} (u_n)_t dx \right)^{-2} \\ &\geq 2C_2(T-t)^{-\frac{\delta-2}{\delta}} \\ &\quad + \int_{\Omega} u^{2-\delta} dx \left[\left(\int_{\Omega} u_n^{1-\delta} (u_n)_t dx \right)^{-1} \right] + 2 \end{aligned}$$

for $n \geq N$ and $t \in [t_0, T)$. It is verified by integral by parts that if $t_0 \leq t < s < T$ and $\delta > 2$ then

$$\begin{aligned} & \frac{\eta}{C_1^2 - \eta} \left[-\delta C_2(T-s)^{\frac{2}{\delta}} + \delta C(T-t)^{\frac{2}{\delta}} \right. \\ &\quad \left. + 2 \left(1 + \int_{\Omega} u(x, t_0)^{2-\delta} dx \right) (s-t) \right] \\ &\geq \delta C_2 \left[-(T-s)^{\frac{2}{\delta}} + (T-t)^{\frac{2}{\delta}} \right] \\ &\quad + \int_{\Omega} u(x, s)^{2-\delta} dx \left(\int_{\Omega} u_n(x, s)^{1-\delta} (u_n)_t(x, s) dx \right)^{-1} \\ &\quad - \int_{\Omega} u(x, t)^{2-\delta} dx \left(\int_{\Omega} u_n(x, t)^{1-\delta} (u_n)_t(x, t) dx \right)^{-1} \\ &\quad + (\delta-2)(s-t) - \frac{\eta(\delta-2)}{\sqrt{C_1^2 - \eta}}(s-t) + 2(s-t) \\ &\geq \delta C_2 \left[-(T-s)^{\frac{2}{\delta}} + (T-t)^{\frac{2}{\delta}} \right] \\ &\quad - \int_{\Omega} u(x, t)^{2-\delta} dx \left(\int_{\Omega} u_n(x, t)^{1-\delta} (u_n)_t(x, t) dx \right)^{-1} \\ &\quad + \delta(s-t) - \frac{\eta(\delta-2)}{\sqrt{C_1^2 - \eta}}(s-t). \end{aligned}$$

Letting $s \nearrow T$, $n \rightarrow \infty$ and $\eta \searrow 0$, we have

$$\begin{aligned} 0 &\geq \delta C_2(T-t)^{\frac{2}{\delta}} - \int_{\Omega} u^{2-\delta} dx \left(\int_{\Omega} u^{1-\delta} u_t dx \right)^{-1} \\ &\quad + \delta(T-t) \\ &\geq \delta C_2(T-t)^{\frac{2}{\delta}} - \int_{\Omega} u^{2-\delta} dx \left(\int_{\Omega} u^{1-\delta} u_t dx \right)^{-1} \end{aligned}$$

for $t \in [t_0, T)$. Therefore, it holds that

$$\delta(T-t)^{\frac{2}{\delta}} \int_{\Omega} u^{1-\delta} u_t dx \leq \frac{1}{C_2} \int_{\Omega} u^{2-\delta} dx$$

which completes this proof. $\square$

Moreover, we can also get the following corollary in the same manner of corollary 2.

Corollary 4. *Let $\delta > 2$ and u be a “Type 2” solution of (1.2) approximated by u_ε solving (1.3). Suppose that*

$u_0 \in C^\infty(\Omega) \cap C(\bar{\Omega})$ satisfies (1.1) and $u_0^{2-\delta} \in L^1(\Omega)$. If (C1) and (C2) holds then there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that $t_k \nearrow T$ as $k \rightarrow \infty$ and $U(x, t) := \frac{u(x, t)}{\|u(\cdot, t)\|_\infty}$ satisfies

$$\frac{\int_{\Omega} |\nabla U(x, t_k)|^2 dx}{\int_{\Omega} U(x, t_k)^2 dx} \rightarrow \lambda \quad \text{as } k \rightarrow \infty,$$

Proof. Since

$$\begin{aligned} \int_{\Omega} u^{1-\delta} u_t dx &= \int_{\Omega} u(\Delta u + \lambda u) dx \\ &= \int_{\Omega} (-|\nabla u|^2 + \lambda u^2) dx, \end{aligned}$$

Theorem 2 implies

$$\int_{\Omega} u(\Delta u + \lambda u) dx \leq \frac{1}{\delta C_2(T-t)^{\frac{2}{\delta}}} \int_{\Omega} u^{2-\delta} dx$$

and then

$$\begin{aligned} \int_{\Omega} (-|\nabla u(x, t)|^2 + \lambda u(x, t)^2) dx \\ \leq \frac{1}{\delta C_2(T-t)^{\frac{2}{\delta}}} \int_{\Omega} u^{2-\delta} dx \end{aligned}$$

for $t \in [t_0, T)$. Hence, $U(x, t) = \frac{u(x, t)}{\|u(\cdot, t)\|_\infty}$ satisfies

$$\begin{aligned} \lambda \int_{\Omega} U(x, t)^2 dx - \frac{1}{\delta C_2(T-t)^{\frac{2}{\delta}} \|u(\cdot, t)\|_\infty^2} \int_{\Omega} u^{2-\delta} dx \\ \leq \int_{\Omega} |\nabla U(x, t)|^2 dx \end{aligned}$$

for $t \in [t_0, T)$. Besides, (C1) implies $\int_{\Omega} u^{2-\delta} dx$ is decreasing in $[t_0, T)$ and

$$\int_{\Omega} |\nabla U(x, t)|^2 dx < \lambda \int_{\Omega} U(x, t)^2 dx \quad \text{for } t \in [t_0, T).$$

Since u is Type 2, there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that

$$t_k \nearrow T \text{ and } (T - t_k)^{\frac{2}{\delta}} \|u(\cdot, t_k)\|_\infty^2 \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

Therefore, we have

$$\frac{\int_{\Omega} |\nabla U(x, t_k)|^2 dx}{\int_{\Omega} U(x, t_k)^2 dx} \rightarrow \lambda \quad \text{as } k \rightarrow \infty. \quad \square$$

Remark 7. If u_0 satisfies $\Delta u_0 + \lambda u_0 \geq 0$ in Ω , then it also holds for $\delta > 2$ that there exists a sequence $\{t_k\}_{k \in \mathbb{N}}$ such that $t_k \nearrow T$ as $k \rightarrow \infty$ and U satisfies

$$\Delta U(x, t_k) + \lambda U(x, t_k) \rightarrow 0 \text{ or } U(x, t_k) \rightarrow 0$$

as $k \rightarrow \infty$ almost all of $x \in \Omega$. We can regard Corollary 4 as a generalization of this profile.

ACKNOWLEDGEMENTS

We would like to thank referees very much for their kind comments and valuable suggestions to improve our paper.

REFERENCES

- [1] Anada, K., Fukuda, I. and Tsutsumi, M.: Regional blow-up and decay of solutions to the Initial-Boundary value problem for $u_t = uu_{xx} - \gamma(u_x)^2 + ku^2$, *Funkcialaj Ekvacioj* **39** (1996) 363–387.
- [2] Anada, K. and Ishiwata, T.: Classification of solutions of a parabolic differential equation by blow-up rates, *Proc. 8th World-Conference on Systemics, Cybernetics and Informatics* Vol. XV (2004) 153–156.
- [3] Angenent, S.: On the formation of singularities in the curve shortening flow, *J. Diff. Geo.* **33** (1991) 601–633.
- [4] Friedman, A. and McLeod, B.: Blow-up of solutions of nonlinear degenerate parabolic equations, *Arch. Rational Mech. Anal.* **96** (1987) 55–80.
- [5] Ishiwata, T. and Tsutsumi, M.: A numerical study of blow-up solutions to $u_t = u^\delta(\Delta u + \mu u)$, *Gakuto International Series, Mathematical Sciences and Applications* **10** (1997), *Nonlinear Waves*, 195–207.
- [6] Tsutsumi, M. and Ishiwata, T.: Regional blow-up of solutions to the initial-boundary value problems for $u_t = u^\delta(\Delta u + u)$, *Proc. Roy. Soc. Edinburgh Sect. A* **127** (1997) 871–887.
- [7] Wiegner, M.: Blow-up for solutions of some degenerate parabolic equations, *Diff. Int. Eq.* **7** (1994) 1641–1647.
- [8] Wiegner, M.: A degenerate diffusion equation with a nonlinear source term, *Nonlinear Anal. T.M.A.* **28** (1997) 1977–1995.
- [9] Winkler, M.: Blow-up of solutions to a degenerate parabolic equation not in divergence form, *J. Diff. Eq.* **192** (2003) 445–474.
- [10] Winkler, M.: Blow-up in a degenerate parabolic equation, *Indiana Univ. Math. J.* **53** (2004) 1415–1442.

Koichi Anada

Waseda University Senior High School, 3-31-1 Kamishakujii Nerima-ku, Tokyo 177-0044, JAPAN.

E-mail: anada-koichi(at)waseda.jp

Tetsuya Ishiwata

Department of Mathematical Sciences, Shibaura Institute of Technology, 307 Fukasaku, Minuma-ku, Saitama 337-8570, JAPAN.

E-mail: tisiwata(at)shibaura-it.ac.jp