On the maximal number of exceptional values of Gauss maps for various classes of surfaces

川上, 裕
九州大学大学院数理学府

<http://hdl.handle.net/2324/21978>

出版情報 : Mathematische Zeitschrift. 274 (3/4), pp.1249-1260, 2012-12. Springer-Verlag
バージョン :
権利関係 : (C) Springer-Verlag Berlin Heidelberg 2012

MI Preprint Series

**Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub**

On the maximal number of exceptional values of Gauss maps for various classes of surfaces

Yu Kawakami

MI 2012-5

(Received May 22, 2012)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

ON THE MAXIMAL NUMBER OF EXCEPTIONAL VALUES OF GAUSS MAPS FOR VARIOUS CLASSES OF SURFACES

YU KAWAKAMI

ABSTRACT. The main goal of this paper is to reveal the geometric meaning of the maximal number of exceptional values of Gauss maps for several classes of immersed surfaces in space forms, for example, complete minimal surfaces in the Euclidean three-space, weakly complete improper affine spheres in the affine three-space and weakly complete flat surfaces in the hyperbolic three-space. For this purpose, we give an effective curvature bound for a specified conformal metric on an open Riemann surface.

INTRODUCTION

The geometric nature of value distribution theory of complex analytic mappings is well-known. One of the most elegant results of the theory is the geometric meaning of the precise maximum “2” for the number of exceptional values of nonconstant meromorphic functions on the complex plane $\mathbf{C}$. In fact, Ahlfors [1] and Chern [5] showed that the maximal number of exceptional values of nonconstant holomorphic maps from $\mathbf{C}$ to a closed Riemann surface coincides with the Euler number of the closed Riemann surface by using Nevanlinna theory (see also [21], [29] and [33]). In particular, for nonconstant meromorphic functions on $\mathbf{C}$, the geometric meaning of the maximal number “2” of exceptional values is the Euler number of the Riemann sphere.

On the other hand, global properties of the Gauss map of complete minimal surfaces in the Euclidean three-space $\mathbf{R}^3$ are closely related to value-distribution-theoretic properties of meromorphic functions on $\mathbf{C}$. In particular, Fujimoto [11] proved that the precise maximum for the number of exceptional values of the Gauss map of a nonflat complete minimal surface in $\mathbf{R}^3$ is “4”, and Osserman [30] showed that the Gauss map of a nonflat algebraic minimal surface can omit at most 3 values (by an algebraic minimal surface, we mean a complete minimal surface with finite total curvature). Recently, the author, Kobayashi and Miyaoka [18] gave an effective upper bound for the number of exceptional values of the Gauss map for a special class of complete minimal surfaces that includes algebraic minimal surfaces (this class is called the pseudo-algebraic minimal surfaces). This also provided a geometric meaning for the Fujimoto and Osserman results for this

2010 *Mathematics Subject Classification.* Primary 30D35, 53C42 ; Secondary 30F45, 53A10, 53A15.

Key words and phrases. Gauss map, exceptional value, conformal metric, minimal surface, front, Bernstein type theorem.

Partly supported by the Grant-in-Aid for Young Scientists (B) No. 24740044, Japan Society for the Promotion of Science.

class, because the upper bound is described in terms of geometric invariants. However, from [18] it was still not possible to understand the geometric meaning for general class.

The author also investigated value-distribution-theoretic properties of Gauss maps for several classes of surfaces which may admit singularities. For instance, by refining the Fujimoto analytic argument, the author and Nakajo [19] showed that the maximal number of exceptional values of the Lagrangian Gauss map of weakly complete improper affine fronts in the affine three-space $\mathbf{R}^3$ is “3”. As an application of this result, a simple proof of the parametric affine Bernstein theorem for improper affine spheres in $\mathbf{R}^3$ was provided. Moreover, the authors [17, 19] proved similar results for flat fronts in the hyperbolic three-space $\mathbf{H}^3$.

The aim of this paper is to reveal the geometric meaning of the precise maximum for the number of exceptional values of Gauss maps for these classes of surfaces. The paper is organized as follows: In Section 1, we give a curvature bound for the conformal metric $ds^2 = (1 + |g|^2)^m |\omega|^2$ on an open Riemann surface Σ , where ω is a holomorphic 1-form and g is a meromorphic function on Σ (Theorem 1.1). The proof is given in Section 2. As a corollary of this theorem, we prove that the precise maximum for the number of exceptional values of the nonconstant meromorphic function g on Σ with the complete conformal metric ds^2 is “ $m + 2$ ” (Corollary 1.2 and Proposition 1.4). We note that the geometric meaning of the “2” in “ $m + 2$ ” is the Euler number of the Riemann sphere (Remark 1.3). In Section 3, we give some applications of the main results. In particular, we give the geometric meaning of the maximal number of exceptional values of Gauss maps for several classes of immersed surfaces in space forms. For instance, the induced metric from $\mathbf{R}^3$ on complete minimal surfaces is $ds^2 = (1 + |g|^2)^2 |\omega|^2$ (i.e. $m = 2$), thereby the maximal number of exceptional values of the Gauss map g of nonflat complete minimal surfaces in $\mathbf{R}^3$ is “4 (= 2 + 2)”. On the other hand, for the Lagrangian Gauss map ν of weakly complete improper affine fronts, since ν is meromorphic, dG is holomorphic and the complete metric is $d\tau^2 = 2(1 + |\nu|^2) |dG|^2$ (i.e. $m = 1$), the maximal number of exceptional values of the Lagrangian Gauss map of weakly complete improper affine fronts in $\mathbf{R}^3$ is “3 (= 1 + 2)”.

Finally, the author would like to thank Professors Junjiro Noguchi, Wayne Rossman, Masaaki Umehara and Kotaro Yamada for their useful advice. In addition, the author would like to express his thanks to Professors Ryoichi Kobayashi, Masatoshi Kokubu, Miyuki Koiso and Reiko Miyaoka for their encouragement of this study.

1. MAIN RESULTS

Now we state the main theorem of this paper.

THEOREM 1.1. *Let Σ be an open Riemann surface with the conformal metric*

$$(1) \quad ds^2 = (1 + |g|^2)^m |\omega|^2,$$

where ω is a holomorphic 1-form, g is a meromorphic function on Σ , and $m \in \mathbf{N}$. Suppose that g omits $q \geq m+3$ distinct values. Then there exists a positive constant C , depending on m and the set of exceptional values, but not the surface Σ , such that for all $p \in \Sigma$ we have

$$(2) \quad |K_{ds^2}(p)|^{1/2} \leq \frac{C}{d(p)},$$

where $K_{ds^2}(p)$ is the Gaussian curvature of the metric ds^2 at p and $d(p)$ is the geodesic distance from p to the boundary of Σ , that is, the infimum of the lengths of the divergent curves in Σ emanating from p .

As a corollary of Theorem 1.1, we give the following Picard-type theorem for the meromorphic function g on Σ with the complete conformal metric $ds^2 = (1 + |g|^2)^m |\omega|^2$.

COROLLARY 1.2. *Let Σ be an open Riemann surface with the conformal metric given by (1). If the metric ds^2 is complete and the meromorphic function g is nonconstant, then g can omit at most $m+2$ distinct values.*

PROOF. By way of contradiction, assume that g omits $m+3$ distinct values. If ds^2 is complete, then we may set $d(p) = \infty$ for all $p \in \Sigma$. By virtue of Theorem 1.1, $K_{ds^2} \equiv 0$ on Σ . On the other hand, the Gaussian curvature of the metric ds^2 is given by

$$(3) \quad K_{ds^2} = -\frac{2m|g'_z|^2}{(1 + |g|^2)^{m+2} |\hat{\omega}_z|^2},$$

where $\omega = \hat{\omega}_z dz$ and $g'_z = dg/dz$. Thus $K_{ds^2} \equiv 0$ if and only if g is constant. This contradicts the assumption that g is nonconstant. $\square$

REMARK 1.3. The geometric meaning of the “2” in “ $m+2$ ” is the Euler number of the Riemann sphere. Indeed, if $m=0$ then the metric $ds^2 = (1 + |g|^2)^0 |\omega|^2 = |\omega|^2$ is flat and complete on Σ . We thus may assume that g is a meromorphic function on $\mathbf{C}$ because g is replaced by $g \circ \pi$, where $\pi: \mathbf{C} \rightarrow \Sigma$ is a holomorphic universal covering map. On the other hand, Ahlfors [1] and Chern [5] showed that the best possible upper bound “2” of the number of exceptional values of nonconstant meromorphic functions on $\mathbf{C}$ coincides with the Euler number of the Riemann sphere. Hence we get the conclusion.

Corollary 1.2 is optimal because there exist the following examples.

PROPOSITION 1.4. *Let Σ be either the complex plane punctured at $q-1$ distinct points $\alpha_1, \dots, \alpha_{q-1}$ or the universal cover of that punctured plane. We set*

$$\omega = \frac{dz}{\prod_{i=1}^{q-1} (z - \alpha_i)}, \quad g = z.$$

Then g omits q distinct values and the metric $ds^2 = (1 + |g|^2)^m |\omega|^2$ is complete if and only if $q \leq m+2$. In particular, there exist examples whose metric ds^2 is complete and g omits $m+2$ distinct values.

PROOF. We can easily show that g omits the q distinct values $\alpha_1, \dots, \alpha_{q-1}$ and ∞ on Σ . A divergent curve Γ in Σ must tend to one of the points $\alpha_1, \dots, \alpha_{q-1}$ or ∞ . Thus we have

$$\int_{\Gamma} ds = \int_{\Gamma} (1 + |g|^2)^{m/2} |\omega| = \int_{\Gamma} \frac{(1 + |z|^2)^{m/2}}{\prod_{i=1}^{q-1} |z - \alpha_i|} |dz| = \infty,$$

when $q \leq m + 2$. □

2. PROOF OF THE MAIN THEOREM

We first recall the notion of chordal distance between two distinct values in the Riemann sphere $\mathbf{C} \cup \{\infty\}$. For two distinct values $\alpha, \beta \in \mathbf{C} \cup \{\infty\}$, we set

$$|\alpha, \beta| := \frac{|\alpha - \beta|}{\sqrt{1 + |\alpha|^2} \sqrt{1 + |\beta|^2}}$$

if $\alpha \neq \infty$ and $\beta \neq \infty$, and $|\alpha, \infty| = |\infty, \alpha| := 1/\sqrt{1 + |\alpha|^2}$. We note that, if we take $v_1, v_2 \in \mathbf{S}^2$ with $\alpha = \varpi(v_1)$ and $\beta = \varpi(v_2)$, we have that $|\alpha, \beta|$ is a half of the chordal distance between v_1 and v_2 , where ϖ denotes the stereographic projection of the 2-sphere $\mathbf{S}^2$ onto $\mathbf{C} \cup \{\infty\}$.

Before proceeding to the proof of Theorem 1.1, we recall two function-theoretical lemmas.

LEMMA 2.1. [13, (8.12) on page 136] *Let g be a nonconstant meromorphic function on $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ ($0 < R \leq +\infty$) which omits q values $\alpha_1, \dots, \alpha_q$. If $q > 2$, then for each positive η with $\eta < (q - 2)/q$, then there exists a positive constant C' , depending on m, q and $L := \min_{i < j} |\alpha_i, \alpha_j|$, such that*

$$(4) \quad \frac{|g'_z|}{(1 + |g|^2) \prod_{j=1}^q |g, \alpha_j|^{1-\eta}} \leq C' \frac{R}{R^2 - |z|^2}.$$

LEMMA 2.2. [12, Lemma 1.6.7] *Let $d\sigma^2$ be a conformal flat metric on an open Riemann surface Σ . Then, for each point $p \in \Sigma$, there exists a local diffeomorphism Φ of a disk $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ ($0 < R \leq +\infty$) onto an open neighborhood of p with $\Phi(0) = p$ such that Φ is a local isometry, that is, the pull-back $\Phi^*(d\sigma^2)$ is equal to the standard Euclidean metric ds_{Euc}^2 on Δ_R and, for a point a_0 with $|a_0| = 1$, the Φ -image Γ_{a_0} of the curve $L_{a_0} = \{w := a_0 s; 0 < s < R\}$ is divergent in Σ .*

Proof of Theorem 1.1. We may assume that $\alpha_q = \infty$. We choose a positive number η with

$$\frac{q - 2(m + 1)}{q} < \eta < \frac{q - (m + 2)}{q}$$

and set $\lambda := m/(q - 2 - q\eta)$. Since $q \geq m + 3$, then $0 < \lambda < 1$ holds.

Now we define a new metric

$$(5) \quad d\sigma^2 = |\hat{\omega}_z|^{2/(1-\lambda)} \left(\frac{1}{|g'_z|} \prod_{j=1}^{q-1} \left(\frac{|g - \alpha_j|}{\sqrt{1 + |\alpha_j|^2}} \right)^{1-\eta} \right)^{2\lambda/(1-\lambda)} |dz|^2$$

on the set $\Gamma' := \{p \in \Sigma; g'_z(p) \neq 0\}$, where $\omega = \hat{\omega}_z dz$ and $g'_z = dg/dz$ with respect to the local complex coordinate z . Take a point $p \in \Sigma'$. Since the metric $d\sigma^2$ is flat, by Lemma 2.2, there exists a local isometry Φ satisfying $\Phi(0) = p$ from a disk $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ ($0 < R \leq +\infty$) with the standard Euclidean metric ds_{Euc}^2 onto an open neighborhood of $p \in \Sigma'$ with the metric $d\sigma^2$, such that, for a point a_0 with $|a_0| = 1$, the Φ -image Γ_{a_0} of the curve $L_{a_0} = \{w := a_0 s; 0 < s < R\}$ is divergent in Σ' . For brevity, we denote the function $g \circ \Phi$ on Δ_R by g in the following. By Lemma 2.1, we get

$$(6) \quad R \leq C' \frac{1 + |g(0)|^2}{|g'(0)|} \prod_{j=1}^q |g(0), \alpha_j|^{1-\eta} < +\infty.$$

Hence

$$L_{d\sigma}(\Gamma_{a_0}) = \int_{\Gamma_{a_0}} d\sigma = R < +\infty,$$

where $L_{d\sigma}(\Gamma_{a_0})$ denotes the length of Γ_{a_0} with respect to the metric $d\sigma^2$. We assume that the Φ -image Γ_{a_0} tends to a point $p_0 \in \Sigma \setminus \Sigma'$ as $s \rightarrow R$. Taking a local complex coordinate ζ in a neighborhood of p_0 with $\zeta(p_0) = 0$, we can write

$$d\sigma^2 = |\zeta|^{-2\lambda/(1-\lambda)} w |d\zeta|^2$$

for some positive smooth function w . Since $\lambda/(1-\lambda) > 1$, we have

$$R = \int_{\Gamma_{a_0}} d\sigma \geq \tilde{C} \int_{\Gamma_{a_0}} \frac{|d\zeta|}{|\zeta|^{\lambda/(1-\lambda)}} = +\infty,$$

which contradicts (6). Thus Γ_{a_0} diverges outside any compact subset of Σ as $s \rightarrow R$.

Since $d\sigma^2 = |dz|^2$, we obtain by (5) that

$$(7) \quad |\hat{\omega}_z| = \left(|g'_z| \prod_{j=1}^{q-1} \left(\frac{\sqrt{1 + |\alpha_j|^2}}{|g - \alpha_j|} \right)^{1-\eta} \right)^\lambda.$$

By Lemma 2.1, we have

$$\begin{aligned} \Phi^* ds &= |\hat{\omega}_z| (1 + |g|^2)^{m/2} |dz| \\ &= \left(|g'_z| (1 + |g|^2)^{m/2\lambda} \prod_{j=1}^{q-1} \left(\frac{\sqrt{1 + |\alpha_j|^2}}{|g - \alpha_j|} \right)^{1-\eta} \right)^\lambda |dz| \\ &= \left(\frac{|g'_z|}{(1 + |g|^2) \prod_{j=1}^q |g, \alpha_j|^{1-\eta}} \right)^\lambda |dz| \\ &\leq (C')^\lambda \left(\frac{R}{R^2 - |z|^2} \right)^\lambda |dz|. \end{aligned}$$

Thus we have

$$d(p) \leq \int_{\Gamma_{a_0}} ds = \int_{L_{a_0}} \Phi^* ds \leq (C')^\lambda \int_{L_{a_0}} \left(\frac{R}{R^2 - |z|^2} \right)^\lambda |dz| \leq (C')^\lambda \frac{R^{1-\lambda}}{1-\lambda} (< +\infty)$$

because $0 < \lambda < 1$. Moreover, by (6), we get that

$$(8) \quad d(p) \leq \frac{C'}{1-\lambda} \left(\frac{1+|g(0)|^2}{|g'(0)|} \prod_{j=1}^q |g(0), \alpha_j|^{1-\eta} \right)^{1-\lambda}.$$

On the other hand, the Gaussian curvature K_{ds^2} of the metric $ds^2 = (1+|g|^2)^m |\omega|^2$ is given by

$$K_{ds^2} = -\frac{2m|g'_z|^2}{(1+|g|^2)^{m+2} |\hat{\omega}_z|^2}.$$

Thus, by (7), we also get that

$$(9) \quad |K_{ds^2}|^{1/2} = \sqrt{2m} \left(\frac{|g'_z|}{1+|g|^2} \prod_{j=1}^q |g - \alpha_j|^{1-\eta} \right)^{1-\lambda}.$$

Since $|g, \alpha_j| \leq 1$ for each j , we obtain that

$$(10) \quad |K_{ds^2}|^{1/2} d(p) \leq \frac{\sqrt{2m}C'}{1-\lambda} =: C.$$

By the definitions of C' and λ , we see that C is positive and depends on m , q and $L := \min_{i < j} |\alpha_i, \alpha_j|$. $\square$

3. APPLICATIONS

In this section, we give several applications of our main results.

3.1. Gauss map of minimal surfaces in the Euclidean 3-space. We briefly recall some basic facts of minimal surfaces in $\mathbf{R}^3$. Details can be found, for example, in [12] and [31]. Let $X = (x^1, x^2, x^3): \Sigma \rightarrow \mathbf{R}^3$ be an oriented minimal surface in $\mathbf{R}^3$. By associating a local complex coordinate $z = u + \sqrt{-1}v$ with each positive isothermal coordinate system (u, v) , Σ is considered as a Riemann surface whose conformal metric is the induced metric ds^2 from $\mathbf{R}^3$. Then

$$(11) \quad \Delta_{ds^2} X = 0$$

holds, that is, each coordinate function x^i is harmonic. With respect to the local complex coordinate $z = u + \sqrt{-1}v$ of the surface, (11) is given by

$$(12) \quad \bar{\partial} \partial X = 0,$$

where $\partial = (\partial/\partial u - \sqrt{-1}\partial/\partial v)/2$ and $\bar{\partial} = (\partial/\partial u + \sqrt{-1}\partial/\partial v)/2$. Hence each $\phi_i := \partial x^i dz$ ($i = 1, 2, 3$) is a holomorphic 1-form on Σ . If we set

$$(13) \quad \omega = \phi_1 - \sqrt{-1}\phi_2, \quad g = \frac{\phi_3}{\phi_1 - \sqrt{-1}\phi_2},$$

then ω is a holomorphic 1-form and g is a meromorphic function on Σ . Moreover, the function g coincides with the composition of the Gauss map and the stereographic projection from $\mathbf{S}^2$ onto $\mathbf{C} \cup \{\infty\}$, and the induced metric ds^2 is given by

$$(14) \quad ds^2 = (1 + |g|^2)^2 |\omega|^2.$$

Applying Theorem 1.1 to the metric ds^2 , we can show the Fujimoto theorem for the Gauss map of minimal surfaces in $\mathbf{R}^3$.

THEOREM 3.1. [11, Theorem I and Corollary 3.4] *Let $X: \Sigma \rightarrow \mathbf{R}^3$ be an oriented minimal surface whose Gauss map $g: \Sigma \rightarrow \mathbf{C} \cup \{\infty\}$ omits more than 4 ($= 2 + 2$) distinct values. Then there exists a positive constant C depending on the set of exceptional values, but not the surface, such that for all $p \in \Sigma$ the inequality (2) holds. In particular, the Gauss map of a nonflat complete minimal surface in $\mathbf{R}^3$ can omit at most 4 ($= 2 + 2$) values.*

3.2. Lorentzian Gauss map of maxfaces in the Lorentz-Minkowski 3-space.

Maximal surfaces in the Lorentz-Minkowski 3-space $\mathbf{R}_1^3$ are closely related to minimal surfaces in $\mathbf{R}^3$. In this subsection we treat maximal surfaces with some admissible singularities, called *maxfaces*, as introduced by Umehara and Yamada [36]. We remark that maxfaces, non-branched generalized maximal surfaces in the sense of [9] and non-branched generalized maximal maps in the sense of [15] are all the same class of maximal surfaces. The Lorentz-Minkowski 3-space $\mathbf{R}_1^3$ is the affine 3-space $\mathbf{R}^3$ with the inner product

$$\langle , \rangle = -(dx^1)^2 + (dx^2)^2 + (dx^3)^2,$$

where (x^1, x^2, x^3) is the canonical coordinate system of $\mathbf{R}^3$. We consider a fibration

$$p_L: \mathbf{C}^3 \ni (\zeta^1, \zeta^2, \zeta^3) \mapsto \operatorname{Re}(-\sqrt{-1}\zeta^3, \zeta^1, \zeta^2) \in \mathbf{R}_1^3.$$

The projection of null holomorphic immersions into $\mathbf{R}_1^3$ by p_L gives maxfaces. Here, a holomorphic map $F = (F_1, F_2, F_3): \Sigma \rightarrow \mathbf{C}^3$ is called *null* if $\{(F_1)'_z\}^2 + \{(F_2)'_z\}^2 + \{(F_3)'_z\}^2$ vanishes identically, where $' = d/dz$ denotes the derivative with respect to a local complex coordinate z of Σ . For maxfaces, an analogue of the Enneper-Weierstrass representation formula is known (see also [20]).

THEOREM 3.2. [36, Theorem 2.6] *Let Σ be a Riemann surface and (g, ω) a pair consisting of a meromorphic function and a holomorphic 1-form on Σ such that*

$$(15) \quad d\sigma^2 := (1 + |g|^2)^2 |\omega|^2$$

gives a (positive definite) Riemannian metric on Σ , and $|g|$ is not identically 1. Assume that

$$\operatorname{Re} \int_{\gamma} (-2g, 1 + g^2, \sqrt{-1}(1 - g^2)) \omega = 0$$

for all loops γ in Σ . Then

$$(16) \quad f = \operatorname{Re} \int_{z_0}^z (-2g, 1 + g^2, \sqrt{-1}(1 - g^2)) \omega$$

is well-defined on Σ and gives a maxface in $\mathbf{R}_1^3$, where $z_0 \in \Sigma$ is a base point. Moreover, all maxfaces are obtained in this manner. The induced metric $ds^2 := f^*\langle \cdot, \cdot \rangle$ is given by

$$ds^2 = (1 - |g|^2)^2 |\omega|^2,$$

and the point $p \in \Sigma$ is a singular point of f if and only if $|g(p)| = 1$.

We call g the *Lorentzian Gauss map* of f . If f has no singularities, then g coincides with the composition of the Gauss map (i.e., (Lorentzian) unit normal vector) $n: \Sigma \rightarrow \mathbf{H}_\pm^2$ into the upper or lower connected component of the two-sheet hyperboloid $\mathbf{H}_\pm^2 = \mathbf{H}_+^2 \cup \mathbf{H}_-^2$ in $\mathbf{R}_1^3$, where

$$\begin{aligned} \mathbf{H}_+^2 &:= \{n = (n^1, n^2, n^3) \in \mathbf{R}_1^3; \langle n, n \rangle = -1, n^1 > 0\}, \\ \mathbf{H}_-^2 &:= \{n = (n^1, n^2, n^3) \in \mathbf{R}_1^3; \langle n, n \rangle = -1, n^1 < 0\}, \end{aligned}$$

and the stereographic projection from the north pole $(0, 0, 1)$ of the hyperboloid onto the Riemann sphere $\mathbf{C} \cup \{\infty\}$ (see [36, Section 1]). A maxface is said to be *weakly complete* if the metric $d\sigma^2$ as in (15) is complete. We note that $(1/2)d\sigma^2$ coincides with the pull-back of the standard metric on $\mathbf{C}^3$ by the null holomorphic immersion of f (see [36, Section 2]).

Applying Theorem 1.1 to the metric $d\sigma^2$, we can get the following theorem.

THEOREM 3.3. *Let $f: \Sigma \rightarrow \mathbf{R}_1^3$ be a maxface whose Lorentzian Gauss map $g: \Sigma \rightarrow \mathbf{C} \cup \{\infty\}$ omits more than 4 ($= 2 + 2$) distinct values. Then there exists a positive constant C depending on the set of exceptional values, but not Σ , such that for all $p \in \Sigma$ we have*

$$|K_{d\sigma^2}(p)|^{1/2} \leq \frac{C}{d(p)},$$

where $K_{d\sigma^2}(p)$ is the Gaussian curvature of the metric $d\sigma^2$ at p and $d(p)$ is the geodesic distance from p to the boundary of Σ . In particular, the Lorentzian Gauss map of a nonflat weakly complete maxface in $\mathbf{R}_1^3$ can omit at most 4 ($= 2 + 2$) values.

As a corollary of this result, we give a simple new proof of the Calabi-Bernstein theorem ([4], [7]) for maximal space-like surfaces in $\mathbf{R}_1^3$ from the viewpoint of value-distribution-theoretic properties of the Lorentzian Gauss map. We remark that Alías and Palmer [2], Estudillo and Romero [8, 9, 10], Osamu Kobayashi [20], Romero [32] and Umehara and Yamada [36] have approached this theorem from other viewpoints.

COROLLARY 3.4. *Any complete maximal space-like surface in $\mathbf{R}_1^3$ must be a plane.*

PROOF. Since a maximal space-like surface has no singularities, the complement of the image of g contains at least the set $\{|g| = 1\} \subset \mathbf{C} \cup \{\infty\}$. On the other hand, we obtain

$$ds^2 = (1 - |g|^2)^2 |\omega|^2 \leq (1 + |g|^2)^2 |\omega|^2 = d\sigma^2.$$

Thus if ds^2 is complete, then $d\sigma^2$ is also complete. By Theorem 3.3, its Lorentzian Gauss map is constant, that is, it is a plane. $\square$

3.3. Lagrangian Gauss map of improper affine fronts in the affine 3-space. Improper affine spheres in the affine 3-space $\mathbf{R}^3$ also have similar properties to minimal surfaces in the Euclidean 3-space. Recently, Martínez [27] discovered the correspondence between improper affine spheres and smooth special Lagrangian immersions in the complex 2-space $\mathbf{C}^2$ and introduced the notion of *improper affine fronts*, that is, a class of (locally strongly convex) improper affine spheres with some admissible singularities in $\mathbf{R}^3$. We note that this class is called “improper affine maps” in [27], but we call this class “improper affine fronts” because Nakajo [28] and Umehara and Yamada [37, 38] showed that all improper affine maps are wave fronts in $\mathbf{R}^3$. The differential geometry of wave fronts is discussed in [34]. Moreover, Martínez gave the following holomorphic representation for this class.

THEOREM 3.5. [27, Theorem 3] *Let Σ be a Riemann surface and (F, G) a pair of holomorphic functions on Σ such that $\operatorname{Re}(FdG)$ is exact and $|dF|^2 + |dG|^2$ is positive definite. Then the induced map $\psi: \Sigma \rightarrow \mathbf{R}^3 = \mathbf{C} \times \mathbf{R}$ given by*

$$\psi := \left(G + \overline{F}, \frac{|G|^2 - |F|^2}{2} + \operatorname{Re} \left(GF - 2 \int FdG \right) \right)$$

is an improper affine front. Conversely, any improper affine front is given in this way. Moreover we set $x := G + \overline{F}$ and $n := \overline{F} - G$. Then $L_\psi := x + \sqrt{-1}n: \Sigma \rightarrow \mathbf{C}^2$ is a special Lagrangian immersion whose induced metric $d\tau^2$ from $\mathbf{C}^2$ is given by

$$d\tau^2 = 2(|dF|^2 + |dG|^2).$$

In addition, the affine metric h of ψ is expressed as $h := |dG|^2 - |dF|^2$ and the singular points of ψ correspond to the points where $|dF| = |dG|$.

The nontrivial part of the Gauss map of $L_\psi: \Sigma \rightarrow \mathbf{C}^2 \simeq \mathbf{R}^4$ (see [6]) is the meromorphic function $\nu: \Sigma \rightarrow \mathbf{C} \cup \{\infty\}$ given by

$$\nu := \frac{dF}{dG},$$

which is called the *Lagrangian Gauss map* of ψ . An improper affine front is said to be *weakly complete* if the induced metric $d\tau^2$ is complete. We note that

$$d\tau^2 = 2(|dF|^2 + |dG|^2) = 2(1 + |\nu|^2)|dG|^2.$$

Applying Theorem 1.1 to the metric $d\tau^2$, we can get the following theorem. This is a generalization of [19, Theorem 3.2].

THEOREM 3.6. *Let $\psi: \Sigma \rightarrow \mathbf{R}^3$ be an improper affine front whose Lagrangian Gauss map $\nu: \Sigma \rightarrow \mathbf{C} \cup \{\infty\}$ omits more than 3 ($= 1 + 2$) distinct values. Then there exists a positive constant C depending on the set of exceptional values, but not Σ , such that for all $p \in \Sigma$ we have*

$$|K_{d\tau^2}(p)|^{1/2} \leq \frac{C}{d(p)},$$

where $K_{d\tau^2}(p)$ is the Gaussian curvature of the metric $d\tau^2$ at p and $d(p)$ is the geodesic distance from p to the boundary of Σ . In particular, if the Lagrangian Gauss map of a weakly complete improper affine front in $\mathbf{R}^3$ is nonconstant, then it can omit at most 3 ($= 1 + 2$) values.

Since the singular points of ψ correspond to the points where $|\nu| = 1$, we can obtain a simple proof of the parametric affine Bernstein theorem ([3], [16]) for improper affine spheres from the viewpoint of value-distribution-theoretic properties of the Lagrangian Gauss map. For the proof, see [19, Corollary 3.6].

COROLLARY 3.7. *Any affine complete improper affine sphere in $\mathbf{R}^3$ must be an elliptic paraboloid.*

3.4. Ratio of canonical forms of flat fronts in the hyperbolic 3-space. We denote by $\mathbf{H}^3$ the hyperbolic 3-space, that is, the simply connected Riemannian 3-manifold with constant sectional curvature -1 , which is represented as

$$\mathbf{H}^3 = SL(2, \mathbf{C})/SU(2) = \{aa^*; a \in SL(2, \mathbf{C})\} \quad (a^* := {}^t\bar{a}).$$

For a holomorphic Legendrian immersion $\mathcal{L}: \Sigma \rightarrow \mathbf{H}^3$ on a simply connected Riemann surface Σ , the projection

$$f := \mathcal{L}\mathcal{L}^*: \Sigma \rightarrow \mathbf{H}^3$$

gives a *flat front* in $\mathbf{H}^3$. Here, flat fronts in $\mathbf{H}^3$ are flat surfaces in $\mathbf{H}^3$ with some admissible singularities (see [23], [26] for the definition of flat fronts in $\mathbf{H}^3$). We call $\mathcal{L}$ the *holomorphic lift* of f . Since $\mathcal{L}$ is a holomorphic Legendrian map, $\mathcal{L}^{-1}d\mathcal{L}$ is off-diagonal (see [14], [25], [26]). If we set

$$\mathcal{L}^{-1}d\mathcal{L} = \begin{pmatrix} 0 & \theta \\ \omega & 0 \end{pmatrix},$$

then the pull-back of the canonical Hermitian metric of $SL(2, \mathbf{C})$ by $\mathcal{L}$ is represented as

$$ds_{\mathcal{L}}^2 := |\omega|^2 + |\theta|^2$$

for holomorphic 1-forms ω and θ on Σ . A flat front f is said to be *weakly complete* if the metric $ds_{\mathcal{L}}^2$ is complete [24, 37]. We define a meromorphic function on Σ by the ratio of canonical forms

$$\rho := \frac{\theta}{\omega}.$$

Then a point $p \in \Sigma$ is a singular point of f if and only if $|\rho(p)| = 1$ [22]. We note that

$$ds_{\mathcal{L}}^2 = |\omega|^2 + |\theta|^2 = (1 + |\rho|^2)|\omega|^2.$$

Applying Theorem 1.1 to the metric $ds_{\mathcal{L}}^2$, we can get the following theorem. This is a generalization of [19, Theorem 4.5].

THEOREM 3.8. *Let $f: \Sigma \rightarrow \mathbf{H}^3$ be a flat front on a simply connected Riemann surface Σ . Suppose that the ratio of canonical forms $\rho: \Sigma \rightarrow \mathbf{C} \cup \{\infty\}$ omits more than 3 ($= 1 + 2$) distinct values. Then there exists a positive constant C depending on the set of exceptional values, but not Σ , such that for all $p \in \Sigma$ we have*

$$|K_{ds_{\mathcal{L}}^2}(p)|^{1/2} \leq \frac{C}{d(p)},$$

where $K_{ds_{\mathcal{L}}^2}(p)$ is the Gaussian curvature of the metric $ds_{\mathcal{L}}^2$ at p and $d(p)$ is the geodesic distance from p to the boundary of Σ . In particular, if the ratio of canonical forms of a weakly complete flat front in $\mathbf{H}^3$ is nonconstant, then it can omit at most 3 ($= 1 + 2$) values.

If Σ is not simply connected, then we consider that ρ is a meromorphic function on its universal covering surface $\tilde{\Sigma}$. As a corollary of Theorem 3.8, we give a simple proof of the classification ([35], [39]) of complete nonsingular flat surfaces in $\mathbf{H}^3$. For the proof, see [17, Corollary 3.5].

COROLLARY 3.9. *Any complete nonsingular flat surface in $\mathbf{H}^3$ must be a horosphere or a hyperbolic cylinder.*

REFERENCES

- [1] L. AHLFORS, Zur Theorie der Überlagerungsflächen, Acta Math. **65** (1935), 157–194.
- [2] L. J. ALÍAS AND B. PALMER, On the Gaussian curvature of maximal surfaces and the Calabi-Bernstein theorem, Bull. London Math. Soc. **33** (2001), 454–458.
- [3] E. CALABI, Improper affine hypersurfaces of convex type and a generalization of a theorem by K. Jörgens, Mich. Math. J. **5** (1958), 108–126.
- [4] E. CALABI, Examples of Bernstein problems for some nonlinear equations, Proc. Sympos. Pure Math. **15** (1970), 223–230.
- [5] S. S. CHERN, Complex analytic mappings of Riemann surfaces I., Amer. J. Math. **82** (1960), 323–337.
- [6] B. Y. CHEN AND J. M. MORVAN, Géométrie des surfaces lagrangiennes de $\mathbf{C}^2$, J. Math. Pures Appl. **66** (1987), 321–335.
- [7] S. Y. CHENG AND S. T. YAU, Maximal spacelike hypersurfaces in the Lorentz-Minkowski spaces, Ann. of Math. **104** (1976), 407–419.
- [8] F. J. M. ESTUDILLO AND A. ROMERO, On maximal surfaces in the n -dimensional Lorentz-Minkowski space, Geom. Dedicata **38** (1991), 167–174.
- [9] F. J. M. ESTUDILLO AND A. ROMERO, Generalized maximal surfaces in Lorentz-Minkowski space L^3 , Math. Proc. Cambridge Philos. Soc. **111** (1992), 515–524.

- [10] F. J. M. ESTUDILLO AND A. ROMERO, On the Gauss curvature of maximal surfaces in the 3-dimensional Lorentz-Minkowski space, *Comment. Math. Helv.* **69** (1994), 1–4.
- [11] H. FUJIMOTO, On the number of exceptional values of the Gauss map of minimal surfaces, *J. Math. Soc. Japan* **40** (1988), 235–247.
- [12] H. FUJIMOTO, Value Distribution Theory of the Gauss Map of Minimal Surfaces in $\mathbf{R}^m$, *Aspects of Mathematics*, E21. Friedr. Vieweg & Sohn, Braunschweig, 1993.
- [13] H. FUJIMOTO, Nevanlinna theory and minimal surfaces, *Geometry V*, 95–151, 267–272, *Encyclopaedia Math. Sci.*, **90**, Springer, Berlin, 1997.
- [14] J. A. GÁLVEZ, A. MARTÍNEZ AND F. MILÁN, Flat surfaces in hyperbolic 3-space, *Math. Ann.* **316** (2000), 419–435.
- [15] T. IMAIZUMI AND S. KATO, Flux of simple ends of maximal surfaces in $\mathbf{R}^{2,1}$, *Hokkaido Math. J.* **37** (2008), 561–610.
- [16] K. JÖRGENS, Über die Lösungen der differentialgleichung $rt - s^2 = 1$, *Math. Ann.* **127** (1954), 130–134.
- [17] Y. KAWAKAMI, A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space, preprint, arXiv:1110.3110.
- [18] Y. KAWAKAMI, R. KOBAYASHI AND R. MIYAOKA, The Gauss map of pseudo-algebraic minimal surfaces, *Forum Math.* **20** (2008), 1055–1069.
- [19] Y. KAWAKAMI AND D. NAKAJO, Value distribution of the Gauss map of improper affine spheres, to appear in *Journal of the Mathematical Society of Japan*, arXiv:1004.1484.
- [20] O. KOBAYASHI, Maximal surfaces in the 3-dimensional Minkowski space $\mathbf{L}^3$, *Tokyo J. Math.* **6** (1983), 297–309.
- [21] R. KOBAYASHI, Toward Nevanlinna theory as a geometric model for Diophantine approximation, *Sugaku Expositions* **16** (2003), 39–79.
- [22] M. KOKUBU, W. ROSSMAN, K. SAJI, M. UMEHARA AND K. YAMADA, Singularities of flat fronts in hyperbolic space, *Pacific J. Math.* **221** (2005), 303–351.
- [23] M. KOKUBU, W. ROSSMAN, M. UMEHARA AND K. YAMADA, Flat fronts in hyperbolic 3-space and their caustics, *J. Math. Soc. Japan* **59** (2007), 265–299.
- [24] M. KOKUBU, W. ROSSMAN, M. UMEHARA AND K. YAMADA, Asymptotic behavior of flat surfaces in hyperbolic 3-space, *J. Math. Soc. Japan* **61** (2009), 799–852.
- [25] M. KOKUBU, M. UMEHARA AND K. YAMADA, An elementary proof of Small’s formula for null curves in $\mathrm{PSL}(2, \mathbf{C})$ and an analogue for Legendrian curves in $\mathrm{PSL}(2, \mathbf{C})$, *Osaka J. Math.* **40** (2003), 697–715.
- [26] M. KOKUBU, M. UMEHARA AND K. YAMADA, Flat fronts in hyperbolic 3-space, *Pacific J. Math.* **216** (2004), 149–175.
- [27] A. MARTÍNEZ, Improper affine maps, *Math. Z.* **249** (2005), 755–766.
- [28] D. NAKAJO, A representation formula for indefinite improper affine spheres, *Results Math.* **55** (2009), 139–159.
- [29] J. NOGUCHI AND T. OCHIAi, *Geometric Function Theory in Several Complex Variables*, *Transl. Math. Monogr.* **80**, Amer. Math. Soc., Providence, RI, 1990.
- [30] R. OSSERMAN, Global properties of minimal surfaces in E^3 and E^n , *Ann. of Math.* **80** (1964), 340–364.
- [31] R. OSSERMAN, *A survey of minimal surfaces*, second edition, Dover Publication Inc., New York, 1986.

- [32] A. ROMERO, Simple proof of Calabi-Bernstein's theorem on maximal surfaces, Proc. Amer. Math. Soc. **124** (1996), 1315–1317.
- [33] M. RU, Nevanlinna theory and its Relation to Diophantine Approximation, World Sci., River Edge, NJ, 2001.
- [34] K. SAJI, M. UMEHARA AND K. YAMADA, The geometry of fronts, Ann. of Math. **169** (2009), 491–529.
- [35] S. SASAKI, On complete flat surfaces in hyperbolic 3-space, Kōdai Math Sem. Rep. **25** (1973), 449–457.
- [36] M. UMEHARA AND K. YAMADA, Maximal surfaces with singularities in Minkowski space, Hokkaido Math. J. **35** (2006), 13–40.
- [37] M. UMEHARA AND K. YAMADA, Applications of a completeness lemma in minimal surface theory to various classes of surfaces, Bull. London Math. Soc. **43** (2011), 191–199.
- [38] M. UMEHARA AND K. YAMADA, CORRIGENDUM: Applications of a completeness lemma in minimal surface theory to various classes of surfaces, to appear in Bulletin of the London Mathematical Society. (doi:10.1112/blms/bds017)
- [39] Y. A. VOLKOV AND S. M. VLADIMIROVA, Isometric immersions of the Euclidean plane in Lobačevskii space (Russian), Mat. Zametki **10** (1971), 327–332.

GRADUATE SCHOOL OF SCIENCE AND ENGINEERING,
YAMAGUCHI UNIVERSITY,
YAMAGUCHI, 753-8512, JAPAN
E-mail address: y-kwkami@yamaguchi-u.ac.jp

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p -adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection

- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments

- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow

- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra
- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions

- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing
- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine

- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency
- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO

MI2012-4 Yasuhide FUKUMOTO & Youich MIE

Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field

MI2012-5 Yu KAWAKAMI

On the maximal number of exceptional values of Gauss maps for various classes of surfaces