

Ability of Endophytic Fungi Isolated from Rice to Inhibit *Pyricularia oryzae*-Induced Rice Blast in Indonesia

Suada, I Ketut

Study Program of Agroecotechnology, Faculty of Agriculture, Udayana University

Suhartini, Desak Made Weda Yati

Study Program of Agroecotechnology, Faculty of Agriculture, Udayana University

Sunariasih, Ni Putu Linda

Study Program of Agroecotechnology, Faculty of Agriculture, Udayana University

Wirawan, I Gede Putu

Study Program of Agroecotechnology, Faculty of Agriculture, Udayana University

他

<https://doi.org/10.5109/22047>

出版情報：九州大学大学院農学研究院紀要. 57 (1), pp.51-53, 2012-02. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：



Ability of Endophytic Fungi Isolated from Rice to Inhibit *Pyricularia oryzae*-Induced Rice Blast in Indonesia

I Ketut SUADA¹, Desak Made Weda Yati SUHARTINI¹, Ni Putu Linda SUNARIASIH¹,
I Gede Putu WIRAWAN¹, Kun Woo CHUN²,
Joo Young CHA^{3*} and Shoji OHGA

Laboratory of Forest Resources Management, Division of Forest Environmental Sciences,
Department of Agro-environmental Sciences, Faculty of Agriculture,
Kyushu University, Fukuoka, 811-2415, Japan
(Received October 31, 2011 and accepted November 9, 2011)

The aim of this research was to measure the ability of endophytic fungi isolated from rice to inhibit the growth of *Pyricularia oryzae*, the pathogen that causes rice blast. *Phialemonium curvatum* had the greatest inhibitory effect against *P. oryzae* (66.6%), followed by *Phaeosphaeriopsis musae* (63.3%), making these species good candidates for biocontrol agents against *P. oryzae*.

Key words: Endophytic Fungi, inhibition ability, *Pyricularia oryzae*, rice

INTRODUCTION

Indonesia, with its tropical climate, is habitat to an enormous number of fungal species including endophytes harbored within the tissues of various plants. Rice (*Oryza sativa* L.) is, on a global scale, one of the most important food crops and performance improvements are continuously being sought in order to increase rice yields.

Endophytic fungi colonize their hosts without causing any external disease symptoms (Carroll, 1998), except under stressful conditions. Studies of the microorganisms associated with tropical plant species are becoming more frequent, especially studies describing biological control and the ability of these microorganisms to biosynthesize pharmacologically important compounds (Azevedo *et al.*, 2000). Endophytic fungi have frequently been reported to be associated with crop plants including wheat (*Triticum aestivum*), wild barley (*Hordeum brevisubulatum* and *Hordeum bogdanii*), soybean (*Glycine max*), and maize (*Zea mays*) (Dingle and McGee, 2003; Istifadah and McGee, 2006; Larran *et al.*, 2010). Some of the endophytic fungi associated with these crops confer upon them resistance to insect or fungal pathogens (Sieber *et al.*, 1988).

The association of arbuscular mycorrhizal fungi and endophytic bacteria with rice has been well documented (Glassop, 2007; Mano and Morisaki, 2008; Mattos *et al.*, 2008). Less, however, is known about rice's fungal endophytes. Fungal endophytes have been detected in cultivated rice (Naik *et al.*, 2009; Yuan *et al.*, 2007), and

antagonistic or plant growth-stimulating properties have been claimed for some of them. For example, endophytic *Fusarium* spp. from cultivated rice roots proved to be an effective biocontrol against a root-knot nematode (Le *et al.*, 2009). The occurrence of mycorrhizal and endophytic fungi in a variety of rice cultivars has also recently been reported (Vallino *et al.*, 2008).

Endophytic fungi, in symbiotic associations with rice plants, provide advantages to the plants such as increased growth and resistance against pathogens such as *Pyricularia oryzae*. The use of endophytic fungi to increase plant resistance to pathogens would be a step toward decreasing the use of fungicides and other synthetic chemicals in rice agriculture. In the present study, we screened endophytic fungi isolated from rice plants *in vitro* for possible biocontrol activity against the rice blast pathogen, *P. oryzae*.

MATERIALS AND METHODS

Isolates of *P. oryzae* and endophytic fungi

For the isolation of *P. oryzae*, diseased rice leaf parts were sterilized with 90% alcohol for 1 min and 70% alcohol for 2 min, and then rinsed with sterile water before being cultured in 2% MA medium (20 g malt extract and 18 g agar in 1000 ml distilled water). After incubation for 5 days, the emerging colony was identified according to its conidia. The 11 endophytic fungi used in this study were isolated from rice seeds and seedlings in Bali (Cha *et al.*, unpublished data).

Test for endophytic fungal species inhibition of *P. oryzae*

Each endophytic fungal isolate was cultured, together with *P. oryzae*, in MA medium to determine its inhibitory activity against the pathogen. Agar plugs, 5 mm in diameter, taken from 7-day-old colonies of the endophyte and pathogen pure cultures were inoculated 3 cm apart on MA medium in 9-cm diameter Petri

¹ Study Program of Agroecotechnology, Faculty of Agriculture, Udayana University, Bali, 80361, Indonesia

² Department of Forest Sciences, College of Forest and Environmental Sciences, Kangwon National University, Chuncheon, 200-701, Korea

³ Field Science Center for Northern Biosphere, Hokkaido University, Nayoro, 096-0071, Japan

* Corresponding author (E-mail: jycha@fsc.hokudai.ac.jp)

dishes. These were incubated for 1 week at 28°C. The pathogen inoculated onto the same media without the endophyte and incubated for 6 weeks at 28°C served as a positive control. The diameters of *P. oryzae* colonies grown with each endophyte isolate were used to calculate the amount of pathogen control, referred to as the percentage of inhibition, as follows:

$$\% \text{ inhibition} = (A-B)/A$$

where A is the diameter of the control pathogen colony and B is the diameter of the pathogen colony grown with the endophyte (Nuangmek *et al.*, 2008).

Data analysis

The colonization frequency (CF) of a single endophyte species was calculated as the number of segments colonized by an endophyte species divided by the total number of segments examined $\times 100\%$. The percentage contribution to an endophyte assemblage by the dominant endophyte (DE) was calculated as the CF of the dominant endophyte divided by the sum of the CF of all the endophytes in an assemblage $\times 100\%$ (Xing *et al.*, 2010).

RESULTS AND DISCUSSION

Most of the endophytic fungi isolated from rice seeds and seedlings, with the exception of *Cladosporium* sp. and *Nodulisporium* sp., showed greater than 50% inhibition of *P. oryzae* in *in vitro* testing. The endophyte that showed the greatest inhibition of the pathogen (66.6%) was *Phialemonium curvatum* W. Gams & W. B. Cooke, which had been isolated from seeds. The next most effective endophyte was *Phaeosphaeriopsis musae* M. Arzanlou & Crous, which inhibited the pathogen by 63.3% (Table 1). Some species of endophyte may be beneficial to the plant in other ways (e.g., nutrient and water uptake), but would provide less benefit as the plant produces fewer defensive chemicals in response to

increasing damage (Ahmad *et al.*, 1985; Cheplick and Clay, 1988). The effect of a single fungal strain on the plant can change when a combination of fungal strains infects the same individual (Lewis and Clements, 1986). These endophytic microorganisms are ubiquitous, increasing plant growth (Varma *et al.*, 1999) and fitness through their ability to improve plant tolerance to heavy metals and drought, and by reducing herbivory and/or phytopathogen settling through the induction of plant self-defense mechanisms (Azevedo *et al.*, 2000). Self-defense may also occur via chemicals produced by endophytic fungi and/or space competition within the plant tissue (Azevedo *et al.*, 2000).

In the future, endophytic microorganisms will be used to increase rice growth through mechanisms such as plant defense against herbivores. Studying these microbes may lead to methods to enhance their ability to improve rice productivity and/or their secondary metabolite production. Many secondary metabolites from endophyte-plant interactions have been isolated and used in raw or derived forms as drugs for the treatment of many conditions (Shennan, 2008). Since plants use the antibiotic qualities of endophytic fungi for defense against pathogens, this mostly chemically-based defense mechanism makes endophytic fungi a perfect group for novel antibiotic research, in the same way that other fungi have yielded such useful drugs as penicillin and streptomycin (Schade, 2007). The endophytic fungi studied here have shown high pathogen inhibition ability and should be considered good candidates for biological control of rice blast caused by *P. oryzae*.

ACKNOWLEDGMENTS

We thank the management of Kertalangu Jogging Track Denpasar Bali and their farmers for their hospitality during this work. This study was carried out with the support of the Forest Science & Technology Projects (project no. S210811L010110), provided by the Korea Forest Service.

REFERENCES

- Ahmad, S., S. Govindarajan, C. R. Funk and J. M. Johnson-Cicalese 1985 Fatality of house crickets on perennial ryegrass infected with fungal endophyte. *Entomol. Exp. Appl.*, **39**: 183–190
- Azevedo, J. L., W. J. Maccheroni, J. O. Pereira, and W. L. Araujo 2000 Endophytic microorganisms: A review of insect control and recent advances on tropical plants. *Electron J. Biotechn.*, **3**: 40–65
- Carroll, G. 1988 Fungal endophytes in stem and leaves from latent pathogen to mutualistic symbiont. *Ecology*, **69**: 2–9
- Cheplick, G. P. and K. Clay 1988 Acquired chemical defenses of grasses: the role of fungal endophytes. *Oikos*, **52**: 309–318
- Dingle, J. and P. A. McGee 2003 Some endophytic fungi reduce the density of pustules of *Puccinia recondita* f. sp. *tritici* in wheat. *Mycol. Res.*, **107**: 310–316
- Glassop, D., R. M. Godwin, S. E. Smith and S. W. Smith 2007 Rice phosphate transporters associated with phosphate uptake in rice roots colonized with arbuscular mycorrhizal fungi. *Can. J. Bot.*, **85**: 644–651
- Istifadah, N. and P. A. McGee 2006 Endophytic *Chaetomium globosum* reduces development of tan spot in wheat caused by

Table 1. Percentage of inhibition ability of endophytic fungal species to *P. oryzae*

Fungal species	Inhibition ability (Value \pm SE) (%)
<i>Cladosporium</i> sp.	46.6 \pm 0.1
<i>Nigrospora oryzae</i>	51.1 \pm 2.1
<i>Nigrospora</i> sp.	51.1 \pm 0.5
<i>Nodulisporium</i> sp.	45.6 \pm 1.4
<i>Penicillium citrinum</i>	60.0 \pm 0.6
<i>Penicillium pinophilum</i>	52.2 \pm 0.5
<i>Penicillium radicum</i>	55.6 \pm 2.2
<i>Phaeosphaeriopsis musae</i>	63.3 \pm 0.1
<i>Phialemonium curvatum</i>	66.6 \pm 0.1
<i>Sarocladium oryzae</i>	61.1 \pm 0.6
<i>Sordariomycetes</i> sp.	58.8 \pm 0.7

- Pyrenophora tritici-repentis*. *Australian Plant Pathol.*, **35**: 411–418
- Larran, S. and C. Monaco 2010 Status and progress of research in endophytes from agricultural crops in Argentina. In: Arya, A. and A. E. Perello (Eds.) *Management of fungal plant pathogens*. pp. 149–151
- Le, H. T. T.; J. L. Padgham and R. A. Sikora 2009 Biological control of the rice root-knot nematode *Meloidogyne graminicola* on rice using endophytic and rhizosphere fungi. *Int. J. Pest Manage.*, **55**: 31–36
- Lewis, G. C. and R. O. Clements 1986 A survey of ryegrass endophyte (*Acremonium loliae*) in the U. K. and its apparent ineffectuality on a seedling pest. *J. Agric. Sci.*, **107**: 633–638
- Mano, H. and H. Morisaki 2008 Endophytic bacteria in rice plant. *Microbes Environ.*, **23**: 109–117
- Mattos, K. A., V. L. Pádua, A. Romeiro, L. F. Hallack, B. C. Neves, T. M. Ulisses, C. F. Barros, A. R. Todeschini, J. O. Previato and L. Mendonça-Previato 2008 Endophytic colonization of rice (*Oryza sativa* L.) by the diazotrophic bacterium *Burkholderia kururiensis* and its ability to enhance plant growth. *An. Acad. Bras. Cienc.*, **80**: 477–493
- Naik, B. S., J. Shashikala and Y. L. Krishnamurthy 2009 Study on the diversity of endophytic communities from rice (*Oryza sativa* L.) and their antagonistic activities *in vitro*. *Microbiol. Res.*, **164**: 290–296
- Nuangmek, W., E. H. C. McKenzie and S. Lumyong 2008 Endophytic fungus from wild banana (*Musa acuminata* Colla) works against anthracnose disease caused by *Colletotrichum musae*. *Jour. of Microbiol.*, **3**(5): 368–374
- Schade, R., F. Andersohn, S. Suissa, W. Haverkamp and E. Garbe 2007 Dopamine agonists and the risk of cardiac-valve regurgitation. *New England Journal of Medicine*, **356**(1): 29–38
- Shennan, C. 2008 Biotic interactions, ecological knowledge and agriculture. *Phil. Trans. R. Soc. Br.*, **363**: 717–739
- Sieber, T. N., T. K. Riesen, E. Muller and P. M. Fried 1988 Endophytic fungi in four winter wheat cultivars (*Triticum aestivum* L.) differing in resistance against *Stagonospora nodorum* (Berk.) Cast. & Germ. = *Septoria nodorum* (Berk.) Berk. *J. Phytopathol.*, **122**: 289–306
- Vallino, M., D. Greppi, M. Novero, P. Bonfante and E. Lupotto 2008 Rice root colonization by mycorrhizal and endophytic fungi in aerobic soil. *Ann. Appl. Biol.*, **154**: 195–204
- Varma A., S. Verma, Sudha, N. Sahay, B. Buetehorn and P. Franken 1999 *Piriformospora indica*, a cultivable plant-growth-promoting root endophyte. *App. Environ. Microbiol.*, **65**: 2741–2744
- Xing, X. K., J. Chen, M. J. Xu, W. H. Lin and S. X. Guo 2010 Fungal endophytes associated with Sonneratia (Sonneratiaceae) mangrove plants on the South Coast of China. *Blackwell Verlag GmbH*: 1–11
- Yuan, Z. L., C. C. Dai, X. Li, L. S. Tian and X. X. Wang 2007 Extensive host range of an endophytic fungus affects the growth and physiological functions in rice (*Oryza sativa* L.). *Symbiosis*, **43**: 21–28