

The transcription factor NKX2-3 mediates p21  
expression and ectodysplasin-A signaling in the  
enamel knot for cusp formation in tooth  
development

韓, 雪

<https://hdl.handle.net/2324/2236143>

---

出版情報 : Kyushu University, 2018, 博士 (歯学) , 課程博士  
バージョン :  
権利関係 :

氏 名 : 韓 雪

論 文 名 :

The transcription factor NKX2-3 mediates p21 expression and ectodysplasin-A signaling in the enamel knot for cusp formation in tooth development

(ホメオボックス転写因子 NKX2-3 は p21 の発現調節および EDA シグナル経路を介して歯の咬頭形成を調節する)

区 分 : 甲

### 論 文 内 容 の 要 旨

Tooth morphogenesis is initiated by reciprocal interactions between the ectoderm and neural crest-derived mesenchyme. During tooth development, tooth cusps are regulated by precise control of proliferation of cell clusters, termed enamel knots, that are present among dental epithelial cells. The interaction of ectodysplasin-A (EDA) with its receptor, EDAR, plays a critical role in cusp formation by these enamel knots, and mutations of these genes is a cause of ectodermal dysplasia. NKX2-3, known as a member of transcription factors of NK homeobox family, critically regulates organ development. A previous study showed that *Nkx2-3* deficiency in tooth resulted in the absence of cusp formation. However, the molecular mechanism of NKX2-3 in tooth morphogenesis remains unclear. In this study, we examined the role of NKX2-3 in tooth development. Using gene microarray analysis in mouse embryos, we found that *Nkx2-3* is highly expressed during tooth development and increased during the tooth morphogenesis, especially during cusp formation. In organ culture, *Nkx2-3* siRNA inhibited cusp formation, suggesting that NKX2-3 is critical for tooth morphogenesis. Cusp formation is known to be regulated by an enamel knot expressing the cyclin-dependent kinase inhibitor p21. We found that NKX2-3 induced the expression in mRNA and protein of p21 when dental epithelial cell line, M3H1 cells were transfected with *Nkx2-3* expression vector. Furthermore, ChIP assay showed that NKX2-3 regulated transcription and expression of p21 by binding to its promoter region, resulting in alteration of cell proliferation. *Nkx2-3* was shown to be significantly induced by EDA, suggesting that *Nkx2-3* is a target transcription factor regulated by the *Eda/Edar* signaling pathway in the enamel knot. Moreover, NKX2-3 activated the bone morphogenetic protein (BMP) signaling pathway by up-regulating expression levels of *Bmp2* and *Bmpr2* in dental epithelium and decreased the expression of the dental epithelial stem cell marker SRY box 2 (SOX2).

These results indicate that NKX2-3 plays roles during the tooth cusp formation via *Eda* signaling and regulating p21 expression.