The Magnitude Of Population And Fishing Rate Of King-Crab, *Paralithodes Camtschatica* (Tilesius)

Aikawa, Hiroaki
Fisheries Laboratory, Department of Agriculture, Kyushu University

<https://doi.org/10.5109/22607>

出版情報 : 九州大学大学院農学研究院紀要. 9 (1), pp.41-49, 1948-03. Kyushu University
バージョン :
権利関係 :

THE MAGNITUDE OF POPULATION AND FISHING
RATE OF KING-CRAB, *PARALITHODES*
CAMTSCHATICA (TILESIIUS)

HIROAKI AIKAWA

The Sakhalin district, extending from both coast of southern Sakhalin to the Okhotsk coast of Hokkaidō, the Nemuro district around the southern Kurile Islands to Kushiro along the Pacific coast of Hokkaidō, and the Kamtchatka district, especially off its west coast, are 3 important fishing grounds for the Japanese King-crab fishery. The crab populations of these districts are respectively of different magnitude and also of different race, being distinguishable by their morphological features. Here, the magnitude and fishing rate are computed statistically for the populations in the Nemuro and Kamtchatka districts.

1. Larval death rate and natural decreasing
rates of matured groups

From the epoch-making investigation of Dr. Marukawa (1933), it is known that the larval death rate is catastrophically as high as 0.97 during the period from the 1st. zoea stage, just after hatched out from berried crab, to the youngest crab stage through glaucothoe stage, so that only 3% of the 1st. zoea larvae can survive, till they attain the smallest adult form. These youngest crabs must undergo furthermore remarkable decrease through 7 or 9 moultings, till they grow 7 mm in carapace width in the next Spring. As the decreasing rate is 0.16 on an average at each

moultage, the total decreasing rate of 0-age group is at least $0.99977 = 1 - (1 - 0.964)(1 - 0.16)^9$

On examining the frequency distributions of carapace width, Marukawa considered the isolated dominant size groups to be of different ages. Thence, the matured crabs of 10 cm or thereabouts in carapace width are supposed to be at least of 8-age. Y. Kajita (1932) however observed in the Nemuro district that the matured crabs were at most of 4-age and already attained the size above 10 cm, and Wang (1937) was also of opinion that in the Sakhalin district the matured groups were of 3- and 4-age, and those of 4-age were larger than 17 cm. Therefore, the same age group is certainly composed of some isolated size groups, which are different in the number of moultings from each other.

From Wang's data, it is known that the minimum size of N-age group is l_n cm in carapace width and that of (N+1)-age group l_{n+1} cm. On the other hand, from Dr. Marukawa's data, the minimum size of (N+x)-age group is nearly equal to l_n cm and that of (N+x+y)-age group also to l_{n+1} cm. Then, $(N+x+y) - (N+x) = y$ is the number of moultings, which N-age group undergoes during a year, and after which it becomes a year older. The natural decreasing rate, mainly due to the moultage, is computable as $1 - (1 - 0.16)^y$. The accuracy of these natural decreasing rates computed must be ascertained with other data.

Table 1. Age, number of moultings, and computed decreasing rate of king-crab, *Paralithodes camtschatica*.

Marukawa (1933)				Wang (1937)				
Age	Size (mm)		No. of moulting ages	Decreasing rate (m)	Age	Size (mm)	No. of moulting ages	Decreasing rate (m)
Zoëa	—		4	0.964				
Glaucothë	—		1	0.16				
Young crab	7		8	0.748				
0-age	17		13	0.998				
1	17		6.5	0.69				
2	25		"	"				
3	34		2.5	0.35				
4	42		"	"				
5	53		2	0.294	0-age	<51	33	0.99977

6-age	70							
7	85			1	101	6	0.649	
8	100							
9-age	130	116	1	0.16				
10	132	125			2	145	4	0.502
11	142	130						
12-age	150	135						
13	157	140						
14	167	141			3	170	4	0.502
15	171	148						
16-age	175	151						
17	179	153						
18	183	155			4	190	4	0.502
19	186	157						
20	189	160						
21-age	191	162						
22	193	164						
23	198	166			5	204	4	0.502
24	201	—						
25	204	—						

Note: Average number of moultings assumed as 4 for the groups older than 1-age.

2. The magnitude of population and fishing rate in the Nemuro district

In the Nemuro district, the production of canned crabs amounted to 51×10^3 cases on an average and showed no remarkable fluctuation recently. Then, the population must be balanced with fishing intensity and the annual recruitment amount might be equal to the annual yield of crabs. 51×10^3 cases correspond to 1200×10^3 male crabs. The female crabs may be caught together with gill-nets, but they are usually liberated at once into water. But, most of them die and cannot survive long, as the gill-nets are lifted a few days after being set in water. So, female crabs which were caught must be added to the decrease in population

due to fishing. As female crabs occupy 36% of the shoal, the loss of female crabs is about 670×10^3 individuals, and, thus, the total decrease is 1870×10^3 individuals, which must be equal to the annual recruitment amount here.

Now, 4-age crabs, ranging from 17 cm to 19 cm in carapace width, are dominant in the Nemuro district and the female crabs of 4-age can produce 200×10^3 zoëa larvae on an average. Then, the number of female crabs (R), which can supply the recruitment with 1870×10^3 crabs after 4 years, is

$$R = \frac{1870 \times 10^3}{200 \times 10^3 (1 - 0.99977) (1 - 0.649) (1 - 0.502)^2}$$

$$= 47 \times 10^3 \text{ individuals (female crabs only)}$$

$47 \times 10^3 / 0.36 = 131 \times 10^3$ individuals (male crabs included) are the total number of crabs, which must be left unfished. Then, at the beginning of the fishing season, 2000×10^3 ($= 1870 \times 10^3 + 131 \times 10^3$) crabs must migrate into the Nemuro district. In addition to this 4-age group, 2- and 3- age groups must migrate here and their amounts are respectively

$$\text{3-age group } 2000 \times 10^3 / (1 - 0.502) = 4016 \times 10^3 \text{ individuals}$$

$$\text{2-age group } 2000 \times 10^3 / (1 - 0.502)^2 = 8064 \times 10^3 \text{ individuals}$$

Then, the total spawning shoal will amount to 14080×10^3 individuals. This is the magnitude of population in the Nemuro district. Then, the fishing rate (f) is

$$f = \frac{F}{S} = \frac{1870 \times 10^3}{14080 \times 10^3} = 0.135$$

In the Nemuro district, 390 tagged crabs were liberated and 22 crabs were recaptured within the same fishing season, then the recapture rate (r) is 0.056 (22/390). For the calculation of fishing rate, the recapture rate must be corrected not only with time factor but also with catch factor. The fishing rate is usually $3.5 \times r$ or thereabouts. Then, the fishing rate will never be smaller than 0.196, which is somewhat larger than the rate calculated statistically, so that the decreasing rate of the matured crabs, considered as 0.502, will be rather smaller than the real rate. Therefore, the magnitude of population will be far smaller than 14000×10^3 individuals here.

3. The magnitude of population and fishing rate in the Kamtchatka district

Here, the yield of King-crabs was rather small from the factories installed on land. While, after the fishing experiment during the years from 1914 to 1920, crab fishery with floating factory has been carried out with success since 1923. From the relation between the annual yield (F) and the number of gill-nets used (n), the following 3 equations are established (see fig. 1)

Fig. 1. Relation between the annual yield and the number of gill-net used (see table 2)

$$\left. \begin{array}{ll} 1924-28 \text{ (1st period)} & F = 462 \times 10^4 + 103 \cdot n \\ 1929-32 \text{ (2nd period)} & F = 3671 \times 10^4 - 104 \cdot n \\ 1934-38 \text{ (3rd period)} & F = -429 \times 10^4 - 114 \cdot n \end{array} \right\} \dots\dots\dots (1)$$

During the 1st period, 1924-'28, the annual yield increased in accordance with the increase in the number of gill-nets used, while, during the 2nd period, 1929-'32, the annual yield decreased inversely with the increase in the number of gill-nets used, but during the 3rd period, 1934-'38, the relation between them became again the same as that during the 1st period.

When k is the fishing intensity of an unit fishing effort of gill-net, S the magnitude of population and F the annual

yield, the following equation can be obtained, as far as the gill net is the competitive gear

$$F = S \{1 - (1 - k)^n\} \quad (2)$$

where n is the number of unit fishing efforts. When n is large enough to be considered that $n \cdot (n-1)$ is nearly equal to n^2 and, in addition, k is very small in value, the equation above is

$$F = S \cdot n \cdot k - S \cdot n^2 \cdot k^2 / 2 \quad (2')$$

From equations 1 and 2', the fishing intensity (k) is computable independently from S and n . The k -value was -0.446×10^{-4} during the 1st. period, when the increase in the magnitude of available population exceeded considerably the decrease due to fishing mortality. On the contrary, k -values became positive in the succeeding periods, being 0.057×10^{-4} during the 2nd. period and 0.531×10^{-4} during the 3rd., as the total population here became available since 1929. The k -values were considerably different from each other, as the magnitudes of population varied. From equation (2'), the fishing rate (f) is

$$f = \frac{F}{S} = k \cdot n - \frac{1}{2} k^2 \cdot n^2 \quad (3)$$

and also the magnitude of population (S) is

$$S = \frac{F}{f} = \frac{F}{k \cdot n - 1/2 k^2 \cdot n^2} \quad (4)$$

To these equations, the known values of n and k are substituted, then, f and S are calculated as shown in table 2, where k -value in the 1st. period is assumed to be similar to that in the 3rd. period.

Table 2. Statistical table for king crab fishery in the Kamtchatka district

Year	No. of float. fact.	No. of unit fish. efforts	Annual yield ($F \times 10^3$)	No. of individuals		Fishing rate (f)	Calculated magnitude ($S \times 10^3$)
				unit fish. effort	per a case		
1924	6	12770	3930	48.05	96.1	0.070	56143
1925	8	35546	7648	31.69	72.5	0.181	42254
1926	12	54251	13873	21.25	63.3	0.260	53358
1927	17	271273	22643	22.00	68.5	0.355	63747
1928	14	249871	20113	14.83	66.2	0.414	48583
mean	—	—	—	27.54	73.3	0.256	52819

1929	15	137005	23718	10.83	64.7	0.476	49828
1930	13	147159	19571	9.83	57.9	0.487	40187
1931	8	233521	14141	7.52	66.4	0.545	25947
1932	6	256572	9186	5.95	68.2	0.373	24627
1933	5	—	6624	—	63.7	—	—
mean	—	—	—	8.53	64.2	0.470	35147
1934	5	118000	7788	6.59	67.5	0.428	18196
1935	6	131000	9800	7.54	69.7	0.443	22124
1936	7	148400	13084	8.80	79.0	0.469	27898
1937	—	152400	14194	9.30	76.2	0.473	30008
1938	—	196400	18055	9.20	75.2	0.500	36110
mean	—	—	—	8.29	73.5	0.463	26867

During the 1st. period, the magnitude of population remained unchanged, being 52819×10^3 individuals on an average, despite that the number of gill-nets used and the fishing rate increased. During the 2nd period, however, the magnitude of population decreased year by year with the increase in the number of gill-nets used and also in the fishing rate. Thence, we must consider that the population is being overfished according to the definition given by E. S. Russel (1939). The catch per unit fishing effort became less than one third of that in the preceding period, although the number of individuals required for a case of canned crabs remained unchanged. In the 3rd. period, the magnitude of population became again large according to the increase in the number of gill-nets used and also in the fishing rate. However, the catch per unit fishing effort and the number of individuals required for a case of canned crabs increased, so that the population would never be replenished with sufficient amount, and it would include smaller crabs within it than before. In other words, the small crabs of 2- and 3- age became more available in the 3rd. period than in the previous period. Accordingly, the fishing was furthermore intensified.

Recently, the magnitude of population may be replenished due to the closure for several years during the World War II, and may attain the previous stationary population, 5000×10^3 individuals, as was seen in the 1st. period.

Among the liberated crabs in 1928, 220 tagged ones were

caught within the same fishing season, 96 in 1929 and lastly 8 in 1930. Then, the total decreasing rate (d) is computed with Tauti's formula

$$1-d = \frac{96+8}{220 \times 2 + 96+8} = 1-0.795$$

As the average fishing rate during the years 1930 is 0.459 (see table 2), the natural decreasing rate (m) is computable by the equation below

$$1-d = (1-f)(1-m)$$

then, $1-m = 1-0.646$. This m -value is approximately similar to that of 2-age group (0.649) and also that of 3- age and older age groups (0.502).

The tagging experiment was tried again in 1929. Tagged crabs were liberated every month during the fishing season and some of them were also recaptured within the same season. When T is the whole duration of fishing season, T_n the duration from the month of liberation to the end of season, F the total catch during T and F_n during T_n , the efficiency of liberation of X_n crabs at T_n is equivalent to that of $X_n \cdot T_n / T \cdot F_n / F = X'_n$ at the beginning of the fishing season. Then the fishing rate (f) is

$$f = \frac{x}{X_n \cdot T_n / T \cdot F_n / F} = \frac{x}{X'} \left(\frac{T}{T_n} \cdot \frac{F}{F_n} \right) = r' \left(\frac{T}{T_n} \cdot \frac{F}{F_n} \right) \dots\dots (5)$$

where x is the number of recaptured crabs, T_n/T the time factor and F_n/F the catch factor. F_n , T_n and $\sum_i^n X'_n (=X')$ are shown in tables 3 and 4. The fishing rate is 0.458 in 1928 and 0.335 in 1929, both being approximately equal to f -values obtained statistically (see table 2).

Table 3. Tagging experiments and factors of correction

Year	Month	No. of liberated individuals	Time factor T/T_n	Catch factor F/F_n	No. of recap. indiv.	Do. corrected (x')	Fishing rate (f)
1928	VI	500	1.71	1.43	109	207	0.458
	VII	500	2.40	2.56	41	251	
	Sum	1000	—	—	150	458	
1929	V	77	1.33	1.09	14	22	0.335
	VI	164	1.71	1.43	17	42	
	VII	171	2.40	2.56	12	74	
	VIII	1	4.00	7.69	0	0	
	Sum	413	—	—	43	138	

Table 4. Montly change in catch

Year	Apr.	May	June	July	Aug.	Sept.
1927	—	9.6	38.6	29.0	22.8	—
1928	2.9	11.2	30.6	48.9	6.0	0.4
1929	—	21.7	17.4	13.0	39.2	8.7
1930	2.9	13.2	30.5	40.7	11.8	0.9
mean	1.5	13.9	29.3	32.9	19.9	2.5

Although the magnitude of population is far larger in the Kamtchatka district than in the Nemuro, the rates of recruitment may be nearly equal in both districts. Assuming that the rate of recruitment is 0.135, the yield will be 600×10^3 crabs of large size with the fishing rate of 0.135, as far as the population remains stationary.

REFERENCES: Kajita, Y., 1932, Rep. Fish. Exper. Sta., Hokkaidō, no. 27; 1942, Kaiyō Gyogyō, vol. 4, no. 4; Marukawa, H., 1933, Rep. Fish. Exp. Sta., Tokyō, no. 4; Suisan Kyoku (Bur. Fish.), 1938, Statistical Table of the northern Deep sea fisheries; Tauti, M., 1936, Bull. Japanese Soc. Sci. Fish., vol. 4, no. 4; Wang, Yi-Kuan, 1937; *ibid.*, vol. 5, no. 6; Russell, E.S., 1939, Rapp. et Proc.-Verb., vol. 110.