Studies On Foodstuffs In Emergencies. XXII. On The Starch Of Lycoris Radiata

Yamafuji, Kazuo
Faculty of Agriculture, Kyushu University

Tono, Tetsuzo

<https://doi.org/10.5109/22640>

出版情報 : 九州大学大学院農学研究院紀要. 9 (4), pp.355-360, 1950-11. Kyushu University
バージョン :
権利関係 :

STUDIES ON FOODSTUFFS IN EMERGENCIES. XXII.
ON THE STARCH OF *LYCORIS RADIATA*

KAZUO YAMAFUJI AND TETSUZO TONO

Lycoris radiata has been generally called "Higanbana" in Japan and its bulb contains a considerable amount of starch. This plant is widely distributed in various fields and forests, but it is unsuitable for eating. There exist several poisonous alkaloids in the bulb of *Lycoris* and the chemical constitution of lycorine, the principal base, was clarified by Kondo et al (1). Since these alkaloids have been utilized as medicine, we already studied the extraction, adsorption (2) and elution (3) of the bases. On the other hand Kihara (4) investigated the nature of polysaccharides in *Lycoris* root, and in this institute we also performed some experiments concerning the alcoholic fermentation (5) of these polyoses. In the present article we shall report the results of investigations on the isolation of starch and its decomposition by amylase.

EXPERIMENTAL

1. **Isolation of starch:** The tubers of Higanbana were gathered, dried at room temperature and ground in a mortar. The analysis of this powder gave the data shown in Table 1.

Table 1. Chemical composition of root powder.

Starch, %	Reducing sugar, %	Total N, %	Fibre, %	Ash, %	Water, %
48.65	1.63	0.61	2.82	3.26	18.07

An example of isolation experiments was carried out as follows: 5 g bulb powder was mixed with 50 g water and after 20 hrs. supernatant liquid decanted. The residue was thoroughly

ground in a glass mortar, filtered through a gauze and mixed with water again. This procedure was repeated 3 times and starch particles deposited were gathered. The yield of starch averaged 60%.

It is assumed that the extraction of starch from tuber is influenced by the hydrogen ion concentration of solution. The tuber powder was immersed into diluted hydrochloric acid and in similar manner starch prepared. The results obtained are given in Table 2.

Table 2. Relation between yield of starch and acidity of solution.

Conc. of HCl, %	0	0.1	0.2	0.3	0.4	0.5	0.6	0.7
Yield of starch, %	45	60	77	88	81	75	65	60

By hydrochloric acid of 0.2—0.5% some constituents other than starch are extracted from the plant material. Therefore, the preparations made by using such acid solutions were impure.

In order to compare the extracting power of alkalis with that of acids 100 cc of solution was added to 1 g of bulb powder, after 20 hrs. at 36°C the mixture filtered and the sugar content in this filtrate determined after the hydrolysis.

Table 3. Extraction of starch by alkali.

Conc. of reagent, %	0.2% NaOH	0.2% Na ₂ CO ₃	H ₂ O	0.2% H ₂ SO ₄	0.2% HCl
Yield of starch, %	22.5	22.3	20.7	21.3	22.6

It can be seen from the data in Table 3 that the amount of starch which was extracted by diluted alkalis is much the same as that obtained by treating with acids. Moreover, the alkaline solution of *Lycoris* root is glutinous and we are unable to isolate from this a starch precipitate.

It appears to us that the yield of starch varies with the time of extraction. 5 g of root powder were immersed into 30 cc of water and after 3 hrs. the mixture was filtered. The residue was ground with water and then it was left stand at 24°C for 1 to 8 days. From these solutions starch particles were gathered by the filtration with a cotton cloth.

Table 4. Relation between time of extraction and yield of starch.

Time of extraction, day	1	2	3	4	5	6	7	8
Yield of starch, %...	40.9	48.3	53.8	56.9	57.2	51.6	47.5	45.5

It will be seen from Table 4 that too short or too long extraction gives rather a low yield of tuber starch. Therefore, the time of immersion must be regulated according to the temperature.

The powder of *Lycoris* bulbs swells in water and this phenomenon seems to have a close connection with the extraction of starch. Accordingly, we selected the particles of the same size out of a large quantity of bulb powder, put them in a graduated tube which has been held in water and after varying time intervals determined the volume of swollen particles. The experiments of this kind resulted that after 1 hr. at 27°C the root powders swelled to 180% and after about 2 hrs. the degree of swelling reached to its maximum, namely 190%.

The figures in Table 5 showed that the increase of particle volume after the immersion in water is affected by the variation of temperature.

Table 5. Variation of swelling by temperature.

Temp. of immersion, °C	15	20	25	30	35	40	45	50	55	60
Degree of swelling, %	164	178	188	208	226	246	258	268	250	200

The degree of swelling is highest at 50°C and over 60°C the peptization takes place. In the experiments mentioned above the time of immersion was 40 min. We further estimated the volume of tuber particles in acid or alkali solutions.

Table 6. Swelling of tuber powder in various solutions.

Temperature of immersion, °C	20	30	40	50	60
Swelling in 1% HCl	210	230	270	215	—
Swelling in 5% NaOH	225	270	300	280	250
Swelling in 1% Na ₂ NO ₃	225	245	250	280	240

As is shown in Table 6, the greatest increase of tuber volume is brought about in sodium hydroxide solution.

In general, it is somewhat difficult to prepare *Lycoris* starch of superior quality in good yield. For example, the glucose determination after acid hydrolysis gave the results represented in Table 7.

Table 7. Purity of *Lycoris* starch preparation.

	Excellent quality	Bad quality
Starch content, %	90.0	73.4

The preparations of inferior quality were made from the residue which remained after the isolation of those of good quality.

2. Action of amylase: There is no report on the enzymatic decomposition of *Lycoris* starch. In the present investigation we used two kinds of enzymes of different origins, viz. malt amylase and Taka diastase. Barley malts were manufactured in this laboratory and dried at 40°C. The extract of malt powder or the suspension of Taka diastase was employed as amylase preparation. For instance, 1 cc of such a enzyme solution was added to the mixture of 25 cc 1% starch solution and 10 cc phosphate buffer. In most experiments the time of reaction was 40 min. and the temperature 55°C.

Table 8. Action of amylase on *Lycoris* starch at various acidities.

pH of solution	4.5	4.8	5.4	5.6	5.9	6.3	6.9	7.2	7.7
Malt; mg glucose	118	122	132	140	136	127	116	89	46
Taka; mg glucose	124	132	135	130	123	121	106	102	82

The data of Table 8 indicate that in alkaline medium the activity of amylase is relatively weak.

In the experiments in Table 9 the acidity of solutions was kept at pH 5.4. The results obtained show that the enzymatic hydrolysis of *Lycoris* starch occurs powerfully even at high temperature.

Table 9. Amylolytic decomposition of starch at different temperatures.

Temp. of reaction	36	45	50	55	60	65	70	75
Malt; mg glucose ...	48	62	66	72	78	79	46	41
Taka; mg glucose ...	52	61	67	65	59	54	36	21

From the data in Table 10 it appears that neutral salts have no influence upon the saccharification of *Lycoris* starch by malt and Taka amylase.

Table 10. Effect of salts upon enzymatic hydrolysis of starch.

Kind of salt	NaCl	NH ₄ Cl	(NH ₄) ₂ SO ₄	KNO ₃	H ₂ O
Malt; mg glucose	67	66	65	62	60
Taka; mg glucose	63	61	58	55	57

The determination of viscosity was carried out with an Ostwald's apparatus. The *Lycoris* starch is easily liquefied by malt amylase, as the figures in Table 11 indicate.

Table 11. Liquefaction of *Lycoris* starch at different hydrogen ion concentrations.

Acidity of solution, pH	5.0	5.4	5.6	5.9	6.2	6.4	6.6
Time of flow, sec.	182	179	177	174	176	178	180

Table 12. Enzymatic liquefaction of starch at various temperatures.

Temp. of reaction, °C.	37	42	47	52	59	65	70
Time of flow, sec.	185	174	180	181	184	187	190

As can be seen from Table 12, the optimum temperature for the liquefaction is 42°C. In these experiments the time of reaction was always 20 min.

SUMMARY

Experiments on the preparation of *Lycoris* starch were carried out under various conditions. It seems somewhat difficult that the starch of excellent quality is isolated from the bulbs in high yield.

Various experiments on the decomposition of *Lycoris* starch by amylase were also performed. It was found that the starch is easily decomposed by malt amylase as well as by Taka diastase.

We wish to express our thanks to Mr. T. Miyakuni and Mr. M. Watayama for their assistance.

REFERENCES

- 1) H. Kondo et al.: Jl. pharm. Soc. Japan, **52**, 433 (1932) **54**, 997 (1934).
- 2) K. Yamafuji and H. Kondo: Jl. Agr. Chem. Soc. Japan, in press.
- 3) T. Akita and T. Nakashima: *ibid.*, in press.
- 4) Y. Kihara: *ibid.*, **9**, 1005 (1933).
- 5) K. Yamafuji and T. Akita: *ibid.*, in press.