The Chrysomelidae of Japan and the Ryukyu Islands. VI : Subfamily Galerucinae I

Kimoto, Shinsaku

Hikosan biological laboratory, Department of Agriculture, Kyushu University

<https://doi.org/10.5109/22724>

出版情報 : 九州大学大学院農学研究院紀要. 13 (2), pp.287-308, 1964-10. Kyushu University
バージョン :
権利関係 :

The Chrysomelidae of Japan and the Ryukyu Islands. VI^{1,2)}
Subfamily Galerucinae I

Shinsaku KIMOTO³⁾

Subfamily GALERUCINAE

Key to Japanese genera of Galerucinae

1. Mesosternum horizontal or inclined, not covered by a process of metasternum 2
 Mesosternum with a somewhat convex, immarginate, long median process
 which is covering the entire horizontal surface of mesosternum 33
2. Antennal insertions generally close, at level of anterior margins of eyes or
 further anterior, occiput and pronotum deeply punctate; last abdominal
 segment of male with a triangular or rounded depression with posterior
 border often emarginate, but never trilobed 3
 Antennal insertions generally separated, situated near, but behind anterior
 borders of eyes; when separated or placed further forward, occiput and
 pronotum not heavily punctate; last abdominal segment of male trilobed,
 with a median lobe always distinct 8
3. Anterior coxal cavities closed behind *Galeruca*
 Anterior coxal cavities open behind 4
4. Seta-bearing pore situated on summit of anterior corner of pronotum, tarsal
 claws bifid or appendiculate, but not differing between different sexes 5
 5
 Seta-bearing pore not situated on summit of anterior corner of pronotum,
 but lateral border near to anterior corner, tarsal claws bifid in male but
 appendiculate in female *Apophyllia*
5. Elytra covered by hairs 6
 Dorsal surface entirely glabrous *Lochmaea*
6. Elytral epipleurae distinct at least on basal 1/2, lateral border of pronotum
 somewhat rounded or sinuate 7
 Elytral epipleurae distinct only on basal 1/4 or less; pronotum subquadrate,
 antennae slender, with second joint elongate but 1/3 as long as third
 Chujoa

¹⁾ Partly supported by a grant from Japan Society, New York City, through
B. P. Bishop Museum, Honolulu, Hawaii, U. S. A.

²⁾ Contribution Ser. 2, no. 210, Entomological Laboratory, Kyushu University.

³⁾ Hikosan Biological Laboratory, Faculty of Agriculture, Kyushu University.

7. Disc of pronotum with a large glabrous space at middle *Galerucella*
 Disc of pronotum entirely covered with hairs, but in some cases anterior
 and lateral margins glabrous *Pyrrhalta*
8. Tarsal claws bifid, anterior coxal cavities opened behind 9
 Tarsal claws appendiculate or simple 11
9. Pronotum without a transverse impression, disc evenly convex.. 10
 Pronotum with a transverse impression, sometimes obsolete at middle
 Aulacophora
10. Body very large and short oval, sides strongly round, elytral epipleurae
 wide, inferior, **recurved** basally *Oides*
 Body oblong, somewhat parallel-sided ; epipleurae of elytra normal, relatively
 narrow and simple *Clerotilia*
11. Anterior coxal cavities opened or half opened behind 12
 Anterior coxal cavities closed behind 29
12. Posterior tibiae without a spine at apex **13**
 Middle and posterior tibiae with a spine at apex 20
13. Anterior and posterior margins of pronotum immarginate 14
 Posterior margin of pronotum marginate 16
14. Elytra without longitudinal carinae 15
 Elytra with two pairs of distinct longitudinal carinae *Japonitata*
15. Pronotum distinctly wider than long; basal margin of elytra without any
 tuberculation *Euliroetis*
 Pronotum almost as long as wide, basal margin of elytra with a pair of
 tuberculations *Cerophysella*
16. Anterior margin of pronotum immarginate 17
 Anterior and posterior margins marginate 19
17. Elytra without distinct *costae* at sides 18
 Elytra with one or two, in some cases three, pairs of longitudinal carinae
 at sides *Haplosomoides*
18. Gena very narrower than transverse diameter of eye, frons not so different
 in both sexes, in male third to four antennal joints reflexed..
 Taumacera
 Gena almost as long as transverse diameter of eye, in male frons with a
 pair of large impressions and with a cylindrical process between antennal
 sockets, antennae almost same in both sexes *Fleutiauxia*
19. Elytral epipleurae rather broad in base, gradually narrowed towards apex ...
 Liroetis
 Elytral epipleurae narrow even in base *Cneorane*
20. Basal margin of pronotum margined; pronotum without a transverse impres-
 sion, in some cases with feeble impressions at sides 21
 Basal margin of pronotum immarginate; pronotum with a distinct transverse
 impression *Paridea*
21. First joint of posterior tarsi shorter than, or subequal to, following two
 joints combined ; anterior margin of pronotum marginate 22
 First joint of posterior tarsi distinctly longer than following two joints
 combined 26
22. Body ovate, very large, convex, pronotum more than twice as wide as long
 Morphosphaera

- Body ovate or oblong ovate, sometime parallel-sided, moderately or scarcely convex; pronotum less than twice as wide as long 23
23. Pygidium punctured uniformly 24
 Basal 1/2 of pygidium shiny, glabrous, finely shagreened, apical 1/2 densely punctured, pubescent; lateral margin of elytra visible from above, body oval, dilated posteriorly, tibiae carinate and finely grooved above
 Agelastica
24. Third antennal joint as long as, or 1 1/2 times as long as second] 25
 Third antennal joint almost, or more than twice as long as second, pygidium uniformly punctate throughout, external margin of elytra not clearly visible throughout from above, oblong-oval, weakly broadened posteriorly ; tibiae not distinctly carinate *Luperus*
25. Prosternal process elevated between coxal cavities, separating them at least anteriorly ; apices of epimera distinct, not extending to infra-coxal lobe; coxal cavities opened or partly opened *Exosoma*
 Prosternal process narrow and not elevated between coxal cavities and not separating them ; apices of epimera distinct ; coxal cavities opened.....
 Calomicrus
26. Dorsal surface glabrous or elytra sparsely covered with short hairs 27
 Dorsal surface thickly covered with fine hairs, densely and finely punctate, elytral epipleurae narrowed near middle *Hesperomorpha*
27. Elytral epipleurae wide at base 28
 Elytral epipleurae narrow, feebly and gradually narrowed from base to apex; pronotum transversely rectangular with straight sides and with distinct sublateral depression, hind femora slightly thickened *Stenoluperus*
28. Frontal tubercles transverse, with anterior angles short; apex of frontal carina inserted between these angles or confluent with them; gena short; antennal insertions at middle of eye; prothorax generally with a transverse setigerous pore a short distance from anterior angle of prothorax; hind tibial spine longer than width of apex of tibiae *Atrachya*
 Frontal tubercles elongate-triangular with anterior angles acute, inserted between antennal insertions and separated by a deep groove; gena slightly shorter than smaller diameter of eye; antennal insertions situated just behind level of anterior borders of eyes; prothorax not broader than long; setigerous pore set back 1/5 to 1/6 length of prothorax from anterior angle; hind tibial spine not longer than width of apex of tibiae *Paralupcrodes*
29. First joint of posterior tarsi shorter than two preceding joints combined 30
 First joint of posterior tarsi longer than two preceding joints combined
 Monolepta
30. Basal border of pronotum not margined except near side 31
 Basal border of pronotum entirely margined 32
31. Pronotum with a pair of distinct impressions laterally *Sermylassa*
 Pronotum without a pair of distinct impression laterally *Hamushia*
32. Posterior tibiae with a single spine at apex; pronotum with a distinct depression laterally ; body oblong *Theopea*
 Posterior tibiae with many short spines at apex, pronotum without deep depression but some times feebly depressed laterally; body oval
 Arthrotus

33. Punctures of elytra distinctly 'and confusedly impressed; pronotum with a transverse furrow which is narrow and runs more than $\frac{3}{4}$ width of pronotum *Agelasa*
 Punctures of elytra partly or largely arranged in longitudinal rows; pronotum with or without a pair of depressions *Gallerucida*

Genus *Galeruca* Geoffroy

Galeruca Geoff r., 1762, Hist. Ins. 1: 251-Latreille, 1810, Considerations Generales, Paris, 432 (type : *Chrysomela tanacetii* Linn.).—Weise, 1886, Ins. Deutschl. 6 (4): 578, 637.—Reitter, 1903, Wien. Ent. Ztg. 22: 133.—Maulik, 1936, Fauna India, Galeruc., 97.—Ogloblin, 1936, Fauna USSR 26, 1: 32, 367, 377.—Gressitt & Kimoto, 1963, Pac. Ins. Man. 1B: 391, 398.

Adimonia Laicharting, 1781, Verz. Tyrol. Ins., 190.—Chapuis, 1875, Gen. Col. 11: 220.

Key to Japanese species of *Galeruca*

1. Anterior corner of pronotum distinctly angulate 2
 Anterior corner of pronotum not angulate but widely rounded; length 11.0 to 12.2 mm *extensa*
2. Lateral margins of pronotum deeply sulcated at anterior half; antennae rather robust, fifth joint distinctly shorter than twice as long as wide; length 9.0 mm *spectabilis*
 Lateral margins of pronotum not or feebly depressed at anterior half; antennae rather slender, fifth joint about three times as long as wide; length 10-11 mm *dahli japonica*

Galeruca ex tensa Motschulsky

Galeruca ex tensa Motschulsky, 1861, Etud. Ent. 10: 22 (Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 162 (Japan).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 399, 401 (Korea, NE China, SE Siberia).

Galeruca bang-haasi Weise, 1894, Deutsche Ent. Zeitschr. 1894: 168 (Japan : Yokohama ; ZMB).—Ogloblin, 1936, Fauna USSR 26, 1: 43, 379, fig. 14b (Ussuri, Manchuria, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 162 (Japan).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 399, 401 (synonymized).

Distribution: SE Siberia, Korea, Manchuria, Japan (3 Kyushu, Honshu, Hokkaido). *Naga* 320: Kamikochi (1 ex., 27. Aug. 1913, M. Suzuki leg.).

Hosts: *Allium* spp. (after Chûjô & Kimoto, 1961).

Galeruca spectabilis Faldermann

Galeruca spectabilis Fald., 1837, Soc. Nat. Moscou, Nouv. Mem. 5: 326, pl. 12, fig. 4 (S. Russia).—Reitter, 1903, Wien. Ent. Ztg. 22 : 136.—Ogloblin, 1936, Fauna USSR 26, 1: 41, 378 (*spectabilis* and *spectabilis orientalis* : S. Russia, Turkey, Iran, Syria).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 399, 403 (Turkey, Syria, Iran, S. Russia, China, Japan).

Galeruca ex tensa : Chûjô & Kimoto, 1961, Pac. Ins. 3 (1) : 162 (Japan).

Distribution: Turkey, Syria, Iran, S. Russia, China, Japan (Honshu).

Yamanashi: Masutomi (1 ex., 26-29. July. 1957, S. Miyamoto leg.). *Fukushima*: Nukemizu, Narahara T, Minamiaizu-gun (1 ex., 5. July. 1947, Y. Kurosawa leg.).

Galeruca dahli japonica Weise

Galeruca dahli var. *japonica* Weise, 1894, Dtsche Ent. Ztschr. 1894: 168 (Japan: Yokohama; ZMB).—Ogloblin, 1936, Fauna USSR 26, 1: 51, 381 (Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 162 (Japan).

Distribution: Japan (Honshu, Hokkaido).

Nagano: Karuizawa. *Yamanashi*: Masutomi. *Saitama*: Urawa City. *Tokyo*: Kamisuwa in Okutama. *Hokkaido*: Engaru in Abashiri; Shikaribetsu in Rubeshibe; Ponkikin in Abashiri; Yufutsu; Ashoro in Tokachi; Obihiro City.

Hosts: *Cirsium* spp.

Genus *Apophylia* Duponchel & Chevrolat

Apophylia hp. & Chev., 1842, in d'Orbigny, 1842, Dict. Univ. d'Hist. Nat. 2: 31.—Allard, 1889, Soc. Ent. Belg., Bull. 33: lxxi.—Maulik, 1936, Fauna India, Galeruc., 78 (type: *Apophylia chloroptera*).—Ogloblin, 1936, Fauna USSR 26, 1: 138, 369.—Chûjô 1962, Phil. Jour. Sci. 91 (1-2): 13, 18.—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 391, 426.

Malaxia Fairmaire, 1878, Soc. Ent. France, Ann. ser. 5, 8: 139 (type: *M. flavovirens* Fairm.).—Allard, 1889, Soc. Ent. Belg., Bull. 33: lxxx.

Glyptolus Jacoby, 1884, Leyden Mus., Notes 6: 62 (type: *G. viridis* Jac.; Sumatra).

Glyceresthis Weise, 1896, Dtsche Ent. Ztschr. 1896: 296 (type: *Auchenia?thalassina* Fald.).

Key to Japanese species of *Apophylia*

1. Legs blackish 3
 Legs almost entirely yellowish brown; black, elytra green; antennae entirely yellowish brown; length 4.6 mm *nigriceps*
2. Prothorax reddish brown; head, except clypeus, stained with reddish brown, antennae and legs black, elytra green; length 6.2-9.0 mm *elongata*
 Prothorax blackish; black, basal joints of antennae and tibiae and apical parts of femora somewhat pale, elytra green; 4.2—5.0 mm. *viridipennis*

Apophylia nigriceps Laboissière

Apophylia nigriceps Lab., 1927, Soc. Ent. France, Ann. 96: 62 (Yunnan-Fou).—Chûjô, 1962, Phil. Jour. Sci. 91 (1/2): 21 (Yunnan, Formosa).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 427, 431 (S. China).

This is the first record of the species from Japan.

Distribution: S. China, Taiwan, Japan (Kyushu).

Oita: Mt. Sobo (1 ex., 27. July 1953, S. Kimoto leg.).

Apophyllia elongata (Jacoby) (Fig. 1)

Malaxia elongata Jæc., 1895, Entomolog. 29: 8 (Loochoos: Oshima).

Apophyllia elongata: Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 157 (Ryukyu Is.).

Distribution: Ryukyu Is. (Amami-Oshima, Okinawa Is.).

Okinawa group: Okinawa Is. (4 exs., 2 -4. Apr. 1955, T. Takara leg.; 1 ex., June. 195-1, G. E. Bohart leg.). *Amami group*: Amami-Oshima (after Nakane & Kimoto, 1959, 1961).

Fig. 1. *Apophyllia elongata* (Jacoby).

Apophyllia viridipennis (Jacoby)

Galerucella viridipennis Jacoby, 1885, 2001. Soc. Lond., Proc. 1885: 744, pl. 46, fig. 9 (Japan: Kurigahara, Wadatoge; BM).

Apophyllia viridipennis: Ogloblin, 1936, Fauna USSR 26, 1: 142, 393 (Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3(1): 157 (Japan).

Distribution: Japan (Honshu).

Yamanashi: Kiyosato (4 exs., 29-31. July. 1957, S. Miyamoto leg.); Shosenkyo (1 es., 22. July. 1936, H. Kamiya leg.).

Genus *Lochmaea* Weise

Lochmaea Ws., 1883, Dtsche Ent. Ztschr. 27: 316 (type: *Chrysomela capreae* L.; Europe); 1886, Ins. Deutschl. 6 (4): 575,

610.—Ogloblin, 1936, Fauna USSR 26, 1: 82, 369, 38-L --Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 392, 416.

Lochmaea capreae (Linnaeus)

Chrysomelacapreae L., 1758, Syst. Nat. ed. 10: 376 (Europe).

Adimon in capreae: Motschulsky, 1866, Soc. Imp. Nat. Moscou, Bull. 39 (1): 175 (Japan).

Lochmaea capreae: Ogloblin, 1936, Fauna USSR 26, 1: 87, 369, 385 (E. Siberia, Manchuria, China, Korea, Japan).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 416 (Europe, Siberia, S. China, Korea, Japan).

Lochmaea capreae cribrata: Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 165 (E. Siberia, Manchuria, China, Korea, Japan).

Oblong oval; pronotum with a pair of lateral and one antero-median and one postero-median depressions, surface rather closely and distinctly punctate; elytra distinctly and more closely punctate than in pronotum; black; pronotum and elytra yellowish brown; scutellum dark reddish brown; length 5.0-6.0 mm..

Distribution: Europe, Siberia, N. China, Korea, Japan (Hokkaido, Honshu).

Ishikawa: Hakusan. *Nagano*: Shirahone; Shimashima. *Gunma*: Sampei Pass; Numata City. *Tochigi*: Nikko. *Aomori*: Yunomata in Shimokita Pen. *Hokkaido*: Ashoro, Nukabira in Takachi; Yukomanbetsu at Mt. Daisetsu; Piuca in Kamikawa; Engaru in Abashiri; Nibushi, Kuccharo at Akan Prov.

Hosts: *Populus* spp., *Salix* spp. (after Chûjô & Kimoto, 1961).

Genus *Chujoa* Gressitt & Kimoto

Chujoa Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 392, 437 (type: *Atysa uetsukii* Chûjô, 1954; Japan).

Chujoa uetsukii (Chûjô)

Atysa uetsukii Chûjô, 1954, Mushi 26 (1): 2, pl. 1, fig. 1 (Japan: Chuka-mura in Okayama Pref.; Chûjô).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 158 (Japan).

Chujoa uetsukii: Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 437 (S. Japan).

Oblong, subparallel-sided; closely covered with fine pubescence; antennae slender, long, about $\frac{3}{4}$ as long as length of body; pronotum subquadrate, with a pair of lateral depressions and a longitudinal furrow; reddish brown with antennae, except basal joints, darker than the other parts of body; length 5.8–8.0 mm.

Distribution: Japan (Honshu, Shikoku, Kyushu).

Oita: Mt. Sobo (8 exs., 24. July–4. Aug. 1955, H. Kamiya leg.). *Kochi*: Nishigoya at Mt. Ishizuchi (1 ex., 31. July. 1952, K. Morimoto leg.).

Host: *Fagus crenata* (after Chûjô & Kimoto, 1961).

Genus *Galerucella* Crotch

Galerucella Crotch, 1873, Acad. Sci. Philad., Proc. 1873: 55.—Weise, 1886, Ins. Deutschl. 6 (4): 575, 622.—Maulik, 1936, Fauna India, Galeruc., 214 (designated *Chrysomela nymphaea* L. as type) (part).—Chûjô, 1962, Phil. Jour. Sci. 91 (1-Z): 13, 37, 39.—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 392, 468.

Hydrogaleruca Laboissiere, 1922, Rev. 2001. Afr. 10: 32 (type: *Chrysomela nymphaea* L.; Europe).—Ogloblin, 1936, Fauna USSR 26, 1: 119, 388 (as a subgenus of *Galerucella*).

Key to Japanese species of *Galerucella*

- Sutural angles of elytra obtuse, rounded; antennae rather slender, seventh to tenth joints more than, or nearly, twice as long as wide; reddish brown; each elytron with an ill-defined submarginal pitchy stripe, which is starting from humerus and running towards apex; antennae black with basal joints paler; vertex and meso- and metathorax, in some cases some parts of abdomen also, blackish; length 3.7–5.2 mm *grisea*
- Sutural angles of elytra subquadrate; antennae rather robust, seventh to tenth joints less than, or nearly, $\frac{1}{2}$ times as long as wide; vertex, antennae, pronotum, meso- and metathorax and elytra largely black, anterior half of head, anterior margin and ventral surface of thorax, lateral margins of elytra and abdomen reddish brown; legs reddish brown with tarsi and outer surface of tibiae black; length 4.8–6.0 mm *22 ipponensis*

Galerucella nipponensis (Laboissière)

Galeruca sagittariae: Baly, 1874, Ent. Soc. London, Trans. 1874:178 (Japan: Hiogo).

Hydrogaleruca nipponensis Lab., Jan. 1, 1922, Rev. Zool. Afrique 10: 120 nota (Japan).

—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1):164 (Japan, Korea, SE Siberia).

Galerucella paludosa Weise, Oct. 12, 1922, Tijdschr. Ent. 65: 68 (Japan: Kyoto).

Galerucella (Hydrogaleruca) nipponensis: Ogloblin, 1936, Fauna USSR 26, 1: 126, 389, fig. 52a, 54 (SE Siberia, Korea, Japan).

Galerucella nipponensis: Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 468, 470 (Japan, Korea, SE Siberia).

Distribution: Korea, SE Siberia, Japan (Honshu, Shikoku, Kyushu).

Fukuoka: Kokubu-cho in Kurume City; Shikano-shima in Kasuya-gun; Najima, Kashii in Fukuoka City. *Miyazaki*: Aoshima.

Hosts: *Brasenia Schreberi*; *Ludwigia ovalis*; *Lycopus lucidus*; *Trapa japonica* (after Chûjô & Kimoto, 1961).

Galerucella grisescens (Joannis)

Galeruca grisescens Joannis, 1866, Abeille 3: 98 (Sicily).

Galleruca vittaticollis Baly, 1874, Ent. Soc. Lond., Trans. 1873:178 (Japan: Nagasaki, Yokohama; BM).

Galleruca distincta Baly, 1874, t.c. (Japan: Nagasaki; BM).

Galerucella (Hydrogaleruca) grisescens: Ogloblin, 1936, Fauna USSR 26, 1: 120, 389 (Japan).

Hydrogaleruca distincta subsp. *yakushimana* Nakane, 1958, Saikyo Univ., Sci. Rep. 2 (5): A308 (Japan: Miyazaki; NSM).

Hydrogaleruca vittaticollis: Chûjô & Kimoto, 1961, Pac. Ins. 3 (1):165 (Japan, Korea, Manchuria, N. China, E. Siberia, Sachalin).

Galerucella grisescens distincta: Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 40 (Japan, Korea, China, Siberia, Formosa).

Galerucella grisescens: Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 468 (Europe, Siberia, China).

Distribution: Taiwan, Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Yakushima), Ryukyu Is. (Ishigaki), Korea, Manchuria, N. China, E. Siberia, Sachalin, Europe.

Sakishimagroup: Ishigaki Is. (2 exs., 20–30. Dec. 1952, G. E. Bohart leg.).

Fukuoka: Mt. Kora in Kurume City; Mt. Wakasugi; Mt. Hiko; Fukuoka City; Kokura City; Tashiro in Yame-gun. *Kumamoto*: Mt. Ichifusa. *Miyazaki*: Aoshima; Hyuga Line; Kiyotake in Miyazaki-gun. *Kagoshima*: Sata-misaki; Shiroyama in Kagoshima City. *Kochi*: Jinzenji in Kochi City. *Nagano*: Karuizawa. *Hokkaido*: Engaru in Abashiri; Aizan-kei at Mt. Daisetsu.

Hosts: *Polygonum* spp.; *Rumex* spp.; *Fragaria chiloensis* var. *ananassa* (after Chûjô & Kimoto, 1961).

Genus *Pyrrhalta* Joannis

Pyrrhalta Joannis, 1866, Abeille 3: 82 (type: *Galeruca viburni* Paykull; Europe).—

Weise, 1886, Ins. Deutschl. 6 (4): 621.—Ogloblin, 1936, Fauna USSR 26, 1: 97, 385.

—Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 37.—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 392, 438.

- Hoplostines* Blackburn, 1890, Linn. Soc. N. S. Wales, Proc. ser. 2, 5 : 361 (type : *Hoplostinesviridipennis* Blackburn, 1890).
- Tricholochmaea* Laboissière, 1932, Mus. Nat. Hist. Nat. Paris, Bull. ser. 2, 4: 963 (type : *L. indica* Lab., India).-Ogloblin, 1936, Fauna USSR 26, 1: 91, 385 (as a subgenus of *Lochmaea*).
- Xanthogaleruca* Lab., 1934, Soc. Ent. France, Ann. 103 : 29, 67 (type : *luteola* Miiller ; Europe).-Ogloblin, 1936, Fauna USSR 26, 1: 100, 386.—Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 37, 38.
- Galerucella*: Maulik, 1936, Fauna India, Galeruc., 214 (part).-Ogloblin, 1936, Fauna USSR 26, 1: 112, 387.
- Clitena*: Ogloblin, 1936, Fauna USSR 26, 1: 129, 369, 391..
- Isshikia* Chûjô, 1961, Ent. Lab., Univ. Osaka Pref., Pub. 6: 87 (type : *I. isshevikii* Chûjô, 1961; Ryukyu Is.) (as a subgenus of *Galeruca*). New **Synonymy**
- Neogalerucella* Chûjô, 1962, Phil. Jour. Sci. 91 (1-2) : 37, 38 (type : *Chrysomelatenella* Linnaeus, 1761; Europe). New **Synonymy**

Key to Japanese species of *Pyrrhalta*

1. Antennae rather robust, fifth to tenth almost, or shorter than, twice as long as wide 2
 Antennae rather slender, fifth to tenth almost, or more than three times as long as wide*.....*..... 9
2. Lateral margin of elytra with a distinct longitudinal convexity which is extending to apex 3
 Lateral margin of elytra without distinct longitudinal convexity which is extending to apex 6
3. Elytral epipleurae distinct on more than basal $3/4$ 4
 Elytral epipleurae distinct on basal $2/3$; brown, vertical area of vertex, middle of pronotum, scutellum, tibiae, tarsi apico-dorsal surface of femora black or blackish ; antennae almost entirely black ; length 5.2-5.3 mm *kawashimai*
4. Abdomen yellowish or reddish; basal margin of pronotum obliquely truncate on its lateral area; smaller than 5.0 mm*5
 Abdomen entirely black; basal margin of pronotum emarginated on its lateral area ; head black with frontal tubercles dark reddish, antennae black with one or two basal joints pale; pronotum, scutellum and elytra reddish brown, with basal area of scutellum infusate; ventral surface and legs black with tibiae and tarsi reddish) brown; length 5.2-6.0 mm *takeii*
5. Elytral epipleurae distinct until apex; dorsal surface entirely yellowish brown ; antennae, especially dorsal surface dark brown ; length 4.0 mm. *konishii*
 Elytral epipleurae not distinct at apical area; dorsal surface largely reddish brown to brown; scutellum black, in many cases pronotum black on middle or entirely black; head, meso- and metathorax and legs variable in coloration, generally ground color of those reddish but in many cases stained with black in various degrees; antennae usually black, in some cases four or five basal joints pale; length 3.0-5.0 mm *semifulva*
6. Third antennal joint subequal to or slightly shorter than fourth 7
 Third antennal joint much longer than fourth8

7. Elytra dark reddish brown with sutural and lateral margins black ; underside black; legs, lower part of head and thorax dark reddish brown ; thorax with three black spots; length 4.2 mm *nigromarginata*
 Elytra dark reddish brown, with a black humeral spot and a lateral stripe situated on same level with humerus; brown, metathorax and abdomen largely black, head with a black spot on vertex, frontal tubercles black; antennae black; prothorax with a large black spot on middle and a pair of broad black stripes situated closely to lateral margin of prothorax; legs blackish, femora and basal part of tibiae reddish brown ; length 6.5 mm . . .
 maculicollis
8. Elytral epipleurae not distinct on their extremities; brown, vertical area of vertex, middle of pronotum, scutellum, humeri, meso- and metathorax, and entire first (visible) to third and middle of fourth segments blackish ; antennae black with basal parts of each joint more or less brownish; length 5.0 mm* *lineola*
 Elytral epipleurae distinct on their extremities; brown, vertical area of vertex, antennae, except two or three basal joints pale, middle of pronotum, scutellum, a broad band running from humerus to subapical area, underside of meso- and metathorax and first to third abdominal segments blackish; length 4.2 mm* (typical form); in some cases brown, antennae except three or four basal joints pale, humeri and underside of meso- and metathorax blackish ; length 4.0 mm (*lythri* type) *calmariensis*

Fig. 2. a, *Pyrrhaltakonishii* Kimoto; b, *P. yasumatsui* n. sp.

9. Apex of scutellum truncate and trapezoidal in form; large in size, larger than 5.5 mm 10
 Apex of scutellum rounded; subtriangular ; brown, antennae dark reddish brown, except three or four basal joints pale; in many cases elytra green-

* These descriptions were prepared by the specimens collected in Germany.

- ish ; length 4.0—4.8 mm *yasumatsui*
10. Elytral punctures very minute and sparsely impressed 11
 Elytral punctures rather strongly and closely impressed 13
11. Elytral epipleurae gradually narrowed towards apex 12
 Elytral epipleurae not narrowed towards apex but subequal in width between preapical and basal areas; yellowish brown ; apical part of femora, dorsal surface of tibiae and tarsi infusate in various degrees; antennae blackish with basal 1/4 to 1/6 of each joint pale; a broad stripe situated closely to lateral margin of prothorax, in some cases a small spot on middle of base also, and a spot on humeri of elytra black ; length 7.0—8.2 mm. *esakii*
12. Third antennal joint more than twice as long as second, a broad stripe situated closely to lateral margin of prothorax, a small spot on middle of base and a spot on humeri of elytra black; length 5.8—6.8 mm
 humeralis
 Third antennal joint one and half times as long as second, a broad stripe situated closely to lateral margin, a small basal spot on middle and a spot on humeri of elytra black; length 7.2-8.8 mm *annulicornis*
13. Elytra entirely fulvous 14
 Elytra dark green with cupreous luster; black, lower part of head, thorax, basal 3/4 of femora and abdomen yellowish brown; thorax with three blackish spots; length 8.0 mm *fuscipennis*
14. Elytra with a distinct longitudinal costa along lateral margin 15
 Elytra without a longitudinal costa along lateral margin; yellowish brown; antennae and tibiae blackish ; length 7.5—8.0 mm *tibialis*
15. Elytra rugosely punctured but those punctures impressed rather shallowly; black, basal part of pronotum and entire elytra yellowish brown; length 7.5 mm *seminigra*
 Elytra distinctly and closely punctured but a diameter of each puncture almost as wide as their interstices, and those punctures rather deeply impressed ; yellowish to reddish brown; antennae, tibiae, tarsi, apex of femora, a marking on vertex, lateral area of pronotum together with a marking before scutellum, and elytral humeri black; length 7.0-8.0 mm...
 issh ikii

Pyrrhalta kawashimai n. sp.

Brown; vertical area of vertex, middle of pronotum, scutellum, tibiae, tarsi and apico-dorsal surface of femora black or blackish; antennae almost entirely black.

Vertex rugosely punctate and covered with fine pubescence; frontal tubercles subquadrate, distinctly raised ; inter-antennal space narrowly carinate. Antennae almost 2/3 as long as length of body, first large and club-shaped; second short, third one and half times as long as second, fourth one and a quarter times as long as third, fourth to seventh subequal in length and shape to each other, eighth to tenth subequal in length and shape to each other, eleventh one and half times as long as tenth and pointed at apex. Pronotum transverse, twice as wide as long, anterior margin almost straight, lateral margin rounded, widest almost at middle, basal margin slightly emarginate at middle and turned obliquo-forward 15 at lateral area, anterior and posterior corners each with a seta-bearing pore;

surface covered with fine pubescence, concaved as a whole, and more deeply concaved on middle and on either side. Scutellum trapezoidal, densely covered with fine pubescence. Elytra broader than prothorax at base, closely and rather distinctly punctured; epipleurae narrowed towards apex and distinct only on basal 2/3.

Length 5.2-5.3 mm.

Holo type: Yugawara, Kanagawa Pref. (2. May. 1955, K. Kawashima leg.) (Entomological Laboratory, Kyushu University).

Paratopotype: 1 ex., same as the holotype.

This species is characteristic in having the elytral epipleurae which are distinct on basal 2/3.

Pyrrhalta takeii (Chûjô) New Combination (Fig. 3b)

Lochmaea (*Tricholochmaea*) *takeii* Chûjô, 1950, Kontyû 18 (5): **114**, 1 fig. (Japan: Numata-cho in Gumma Pref.; CHUJO).

Tricholochmaea takeii: Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 169 (Japan).

Distribution: Japan (Honshu).

Tottori: Hoki-Daisen (1 ex., 24. May. 1954, S. Kimoto leg.).

Pyrrhalta konishii Kimoto (Figs. 2a, 3a)

Pyrrhalta konishii Kimoto, 1963, Fragm. Col., Kyoto (4): 17 (Maruyama in Hokkaido; KU).

Distribution: Japan (Hokkaido).

Fig. 3. a and cl, male genitalia, b and c, pronotum : a, *Pyrrhalta konishii* Kimoto; b, *P. takeii* (Chûjô); c d, *P. semifulva* (Jacoby).

Pyrrhalta semifulva (Jacoby) New Combination (Figs. 3c-d)

Galerucella semifulva Jacoby, 1885, Zool. Soc. Lond., Proc. 1885 : 745, pl. 46, fig. 11 (Japan: Kiga; BM).

Galerucella modesta Jacoby, 1885, loc. cit. (Japan: Nikko; BM).

Lochmaea japonica Weise, 1922, Tijdschr. Ent. 65 : 67 (Japan: Kyoto).

Lochmaea (*Tricholochmaea*) *semifulva* : Ogloblin, 1936, Fauna USSR 26, 1: 91, 385, fig. 38 (E. Siberia, Manchuria, China, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 166 (E. Siberia, Manchuria, China, Japan).

The verification of the record of the species from outside of Japan may be required. *P. limbata* Chen was first described as *semifulva* subsp. *limbata* but Gressitt & Kimoto (1963) treated as an independent species in having characteristic male organ among the species group.

Distribution: E. Siberia, Manchuria, Japan (Hokkaido, Honshu, Sado I., Shikoku, Kyushu).

Fukuoka: Nakatashiro, Tashiro in Yame-gun; Mikazuki-yama in Fukuoka City; Mt. Fukuchi; Kokura City; Mt. Hiko. *Oita*: Mt. Sobo. *Kagoshima*: Shiro-yama in Kagoshima City. *Kochi*: Jinzenji in Kochi City; Kuroson. *Tokushima*: Jinryomura in Myosai-gun. *Osaka*: Minoo. *Nara*: Mt. Yoshino; Mt. Kasuga. *Wakayama*: Koya-san. *Fukui*: Mt. Murakuni in Takefu City. *Nagano*: Asama-Onsen; Wada-toge; Karuizawa. *Yamanashi*: Shosenkyo. *Tochigi*: Nikko.

Hosts: Cherry; *Corylopsis Gotoana*; *Sorbus japonica* (after Chûjô & Kimoto, 1961).

Pyrrhalta nigromarginata (Jacoby)

Galerucella nigromarginata Jacoby, 1885, Zool. Soc. Lond., Proc. 1885: 743, pl. 46, fig. 10 (Japan; BM).

Galerucella (*Xanthogaleruca*) *nigromarginata* : Ogloblin, 1936, Fauna USSR 26, 1: 112, 386 (Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 163 (Japan).

Pyrrhalta nigromarginata : Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 439, 459 (Japan, Hainan Is.).

I have not seen any specimens collected in Japan besides the type. This seems to be a very rare species.

Distribution: Japan, Hainan Is.

Pyrrhalta maculicollis (Motschulsky)

Galeruca maculicollis Motschulsky, 1853, Etud. Ent. 2: 49 (N. China).—Baly, 1874, Ent. Soc. Lond., Trans. 1874: 177 (Nagasaki, Yokohama).

Gallerucella (*Xanthogaleruca*) *maculicollis* : Oglobin, 1936, Fauna USSR 26, 1: 104, 387, fig. 41 (Siberia, Manchuria, China, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 162 (Japan, Korea, Manchuria, N. China, E. Siberia).

Pyrrhalta maculicollis : Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 439, 457 (SE Siberia, E. China).

Distribution: SE Siberia, E. China, Manchuria, Korea, Japan (Hokkaido, Honshu, Sado I., Shikoku, Kyushu).

Tokushima: Kamiyama-mura in Myosai-gun. *Nagano*: Utsukushigahara ; Asama-Onsen. *Yamanashi*: Atago-yama in Kofu City. *Hokkaido*: Yubari in Sorachi ; Nibushi at Akan Prov.; Mt. Daisetsu ; Jozankei nr-. Sapporo City; Ashoro in Tokachi; Obihiro City; Sapporo City.

Hosts: *Ulmus* spp., *Zelkova serrata* (after Chûjô & Kimoto, 1961).

Pyrrhalta lineola (Fabricius) (Fig. 4a)

Chrysomela lineola F., 1781, Spec. Ins. 1: 149 (Europe).

Galerucella lineola: Ogloblin, 1936, Fauna USSR 26, 1: 114, 388 (Europe to Ussuri).

—Chûjô, 1959, Kagawa Univ., Mem. Fac. Lib. Arts & Educ. 2 (81): 11 (Japan : Rishiri-to, Mt. Tokachidake in rang of Mt. Daisetsu-san Nat. Park in Hokkaido).

—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1) 162 (Palearctic region, Japan).

Pyrrhalta lineola: Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B:442, 455 (Europe, N. Asia, NE China, Sachalin, N. Japan).

The illustration of male aedeagus of the species was prepared by a German specimen.

Distribution: Europe, Siberia, Manchuria, Sachalin, Japan (Hokkaido).

Hosts: *Alnus* spp.; *Corylus* spp.; *Salix* spp. (after Chûjô & Kimoto, 1961).

Fig. 4. Male genitalia: a, *Pyrrhalta lineola* (Fabricius) ; b, *P. calmariensis* (Linnaeus).

Pyrrhalta calmariensis (Linnaeus) (Fig. 4b)

Chrysomela calmariensis L., 1767, Syst. Nat. ed. 12: 600 (Europe).

Galerucella lineatipes Takei, 1916, Konchû Sekai, Gifu 20 (226) : 253 (Japan : Gumma Pref.).

Galerucella calvariensis : Ogloblin, 1936, Fauna USSR 26, 1: 115, 388, figs. 46, 47b, 48.—Chûjô, 1956, Kagawa Univ., Mem. Fac. Lib. Arts & Educ. 2 (31): 8, fig. 4 (Japan : Odayama and Wakamatsushi in Fukushima Pref.).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 162 (Palearctic region, Japan).

Pyrrhalta calvariensis : Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B:447 (Europe, N. China).

Distribution : Palearctic region ; Japan (Honshu, Hokkaido).

Hokkaido : Tomakomai City (2 exs., 5. Aug. 1952, S. Ishii leg.).

Hosts : Tn Europe, *Lythrum Salicaria* ; *Stachys palustris*.

Pyrrhalta yasumatsui n. sp. (Fig. 2b)

Brown; antennae dark reddish brown with three or four basal joints pale; elytra in many cases somewhat greenish.

Vertex rugosely punctate and covered with fine pubescence; frontal tubercles convex, subquadrate, separated to each other by a longitudinal groove, inter-antennal space extremely narrow but carinate. Antennae almost 2/3 as long as body length, first joint long and club-shaped, second slender, about 2/3 of first in length, third long, nearly twice as long as second; fourth slightly shorter than third, fifth slightly shorter than fourth and subequal in. length to first. Thorax transverse, twice as wide as long, anterior margin slightly emarginate, lateral margin widest almost at middle, almost straightly narrowed anteriorly and posteriorly, basal margin slightly emarginate at middle and turned obliquo-forwardly at lateral area, anterior and posterior corners each with a seta-bearing pore, surface rugosely punctate and covered with fine pubescence, longitudinally grooved on middle and strongly depressed on each side. Scutellum triangular, apex rounded, covered with fine pubescence. Elytra broader than thorax at base, parallel-sided, lateral margin narrowly marginate, surface closely punctate and closely covered with pubescence ; elytral epipleurae gradually narrowed towards apex.

Length 4.0 -4.8 mm.

Distribution : Ryukyu Is. (Okinawa, Amami-Oshima).

Holotype : Okinawa, Ryukyu Is. (7. May. 1958, T. Takara leg.) (Entomological Laboratory, Kyushu University).

Pam types : 3 exs., Okinawa, Ryukyu Is. (3. May. 1958, T. Takara leg.); 1 ex., Yakkachi (Sumiyo-mura), Amami-Oshima (18. July. 1933, T. Esaki & K. Yasumatsu leg.); 1 ex., Gusuku-Nishinakama (Santarotoge), Amami-Oshima (15. July. 1933, T. Esaki & K. Yasumatsu leg.).

This species is characteristic in having a rounded apex of scutellum and being small in size.

Pyrrhalta esakii Kimoto

Pyrrhalta esakii Kimoto, 1963, Fragm. Col., Kyoto, (4): 17 (Mt. Takao in Tokyo, Shirahone in Nagano Pref., Mt. Yoshino in Nara Pref., Gifu City, Kibune in Osaka, Odaigahara, Hanazono near Kyoto, Katsuragawa near Kyoto; KU).

Distribution : Japan (Honshu),

Pyrrhalta humeralis (Chen)

Galerucella humeralis Chen, 1942, Notes d'Ent. Chinoise 9: 17 (China).

Pyrrhalta humeralis : Nakane & Kimoto, 1961, Kontyû 29 (1) : 21 (Hentona in Okinawa ; Japan).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 439, 451 (China).

Distribution: China, Ryukyu Is. (Okinawa Is.), Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Okinawa group: Okinawa Is. (after Nakane & Kimoto, 1961).

Fukuoka: Mt. Hiko ; Kokura City. *Miyazaki*: Miyazaki City ; Aoidake. *Toku-shima*: Jinryo-mura in Myosai-gun. *Okayama*: Kamo-cho in Tomata-gun. *Fukui*: Mt. Murakuni in Takefu City. *Nagano*: Ontake in Kiso-Fukushima ; Karuizawa ; Wadatoge. *Aomori*: Yunomata in Shimokita Pen. *Hokkaido*: Nukabira in Tokachi ; Mt. Hakodate in Oshima Pen.

Hosts: In China, *Viburnum* sp., *Salix* sp. (after Gressitt & Kimoto, 1963).

Pyrrhalta annulicornis (Baly)

Galleruca annulicornis Baly, 1874, Ent. Soc. Lond., Trans. 1874: 177 (Japan: Hiogo ; BM).

Galerucella (Pyrrhalta) viburni annulicornis : Ogloblin, 1936, Fauna USSR 26, 1: 100, 385 (Manchuria, China, Japan).

Pyrrhalta annulicornis : Chûjô & Kimoto, 1961, Pac. Ins. 3 (1) : 168 (Japan, Ryukyu Is., Korea, N. China, Manchuria, Sachalin).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 439, 445 (Japan, NE China, SE Siberia).

Distribution: Ryukyu Is. (3 Okinawa), Japan (Honshu, Shikoku, Kyushu, Tsushima), Korea, Manchuria, SE Siberia, Sachalin.

Fukuoka: Mt. Hiko. *Oita*: Mt. Kuju ; Mt. Sobo. *Tokushima*: Jinryo-mura in Myosai-gun ; Mt. Kenzan. *Kochi*: Nishigoya at Mt. Ishizuchi. *Tokyo*: Mt. Takao.

Pyrrhalta fuscipennis (Jacoby)

Galerucella fuscipennis Jacoby, 1885, Zool. Soc. Lond., Proc. 1885: 746 (Japan: Aomori ; BM).

Clitena fuscipennis : Ogloblin, 1936, Fauna USSR 26, 1: 132, 392, fig. 56 (Amur, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 160 (E. Siberia, Korea, Japan).

Pyrrhalta fuscipennis : Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 450 (Japan, Korea, E. Siberia).

Distribution: Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yakushima), Korea, E. Siberia.

Kochi: Mt. Kuishi (1 ex., 18. Aug. 1950, K. Morimoto leg.).

Hosts: *Acer* spp. ; *Populus* spp. (after Chûjô & Kimoto, 1961).

Pyrrhalta tibialis (Baly)

Gallerucata tibialis Baly, 1874, Ent. Soc. Lond., Trans. 1874: 176 (Japan : Nagasaki ; BM).

Clitena tibialis : Ogloblin, 1936, Fauna USSR 26, 1: 131, 392 (China, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1) : 161 (Japan, N. China).

Pyrrhalta tibialis : Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 442, 466 (C. China).

Distribution: Japan (Honshu, Shikoku, Kyushu), C. China.

Fukuoka: Shira-shima in Wakamatsu City (1 ex., 23. Sept. 1954, T. Yoshida leg.) ; Sarakura-yama in Yahata City (1 ex., 30. July. 1953, T. Yoshida leg.). *Kyoto*: Daihi (1 ex., 27. May. 1950); Niomon (1 ex., 12. July. 1951, T. Horio leg.).

Host: *Celtis sinensis* var. *japonica* (after Chûjô & Kimoto, 1961).

Pyrrhalta seminigra (Jacoby)

Galerucella seminigra Jacoby, 1885, 2001. Soc. Lond., Proc. 1885 : 744 (Japan : Yagohara ; BM).

Galerucella (*Xanthogaleruca*) *seminigra* : Ogloblin, 1936, Fauna USSR 26, 1: 111, 386 (Japan),

Clitena seminigra : Chûjô & Kimoto, 1961, Pac. Ins. 3 (1) : 160 (Japan).

Pyrrhalta seminigra: Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 411 (Japan).

Distribution: Japan (Hokkaido, Honshu, Shikoku, Kyushu).

Oito: Mt. Sobo (1 ex., 6. Aug. 1954, H. Kamiya leg.). *Tokushima*: Jinryo-mura in Myosai-gun (1 ex., 20. July. 1952, I. Hiura leg.). *Aomori*: Yunomata in Shimokita Pen. (1 ex., 20. July. 1956, K. Morimoto leg.). *Hokkaido*: Mt. Hakodate in Oshima Pen. (1 ex., 31. July. 1959, H. Kamiya leg.).

Pyrrhalta isschikii (Chûjô) New Combination

Galeruca (*Isshikia*) *isschikii* Chûjô, 1961, Ent. Lab., Univ. Osaka Pref., Pub. 6: 86 (Naze in Amami-Oshima ; CHUJO).

Pyrrhalta nigricornis Ohno, 1962, Kontyû 30 (1) : 27 (Amami-Oshima ; OHNO). New **Synonymy**

Distribution: Ryukyu Is. (Amami-Oshima).

Amami group: Amami-Oshima (4 ex., K. Matsui leg.).

Hosts: *Viburnum suspensum*, *V. Awabuki* (after Ohno, 1962).

Genus *Aulacophora* Chevrolat

Aulacophora Chevr., 1837, 'in Dejean, Cat. Col. ed. 2, 378, ed. 3, 402.—Duponchel & Chevrolat, 1842, in d'Orbigny, Dict. Univ. d'Hist. Nat. 2: 337 (type : *Galleruca quadraria* Olivier ; Europe).—Chapuis, 1875, Gen. Col. 11: 158.—Weise, 1886, Ins. Deutschl. 6 (4) : 574.—Laboissière, 1929, Soc. Ent. France, Ann. 98 (3): 256.—Maulik, 1936, Fauna India, Galeruc., 169.—Hincks, 1949, Ann. Mag. Nat. Hist. ser. 12, 2 : 609; 1950, *op. cit.* 3 : 87.—Chen & Kung, 1959, Acta Ent. Sinica 9 (4) : 273.—Chûjô, 1962, Philip. Jour. Sci. 91 (1-2): 14, 74.—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 393, 482.

Raphidopalpa Chevr., 1845, in d'Orbigny, Dict. Univ. d'Hist. Nat. 6: 5 (type : *Crioceris abdominalis* Fab. ; Europe).—Hincks, 1949, Ann. Mag. Nat. Hist. ser. 12, 2 : 620; 1950, *op. cit.* 9: 88.

Raphidopalpa: Rosenhauer, 1856, Tiere Andalus., 325 (type : *Galleruca foveicollis* Lucas ; Europe).—Ogloblin, 1936, Fauna USSR 26, 1: 19, 150, 370, 396.—Hincks, 1949, Ann. Mag. Nat. Hist. ser. 12, 2 : 621.

Ceratia Chapuis, 1876, Soc. Ent. Belg., CR. 19: c (type: *Aulacophora marginalis* Chapuis).—Laboissière, 1929, Soc. Ent. France, Ann. 98 (3): 257.

Triplatys Fairmaire, 1877, Petites Nouvelles Ent. 2 (185) : 186 ; 1879, Mus. Godeffroy, Jour. 14: 118.

Orthaulaca Weise, 1892, Dtsch. Ent. Ztschr. 1892: 392 (type : *Galeruca similis* Oliv.; SE Asia).

Cerania Weise, 1892, Dtsch. Ent. Ztschr. 1892: 392 (type: *Aulacophora cornuta* Baly; SE Asia). —Laboissière, 1929, Soc. Ent. France, Ann. 98 (3): 256.

Spaerarthra Weise, 1892, *t. c.* 392 (type : *Aulacophora cyanoptera* Boisduval ; New Guinea).—Laboissière, 1929, Soc. Ent. France, Ann. 98 (3): 257.

Key to Japanese species of *Aulacophora*

1. Elytra largely yellowish or reddish brown?.....2.
 Elytra black or blue or green 3
2. In male elytra with a tuft of hairs on humeri, and first antennal joint robuster than in female, fifth (visible) abdominal segment trilobed and median lobe deeply excavated on middle and clearly longer than lateral one; in female fifth (visible) abdominal segment sharply notched at apex and its lateral portion clearly depressed ; reddish or yellowish brown; meso- and metathorax and middle and hind legs black, antennae reddish brown but in many cases preceding joints darkened ; rather smaller in size ; length 5.6-7.3 mm *femoralis*
- In male elytra without a tuft of hairs on humeri, first antennal joint normal, fifth (visible) abdominal segment trilobed and median lobe with almost flat surface and not longer than lateral one; in female fifth (visible) abdominal segment entire and rounded at apex; reddish brown, in some specimens each elytron with two or three pairs of black spots, of which one situated near scutellum, one at humeri and another behind middle, abdomen and both of middle and hind legs black, in some cases preceding joints of antennae darkened; rather large in size; length 7.6-8.3 mm *bicolor*
3. Ventral surface of meso- and metathorax black; elytra black or blue or greenish blue ; head, prothorax and abdomen reddish brown, antennae and legs black ; length 5.8-6.3 mm4.
 Ventral surface entirely reddish or yellowish brown 5
4. Elytra black.....*nigripennis nigripennis*
 Elytra blue or greenish blue*nigripennis nitidipennis*
5. In male third antennal joint robuster than female, and second joint slightly robuster than female and its apical portion produced laterally, fifth (visible) abdominal segment trilobed, and its median lobe very deeply excavated on middle; in female third to fifth antennal joints subequal to each other in length ; elytra bluish green ; head, pronotum and abdomen yellowish brown ; antennae and legs largely black ; length 7.5-9.0 mm *loochooensis*
- In male third to sixth antennal joints slightly robuster than female, second joint normal ; in female third antennal joint slightly longer than fourth, the latter distinctly longer than fifth; head, pronotum, abdomen, antennae and legs yellowish brown; elytra black; length 5.3-6.0 mm *lewisi*

Aulacophora bicolor (Weber)

Galleruca bicolor Weber, 1801, Qbs. Ent., 56 (Sumatra).

Galeruca sexpunctata Oliv., 1808, Ent. 6: 627, pl. 2, fig. 29 (Timor).

Aulacophora 6-punctata: Schönfeldt, 1890, Ent. Nachr. 16 (11): 174 (Loochoos : Oshima).

Aulacophora bicolor: Chûjô, 1935, Nat. Hist. Soc. Formosa, Trans. 24: 205 (Loochoos: Iriomote, Ishigaki).— Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 158 (Sunda

I., Ceylon, Philippines, Taiwan, Ryukyu Is.).—Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 77, 86 (India, Andaman Is., Nicobar Is., Sumatra, Java, Timor, Philippines, Formosa, Loochoos).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 483, 484 (China, Taiwan, Ryukyu Is., Philippines, SE Asia, Indonesia).

The specimens taken from Okinawa Is. and Amami-Oshima are all f. *sexpunctata* and those from Sakishima group are all the ninate form.

Distribution: SE Asia, Taiwan, Ryukyu Is. (Iriomote, Ishigaki, Okinawa, Amami-Oshima).

Sakishima group: Ishigaki Is., Iriomote Is. (after Nakane & Kimoto, 1961).

Okinawa group: Okinawa Is. (after Nakane & Kimoto, 1961). *Amami group*: Amami-Oshima (after Nakane & Kimoto, 1961); Kakeroma Is.

Hosts: *Cucurbitaceae* spp. (after Chûjô & Kimoto, 1961).

Aulacophora lewisii Baly

Aulacophora lewisii Baly, 1886, Linn. Soc. Lond., Jour. 20: 5, 24♀ (China ;BM).—Chen, 1959, Acta Ent. Sin. 9 (4): 375, 378 (India, Hongkong).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 484, 488 (India, Ceylon, SE Asia, China, Loochoos, Japan).

Orthaulaca (Ceratia) cattigarensis Ws., 1892, Dtsch. Ent. Ztschr. 1892: 397 (Japan ; Shanghai).

Aulacophora cattigarensis: Ogloblin, 1936, Fauna USSR 26, 1: 156, 397, fig. 65a (Japan, China, Annam, Tonkin, Cochinchina).—Chûjô, 1957, Kontyû 25 (1): 15 (Loochoos: Amami-Oshima).—Chen, 1959, Acta Ent. Sin. 9 (4): 375, 379, figs. 6-7 (China, India, Vietnam, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 158 (Japan, Ryukyu, S. China, Indo-China).—Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 78, 99 (China, Indo-China, Japan, Loochoos, Formosa, incl. Percadore, Samosana).

Distribution: India, Ceylon, SE Asia, Japan (Yakushima), Ryukyu Is. (Iriomote, Ishigaki, Miyako, Okinawa, Amami-Oshima), Taiwan.

Okinawa group: Okinawa Is. (after Nakane & Kimoto, 1961). *Amami group*: Amami-Oshima (after Nakane & Kimoto, 1961).

Hosts: *Cucurbitaceae* spp. (after Chûjô & Kimoto, 1961).

Aulacophora femoralis (Motschulsky)

Rhaphidoplaba femoralis Mots., 1857, Etud. Ent., 6: 37 (Japan).—Ogloblin, 1936, Fauna USSR 26, 1: 154, 396 (Korea, ? China, Japan).

Aulacophora femoralis: Chen & Kung, 1959, Acta Ent. Sin. 9 (4): 375, 385 (Taiwan, Japan, ? Korea).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 158 (Japan, Ryukyu Is., Korea, N. China).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 483, 486 (E. Siberia, China, Hainan, Vietnam, Korea, Japan).

Distribution: Japan (Honshu, Aogashima, Hachijo-jima, Hachijo-kojima, Awashima, Sado I., Oki I., Shikoku, Okinoshima, Kyushu, Tsushima, Yakushima, Kuchinoerabu-jima), Ryukyu Is. (Tokara, Amami-Oshima, Kita & Minami-Daito-jima, Okinawa, Taketomi, Miyako, Ishigaki, Iriomote), Korea, China, Taiwan.

Sakishima group: Miyako Is. Iriomote Is. & Ishigaki Is. (after Nakane & Kimoto, 1961). *Okinawa group*: Okinawa Is. (after Nakane & Kimoto, 1961). *Amami group*: Amami-Oshima (after Nakane & Kimoto, 1961). *Tokara group*: Nakanoshima & Takarajima (after Nakane & Kimoto, 1961).

Fukuoka: Fukuoka City. *Kumamoto*: Tatsuta-yama in Kumamoto City. *Yaku-*

shima : Amboo. *Kochi* : Kuroson. *Okayama* : Kamo-cho in Tomata-gun. *Kyoto* : Yoshida in Kyoto City. *Fukui* : Mt. Murakuni in Takefu City. *Yamanashi* : Shosenkyo; Komagatake. *Miyagi* : Sendai City.

Hosts : *Cucurbitaceae* spp. (after Chûjô & Kimoto, 1961).

Aulacophora loochooensis Chûjô

Aulacophora loochooensis Chûjô, 1957, Kagawa Univ., Mem. Fac. Lib. Arts & Educ. 2 (52): 2 (Loochoos : Shinmura-Akatsuchiyama in Amami-Oshima ; CHUJO).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 158 (Ryukyu Is.).

Distribution : Ryukyu Is. (Amami-Oshima).

A mami group : Yuwan in Amami-Oshima (2 exs., 5. Apr. 1958, M. Takahashi leg.); Shinmura in Amami-Oshima (1 ex., 2. Apr. 1958, M. Takahashi leg.).

Hosts : *Cucurbitaceae* spp. (after Chûjô & Kimoto, 1961).

Aulacophora nigripennis nigripennis Motschulsky

Aulacophora nigripennis Mots., 1857, Etud. Ent. 6: 38 (Japan).—Ogloblin, 1936, Fauna USSR 26, 1: 156, 397, figs. 64, 65b (Amur, China, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 159 (Japan, Korea, E. Siberia, Manchuria, China, Formosa).—Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 79, 91 (Japan, Loochoos, Formosa, Korea, Manchuria, China, Amur, Ussuri).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B:484, 489 (E. Siberia, China, Korea, Japan).

Distribution : E. Siberia, Manchuria, China, Taiwan, Korea, Japan (Honshu, Awa-shima, Sado I., Oki Is., Shikoku, Okinoshima, Kyushu, Tsushima, Goto Is., Yakushima).

Fukuoka : Magaribuchi in Sawara-gun; Tashiro in Yame-gun; Mt. Wakasugi; Mt. Hiko ; Nishi-koen, Hirao in Fukuoka City; Kokura City ; Mt. Kora in Kurume City ; Shikanoshima in Kasuya-gun ; Mt. Fukuchi; Shirashima in Wakamatsu City. *Kagoshima* : Sata-misaki. *Miyazaki* : Mt. Osuzu. *Yakushima* : Amboo. *Kochi* : Ashizuri-misaki. *Kyoto* : Kibune in Kyoto City. *Nara* : Mt. Yoshino. *Kanagawa* : Yugawara. *Miyagi* : Sendai City.

Hosts : *Cucurbitaceae* spp. (after Chûjô & Kimoto, 1961).

Aulacophora nigripennis nitidipennis Chûjô

Aulacophora (Ceratia) nitidipennis Chûjô, 1935, Nat. Hist. Soc. Formosa, Trans. 25: 82 (Loochoos: Nago in Okinawa; Naze in Amami-Oshima; Iriomote I.).

Aulacophora nigripennis nitidipennis : Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 159 (Ryukyu Is.).

Distribution : Ryukyu Is. (Amami-Oshima, Okinawa, Ishigaki, Iriomote).

Sakishima group : Ishigaki Is. (after Nakane & Kimoto, 1961). *Okinawa group* : Okinawa Is. (after Nakane & Kimoto, 1961). *Amami group* : Amami-Oshima (after Nakane & Kimoto, 1961).

Hosts : *Cucurbitaceae* spp. (after Chûjô & Kimoto, 1961).

Genus *Oides* Weber

Oides Weber, 1801, Obs. Ent., 26 (type : *Chrysomela bipunctata* F.).—Laboissiere, 1922, Soc. Ent. France, Ann. 90: 194.—Maulik, 1935, Fauna India, Galeruc., 105.

- Ogloblin, 1936, Fauna USSR 26, 1: 18, 145, 395.—Hincks, 1949, Ann. Mag. Nat. Hist. ser. 12, 2: 617.—Chûjô, 1962, Phil. Jour. Sci. 91 (1-2): 14, 61.—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 392, 474.
- Adorium* Fabricius, 1801, Syst. Ent. 1: 409 (type : *Chrysomela bipunctata* F.).—Chapuis, 1875, Gen. Col. 11: 156.
- Ochrælea* Chevrolat, 1837, in Dejean, Cat. Col. ed. 2, 375, ed. 3, 399 (type: *Adorium flavum* Olivier, 1807; monobasic); 1845, in d'Orbigny, Dict. Univ. d'Hist. Nat. 6: 5; 1846, loc. cit. 8: 713.—Hincks, 1949, Ann. Mag. Nat. Hist. ser. 12, 2: 617.
- Rhombopalpa* Chevrolat, 1837, in Dejean, Cat. Col. ed. 2, 375, ed. 3, 399 (type : *Adorium decempunctata* Billberg; monobasic); 1845, in d'Orbigny, Dict. Univ. d'Hist. Nat. 6: 5.—Hincks, 1949, Ann. Mag. Nat. Hist. ser. 42, 2: 617.
- Botanctona* Fairmaire, 1877, Petites Nouvelles Ent. 2 (185): 185; 1897, Mus. Godefroy, Jour. 14: 113.
- Boisduvallia* Montrouzier, 1885, Soc. Agr. Lyon, Ann. 7: 72.

Oides bow ringii (Baly)

- Adorium bowringii* Baly, 1863, Ent. Soc. Lond., Trans. ser. 3, 1: 623 (China : Hong Kong; BM).
- Oides bowringi* : Seki, 1933, Konchu-kai 1 (5): 494 (Japan : Kobe).—Ogloblin, 1936, Fauna USSR 26, 1: 149, 396 (Korea, China, Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 167 (China, Korea, Japan).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 475, 476 (S. China, Korea, Japan).
- Large in size, round, strongly convex; yellowish brown; elytra largely bluish black with lateral and apical margins broadly and sutural margin narrowly yellowish brown; antennae yellowish brown with four apical joints blackish; legs yellowish brown with tarsi and apex of tibiae blackish; length 13-15 mm.
- Distribution*: S. China, Korea, Japan (Honshu).
- Hyogo*: Karasubara in Kobe City (1 ex., 4 May. 1937, M. Uno leg.); Mt. Rokko in Kobe City (1 ex., 12 Aug. 1951, Y. Wada leg.).
- Hosts*: *Kadsura japonica*; *Schisandra nigra* (after Chûjô & Kimoto, 1961).

Genus *Clerotilia* Jacoby

- Clerotilia* Jac., 1885, 2001. Soc. Lond., Proc. 1885: 751 (type : *C. flavomarginata* Jac.; Japan).—Ogloblin, 1936, Fauna USSR 26, 1: 217, 370.—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 393, 479.

Clerotilia flavomarginata Jacoby

- Clerotilia flavomarginata* Jacoby, 1885, 2001. Soc. Lond., Proc. 1885: 751, pl. 46, fig. 12 (Japan : Tsumago, Nara; BM).—Ogloblin, 1936, Fauna USSR 26, 1: 218, 407 (Japan).—Chûjô & Kimoto, 1961, Pac. Ins. 3 (1): 160 (Japan).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 480, 482 (Japan, E. China).
- Oblong, subparallel-sided, slightly convex; antennae long, slender but in male third to eighth joints robuster than in female and strongly incrassate at apex; pronotum smooth; elytra finely granulate, rather distinctly and closely punctate; reddish brown to dark reddish brown; elytra green with slight cupreous luster, and its lateral and apical margins narrowly brownish; antennae piceous to black; length 4.3-5.8 mm.

Distribution : E. China, Japan (Honshu, Shikoku, Kyushu).

Fukuoka : Mt. Hiko; Mt. Fukuchi; Mt. Wakasugi; Hirao in Fukuoka City.
Kochi : Jinzenji in Kochi City. *Nagano* : Karuizawa ; Ontake in Kiso-Fukushima.
Yamanashi : Obinayama, Atago-yama in Kofu City.

Host : *Rhamnella fraguloides* (after Chûjô & Kimoto, 1961).

Genus *Japonitata* Strand

Japonia Weise, 1922 (nec Gould), Tijdschr. Ent. 65: 70 (type : *Phyllobrotica nigrita* Jacoby ; Japan).-Ogloblin, 1936, Fauna USSR 26, 1: 196, 371.

Japonitata Strand, 1935, Folia Zool. Hydrobiol. 7: 294 (new name for *Japonia* Weise, nec Gould, 1859).—Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B:394,501.

Japonitata nigrita (Jacoby)

Phyllobrotica nigrita Jacoby, 1885, 2001. Soc.Lond.,Proc. 1885 :742, pl. 46, fig. 3 (Japan : Osaka ; BM).

Japonin nigrita : Ogloblin, 1936, Fauna USSR 26, 1: 196, 403 (Japan).

Japonitata nigrita : Chûjô & Kimoto, 1963, Pac. Ins. 3 (1): 165 (Japan).---Gressitt & Kimoto, 1963, Pac. Ins. Mon. 1B: 501 (Japan).

Oblong ovate, convex; pronotum smooth, impunctate and with a pair of depressions laterally; elytra with two pairs of lateral costae, which start from humerus and of which interior one is interrupted at base; shining black, abdomen yellowish ; length 4.0—5.8 mm.

Distribution : Japan (Honshu, Shikoku, Kyushu).

Fukuoka : Mt. Hiko (1 ex., 15. Aug. 1954, T. Okuma leg.). *Kumamoto* : Mt. Ichifusa (1 ex., 26. July. 1952, S. Kimoto leg.). *Kochi* : Ashizuri-misaki (1 ex., 27. Apr. 1956, Y. Wake leg.). *Kyoto* : Seriu (1 ex., 14. June. 1954, M. Nakayama leg.).