On the bionomics of *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) (Anthocoridae, Heteroptera)

Chu, Yau-i
Entomological Laboratory, Department of Agriculture, Kyushu University

<https://doi.org/10.5109/22783>

出版情報：九州大学大学院農学研究院紀要. 15 (1), pp.1-136, 1969-03. Kyushu University
バージョン：
権利関係：

On the bionomics of *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) (Anthocoridae, Heteroptera) *

Yau-i CHU

Contents

1	Introduction	1
2	Historical review	2
3	Economic importance of Anthocorid bugs as natural enemies	3
4	Systematics and descriptions	6
5	Habitat	25
6	Laboratory observation on the life history	26
7	Field observations	41
8	Preying habits	47
9	Effects of food on the fecundity	68
10	Rearing with artificial media	75
11	Ovarian development under certain nutritional conditions	83
12	Effect of population density on the survival, growth and fecundity	87
13	Overwintering and cold-hardiness	92
14	Relative toxicity of pesticides	95
15	Inter-specific competition between <i>L. beneficus</i> and <i>X. galactinus</i>	104
16	Co-existence with <i>Apanteles chilonis</i> [Munakata]	108
17	Approach to the mass production	110
18	Attack of bugs to man	113
19	Conclusion-Evaluation of bugs as predators against the rice stem borer...	114
20	Summary	119
21	Acknowledgement	123
22	Literature cited	124

Introduction

In certain rice growing countries, the rice stem borer (*Chilo suppressalis* (Walker)) is causing serious damage to rice, and to combat the borer, an extensive use of insecticides is necessary to obtain a standard yield. On the other hand, the feasibility of biological control of the borers has been recently re-emphasized, and voluminous papers have been published in these

* Contribution Ser. 2, No. 288, Entomological Laboratory, Kyushu University.
Special Project Research supported by the Ministry of Education "Studies on the
Dynamic Status of Biosphere." This study was carried out as part of JIBP.

several years. The list of natural enemies of rice stem borers was compiled by Nickel (1964), Rao and Chawla (1964), Yasumatsu (1967), Yasumatsu and Torii (1968) and others. Yasumatsu recorded 79 species of parasites and 20 species of insect predators from Asian area. Up to the present, many prominent works have been conducted on some parasitic insects, while little is known on the predatory natural enemies of the borers, although they also contain very effective members. For example, the Anthocorid bug, *Lyctocoris beneficus* (Hiura), sometimes occurs abundantly in the straw piles and barns, attacking overwintering larvae and the newly emerged adult moths of rice stem borer in the spring. According to Oho's investigation, during the spring of 1953, more than 90 percent newly emerged adults of the rice stem borer were attacked by this bug in the straws in Kyushu, Japan (Oho 1955a, b). The importance of this bug as a controlling agent against the overwintering rice stem borer during the colder season was also emphasized by some workers (Kobayashi 1958, Bess 1967, Yasumatsu 1967, etc.). Although some fragmental informations on this Anthocorid predator have been recorded by some workers, it seems extremely necessary to make a detailed study of this bug before the practical trial of this Anthocorid as a biological control agent against the rice stem borer. The purpose of the present work is to get fundamental informations to evaluate the availability of *L. beneficus* and *Xylocoris galactinus* (Fieber), another Anthocorid, which is often more predominantly found than *L. beneficus* in the similar environment, as natural enemies against the rice stem borer.

Historical review

Lyctocoris beneficus (Hiura) was at first described by Hiura (1957) as a new species of *Euspudaeus*, but the history of the finding of this species goes back to 1953. In this year great numbers of this bug were discovered in the giant straw piles at Saga Paper Board Co. in Kyushu, Japan, and they were considered to be an effective predator in checking the emergence of rice stem borer moths in late spring. On the other hand, the high mortality of this bug against BHC was reported (Anonym 1953). In the next year, the susceptibility of this bug against several insecticides was tested in Saga Agricultural Experimental Station, and at the same time some brief biological notes were recorded (Anonym 1954). Oho (1955a) made a preliminary report of this bug under the name of *Euspudaeus* sp. and recorded fragmental ecological notes as well as a highly susceptibility against BHC. This same preliminary report was also published in another magazine (Oho 1955b). In the same year, he published the feeding habit and life history of this bug as the first report of the study of this bug (Oho 1955c), and some illustrations relating to this bug were also published (Mizukami and Oho 1955). In 1957, Hiura described the adult as a new species of *Euspudaeus*, and gave the description of each stage of the nymphs of this bug together

with its biological data. His materials were collected from 31 localities which were covering the entire Japan except Hokkaido. The anatomical studies on the ovariole and alimental canal of this bug as well as *Xylocoris galactinus* (Fieber) were conducted by Miyamoto (1957, 1961). The generic name of this bug was later revised to *Lyctocoris* by I-Iiura (1966). The distribution of this bug is now confined to Japan and Quelpart Island in Korea (Miyamoto and Lee 1966).

On the contrary, *Xylocoris galactinus* (Fieber) was discovered more than 130 years ago. At first, Fieber (1836) described the species under the name of *Anthocoris galactinus*. Since then, it has long been known as *Piezostethus galactinus* (Fieber). Its taxonomical, morphological and biological studies as well as collecting data have been accumulated by many workers (Reuter 1884, Poppius 1909, Butler 1923, Blatchley 1926, Carayon 1953, Southwood and Les-ton 1959, Hiura 1966, etc.). Contrary to the long year studies of this species, the nymphal morphology was first described by Hall (1951), and he also studied its ecology in the grain stock yard. The biology under the same environment was later discussed by Sinha (1961). Thus, the bug is known as a cosmopolitan species and recorded from Europe, North and South America, Asia, North Africa and Micronesia etc., and Herring (1966) attributed its wide distribution to the transportation of bug-infested food-stuff.

Economic importance of Anthocorid bugs as natural enemies

Both *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) belong to the family Anthocoridae, which include numerous small and inconspicuous species living in flower and other more or less hidden situations, and commonly known as the name of "flower bug" or "minute pirate bug." A number of Anthocorid bugs were recorded as the predators of plant mites, Lepidopterous larvae and eggs, thrips, leafhoppers, psyllids, scale insects, whiteflies and similar small or soft sclerotized insects. However, their predatory behaviour had been almost overlooked because of their small size and inconspicuous habit. The importance of these bugs as the biological control agents has been pointed out by some workers (Massee and Steer 1926, Ishihara 1941, Carayon 1961, etc.). For example, *Orius insidiosus* (Say) which preys on more than 40 species of insects and mites was recognized as an important predator of *Tetranychus bimaculatus* Harvey (McGregor 1913), *T. pacificus* McGregor (Michelbacher et al. 1952), *Paratetranychus pilosus* C.-F. (Garman and Jewett 1938, Lord 1949, Johansen and Brekey 1949), *Taeniothrips gladioli* M.-S. (Richardson 1933), *Blissus leucopterus* (Say) (Flint 1918), *Thrips tabaci* Lindeman (Ogilvie 1927), *Heliothis obsoleta* (Fabricius) (Phillips and Barber 1933, Barber 1936a, b, Hanison 1960), *H. zea* (Boddie) (Bell and Whitecomb 1964), *Psallus seriatus* Reuter and *Alabama argillacea* Hiibner (McGregor

1942). In addition to these, some experimental works on the biological control of insect pests or mites using Anthocorid bugs, such as *Paratriphleps laeviusculus* (Champion), *Orius insidiosus* (Say), *O. tristictus* (White), *Montandoniola moraguesi* Puton, *Tetraphleps* sp. etc. have been undertaken by several American entomologists (Annand 1942a, b, Weber 1953, Sweetman 1958, Davis and Krause 1963, 1964, 1965a, b, Funasaki 1966, Mitchell and Wright 1967).

In Japan, the studies on the Anthocorid bugs as the natural enemies were rather few and all the records on the bugs with exact specific names were compiled by Yasumatsu and Watanabe (1964). Besides, Ito et al. (1960) and Oku and Kobayashi (1966) reported the predation of *Orius* spp. on *Pieris rapae crucivora* Boisduval and some species of aphids in soybean field respectively.

Before giving the author's observation on the predation of *L. beneficus* and *X. galactinus*, the author gives the world records of data which have been observed by many authors on the Lepidopterous hosts preyed by various species of Anthocorids. As mentioned previously, Lepidopterous larvae are actually the most favourable hosts of the Anthocorid bugs, and sometimes their eggs become more attractive food for the bugs. The oldest preying data was introduced by Douglas and Scott (1865) who reported the preying of *Anthocoris nemoralis* (Fabricius) and *A. nemorum* (Linné) on the *Lithocolletis* spp. (Lithocolletidae). Since then, 11 species of predators and 19 species of prey have been recorded by several workers.

Although *Lyctocoris campestris* (Fabricius) and *Xylocoris cruditans* (Fallen) were described far back to 1794 and 1807 respectively, studies on their preying habits have been neglected. The first preying record of *Lyctocoris* bug was made by Corti (1921) who observed the attack of *L. campestris* on the silkworm larvae as well as prepupa through the wall of its cocoon, but later Malenotti (1925) opposed Corti's observation and wrote that the bug does not prey upon the prepupa through its cocoon. Two other preying records of *L. campestris* were made on the potato tuber moth (*Phthorimaea terrella* Zeller) and Tussock moth (*Hemerocampa vetusta* Boisduval) as given in Table 1. The association of *L. campestris* with stored products was reported by Howe (1952) who also indicated the predation of this bug on Khapra beetle (*Trogoderma granarium* Evert). The preying of another *Lyctocoris* bug was observed by Fronk (1947). He observed that *L. elongatus* (Reuter) was feeding on the pine beetle *Dendroctonus frontalis* Zimmerman.

The first preying data of *Xylocoris* bug was recorded by Trägårdh (1914) who found *X. cruditans* in the burrows of bark beetles and confirmed the predation of the bugs on their larvae. The preying data on another bark beetle, *Ips typographus* Linné, was reported by another worker (Mokrzecki 1923). The attack on *Tribolium confusum* Duval by *X. cruditans* was observed by Good (1936). Nutting and Gerhardt (1964) found that *X. flavipes* (Reuter) was abundant in grain storage which was heavily infested by Khapra beetle,

Table 1. Preying record of Anthocorid bugs on Lepidopterous insects
(Arranged in chronological order).

Predator	Host	Locality	Reference
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Garman et al. 1914
<i>Orius</i> sp.	<i>Spilonota ocellana</i>	U.S.A.	Wilson et al. 1915
<i>Orius</i> sp.	<i>Pectinophora gossypiella</i>	Egypt	Willcocks 1916
<i>Orius</i> sp.	<i>Spilonota ocellana</i>	Canada	Duport 1917
<i>Lyctocoris campestris</i>	<i>Bombix mori</i>	Italy	Corti 1921
<i>Orius tantilus</i>	<i>Pectinophora gossypiella</i>	India	Ballard 1921
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Phillip et al. 1923
<i>Orius australis</i>	<i>Heliothis obsoleta</i>	Australia	China 1926
<i>Lyctocoris campestris</i>	<i>Phthorimaea terrella</i>	U.S.A.	Underhill 1926
<i>Orius australis</i>	<i>Heliothis obsoleta</i>	Australia	Veitch 1927
<i>Orius insidiosus</i>	<i>Laphygma frugiperda</i>	U.S.A.	Luginbill 1928
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Winburn et al. 1932
<i>Orius</i> sp.	<i>Heliothis 'obsoleta</i>	S. Africa	Parsons et al. 1934
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Barber 1936a, b.
<i>Anthocoris musculus</i>	<i>Eulia mariana</i>	Canada	Gilliatt 1937
<i>Orius sauteri</i>	<i>Ostrinia nubilalis</i>	Japan	Koo 1940
<i>Scoloposcelis parallelus</i>	<i>Scirpophaga nivella</i>	Java	Diakonoff 1940
"	<i>Scirpophaga auriflua</i>	"	"
"	<i>Proceras venosatus</i>	"	"
"	<i>Chilotraea auricilia</i>	"	"
"	<i>Ostrinia nubilalis</i>	"	"
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Marcovitch et al. 1941
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Annand 1942a
<i>Paratriphleps laeviusculus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Annand 1942b
<i>Orius</i> sp.	<i>Heliothis virescens</i>	Peru	Anonym 1942
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Fletcher et al. 1943
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Ewing et al. 1943
<i>Orius insidiosus</i>	<i>Heliothis obsoleta</i>	U.S.A.	Barber 1943
<i>Paratriphleps laeviusculus</i>	<i>Heliothis virescens</i>	Peru	Hambleton 1944
<i>Orius albidipennis</i>	<i>Prodenia litura</i>	Egypt	Kamel 1951
<i>Paratriphleps laeviusculus</i>	<i>Heliothis virescens</i>	Peru	Wille 1951
<i>Orius tristicolar</i>	"	"	"
<i>Paratriphleps laeviusculus</i>	<i>Heliothis virescens</i>	Peru	Simon 1954
<i>Orius tristicolar</i>	"	"	"
<i>Orius insidiosus</i>	<i>Ostrinia nubilalis</i>	U.S.A.	Bartholomai 1954
<i>Anthocoris musculus</i>	<i>Spilonota ocellana</i>	Canada	Stultz 1955
<i>Orius tristicolar</i>	<i>Platynota stultana</i>	U.S.A.	Atkins et al. 1957
<i>Lyctocoris campestris</i>	<i>Hemerocampa vetusta</i>	U.S.A.	Atkins 1958

Predator	Host	Locality	Reference
Orius sp.	Prodenia sp.	Egypt	Hassan et al. 1960
Orius sp.	<i>Pieris rapae crucivora</i>	Japan	Ito et al. 1960
Orius insidiosus	<i>Heliothis obsoleta</i>	U.S.A.	Hanison 1960
Orius sp.	Bucculatrix thurberialla	U.S.A.	Tuttle et al. 1961
Orius sp.	<i>Heliothis zea</i>	U.S.A.	Wene et al. 1962
Orius insidiosus	<i>Heliothis zea</i>	U.S.A.	Bell et al. 1964

though its feeding habit was not observed. But they listed the bug as its predator. The beneficence of *X. flavipes* and *L. campestris* was postulated by Bosvine (1951) writing as follow : "Those bugs may be encountered in food stores, especially large ones such as warehouse and granaries. — they are quite harmless to man and even beneficial, since they live sucking the juice of the beetles and moths or sometimes of mites." A Tingid bug, *Gargaphia bimaculata* Parshley added to a prey list of *Xylocoris* bug by Bondar (1924). Besides, some predatory observations of *X. galactinus* were recorded by Hall (1951).

Systematics and description

The family Anthocoridae occurs in almost all parts of the world and includes more than 400 described species. The character of this family was summarized by Blatchley (1926) as follows : "very small oval or oblong-oval sub-depressed bugs having the head long, porrect, inserted in thorax to or almost to eyes; tylus stout, prominent, protruding between and in front of bases of antennae, its apex blunt ; ocelli present ; bucculae wanting ; beak 3-segmented, its apex acuminate, the first segment usually shorter than head ; antennae 4-segmented ; pronotum short, more or less trapezoidal ; scutellum small, triangular ; mesonotum in part or wholly visible ; elytra usually present, covering the abdomen and with a distinct cuneus and embolium, membrane without closed cells, its veins few or wanting ; legs short, subequal in length, the front ones not raptorial, tarsi 3-segmented; meta-sternum with a more or less distinct, osteolar channel connecting with the scent glands. Males with a single asymmetrical genital plate ; genitals of female with a narrow medial sheath flanked on each with two larger triangular plates."

The family was divided into three subfamilies by Blatchley (1926) and China and Miller (1955, 1959).

Family Anthocoridae (Amyot & Serville)

Amyot & Serville, 1843. Hist. Nat. Hemipt., pp. 37, 262 (Anthocorides).

Fieber, 1851. Genera Hydroc., p. 9 (Anthocoridae).

Subfamily Lyctocorinae (Reuter)

- Reuter, 1884. Monog. Anthoc., p. 4. Div. Lyctocorina.
Champion, 1900. Biol. Centr. Am., Heter., 2: 306. Div. Lyctocoraria.
Oshanin, 1909. Verz. palae. Hemip., 1 : 613. Div. Lyctocoraria.
Oshanin, 1912. Kat. palae. Hemip., p. 58. Tribe Lyctocoraria.
Van Duzee, 1917. Cat. Hemip. Am. N. Mexico, p. 288. Subfam. Lyctocorinae.

Subfamily Anthocorinae (Reuter)

- Reuter, 1884. Monog. Anthoc., p. 4. Div. Anthocoraria.
Champion, 1900. Biol. Centr. Am., Heter., 2: 319. Div. Anthocoraria.
Oshanin, 1909. Verz. palae. Hemip., 1: 613. Div. Anthocoraria.
Oshanin, 1912. Kat. palae. Hemip., p. 56. Tribe Anthocoraria.
Van Duzee, 1917. Cat. Hemip. Am. N. Mexico, p. 292. Subfam. Anthocorinae.

Subfamily Dufouriellinae Van Duzee

- Reuter, 1884. Monog. Anthoc., p. 4. Div. Xylocoraria.
Champion, 1900. Biol. Centr. Am., Heter., 2: 330. Div. Xylocoraria.
Oshanin, 1909. Verz. palae. Hemip., 1: 633. Div. Xylocoraria.
Van Duzee, 1916. Check list Hemip., p. 35. Subfam. Dufouriellinae.
Van Duzee, 1917. Cat. Hemip. Am. N. Mexico, p. 295. Subfam. Dufouriellinae.

These subfamilies are easily distinguished by the following key (China and Miller 1959).

- 1a Basal margin of pronotum not deeply emarginate ; scutellum not transversely sulcate ; cell of the hind wings with a hamus..... 2
- b Basal margin of pronotum deeply emarginate; scutellum transversely sulcate ; cell of the hind wings without a hamus.
..... Subfamily Dufouriellinae
- 2a Third and fourth segments of antennae slender, linear and beset with numerous long hairs ; hamus with a connecting vein; front femora usually more or less swollen. Subfamily Lyctocorinae
- b Third and fourth segments of antennae short, fusiform, with hairs very short ; hamus without a connecting vein and run from subtended or decurrent vein; frontal femora never swollen.
..... Subfamily Anthocorinae

Subfamily Lyctocorinae was firstly established by Reuter (1884) by the name of Division Lyctocoraria and this name was later adopted by some authors and finally changed to the present name by Van Duzee and Edward (1916). The species of this subfamily have the third and fourth antennal segments much more slender than the preceding segments, and longer than the diameter of the segments. The hairs on the third and fourth antennal segments are long and remarkable. In hind-wings, hamus runs from the connecting vein, except in the case of the genus *Plochicoris*. The subfamily includes 14 genera and they are:

1. ***Asthenidea*** Reuter
Reuter, 1884. Monog. Anthoc. p. 602.
2. ***Eulasiocolpus*** Champion
Champion, 1900. Biol. Centr. Am., Heter., 2 : 313.
3. ***Euspudaeus*** Reuter
Reuter, 1884. Monog. Anthoc., p. 565.
4. ***Iella*** Carayon
Carayon, 1958. Memo. Inst. Sci. Madag., (E) 9 : 335.
5. ***Lasiochiloides*** Champion
Champion, 1900. Biol. Centr. Am., Heter., 2 : 311.
6. ***Lasiochilus*** Reuter
Reuter, 1871. Öf. Vet. Aka. Förh., 28: 562.
7. ***Lasiocolpoides*** Champion
Champion, 1900. Biol. Centr. Am., Heter., 2 : 313.
8. ***Lasiocolpus*** Reuter
Reuter, 1884. Monog. Anthoc., p. 581.
9. ***Lepidophorella*** Poppius
Poppius, 1909. Acta Soc. Sci. Fenni., 37 (9) : 43.
10. ***Lilia*** White
White, 1879. Ent. mon. Mag., 16 : 147.
11. ***Lyctocoris*** Hahn
Hahn, 1835. Wanz. Ins., 3 : 19.
12. ***Oplobates*** Reuter
Reuter, 1895. Ent. mon. Mag., 31 : 171.
13. ***Plochiocoris*** Champion
Champion, 1900. Biol. Centr. Am., Heter., 2 : 311.
14. ***Xylocoris*** Dufour
Dufour, 1831. Ann. Sci. Nat. Paris, 22: 423.

These genera may be distinguished by the following key:

- 1a Wings without hamus in the cell. *Plochiocoris* Champion
- b Wings with hamus in the cell. 2
- 2a Sides of pronotum marginate and explanate, not setose: fore femora with a tooth beneath at about distal one-fourth. *Lilia* White
- b Sides of pronotum not or narrowly or incompletely explanate, always distinctly setose, fore femora not so armed. 3
- 3a Head about twice as long in front of eyes as length of an eye. 4
- b Head measured from above, not conspicuously longer in front of eye than length of an eye. 10
- 4a Osteolar channel on metapleuron short or long, forwardly curved. 5
- b Osteolar channel on metapleuron short, backwardly curved. 7
- 5a Osteolar channel short.*..... *Lasiocolpoides* Champion

- b Osteolar channel long. 6
- 6a Membrane of hemielytra with a single distinct vein, abdominal apex with several long bristle-like hairs. **Lasiochiloides** Champion
- b Membrane of hemielytra with 4 distinct veins, abdominal apex ciliate. **Oplobates** Reuter
- 7a Sides of hemielytra serrately emarginate, rostrum reaching to metacoxae. **Lasiocolpus** Reuter
- b Sides of hemielytra not emarginate, rostrum extending to mesocoxae. 8
- 8a Notum covered with yellowish normal hairs. **Lasiochilus** Reuter
- b Notum covered with scale-like hairs. 9
- 9a Hairs on hemielytra composing tufts. **Iella** Carayon
- b Hairs on hemielytra not composing tufts. **Lepidophorella** Poppius
- 10a Osteolar channel curved forward or backward, with a rounded angle. 11
- b Osteolar channel nearly straight, joining at a sharp right angle. 13
- 11a Frontal femora more or less swollen, clavus impunctate. **Xylocoris** Dufour
- b Frontal femora long and slender, clavus with two or three rows of punctures. ... 12
- 12a Eyes very large and oval, without a neck, sides of pronotum strongly sinuate. **Eulasiocolpus** Champion
- b Eyes not large, with a stout and cylindrical neck behind eyes, sides of pronotum not sinuate. **Asthenidea** Reuter
- 13a Sides of pronotum distinctly marginate, the apical angles wide and dull, not flattened, hemielytra with or without luster, thickly punctate. **Lytcocoris** Hahn
- b Sides of pronotum sinuate and not marginate, the apical angles flattened, hemielytra shining, coarsely punctate. **Euspodaes** Reuter

The genus *Lytcocoris* Hahn was erected to contain a single species, *domesticus* (Hahn 1835). Subsequently, Reuter (1871a) placed ***domesticus*** as a synonym of *Acanthia campestris* Fabricius. Reuter (1871b) erected the genus ***Dolichomerus*** with two included species, but later considered it as a synonym of *Lytcocoris* Hahn (Reuter 1884) and the genus remains taxonomically stable to the present.

Genus ***Lytcocoris*** Hahn

Haplotype, ***domesticus*** Hahn= ***campestris*** (Fabr.)

Hahn, 1835. Wanz. Ins., 3: 19.

Fieber, 1869. Wien Ent. Monat., 3: 264.

Flor, 1860. Rhyn. Livl., 1 : 665 (Subgenus of ***Xylocoris***).

Fieber, 1861. Europ. Hemip., p. 38.

Douglas & Scott, 1865. Brit. Hemip., p. 498.

Reuter, 1871. Öf. Vet. Aka. Förh., 27 : 409 (***Dolichomerus***).

Reuter, 1884. Monog. Anthoc., p. 560.

Champion, 1900. Biol. Centr. Am., Heter., 2: 309.

Provancher, 1886. Pet. Fauna Ent. Can., 3: 90 (***Tetraphleps***).

(Haplotype=*canadensis* Prov.).

Van Duzee, 1917. Cat. Hemip. Am. N. Mexico, p. **288**.

Blatchley, 1926. Heter. E-N. Am., p. 623.

Zimmerman, 1948. Ins. Hawaii 3: 174.

Kelton, 1962. Can. Ent. 94 : 1302.

Kelton, 1967. Can. Ent. 99 : 807,

This genus may be known by the following characters:

Body oblong or oblong-oval, usually subopaque almost glabrous species having the head across as wide as long, narrower than the front of pronotum, eyes large, placed close to the base of head. Ocelli situated near the inner margins of the eyes. Tylus stout, its apex truncate. Antennae: 1st and 2nd segments thick, 1st cylindrical, reaching to the end of the face ; 2nd gradually clavate, about $2\frac{1}{2}$ times as long as 1st ; 3rd and 4th thin, filiform, subequal in length, with prominent bristles. Rostrum long, reaching to the apex of the metasternum; 1st segment scarcely so long as the head, 2nd $1\frac{2}{3}$ times as long as the 1st. Pronotum subtrapezoidal, with its anterior margin sinuate, wider than the head across the eyes; anterior angles depressed, rounded ; side margins thick, raised, with a slight linear impression within; posterior margin concave. Scutellum large, impressed at the middle. Hemelytra long oval, reaching apex of abdomen, clavus concave; claval suture depressed, corium flat ; anterior margin reflexed; cuneus deflected; membrane with 1 vein at the corner of the cell. Mesosternum short, very slightly convex, posteriorly with a broad flat depression. Metasternum right angled, base convex, end depressed. Legs all alike, slender, while front coxae short, subcontiguous; front femora elongate, fusiform, but slightly swollen, unarmed. Osteolar channel nearly straight jointing at a sharp right angle a very fine carina which is extending to the front margin of metasternal plate. Males with a short spongy fossa at the apex. According to Kelton (1967), this genus is easily distinguished from the other Anthocoridae by the shape of 5 following points : 1) the osteolar channel, 2) 3rd and 4th antennal segments, 3) antero-lateral margins of the pronotum, 4) male genitalia and 5) female abdominal segments.

This genus includes 24 species, 5 from Palaearctic, 8 from Nearctic, 6 from Ethiopian, 6 from Neotropical, 2 from Oriental and Australasian Regions. Among them one species is world-wide in distribution.

1. *Lyctocoris albifer* Walker

Lyctocoris albifer Walker, 1872. Cat. Heter. Brit. Mus. 5: 154.

Madeira.

2. *Lyctocoris beneficus* (Hiura)

Euspodaeus beneficus Hiura, 1957. Sci. Bull. Fac. Agr. Kyushu Univ. 16 (1): 31.

Lyctocoris beneficus Hiura, 1966. Bull. Osaka Mus. Nat. Hist. 19 : 33.

Japan, Korea.

3. *Lyctocoris campestris* (Fabricius)

Acanthia campestris Fabricius, 1794. Ent. Syst. 4: 75.

Salda campestris Fabricius, 1803. Syst. Rhyn. p. 116.

Anthocoris campestris : Stål, 1868. Hemip. Fabr. 1 : 90.

Lyctocoris carnepestris : Reuter, 1871. Öf. Vet. Aka. Förh. 28: 409.

Cimex pallidus Rossi, 1794. Mant. Ins. 2: 55.

Lygaeus arvicola Latreille, 1804. Hist. Nat. Crust. Ins. 12: 220.

Phytocoris pallens Fall&, 1829. Hemip. Suec., Cimic. p. 103.

Cimex domesticus Schilling, 1834. Isis p. 738.

Lyctocoris domesticus Hahn, 1835. Wanz. Ins. 3 : 20.

Anthocoris domesticus : Herrich-Schaeffer, 1853. Wanz. Ins. 9 : 228.
Anthocoris bicuspis Herrich-Schaeffer, 1835. Nomen. Ent. 1 : 60.
Xylocoris dimidiata Spinola, 1840. Ess. Hemp. p. 236.
Xylocoris parisiensis Amyot & Serville, 1843. Hemip., p. 264.
Xylocoris americanus Dallas, 1852. List Hemip. 2 : 589.
Cardiastethus currax Garbiglietti, 1869. Bull. Ent. Ital. 1 : 123.
Lyctocoris fitchii Reuter, 1871. Öf. Vet. Aka. Förh. 28 : 557.

Cosmopolitan.

4. ***Lyctocoris canadensis*** Kelton

Lyctocoris canadensis Kelton, 1967. Can. Ent. 99 (8) : 807.

Canada.

5. ***Lyctocoris cohici*** Delamare

Lyctocoris cohici Delamare, 1952. Encyl. Biogeogr. ecol. 8: 81.

Ivory coast.

6. ***Lyctocoris doris*** Van Duzee

Lyctocoris doris Van Duzee, 1921. Proc. Calif. Acad. Sci. 11 : 137.

Canada, U. S. A.

7. ***Lyctocoris dorni*** E. Wagner

Lyctocoris dorni E. Wagner, 1941. Gulde, Wanz. 8: 223.

Europe, Syria.

8. ***Lyctocoris elongatus*** (Reuter)

Dolichomerus elongatus Reuter, 1871. Öf. Vet. Aka. Förh. 28: 558.

Lyctocoris elongatus Reuter, 1884. Monog. Anthoc. p. 565.

u. s. A.

9. ***Lyctocoris hasegawai*** Hiura

Lyctocoris hasegawai Hiura, 1966. Bull. Osaka Mus. Nat. Hist. 19 : 36.

Formosa.

10. ***Lyctocoris hawaiiensis*** (Kirkaldy)

Nesidiocheilus hawaiiensis Kirkaldy, 1902. Fauna Hawaii. 3 (2) : 93.

Lyctocoris hawaiiensis : Zimmerman, 1948. Ins. Hawaii. 3: 174.

Hawaii.

11. ***Lyctocoris latus*** Poppius

Lyctocoris Zatus Poppius, 1909. Acta Soc. Sci. Fenn. 37 (9) : 42.

Peru.

12. ***Lyctocoris longirostris*** Horváth

Lyctocoris longirostris Horváth, 1911. Bull. Mus. Paris p. 216.

Dahomey.

13. ***Lyctocoris lugubris*** Poppius

Lyctocoris lugubris Poppius, 1909. Acta Soc. Sci. Fenn. 37 (9) : 42.

West Africa.

14. ***Lyctocoris mexicanus*** Kelton

Lyctocoris mexicanus Kelton, 1966. Can. Ent. 98 (3) : 320.

Mexico.

15. ***Lyctocoris nidicola*** E. Wagner

Lyctocoris nidicola E. Wagner, 1955. Notul. Ent. 35 (2): 61.

Finland.

16. ***Lyctocoris obsoletus*** (Blanchard)

Anthocoris obsoletus Blanchard, 1852. Illist de Chile 2001. 7: 140.

Lyctocoris obsoletus: Signoret, 1863. Ann. Soc. Ent. Fr. p. 566.

Chile.

17. ***Lyctocoris okanaganus*** Kelton & Anderson

Lyctocoris okanaganus Kelton & Anderson, 1962. Can. Ent. 94 (12): 1302.

Canada, U. S. A.

18. ***Lyctocoris rostratus*** Kelton & Anderson

Lyctocoris rostratus Kelton & Anderson, 1962. Can. Ent. 94 (12) : 1302.

Canada, U. S. A.

19. ***Lyctocoris signoreti*** Reuter

Lyctocoris signoreti Reuter, 1884. Monog. Anthoc. p. 563.

Venezuela.

20. ***Lyctocoris spangbergi*** Reuter

Lyctocoris spangbergi Reuter, 1884. Monog. Anthoc. p. 562.

Venezuela.

21. ***Lyctocoris stalii*** (Reuter)

Dolichomerus stalii Reuter, 1871. Öf. Vet. Aka. Förh. 28 : 558.

Dolichomerus reuteri White, 1879. Ent. mon. Mag. 16: 146.

Lyctocoris stalii Reuter, 1884. Monog. Anthoc. p. 564.

Canada, U. S. A.

22. ***Lyctocoris subelegans*** Breddin

Lyctocoris subelegans Breddin, 1928. Denskschr. Med. Ges. Naturw. Jena 17: 81.

South Africa.

23. ***Lyctocoris tuberosus*** Kelton & Anderson

Lyctocoris tuberosus Kelton & Anderson, 1962. Can. Ent. 94 (12) : 1302.

Canada, U. S. A.

24. ***Lyctocoris uyttendoogaati*** Blöte

Lyctocoris nyttendoogaati Blöte, 1926. Tidsk. Ent. 72: 163.

Canary Is.

The Palaearctic and Oriental species are characterized by the following

key (almost followed Wagner (1955)).

- 1a Vertex narrower than twice of eye width. *L. hasegawai*
- b Vertex wider than twice of eye width. 2
- 2a Vertex 2-2.2 times as wide as an eye. *L. dorni*
- b Vertex 2.6-3.1 times as wide as an eye. 3
- 3a Second segment of antennae distinctly longer than width of head. 4
- b Second segment of antennae as long as width of head. 5
- 4a Lateral margins of pronotum marginated, cuneus shallowly punctate.
..... *L. campestris*
- b Lateral margins of pronotum not marginated, cuneus rugose not punctuated.
..... *L. beneficus*
- 5a Vertex 2.9-3.0 times as wide as eye, length of posterior margin of pronotum about
1.5 times as long as anterior margin, *L. nidicola*
- b Vertex 2.6-2.7 times as wide as eye, length of posterior margin of pronotum about
2.0 times as long as anterior margin. *L. uytenboogaati*

The taxonomical studies of *L. beneficus* was first made by Hiura (1957, 1966), and his former paper contains the descriptions of adult and nymphs.

Description of *L. beneficus* :

Egg. The egg of *L. beneficus* is elongate and slightly curved. It is rounded at one end and provided with chalky white operculum at the opposite end. The operculum is roughly circular in outline and divided into two distinct zones. The marginal zone consists of a ring of radially elongate "cells" roughly rectangular in shape, such columnar system composed of about 27 cells. In width this zone occupies approximately half the diameter of the operculum. The second or inner zone is slightly convex provided with 15-20 rounded processes. The operculum is normally the only portion exposed on the filter paper in laboratory condition, and chalky white in appearance in a newly oviposited egg, while the colour changes to pinkish red hue after several hours of oviposition.

Average 1.15 mm in length (range 1.13-1.18), and 0.35 mm in width (range 0.32-0.37).

First instar. Immediately after the eclosion, the nymph is pale yellow in coloration. Head : pale yellow with eyes deeper brilliant red colour, sutures obscurely visible; antennal segments I-III darker in colour, IV nearly colourless or very pale straw-coloured, diameters of segments subequal; rostrum pale orange in colour while dark grey in distal part, and much shorter than antennae or about three-fourths as long as antennae. Thorax : in general colour more dusky than head ; pronotum pale yellowish, rectangular in shape; mesonotum with a pale yellowish widening posteriorly; mesonotum pale yellow, paler than the preceding, lateral parts of thoracic segments deeply tinged. Legs : pale yellow, femora slightly darker than tibiae and tarsi. Abdomen : orange-red with two long ciliae on the last segment.

Second instar. Head : brownish yellow, eyes brownish red, suture pale and well marked ; antennae pale, suffused with dusky brown, especially on distal part of segment I, entire part of II and base of IV, III and IV more slender than I and II in diameter, III and IV with prominent hairs; rostrum brownish, deeper on distal part. Thorax: all segments yellowish with slightly brownish tinge, margins with reddish tinge; wing-pads slightly

observable. Legs yellow with dusky suffusion. Abdomen: segments I and II yellow as in thorax; remainder pale, with dark internal organs visible through the hinder part of integument, posterior margin of the last segment pinky tinged, with two developed long ciliae.

Third instar. Head : brownish yellow with pale sutures; eyes red, surrounded by a dirty brown green area; antennal segments I-III dusky grey, with pale distal parts on segments II and III, the basal one-third of segment IV dull grey; rostrum yellowish brown, distal part deep brown. Thorax: dirty brown yellow with a pair of large dirty yellow-green areas on the sides of a longitudinal central line, tinge of the area slightly lighter on mesonotum; wing-pads nearly extending to the posterior margin of the first abdominal segment. The anterior margin of segment II approaching to wing-pads with pinky hue. Legs: dusky grey, similar in colour to basal part of pronotum, dusky grey in the antero-dorsal part of each femur. Abdomen : segments I-III dark yellowish brown, remaining segments deep in colour with dirty brown area at the central part, terminal cilia absent.

Fourth instar. Head : colour as in the preceding instar, but more brownish, suffused with dark tinged; eyes red; antennal segments I-III dark grey to black with distal one-eighth of II, III pale cream coloured, IV pale coloured, slightly dark distally. Thorax: colour as in the third instar but more brown tinged; wing-pads more developed, exceeding beyond the posterior margin of first abdominal segment. Legs: pale testaceous with grey-black suffusion. Abdomen: colour as in the third instar.

Fifth instar. Head: deep dark brown coloured; antennal segments, I-III dark grey to black, with distal extremities narrowly pale creamy white, greyish hue most distinct on III, IV dark grey on basal part, remainder brownish-yellow; rostrum darker at median part, with one pinkish area on each lateral side of the base. Thorax : pronotum deep brownish black coloured, becoming darker laterally: mesonotum and metanotum brown coloured medially, dark red-brown on antero-lateral regions, becoming yellowish-brown posteriorly on wing-pads. Legs: testaceous with dark grey suffusion, especially on femora and fore-tibia. Abdomen: yellowish brown with large black area except the last segment, the area on segments VII, VIII and VIII separated in mesial part, segment I very faintly pink. Fore wing-pads entirely covering hind wing-pads, extending to middle of segment II.

Adult. Head: black, distal part pale brown, upper surface coarsely punctate, except oral area and around eyes, ventral surface reddish black; eyes deep brown; antennae with all segments dark brown except basal part of II which is yellow brown, with distinct hairs, hairs on III and IV longer than diameter of the segments; rostrum brown in colour, reaching the posterior margin of mesonotum. Thorax: pronotum subtrapezoidal, with a longitudinal median canal, anteriorly levigate, with two large triangular depressions posteriorly, anterior margin curved a little, posterior margin markedly sinuate, about three times as wide as anterior margin, lateral margins gently curved, scutellum wider than long, 4 : 3 in proportion, divided by a transverse sulcus into an anterior raised and rounded portion and a posterior portion with transverse striations. Osteolar channel curved forwardly at a right angle, running parallel with outer margin and extending to anterior margin of pleuron. Elytra coarsely punctate, cuneus smooth, clavus and cuneus wide, embolium comparatively narrow, membrane with four subparallel veins, anterior one distinct. Legs: very dark brown apex of fore- and middle-tibia each with a distinct small spongy furrow, tarsi dark brown or black, three-segmented.

The measurements of body parts of each stage are given in Tables 2-4.

Table 2. *Lyctocoris beneficus*; dimensions of nymphs and adult in millimeters. (I)

Instar No.	Head length	Head width	Pronotum length	Pronotum width	Abdomen length	Total length
1	0.24	0.22	0.13	0.25	0.45	1.10
2	0.27	0.23	0.18	0.37	0.85	1.86
3	0.38	0.36	0.32	0.63	1.10	2.53
4	0.39	0.36	0.36	0.74	1.62	2.92
5	0.41	0.39	0.44	0.91	2.08	3.72
Adult	0.46	0.54	0.69	1.27	2.12	3.84

(Avg of 5 individuals).

Table 3. *Lyctocoris beneficus*; dimensions of nymphs and adult in millimeters. (II)

Instar No.	Antennal segments					Rostral segments				
	I	II	III	IV	total	I	II	III	IV	total
1	0.06	0.18	0.15	0.16	0.55	0.03	0.10	0.18	0.13	0.44
2	0.09	0.27	0.21	0.24	0.81	0.05	0.15	0.27	0.18	0.65
3	0.12	0.32	0.25	0.30	0.99	0.08	0.18	0.36	0.24	0.86
4	0.16	0.36	0.26	0.34	1.12	0.09	0.21	0.40	0.26	0.96
5	0.17	0.51	0.32	0.35	1.30	0.11	0.26	0.48	0.30	1.15
Adult	0.19	0.58	0.39	0.39	1.54	0.11	0.35	0.54	0.39	1.38

(Avg of 5 individuals).

Table 4. *Lyctocoris beneficus*; dimensions of nymphs and adult in millimeters. (III)

Instar No.	Fore leg			Middle leg			Hind leg		
	Femur	Tibia	Tarsus	Femur	Tibia	Tarsus	Femur	Tibia	Tarsus
1	0.21	0.16	0.11	0.20	0.10	0.04	0.25	0.20	0.08
2	0.38	0.33	0.18	0.34	0.37	0.13	0.51	0.56	0.21
3	0.53	0.47	0.25	0.52	0.48	0.23	0.72	0.75	0.24
4	0.61	0.54	0.26	0.55	0.58	0.26	0.76	0.87	0.25
5	0.77	0.73	0.27	0.68	0.64	0.27	1.01	1.04	0.26
Adult	0.77	0.77	0.28	0.92	0.65	0.28	1.15	1.09	0.27

(Avg of 5 individuals).

Each instar of *L. beneficus* nymphs is easily distinguishable by the following key.

- 1a Posterior margin of abdomen with cilia. 2
- b Posterior margin of abdomen without cilia. 3
- 2a Wing-pads not visible. 1st instar
- b Wing-pads slightly visible. 2nd instar
- 3a Wing-pads not exceeding posterior margin of the first abdominal segment. 3rd instar
- b Wing-pads exceeding posterior margin of the first abdominal segment. 4
- 4a Hind wing-pads partly covered with fore ones, without black brown areas on dorsal surface of 1st and 2nd abdominal segments. 4th instar
- b Hind wing-pads entirely covered with fore ones, with black brown area on dorsal surface of abdomen, especially prominent on 1st and 2nd segments. 5th instar

The genus *Xylocoris* Dufour was erected to contain a single species *rufipennis* by Dufour (1831). In 1833, Dufour added the second species, *X. ater* which was later moved to the genus *Dufouriella* by Kirkaldy. Since then, many species had been included in this genus. On the other hand, Fieber (1861) erected a new genus *Piezostethus* designating *P. cursitans* as the haplotype. Later *Piezostethus* was proved to be a synonym of *Xylocoris* by Kirkaldy (1906).

Genus *Xylocoris* Dufour

Haplotype *rufipennis* Duf. = *cursitans* (Fall&).

- Dufour, 1831. Ann. Sci. Nat. Paris 22: 423.
 Dufour, 1833. Ann. Soc. Ent. Fr. 1 (2) : 106.
 Burmeister, 1835. Handb. d'Ent. 2 : 289.
 Spinola, 1837. Ess. Hemip. p. 235.
 Amyot & Serville, 1843. Hemip. p. 235.
 Herrich-Schaeffer, 1853. Wanz. Ins. 9 : 170.
 Stål, 1865. Hemip. Afr. 3 : 23.
 Kirkaldy, 1906. Trans. Am. Ent. Soc. 32 : 119.
 Reuter, 1912. Öf. Fin. Vet. Soc. Fork. Liv. Afd. (A) 7: 25.
 Van Duzee, 1917. Cat. Hemip. Am. N. Mexico p. 290.
 Blatchley, 1926. Heter. E-N. Am. p. 627.
 Zimmerman, 1946. Ins. Hawaii. 3: 175.
 Herring, 1967. Ins. Micronesia 7 (8) : 142.

Genus *Piezostethus* Fieber

- Fieber, 1869. Wien Ent. Monat. 4: 365.
 Fieber, 1861. Europ. Hemip. pp. 38, 139.
 Douglas & Scott, 1865. Brit. Hemip. p. 500.
 Reuter, 1871. Öf. Vet. Aka. Förh. 28: 410.
 Reuter, 1884. Monog. Anthoc. p. 29.
 Champion, 1900. Biol. Centr. Am. Heter. 2: 315.

Oshanin, 1909. Verz. palae. Hemip. 1: 615.

Banks, 1910. Cat. Nearc. Hemip. p. 24.

The genus *Xylocoris* is characterized by the following features:

Small, oval or oblong-oval, subglabrous shining, having the head about as long as wide across the eyes, hidden in thorax almost to eyes, eyes large, ocellus distinct and situated close to the base of eyes. Rostrum reaching to or rather beyond the base of mesonotum, 1st segment about $1/2$ the length of head; 2nd about $2 \frac{1}{2}$ times longer than the 1st; 3rd about as long as the 1st. First segment of antennae cylindrical, very short, not reaching to the apex of head, 2nd gradually clavate, about $2 \frac{1}{2}$ times as long as the 1st; 3rd and 4th thin, filiform, subequal in length, taken together much longer than the second. Pronotum subtrapezoidal, without an apical constriction, smooth and shining, hind margin feebly sinuate or subtruncate, with a transverse post-median impression, sides in front of middle strongly deflexed, not margined or ciliated, base sinuate, scutellum nearly flat, slightly concave. Hemelytra usually dimorphic, in macropterous form elongate, parallel-sided, membrane with distinct veins, reaching tip of abdomen, in brachypterous form not further back than fourth abdominal segment, legs short, femora slightly incrassate, especially on fore legs, front tibiae of males much enlarged near apex, obliquely truncate, spinose beneath. Osteolar channel long, curved forwards from middle, extending almost or quite anterior margin of metapleura.

This genus includes 38 species, 23 from Palaearctic, 9 from Nearctic, 5 from Neotropical, 5 from Australasian, 4 from Oriental, and 8 from Ethiopian Regions. Among them 2 species are known as cosmopolitan species.

1. *Xylocoris albonotatus* (Champion)

Piezostethus albonotatus Champion, 1900. Biol. Centr. Am. Heter, 2:316.

Guatemala, Panama.

2. *Xylocoris afer* (Reuter)

Piezostethus afer Reuter, 1884. Monog. Anthoc. p. 592.

Tunisia, South Africa, Egypte.

3. *Xylocoris antaoensis* E. Wagner

Xylocoris antaoensis E. Wagner, 1957. Comm. Biol. Helsing. 16 (2) : 3.

Cape Verde Is.

4. *Xylocoris balteatus* Walker

Xylocoris balteatus Walker, 1872. Cat. Heter. Brit. Mus. 5: 159.

Piezostethus balteatus: Reuter, 1884. Monog. Anthoc. p. 714.

Madeira.

5. *Xylocoris betulinus* Drake & Harris

Xylocoris betulinus Drake & Harris, 1926. Proc. Biol. Soc. Wash. 39 : 37.

U. S. A.

6. *Xylocoris bimaculatus* (Champion)

Piezostethus bimaculatus Champion, 1900. Biol. Centr. Am. Heter. 2: 316.

Guatemala.

7. *Xylocoris californicus* (Reuter)

Piezostethus californicus Reuter, 1884. Monog. Anthoc. p. 600.

Canada, U. S. A.

8. *Xylocoris canariensis* E. Wagner

Xylocoris canariensis E. Wagner, 1955. Comm. Biol. Helsing. 14 (2) : 27.

Canary Is.

9. *Xylocoris ciliatus* (Jakovlev)

Piezostethus ciliatus Jakovlev, 1877. Bull. Soc. Mosc. 52 (2) : 300.

U. S. S. R.

10. *Xylocoris congoensis* (Bergroth)

Piezostethus congoensis Bergroth, 1905. Ann. Soc. Ent. Belg. (1905) : 386.

Congo.

11. *Xylocoris contiguus* E. Wagner

Xylocoris contiguus E. Wagner, 1955. Comm. Biol. Helsing. 14 (2) : 24.

Canary Is.

12. *Xylocoris cursitans* (Fallén)

Lygaeus cursitans Fallén, 1807. Mon. Cimic. Suec. p. 74.

Anthocoris cursitans Fallén, 1829. Hem. Suec. p. 69.

Xylocoris cursitans : Sahlberg, 1848. Monog. Geor. Fenn. p. 80.

Piezostethus rufipennis Dufour, 1833. Ann. Soc. Ent. Fr. 2: 106.

Lyctocoris corticalis Hahn, 1835. Wanz. Ins. 3 : 21.

Xylocoris bicolor Schultz, 1846. Uebers Schles. Ges. (1846) : 116.

Xylocoris latior Mulsant, 1852. Ann. Soc. Linn. Lyon p. 160.

Piezostethus cursitans : Reuter, 1871. Öf. Vet. Aka. Förh. p. 411.

Cosmopolitan.

13. *Xylocoris deserti* Villiers

Xylocoris deserti Villiers, 1918. Bull. Inst. Franz. Afr. Noir (A) 18: 838.

Mauritania.

14. *Xylocoris discalis* (Van Duzee)

Scoloposcelis discalis Van Duzee, 1914. Trans. San Diego Soc. Nat. Hist. 2: 15.

Hawaii, U. S. A.

15. *Xylocoris dybasi* Herring

Xylocoris dybasi Herring, 1967. Ins. Micronesia 7 (8) : 413.

Mariana Is.

16. *Xylocoris flaccidus* (Van Duzee)

Piezostethus flaccidus Van Duzee, 1914. Trans. San Diego Soc. Nat. Hist. 2: 14.

U. S. A.

17. *Xylocoris flavipes* (Reuter)

Piezostethus flavipes Reuter, 1875. Sv. Aka. Handl. 3 (1) : 65.

Triphleps frumenti Zacher, 1922. Arb. Biol. Reichanst. Land und Forst Wirt. 12 : 236.

Triphleps sinui Narayanan & Chatterji, 1952. Proc. 2001. Soc. Bengal 5 (2): 163,

Triphleps ramae Narayanan & Chatterji, 1953. Proc. 2001. Soc. Bengal 6 (2) : 121.

Europe, Algeria, South Africa, India.

18. *Xylocoris formicetorum* (Boheman)

Anthocoris formicetorum Boheman, 1844. Öf. Aka. Förh. 1: 158.

Xylocoris formiceticola Sahlberg, 1848. Mon. Geoc. Fenn. p. 82.

Xylocoris coenomyces Baerensprung, 1858. Berl. Ent. Zeit. 2 : 195.

Piezostethus formicetrum : Fieber. 1861. Europ. Hemip. 139 : 2.

Europe, Algeria.

19. *Xylocoris galactinus* (Fieber)

Anthocoris galactinus Fieber, 1836. Weit. Beit, 1 (107) : 7.

Anthocoris pulchellus Fieber, 1840. Zett. Ins. Lap. 265: 3.

Piezostethus albipennis Herrich-Schaeffer, 1853. Wanz. Ins. 9 : 173.

Piezostethus galactinus Fieber, 1861. Europ. Hemip. 139 : 1.

Cosmopolitan.

20. *Xylocoris heluanensis* E. Wagner

Xylocoris heluanensis E. Wagner, 1961. Bull. Soc. Ent. Egypte 45: 301.

Egypt.

21. *Xylocoris hirsutus* Carayon

Xylocoris hirsutus Carayon, 1961. S. Afr. Anim. life 3 : 550.

South Africa, Kenya.

22. *Xylocoris jeanneli* (Poppius)

Piezostethus jeanneli Poppius, 1920. Voyage Alluaud Hemip. 4: 322.

East Africa.

23. *Xylocoris lativentris* (Sahlberg)

Piezostethus lativentris Sahlberg, 1861. Not. Pro. Faun. Flor. Fenn. 9: 287.

North Europe, Bulgaria.

24. *Xylocoris machadoi* (Carvalho)

Piezostethus machadoi Carvalho, 1952. Publ. Cult. Comp. Diam. Angola 15: 19.

Angola.

25. *Xylocoris maculipennis* Baerensprung

Xylocoris maculipennis Baerensprung, 1858. Berl. Ent. Zeit. 2 : 197.

Piezostethus maculipennis: Reuter, 1883. Ent. Tidsk. 4: 137.

North Africa, Syria, South Europe.

26. *Xylocoris nigrutilus* (Reuter)

Piezostethus nigrutilus Reuter, 1877. Öf. Finsk. Vet. Soc. 21 (40) : **19.**

Piezostethus nigricans Reuter, **1877.** Öf. Finsk. Vet. Soc. 21 (40) : 19.

Xylocoris formicetorum : Baerensprung, 1858. Berl. Ent. Zeit. 2 : 195.

Germany.

27. ***Xylocoris obliquus*** Costa

Xylocoris obliquus Costa, 1852. Cent. 7 (241) : 28.

Piezostethus signatus Jakovlev, 1882. Troudy Ent. Ross. 12: 148.

Piezotethus obliquus : Reuter, 1884. Monog. Anthoc. p. 591.

var. ***orientalis*** Reuter, 1884. Monog. Anthoc. p. 591.

Europe, Syria.

Morocco, Syria, Persia. (var. *orientalis*)

28. ***Xylocoris pallidipes*** (Poppius)

Piezostethus pallidipes Poppius, 1920. Voyage Alluaud Hemip. 4 : 3.

East Africa.

29. ***Xylocoris parvulus*** (Reuter)

Piezostethus parvulus Reuter, 1871. Notis. Skpts. pro. Faun. et Flora Fenn. 11: 321.

Finland, Holland, Hungary.

30. ***Xylocoris piceus*** (Reuter)

Piezostethus piceus Reuter, 1884. Monog. Anthoc. p. 590.

Siberia.

31. ***Xylocoris queenslandicus*** Gross

Xylocoris queenslandicus Gross, 1956. Rec. S. Aust. Mus. 11: 151.

Australia.

32. ***Xylocoris spangnicola*** (Reuter)

Piezostethus spangnicola Reuter, 1883. Ent. Tidsk. 4 : 135.

Finland.

33. ***Xylocoris sordidus*** (Reuter)

Piezostethus sordidus Reuter, 1871. Öf. Vet. Aka. Förh. 28: 560.

Piezostethus binotatus Reuter, 1871. Öf. Vet. Aka. Förh. 28: 560.

U. S. A., Central and South America.

34. ***Xylocoris terricola*** (Reuter)

Piezostethus terricola Reuter, 1902. Ent. mon. Mag. 38 : 102.

Spain.

35. ***Xylocoris thomsoni*** (Reuter)

Piezostethus thomsoni Reuter, 1883. Ent. Tidsk. 4 : 137.

Sweden, U. S. S. R.

36. ***Xylocoris transversus*** E. Wagner

Xylocoris transversus E. Wagner, 1955. Comm. Biol. Helsing. 14 (2) : 22.

Canary Is.

37. *Xylocoris umbrinus* Van Duzee

Xylocoris umbrinus Van Duzee, 1921. Proc. Calif. Acad. 4 (11): 137.

Canada, U. S. A.

38. *Xylocoris vicarius* (Reuter)

Xylocoris vicarius Reuter, 1884. Monog. Anthoc. p. 599.

Canada, U. S. A.

The Palaearctic and Oriental species are characterized by the following key.

- 1a Base of pronotum strongly sinuate, sides of pronotum anteriorly marginate. 2
- b Base of pronotum very widely and slightly sinuate or subtruncate, not marginate at sides. 3
- 2a First segment of rostrum extending to base of eye, antennal segment III as long as segment II. *X. maculipennis*
- b First segment of rostrum only reaching to the middle of eye, antennal segment III shorter than segment II. *X. lativentris*
- 3a Apex of abdomen free from long protruded hair, legs somewhat long, fairly slender, entirely pale yellow. 4
- b Apex of abdomen furnished with long protruding hairs, legs somewhat short, fairly robust, femora mostly pitchy-black or brown. 5
- 4a Osteolar channel reaching to anterior margin of pleura. *X. flavipes*
- b Osteolar channel not reaching to anterior margin but to lateral margin of pleura. *X. transversus*
- 5a Osteolar channel very long, forming a carinate and acuminate apex which is reaching to basal margin of pleura. 6
- b Osteolar channel not strongly and acuminately prolonged near apex, ending a little below of basal margin of pleura. 10
- 6a Posterior tibiae with long fine white hairs..... *X. parvulus*
- b Posterior tibiae with short hairs. 7
- 7a Large-sized species at least 2 ½ mm in length. 8
- b Small-sized species at most 2 mm. 9
- 8a Hemelytra whitish, partly brownish. *X. galactinus*
- b Hemelytra except interior angle brown or pitch black..... *X. afer*
- 9a Antenna entirely pale yellow testaceous, legs entirely yellow testaceous. *X. formicetorum*
- b Antenna pitch-black, femora except apex blackish..... *X. spangnicola*
- 10a Hemelytra entirely fuscous. 11
- b Hemelytra strongly shining. 12
- 11a Hind tibiae with marked long hairs..... *X. contiguus*
- b Hind tibiae without long hairs. *X. piceus*
- 12a Hemelytra pitch black, corium except for apex and frequently embolium blue whitish. 13
- b Hemelytra unicolous, strongly shortened, its apex truncate. 16
- 13a Body size less than 1.8 mm, apex of mesocorium black..... 14
- b Body size more than 2.4 mm, apex of mesocorium shining..... 15

- 14a Small-sized species **0.5-1.4** mm, with distinct tibial hairs. *X. terricola*
 b Large-sized species 1.4-1.8 mm, without distinct tibial hairs. *X. heluanensis*
- 15a Pronotum strongly slender anteriorly, lateral margin of pronotum linear.
 * * *X. obliquus*
 b Pronotum not markedly slender anteriorly, lateral margin of pronotum widely sinuate. * * *X. canariensis*
- 16a Large-sized species, oblong oval, at least 2 ½ mm in length, abdomen not strongly dilated. *X. cursitans*
 b Small-sized species, short-oval, at most 1½ mm in length, abdomen strongly dilated. *X. thomsoni*

Note: From the above key the following six species are excluded.

- X. antaensis* E. Wagner, 1957. Comm. Biol. Helsing. 16 (2) : 3.
X. balteatus Walker, 1872. Cat. Heter. Brit. Mus. 5 : 159.
X. ciliatus (Jakovlev, 1877). Bull. Soc. Mosc. 52 (2) : 300.
X. deserti Villiers, 1918. Bull. Inst. Franz. Afr. Noir (A) 18 : 838.
X. machadoi (Carvalho, 1952). Publ. Cult. Comp. Diam. Angola 15 : 19.
X. nigritulus (Reuter, 1877). Öf. Finsk. Vet. Soc. 21 (40) : 19.

Description of the adult of *X. galactinus* was first treated by Fieber (1836). Since then, redescriptions have been made by Reuter (1884), Blatchley (1926), Carayon (1953) etc., and works on nymphal stages were done by Hall (1951).

Description of *X. galactinus* :

Egg. The egg of *X. galactinus* is elongate and slightly curved. It is rounded at one end and provided with an operculum at the opposite end. The operculum is roughly circular in outline, and divided into two distinct zones as in the case of many Anthocorid bugs. The marginal zones consists of about 22 rectangular cells which are arranged radially around the inner zone, but varying somewhat in detail. In width this zone is approximately half the diameter of operculum. The second or inner zone is slightly convex and composed of flattened area with a number of fine protuberances. The operculum is chalky white in coloration and the main portion is exposed on the filter paper in which the egg is deposited in laboratory condition. The sculpture of operculum differs distinctly from that of *L. beneficus*.

Average 0.66 mm in length (range 0.64-0.69), and 0.32 mm in width (range 0.29-0.34).

First instar. Immediately after the eclosion the nymph is pale pinky colour but soon becomes darker. Head : pale yellow with dusky suffusion, eyes deeper red colour, sutures clearly visible ; antennal segments I-III with grey tinge, IV translucent. Thorax : generally more dusky than head; pronotum rectangular in shape; mesonotum widening posteriorly, anterior margin of pro- and mesonota with pinky tinge; metanotum paler than aforementioned. Legs: almost translucent with femora slightly darker than tibiae and tarsi. Abdomen: pale yellowish, lateral parts of anterior half pink tinged, mesial part of segments III-IV with distinct pink speckles, with 4 long setae on abdominal tip.

Second instar. Head : reddish orange, eyes red, suture pale and well marked ; antennae translucent, suffused with dusky grey except for segment IV; rostrum yellowish, trans-

lucent, slightly dusky grey on distal part. Thorax : all segments reddish, suffused with dirty grey. Legs : coloured as in the first instar, while slightly more brownish, hind tibiae with developed setae. Abdomen: reddish brown, pinkish tinge on lateral parts more distinct, speckles on segments III-VI red brown, terminal setae present.

Third instar. Head : dark brown with pinkish sutures, ventral surface reddish ; antennal segments I-III dusky grey with pale distal zones, distal part of segment IV yellowish, rostrum; dark brown, especially on basal part. Thorax: yellowish brown suffused with dusky grey hue, pronotum dark brown especially on shoulder parts, anterior part of mesonotum lighter in colour ; wing-pads just visible, showing dark margins, extending to middle of abdominal segment I. Legs : slightly brown, fore femora distinctly swollen, antero-lateral parts of femora greyish. Abdomen : segments I-II yellowish brown, remainder deep brown, especially on posterior part of each segment, speckles disappeared, setae existing.

Fourth instar. Head : colour as described for the third instar; antennal segments I-III dark grey to black with distal one-eighth of II cream coloured, IV yellowish, slightly brownish distally. Thorax : colour as in the third instar, but more fuscous; wing-pads further developed. Legs : pale testaceous with grey-black suffusion. Abdomen : colour as in the third instar.

Fifth instar. Head: deep brown and pinkish red on dorsal and ventral surfaces respectively, eyes deep red ; antennal segments I-III dark grey to black, with distal extremities narrowly pale creamy white, IV yellowish, translucent on distal part ; rostrum brown, darker at tip and ventral surface. Thorax : pronotum deep brown, becoming darker laterally, mesonotum brown ; metanotum brown tinged on anterior portion, yellowish posteriorly with pinkish brown along posterior margin. Wing-pads exceeding posterior margin of first abdominal segment and fore wing-pads entirely covering hind ones. Legs : yellowish brown with dark grey suffusion. Abdomen : segments I and II yellowish in colour, with pinky suffusion, remainder of abdomen dark brown or rather approaching to black, terminal setae present.

Adult. Head : brownish black, thinly clothed with short fine yellowish hairs, compound eyes deep brown, ocelli somewhat brilliant red; antennal segments I dark brown, II yellowish brown, the tip fuscous, III and IV yellowish-brown, slender, subequally in length with prominent hairs which are longer than the diameter of the segments ; rostrum yellowishbrown, long and extending to the fore coxae. Thorax : pronotum black brown, subtrapezoidal, with base about one third as wide as apex; postmedian transverse impression distinct mesially, subobsolete on sides; collar almost smooth and minutely rugose; side margins provided with two erect slender setae; scutellum with a triangle-shaped median depression wide and deep, both pronotum and scutellum covered with fine yellow hairs. Hemelytra very finely and sparsely pubescent, uniformly straw-yellow, with at most the tips of clavus and cuneus and other margins of membrane fuscous. Legs : femora very dark brown, tibiae yellowish brown, tarsi paler, 3-segmented, bristles on tibiae very prominent. Abdomen as pronotum in coloration, with terminal setae.

The dimension of each stage is given in Tables 5-7.

Each instar of *X. galactinus* nymphs is easily distinguishable by the following key.

Head longer than pronotum,	2
Head shorter than pronotum,	3

Table 5. *Xylocoris galactinus* ; dimensions of nymphs and adult in millimeters. (I)

Instar No.	Head length	Head width	Pronotum length	Pronotum width	Abdomen length	Total length
1	0.17	0.17	0.15	0.28	0.48	0.88
2	0.21	0.20	0.18	0.48	0.72	1.22
3	0.23	0.21	0.32	0.60	1.12	1.86
4	0.25	0.23	0.44	0.68	1.21	2.60
5	0.28	0.26	0.48	0.80	1.30	2.81
Adult	0.31	0.42	0.50	0.89	1.46	2.92

(Avg of 5 individuals).

Table 6. *Xylocoris galactinus* ; dimensions of nymphs and adult in millimeters. (II)

Instar No.	Antennal segments					Rostral segments				
	I	II	III	IV	total	I	II	III	IV	total
1	0.03	0.10	0.10	0.14	0.37	0.01	0.06	0.12	0.08	0.27
2	0.07	0.15	0.16	0.18	0.56	0.03	0.11	0.19	0.10	0.43
3	0.10	0.24	0.24	0.21	0.79	0.05	0.13	0.29	0.13	0.60
4	0.13	0.32	0.31	0.24	1.00	0.06	0.15	0.31	0.14	0.65
5	0.15	0.35	0.32	0.26	1.08	0.08	0.17	0.32	0.14	0.71
Adult	0.19	0.36	0.32	0.29	1.17	0.12	0.19	0.34	0.15	0.81

(Avg of 5 individuals).

Table 7. *Xylocoris galactinus* ; dimensions of nymphs and adult in millimeter. (III)

Instar No.	Fore leg			Middle leg			Hind leg		
	Femur	Tibia	Tarsus	Femur	Tibia	Tarsus	Femur	Tibia	Tarsus
1	0.13	0.12	0.02	0.13	0.12	0.02	0.13	0.14	0.04
2	0.25	0.21	0.09	0.21	0.20	0.06	0.25	0.23	0.08
3	0.38	0.25	0.12	0.36	0.33	0.13	0.39	0.38	0.14
4	0.43	0.32	0.13	0.44	0.43	0.17	0.54	0.59	0.18
5	0.54	0.44	0.22	0.48	0.46	0.22	0.66	0.71	0.24
Adult	0.58	0.50	0.27	0.62	0.58	0.27	0.73	0.88	0.35

(Avg of 5 individuals).

2a All antennal segments almost with similar diameter.	1st instar
b Segments I-II thicker than segments III-IV.	2nd instar
3a Hind wing-pads entirely covered with fore ones.	5th instar
b Hind wing-pads not entirely covered with fore ones.	4
4a Wing-pads just visible.	3rd instar
b Wing-pads developed and extending to the posterior margin of the first abdominal segment.	4th instar

The adults of *L. beneficus* and *X. galactinus* are easily discriminated by the following characters : 1) Larger body size, viz. about 3.8 mm in the former species, and less than 3.0 mm in the latter. 2) Distinguished cilia on the posterior margin of the abdomen in the latter species. 3) The front femora and tibiae more or less swollen in the latter species but not in *L. beneficus*. The discrimination of the nymphal stage is somewhat difficult in the newly hatched nymphs between the two species. It is only distinguishable by the body size which are about 1.1 and 0.9 mm in *L. beneficus* and *X. galactinus* respectively. To distinguish the newly hatched nymphs more than one day old and the following instar nymphs, following characters are available.

- 1) Body size, as shown in Tables 2 and 5, larger in *L. beneficus*.
- 2) Coloration, light yellow to brown yellow in *L. beneficus*, more or less brilliant red-brown in *X. galactinus*.
- 3) In *L. beneficus*, abdominal segment apparently dilated on the sides of IV-VI segments, thus making sub-spatulate in form. The dilation in *X. galactinus* is not prominent and rather semi-ellipsoidally shaped.
- 4) In all nymphal stages of *X. galactinus*, the posterior margin of abdomen is provided with terminal ciliae while the ciliae are only present on the 1st and 2nd instars in *L. beneficus*.

Habitat

According to the previous records, *Lyctocoris beneficus* (Hiura) was known to be abundant in a giant pile at Saga Paper Board Co. in Kyushu (Oho 1955a, b, Hiura 1957). Besides, Hiura (1957, 1966) examined the specimens of this species which were collected from the following habitats: farm store, thatch, piles of fire wood, grass (ex. *Miscanthus sinensis*), and harvested plants such as wheat, sweet potato, vine, broad-bean stem. Also some specimens were collected at the lamp during July to September.

Xylocoris galactinus (Fieber) is commonly known under the name of "hot-bed bug", and in Europe it was found in manure-heaps, hot-bed, stable straw, grain bins, marsh haystacks and was also recorded on oak and beech and under the bark of other trees (Morley 1905, Butler 1923, Hall 1951, Southwood and Leston 1959, Sinha 1961 etc.). As the habitat of the bug, Reuter (1871a) wrote as follows: " Amongst decaying grass, on dung (Schiödte), with *Myrmica calspitum* (Bellevoye), on oak and beech, and under

the bark of tree." Further, Fieber (1860) recorded "on plants in meadows." Of the habitat in Japan, Hiura (1966) recorded that "the bug lives in farm store of wheat straw, sometimes in the open air stacks of rice straw. Four specimens were obtained under the pile of dead sea grass on the beach. Often appearing at the lamp." In Micronesia, this species was collected from decaying coconuts (Herring 1967).

During the author's work, some individuals of both species of bugs were collected at the farm store in winter, while in the warmer season a great number of them were found in the giant straw piles at Saga Paper Board Co. Sometimes, the bugs were extremely abundant in paddy or wheat straw just transported to the Company yard from the farmer's stock yard or field piled thatch. *L. beneficus* was often much more predominant than *X. galactinus* on wheat straw which is generally more parched than paddy straw. The latter species was in almost all cases more prevalent in paddy straw especially of somewhat moist condition. The trials to collect the bugs in the paddy field and the grass meadow adjacent to the giant straw pile at Saga with a suction apparatus were failed. In August, many specimens of both species including nymphs were collected at the just harvested paddy straw at Kumamoto, in Kyushu, and this fact may indicate the occurrence of these bugs in the paddy field and the meadow nearby. During July to September many specimens of both species were also collected by the lamp. The nocturnal activities of both species will be given in detail in the later part of this report.

In conclusion, the author's observation indicates that *L. beneficus* is abundant in relatively dried places, while *X. galactinus* prefers rather marsh environment.

Laboratory observation on the life history

I. Egg.

Judging from their habitat in the field condition, paddy straw seems to be an important ovipositional substratum both for *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber). Under the laboratory condition, eggs are actually laid in the moist paddy straw which is provided for oviposition. On the other hand, eggs are also laid to the other materials such as the cherry twig, moist filter paper, molded cracked corn, moist vermiculite, etc. The oviposition preference by the adult females to these materials will be discussed in another column of this chapter. On the filter paper, eggs are inserted horizontally, causing the paper to bulge into a small but clearly visible blister which facilitates the location and counting of eggs. The operculum tends to face obliquely upwards because of the slight curvature of the egg. When the cherry twig is offered for the ovipositional substratum, the egg is oviposited just below the epidermis, exposing only

a white sculptured operculum. When the twig is provided in an upright position, the adult female takes the position with the head upward in most cases, and lays eggs. Consequently, the egg operculum directs upwards. But rarely the egg was seen to be deposited the operculum downwards.

The range of incubation period is certainly affected by the temperature (Figs. 1 & 2). Hiura (1957) and Oho (1955c) indicated the incubation period

Fig. 1. Frequency distribution histograms of the incubation period of *Lyctocoris beneficus*.

Abscissa: time in days, **Ordinate:** Number of observations.

Fig. 2. Frequency distribution histograms of the incubation period of *Xyllocoris galactinus*.

Abscissa : time in days, **Ordinate :** Number of observations.

of *L. beneficus* for 7 and 6 days respectively. In the present work, majority of eggs of *L. beneficus* hatched after the incubation period of approximately 4 days under 30°C condition, and that of *X. galactinus* was 3 days. No significant effect was observed on the incubation period under the range of 53-100 % of relative humidity. However, the water content of ovipositional substrata gives a great influence on the oviposition preference as well as the development of eggs. Namely, the adult does not lay any eggs on the dry substance. Moreover, if the egg supporting media dry up after the oviposition, the eggs fail to develop. Even if moderate humidity is supplied to the parched egg supporting media, few eggs resume their development and the rate of hatching is very low. The egg mortality in the dry substrata may be caused by the combination of two actions, one is the drought of egg itself, and another is the compression of egg by the desiccated substratum.

During the embryonic development, marked colour changes may be seen in the eggs of both species. The process of colour change is very similar in both species. But the change of colour in *X. galactinus* is slightly faster than that of *L. beneficus* owing to the shorter incubation period, and in the former species reddish tinge was always stronger than the latter species. The following is the description of colour change of the egg of *L. beneficus* under 30°C condition. At first, the egg is almost colourless and opalescent, showing slight pale orange tinge. This tinge becomes stronger and more extensive within 24 hours after oviposition. At the beginning of the second day, the posterior two-thirds of the egg becomes milky orange. At the end of the same day, the egg becomes entirely orange coloured except the anterior third which is still much paler than the rest. The egg is deep orange coloured at about one third of its length toward the operculum and also at the extreme tip. Later these concentrations may spread so that about half of the egg is tinged with orange. By the third day, they form two distinct zones, one occupying the posterior half of the egg and another forming a zone just behind the operculum from which it is separated by a narrow pale zone ; a broader pale orange zone separates the two deeper orange zones from each other. Toward that day, the entire egg becomes orange in colour, and orange tinge becomes deeper later, then reddish zones appear at a sub-operculum and mid-ventral part. At this stage the pigmented eyes are visible through the chorion. In this condition, the egg is ready to hatch.

Oviposited eggs are often pierced and sucked by their adults, thus leaving only the shrivelled and transparent chorions. This phenomenon occurs even in the presence of abundant food, but is more frequently seen under the dry condition. Contrarily, Hill (1957, 1961) indicated that *Anihocoris nemorum* (Linné) and *A. sarothamni* Douglas et Scott sucked their eggs in spite of an abundance of moisture. In any case a long time exposure of ovipos-

ited filter paper with the adult is harmful to the eggs. The author's experiment shows that the rate of hatching is very low from a filter paper which remained in the container with the adults for more than 48 hours. Though the ratio of pierced egg deposited on the other substrata was not observed in the present work, Hill (1965) indicated that the eggs of *Anthocoris confusus* (Reuter) laid on the filter paper were easier to be sucked by the adults than the eggs on plant material. The egg-piercing habit is more pronounced in *X. galactinus* which prefers moist circumstance than *L. beneficus*. Therefore, it is presumed that the eggs may serve as somewhat suitable water supply for the adult bug under a certain parched condition. Under the favorable condition, the rate of hatching is apparently very high, being more than 95 %. While in some cases, a low percentage of hatchability was observed. As to the cause of low rate of hatching, the following three cases are considered, 1) dryness of oviposited material, 2) egg-piercing habit by adult and 3) age of female adult. The third case is discussed in the later part of this chapter.

II. Nymph.

The eclosion occurs mainly at 8-10 am in both species, and only a few instances were observed in the night. The average time taken for eclosion by *L. beneficus* and *X. galactinus* under 30°C condition were 9 and 7 minutes respectively. The process of eclosion is very similar in both species. Herewith is described the eclosion process of *L. beneficus*. Eclosion activity was at first begun with the vigorous movement of embryo. Through this movement the chorion which was imbedded into the filter paper became removable, resulting the position of egg vertical or perpendicular to the surface of paper. At this stage, the operculum still remained the same position, but after 4-5 min the nymph before hatching began slow bending or dorso-ventral levering movement and it dropped off suddenly, after then the embryo came to lie at an obtuse angle to the paper and began violent movement to liberate appendages from the chorion. After 2 minutes the antennae were extended, and it took only 30 seconds to stretch all the legs. The continuous muscular movement freed the abdomen from the chorion. The whole process of eclosion was accomplished by the nymphal bending forward and grasping the paper by the front legs and, only a few seconds were needed to complete the last stage of eclosion.

As many other species of Anthocorid bugs, five nymphal instars are recognized in both species. Under the condition of 30°C the total period from eclosion to adult ranged approximately from 17 to 22 days and 11 to 15 days in *L. beneficus* and *X. galactinus* respectively. In each test, complete records were made on 20-30 individuals to observe the duration of nymphal instars under various temperature conditions (Tables 8 & 9).

It is evident that within the range of temperatures tested the nymphal development becomes more speedy in warmer conditions. The acceleration

Table 8. Duration of nymphal instars of *Lyctocoris benejcus* at different temperature conditions (in days).

Temp	1	2	Instar 3	4	5	Total
25°C	2-2.3-4	2-3.2-5	4-4.8-8	4-5.6-8	5-6.9-8	20-22.8-25
30°C	2-2.2-4	2-2.6-4	3-4.7-6	4-5.1-7	4-5.1-7	17-18.7-22
33°C	2-2.2-4	2-2.5-4	3-4.1-6	3-4.4-6	4-4.7-7	16-17.9-22
Room temp (28-33°C)	2-2.4-4	2-2.7-4	3-4.3-6	4-4.8-7	4-5.2-8	18-19.4-24

Table 9. Duration of nymphal instars of *Xylocoris galactinus* at different temperature condition (in days).

Temp	1	2	Instar 3	4	5	Total
25°C	2-2.7-4	2-3.0-5	2-3.3-5	2-3.7-5	5-6.7-9	17-19.4-22
30°C	1-1.8-3	1-1.9-3	2-2.6-4	2-2.2-3	3-4.5-7	11-13.6-15
33°C	1-1.7-3	1-1.7-3	2-2.4-3	2-2.1-3	3-4.3-6	11-12.4-14
Room temp (28-33°C)	1-2.0-3	1-2.2-3	2-2.7-4	2-2.4-4	3-4.8-7	12-14.1-16

of nymphal development under certain higher temperature conditions is generally known in many insects including another Anthocorid bug (Butler 1966). Among 5 stadia, the fifth stadium was the longest and the first and second stadia were the shortest in both species. Relative humidity ranging between 53-100 % was not influential on the duration of nymphal period. Further, there was no significant difference between the rate of nymphal development of male and female individuals.

III. Adult.

1. Emergence. When a nymph accomplishes nymphal growth, emergence of both species takes place in the morning as indicated in Figs. 3 and 4.

Fig. 3. Frequency distribution histogram of the time of emergence observed in *Xylocoris galactinus*. *Abscissa* : time in o'clock, *Ordinate* : Number of observations.

Fig. 4. Frequency distribution histogram of the time of emergence observed in *Lyctocoris benejcus*. *Abscissa* : time in o'clock, *Ordinate* : Number of observations.

The body of newly emerged adult is milky pinkish in colour and the wings are whitish. In *L. beneficus*, it took 15 minutes before the clavus starts to get its dark colour. Darkening of the eyes also took about the same minutes. Gradually, the pale pink colour of the adult changed to the full coloration 4 to 5 hours after emergence. The process of colour change of newly emerged adult of *X. galactinus* was similar to that of *L. beneficus*, but it took only 100 to 120 minutes to accomplish the whole process pigmentation.

2. Copulation. If the male and female are placed together in the rearing tube, copulation or attempt to copulate may occur within a few seconds at any time of the day or night. Mating was even observed in the insect aspirator among the just collected individuals from the field. The pre-copulatory period of male adult in *X. galactinus* was very short and mating behaviour occurred after several hours since the darkening of colour was finished, and that of *L. beneficus* was somewhat similar to *X. galactinus*. This fact nearly indicates that in the newly emerged male adult the reproductive organs already attained to maturity. Adult female of any age including mature and newly emerged pink coloured individuals may become the object of copulation for the male. The fertilized females very frequently refuse the males and struggle against the males vigorously. Virgin females, on the other hand, readily accept the males. The male mounts on the back of the female, keeping his body dorso-laterally to the body of female and to her right side. Male holds her with the help of numerous spines found on the inner margin of the fore tibiae. In this case, his three left legs are placed (1) on her pronotum, (2) at the basal portion of the hemielytron and (3) at the cuneus of the left hemielytron. Right fore-leg is lifted up while the right mid-leg is touching the substratum, thus supporting the weight of the male. During the course of copulation, the antennae of male are making back and forth movements. Those of the female are always stretched forward and very rarely touching the male antennae. The female remains motionless during the entire course of copulation, stretching her rostrum forward. In a very few instances, copulating females were observed depressing her head and raising the tip of abdomen and walking with the mate on her back. The left ventro-lateral surface of the male's body was in contact with the right ventro-lateral surface of the female's body, thus allowing the insertion of the left paramere of the male to the female genital opening without any difficulty. The average time consumed for the copulation of *L. beneficus* and *X. galactinus* was 2.5 and 2.0 minutes respectively. One mating is seemed to be almost enough to supply sufficient spermatozoa to the female to lay fertilized eggs throughout her life.

3. Oviposition. The pre-oviposition period is very much variable depending upon the temperature conditions and the records of experimental data are indicated in Tables 12-17. It was 15.2, 7.3 and 6.9 days under 25°, 30° and 33°C respectively in *L. beneficus* and 12.8, 4.1 and 3.8 days in *X. galactinus* on

an average. Within the range of such temperature conditions, relative humidity between 53–100% did not give any influences upon the pre-oviposition period. If the temperature decreases during the earlier life of adult, the period may be prolonged. If a 2 day old female adult was placed to 25°C condition from 30°C, her pre-oviposition period was prolonged to 17.6 and 15.3 days in *L. beneficus* and *X. galactinus* respectively. This data were apparently longer than the period shown by the adult reared under the constant temperature condition. While the inverse transfer of adults, namely low to high, did not shorten the pre-oviposition period.

The oviposition is conducted at any time of the day or night, although somewhat more eggs are produced during the daytime. But no significant diurnal ovipositional rhythm is observed.

When the female is ready to oviposit, she moves slowly on the substratum examining it every few seconds with the tip of her rostrum while she is vibrating her antennae rapidly. Quite suddenly, the female stands still, raises up her body high and bends the tip of her abdomen so that it touches the surface of paper. Then, her ovipositor is stretched vertically to be inserted in the substratum. When the insertion of ovipositor is finished, the female takes the normal position with the antennae stretched forward, and with the abdomen showing rhythmical contracting movements. By these movements, the ovipositor is deeply inserted into the substratum. The final passage of the egg into the substratum is carried out without any perceptible extra effort and the ovipositor is at once removed. The average times taken for oviposition process in *L. beneficus* and *X. galactinus* were 80 and 50 seconds respectively. The eggs are not laid aggressively but placed singly.

It is almost evident that especially in the field the barn straw is the most important oviposition substratum. Under the laboratory condition, both species actively deposited eggs on the moist straw. In addition to straw, the cherry twig, moist filter paper, moist vermiculite and slightly molded cracked corn are excellent substrata for the oviposition of both species. A few eggs are also deposited on the moist wood chops (*Acacia* sp.). As the optimal condition for the oviposition substratum, suitable moisture and softness of materials are demanded. Although Hill (1957, 1961) and Tawfik and Nagui (1965) observed the oviposition of other Anthocorid bugs on the wall of rearing glass tube, no egg was observed on the wall of a glass tube in the present experiment.

To compare the suitability of these materials as the ovipositional substratum, 20 adult females were kept in the 100 ml beaker with one of these materials and, thus six beakers of this kind were prepared and the number of newly hatched nymphs were counted after several days in each beaker. As the counting of oviposited eggs on such materials as vermiculite, corn, straw and wood chop was actually difficult, the number of hatched nymphs

was taken as the index of oviposition suitability and preference. Therefore, in this test, it was possible to compare the suitability both for oviposition and egg development at the same time. This work was conducted for 5 days, renewing the substratum every day (Table 10). In another series of test, the combination of more than 2 kinds of oviposition substrata were used in each beaker containing 20 adult females and examined their oviposition preference. In this connection, it must be noted that the vermiculite used contained some water (vermiculite : water=100 : 15 w/w) (Table 11).

Table 10. Number of newly hatched nymphs from the various ovipositional substrata per day, each substratum provided with 20 adult females of *Lyctocoris beneficus* and *Xylocoris galactinus* at the beginning of the experiment.

Substratum	<i>L. beneficus</i>	<i>X. galactinus</i>
Moist straw	17.2	21.0
Moist filter paper	24.4	25.8
Moist vermiculite	18.5	24.3
Cherry twig	32.0	21.7
Molded cracked corn	21.6	27.5
Wood chop (<i>Acacia</i> sp.)	5.1	8.3

Table 11. Number of newly hatched nymphs from the various ovipositional substrata per day, each combination provided with 20 adult females of *Lyctocoris beneficus* and *Xylocoris galactinus* at the beginning of the experiment.

Combination	Substrataum	No. of nymphs hatched	
		<i>L. beneficus</i>	<i>X. galactinus</i>
A	Straw	8.2	6.8
	Filter paper	9.4	9.4
	Cherry twig	18.2	13.2
	Cracked corn	10.6	14.6
B	Straw	8.4	4.4
	Filter paper	11.6	8.2
	Cherry twig	14.4	9.8
	Vermiculite	5.0	10.4
C	Straw	8.2	5.4
	Filter paper	14.6	13.6
	Cherry twig	16.4	6.6
	Wood chop	1.8	1.2

As the data of Tables 10 and 11 show, cherry twig was the most preferred oviposition substratum for *L. beneficus*. Filter paper and molded cracked corn were also an excellent ones for this species, but dry cracked corn was not suitable for the oviposition of both species. In the case of *X. galactinus*, majority of newly hatched nymphs were from filter paper, molded cracked corn and vermiculite. Wood chop was the worst in both species. Actually the oviposition was influenced by the kind of substratum. The influence of substrata on the oviposition behaviour was also observed in some groups of insects by Chalfant and Mitchell (1967). In conclusion, the following order may be established from Tables 10 and 11 on the suitability and preference of oviposition in each species.

- L. beneficus.* cherry twig > cracked corn > filter paper > straw > vermiculite > wood chop.
- X. galactinus.* cracked corn > vermiculite > filter paper > cherry twig > straw > wood chop.

The oviposition preference between the two species is apparently variable not only with the kind of substratum but also with the condition of substratum, especially relating to its water content. In any cases, adults avoid to deposit their eggs on the parched or extremely moist substrata.

The oviposition period, number of eggs per female and longevity were examined by the use of fertile females which were kept separately in a 9 x 2.5 cm (length x diameter) flat-bottomed corked glass tube. And a sheet of filter paper cut in 1 x 4 cm square was provided in the tube for the supply of water as well as oviposition substratum. Adult females readily accepted the moist paper for oviposition, and the larvae of red flour beetle (*Tribolium castaneum* (Herbst)) were supplied as food. The tubes were examined daily. The filter papers were renewed and the eggs deposited on the paper were counted daily. The tests were carried out under 25°, 30° and 33°C temperature conditions. The detailed results are given in Tables 12-17 and the summary is shown in Tables 18 & 19.

It is evident that the individual variation is extremely large both in longevity and egg-production activity. Actually these are easily influenced by some factors as nutritional condition, population density, etc., and the affects of these factors are discussed in the later chapters. So far as the data in these tables are concerned, the longevity of adult female was greatly affected by temperature. In *L. beneficus*, it showed 39.2, 31.0 and 24.5 days on an average under the respective temperature condition. When the adults were kept under 25°C condition, the minimum longevity was 23 days while the maximum one was 53 days. The average longevity of *X. galactinus* was 36.9, 26.4 and 22.6 days for 25°C, 30°C and 33°C conditions respectively, and it was evidently shorter than that of *L. beneficus*. In both species, high temperature gave bad effect on the longevity as well as on the egg-

Table 13. Length of life and egg-laying activity of *Lyctocoris beneficus* under 25°C condition.

Female No.	Pre-oviposition period(day)	Oviposition period (day)	Post-oviposition period(day)	Longevity (day)	No. of eggs Jaid per week (from beginning of oviposition)						Total of eggs(A)	Actual days of oviposition(B)	Eggs per day (A/B)
					1	2	3	4	5	6			
2	14	23	0	37	13	18	18	8	0	-	39	8	7.56
3	14	23	0	37	13	18	18	8	0	-	39	8	7.56
4	13	11	8	32	16	8	0	-	-	-	28	4	7.0
5	16	33	0	49	11	9	7	0	5	-	32	6	6.3
6	14	38	0	52	11	20	17	8	2	-	58	8	7.3
8	15	17	21	53	22	3	0	0	0	0	14	2	7.0
9	16	16	0	32	18	21	4	-	-	-	47	6	7.8
10	14	29	3	46	12	21	16	5	8	-	62	8	7.8
Avg	14.8	20.1	4.3	39.2	15.8	13.8	7.1	2.4	1.5	0.0	40.6	5.7	7.3

Table 13. Length of life and egg-laying activity of *Lyctocoris beneficus* under 30°C condition.

Female No.	Pre-oviposition period(day)	Oviposition period (day)	Post-oviposition period(day)	Longevity (day)	No. of eggs laid per week (from beginning of oviposition)						Total of eggs(A)	Actual days of oviposition (B)	Eggs per day (A/B)
					1	2	3	4	5	6			
1	6	16	9	31	21	12	3	0	-	-	36	5	7.2
2	9	15	4	28	2	25	5	-	-	-	46	10	4.2
3	7	8	5	20	8	2	-	-	-	-	10	2	5.2
4	6	15	18	39	17	7	3	0	0	1	27	5	5.4
5	8	16	0	24	4	15	2	-	-	-	21	5	4.2
6	6	23	8	37	18	19	11	8	0	-	56	8	7.0
7	10	15	3	28	10	8	2	-	-	-	20	5	4.0
8	7	24	0	31	17	12	13	11	0	-	53	6	8.3
9	7	17	9	33	28	18	18	0	0	1	65	7	9.3
10	7	22	10	39	21	8	0	2	0	-	31	6	5.2
Avg	7.3	17.1	6.6	31.0	16.4	12.2	5.7	2.1	0.0	0.1	36.5	5.9	6.0

Table 14. Length of life and egg-laying activity of *Lyctocoris beneficus* under 33°C condition.

Female No.	Pre-oviposition period(day)	Oviposition period (day)	Post-oviposition period(day)	Longevity (day)	No. of eggs laid per week (from beginning of oviposition)						Total of eggs(A)	Actual days of oviposition (B)	Eggs per day (A/B)
					1	2	3	4	5	6			
1	7	13	0	20	12	8	-	-	-	-	20	5	4.0
2	6	15	2	29	13	16	8	0	1	1	37	4	9.3
3	8	12	17	25	9	7	0	-	-	-	16	4	4.0
4	7	29	0	36	24	15	2	3	7	-	51	9	5.7
5	6	1	11	18	11	0	-	-	-	-	11	1	11.0
6	6	9	2	17	31	4	-	-	-	-	35	7	5.0
7	10	12	1	23	2	7	-	-	-	-	9	3	3.0
8	7	17	3	27	13	8	3	-	-	-	24	5	4.8
9	6	18	2	26	21	12	4	-	-	-	37	6	6.2
10	6	18	0	24	17	6	2	-	-	-	25	5	5.0
Avg	6.9	14.4	3.8	24.5	15.3	8.3	1.9	0.3	0.7	-	26.5	4.9	5.8

Table 15. Length of life and egg-laying activity of *Xylocoris galactinus* under 25°C condition.

Female No.	Pre-oviposition period(day)	Oviposition period (day)	Post-oviposition period(day)	Longevity (day)	No. of eggs laid per week (from beginning of oviposition)						Total of eggs(A)	Actual days of oviposition (B)	Eggs per day (A/B)
					1	2	3	4	5	6			
1	14	22	1	37	21	13	6	2	-	-	42	8	5.3
2	15	16	0	31	18	4	5	-	-	-	27	5	5.4
3	14	22	1	39	24	21	18	12	-	-	75	12	6.3
4	10	37	2	49	16	14	15	16	12	7	80	9	8.7
5	10	36	2	48	14	23	16	8	8	1	70	13	5.4
6	11	22	2	35	6	14	3	4	-	-	27	6	4.5
7	13	16	3	32	19	16	14	-	-	-	49	7	7.0
8	13	22	4	39	8	9	3	2	-	-	22	5	4.4
9	14	13	0	27	6	19	-	-	-	-	25	3	8.3
10	14	17	1	32	11	20	16	-	-	-	47	6	7.8
Avg	12.8	22.3	1.6	36.9	18.8	16.9	7.1	2.2	2.0	0.8	46.4	7.4	6.3

Table 16. Length of life and egg-laying activity of *Xylocoris galactinus* under 30°C condition.

Female No.	Pre-oviposition period(day)	Oviposition period (day)	Post-oviposition period(day)	Longevity (day)	No. of eggs laid per week (from beginning of oviposition)						Total of eggs(A)	Actual days of oviposition (B)	Eggs per day (A/B)
					1	2	3	4	5	6			
1	4	9	3	16	18	13	-	-	-	-	31	4	7.8
2	5	23	10	38	17	10	6	7	0	-	40	5	8.0
3	3	29	5	37	25	19	11	5	2	-	62	10	6.0
4	4	23	1	28	30	19	8	6	-	-	63	8	7.9
5	5	8	0	13	21	3	-	-	-	-	24	3	8.0
6	3	30	8	41	11	21	13	8	8	0	61	6	10.2
7	3	15	4	22	30	12	7	-	-	-	49	5	9.8
8	5	15	0	20	18	6	2	-	-	-	26	4	6.5
9	4	22	5	31	26	17	8	0	-	-	51	6	8.5
10	5	10	3	18	24	16	-	-	-	-	40	6	6.7
Avg	4.1	18.4	3.8	26.4	22.0	13.6	5.0	2.6	1.0	0.0	44.7	5.7	8.0

Table 17. Length of life and egg-laying activity of *Xylocoris galactinus* under 33°C condition.

Female No.	Pre-oviposition period(day)	Oviposition period (day)	Post-oviposition period (day)	Longevity (day)	No. of eggs laid per week (from beginning of oviposition)						Total of eggs(A)	Actual days of oviposition(B)	Eggs per day (A/B)
					1	2	3	4	5	6			
1	4	16	8	28	27	14	2	0	-	-	43	6	7.1
2	3	10	2	15	18	8	-	-	-	-	26	3	8.7
3	4	16	4	24	16	14	7	-	-	-	37	4	9.3
4	4	12	2	18	28	17	-	-	-	-	45	5	9.0
5	5	16	6	27	22	10	4	0	I	-	36	5	7.2
6	3	15	2	20	23	16	9	-	-	-	48	6	8.0
7	4	23	5	32	17	14	6	2	-	-	39	6	6.5
8	4	12	3	19	19	15	0	-	-	-	34	5	6.8
9	4	16	6	26	17	15	6	0	-	-	38	5	7.6
10	4	8	5	17	24	6	-	-	-	-	30	3	10.0
Avg	3.9	14.4	4.3	22.6	21.1	12.9	3.4	0.2	-	-	37.6	4.8	8.0

Table 18. Oviposition of adult *Lytocoris beneficus* under various temperature conditions.

Temp.	Oviposition period (day)	No. of eggs laid per week*						Eggs per female (A)	Days of oviposition (B)	No. of eggs per day (A/B)
		1	2	3	4	5	6			
25°C	20.1	15.8	13.8	7.1	2.4	1.5	0.0	40.6	5.7	7.3
30°C	17.1	16.4	12.2	5.7	2.1	0.0	0.1	36.5	5.9	6.0
33°C	14.4	15.3	8.3	1.9	0.3	0.7	0.0	26.5	4.9	5.8

Table 19. Oviposition of adult *Xylocoris galactinus* under various temperature conditions.

Temp.	Oviposition period (day)	No. of eggs laid per week*						Eggs per female (A)	Days of oviposition (B)	No. of eggs per day (A/B)
		1	2	3	4	5	6			
25°C	22.3	18.8	16.9	7.1	2.2	0.7	0.7	46.4	7.4	6.3
30°C	18.4	22.0	13.6	5.5	2.4	1.0	0.2	44.7	5.7	7.9
33°C	14.4	21.1	12.9	3.4	0.2	0.0	0.0	37.6	4.8	8.0

* from the day of oviposition started.

production capacity. However, shorter pre-oviposition period was observed in the 33°C reared groups, and the production of egg was clearly decreased. The growth was postponed in the individuals which were reared under the condition of low temperature, but the number of eggs oviposited per female was the highest in these individuals. The number of eggs laid by *L. beneficus* and *X. galactinus* were 40.6 and 46.4 respectively, making an apparent contrast with the fewer number 26.5 and 37.6 for the respective species which were reared under the condition of 33°C. The post-oviposition period was longer in the groups which were reared under the higher temperature condition. Though not shown in the Tables, the longest record of post-oviposition period were 36 days in *L. beneficus* and 23 days in *X. galactinus*, and these records were obtained when the adults were kept under the condition of 25°C condition.

The longevity of adult male was given in Table 20, and it seems somewhat shorter than that of the female in both species.

Table 20. Longevity (min-avg-max) of male adult of *Lyctocoris beneficus* and *Xylocoris galactinus* at different temperature conditions (in days).

Temperature	<i>L. beneficus</i>	<i>X. galactinus</i>
25°C	18-33.0-48	13-29.6-45
30°C	15-23.5-42	11-21.2-38
33°C	U-23.3-39	12-20.5-36

High temperature over 33°C was not preferable for both species, and the bad effect of high temperature was more evident in *L. beneficus*. It seems that *X. galactinus* is more adapted to high temperature than *L. beneficus*. This suggestion was also proved by the cold-hardiness test of these bugs as mentioned later.

IV. Effect of maternal age on the vitality of eggs.

It is well-known that egg production of various insects may be greatly affected by environmental conditions and the past history of the mother. A considerable amount of literature on the influence of parental age has been accumulated for some groups of organisms by Strong (1954), and the studies in this field were conducted on some groups of insects (Durrant 1955, Sang 1956, Richard and Kolderie 1957). The author investigated the effect of parental age on the vitality of offspring in both species. For this purpose, 20 couples of adults which were emerged on the same day were kept in 9x6 cm petri-dish (diameter x height), and the eggs deposited on the moist filter paper were isolated from the adult containers every day. The vitality of offspring was measured by the rate of hatching, length of nymphal stage,

percent emergence and the body weight of new adult (Tables 21 & 22). As discussed in the later chapter, the length of nymphal stage, percent emergence, and body weight were apparently influenced by the number of rearing insects per unit space. But the effect of population density on these factors was not considered in this work. The work was carried out under the room temperature condition of 28-33°C.

Table 21. The rate of hatching and the nymphal development of *Lyctocoris beneficus* in relation to parental age.

Parental age (in weeks)	No. of eggs laid	Rate of hatching (%)	Nymphal stage (in days)	Percent emergence (%)	Weight of new adult (mg)	
					♂	♀
1	12	75.0	20.5	77.8	2.2	2.3
2	343	92.7	19.2	85.5	2.4	2.3
3	286	74.5	19.3	86.0	2.4	2.4
4	139	62.6	21.7	60.9	2.3	2.4
5	55	43.6	22.5	58.3	2.3	2.4
6	27	44.4	23.8	50.0	2.3	2.3

Table 22. The rate of hatching and the nymphal development of *Xylocoris galactinus* in relation to parental age.

Parental age (in weeks)	No. of eggs laid	Rate of hatching (%)	Nymphal stage (in days)	Percent emergence (%)	Weight of new adult (mg)	
					♂	♀
1	103	84.5	13.8	62.1	1.3	1.3
2	421	74.6	13.4	83.8	1.3	1.4
3	336	83.6	13.2	81.9	1.3	1.4
4	37	35.1	14.8	46.2	1.2	1.3

As seen in Tables 21 and 22, both *L. beneficus* and *X. galactinus* showed the same tendency as to the effect of maternal age. The egg production was low in the first week especially in *L. beneficus*. It was no doubt due to the fact that most of the adults were still in the pre-oviposition period in this week. The highest percentage of egg-hatching and adult emergence were obtained from the offspring produced by the second and third week old parents. Also the shortening of the duration of nymphal stage was observed in such offsprings. The off spring production was suddenly decreased in the four week old parents, being about half and one third of the younger stage parents in *L. beneficus* and *X. galactinus* respectively. And the vitality of offspring produced by the adults of age later than the fourth week was clearly inferior to that of the younger individuals. No correlation was

seen between the body weight of newly emerged and older adults.

In the laboratory condition under the constant temperature of 30°C, about 54 days are necessary to complete one generation in the case of *L. beneficus*, while 43 days are needed in *X. galactinus*. On the other hand, the duration from "egg to egg" takes only about 31 days in the former and 21 days in the latter species. It is supposed that actually under the optimal artificial condition, both species may have more than 10 generations a year.

Field observations

Field observations on *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) were mainly conducted on the giant straw piles at Saga Paper Board Co., under the various conditions of straws which were transported to the stock yard of the company. Life history in natural condition was also observed at the artificial small straw heap which was piled up in the campus of Kyushu University.

In the investigation at Saga both species were not collected in the months of January, February and March, while from the end of April, small numbers of adults of *X. galactinus* began to appear, and the adults became very abundant in the middle of May. Towards the end of this month, large numbers of both adults and various stages of nymphs were observed, and the abundance of these individuals remained through June. The spring emergence of *L. beneficus* was about 2-3 weeks later than that of *X. galactinus*. At the beginning of July, the population of both species decreased suddenly from the outer part of straw piles, and the low densities of the bugs were maintained till the middle of August. This is probably due to the bugs avoiding the high temperature of summer and emigrate deeply into the interior part of the straw pile, and this phenomenon may be caused owing to the decrease of absolute population by summer heat. This was proved by the fact that in this season many individuals of both species were found in the straw transported from the farmer's stock yard or in newly harvested paddy straws.

During September through October, various stages of both species reappeared in the straw piles. In this season both adults and various stage nymphs of *L. beneficus* were more abundant as compared with the spring emergence, while the population of *X. galactinus* nymphs decreased. To the end of October the nymphs of the latter species were scarcely found. In November, both species almost disappeared in the straw piles, but some were still collected in the straws just transported from the farmer's barn. The materials collected in this season were almost composed of adults of both species. This fact suggests that when the climate becomes cooler, they stop oviposition, and overwinter in the adult stage. The decrease of population in the giant piles is rather reasonably explained by the fact that

the bug emigrates into the internal part of the piles, because the low temperature in November is not enough to kill the bug (cf. Chap. 13).

The seasonal variation of the number of both sexes in field collected adults is given in Table 23. From this table it is proved that the sex ratio ($\frac{\text{♀}}{\text{♂} + \text{♀}}$) in the field condition remains nearly 0.5 through May to November.

Table 23. The seasonal variation of the number of both sexes in field collected *Lyctocoris beneficus* and *Xylocoris galactinus* (♂:♀).

		<i>L. beneficus</i>	<i>X. galactinus</i>
April	B.	—	—
	M.	—	—
	E.	—	13:16(7)
May	B.	4: 2(0)	24: 25(12)
	M.	9: 11(2)	18: 10(7)
	E.	12: 14(4)	23: 18(11)
June	B.	22: 13(8)	19: 16(10)
	M.	13: 12(8)	30: 25(18)
	E.	16: 21(11)	13: 9(5)
July	B.	12: 15(12)	41: 37(26)
	M.	18: 14(10)	23: 34(27)
	E.	19: 21(11)	19: 6(4)
Aug.	B.	5: 10(6)	7: 12(7)
	M.	3: 5(2)	14: 16(9)
	E.	8: 14(9)	30: 29(16)
Sept.	B.	12: 8(6)	34: 42(25)
	M.	19: 24(17)	28: 19(17)
	E.	32: 27(21)	17: 21(15)
Oct.	B.	11: 13(4)	5: 6(2)
	M.	14: 12(7)	4: 8(5)
	E.	8: 11(2)	3: 7(3)
Nov.	B.	2: 5(3)	1: 3(1)
	M.	—	3: 5(3)
	E.	0: 2(1)	—

Figure in parenthesis shows the number of females with mature ovaries.

Generally speaking, in the giant straw piles *X. galactinus* was much more prevalent than *L. beneficus*. While in the straws from the farmer's barn, especially in wheat straws, *L. beneficus* was more prevalent. *X. galactinus* was seemed to prefer somewhat moist straw, although in some cases many

individuals of this species were found in the very dry straw. *L. beneficus* was almost decidedly predominant in somewhat dry paddy straws (Table 24). However, in some cases, *X. galactinus* was found in the dry straw which was generally fitted for the life of *L. beneficus*. In such cases the latter species was never observed. Therefore, we can see a marked habitat segregation between both species. The competition and habitat segregation between these bugs will be well discussed in Chapter 15.

Table 24. Predominancy of *Lyctocoris beneficus* and *Xylocoris galactinus* in relation to the straw condition.

	Condition and kind of straw			
	Dry paddy straw	Moderately dry paddy straw	Wet paddy straw	Dry wheat straw
<i>L. beneficus</i>	++ ~ ++	+	+ ~ -	+++
<i>X. galactinus</i>	+	+++	+++	+

The insect fauna in the straw pile is often comparatively complicated, and seems to be apparently influenced by the condition of straws. Especially humidity of straws seems to play an important role on the occurrence of Anthocorid bugs. The prevalent insects and other creatures in the straw pile are listed in Table 25.

The methods applied to the determination of prey in the field collected insects contain the microscopic examination of gut contents or excreta (Putman 1962, etc.), precipitin tests (Dempster 1960, 1963, Laughton et al. 1955), paper chromatographic analysis (Putman 1965), etc. These methods have proved useful in some circumstances, but all have certain limitations. The author at first tried the paper chromatographic analysis using Metcalf and Newell's solvent system (Metcalf and Newell 1962). But it was proved that this method can not be applicable at least against the author's materials. The results of both circular and ascending developments failed to the separation of pigments, and in addition to it, the amount of fluorescent substances were too small to be detected in the chromatographic technique. In the present work, one of the field collected insects was reared in the laboratory together with the Anthocorid bugs, and the prey-predator relationship was observed (Table 26).

The results of feeding test indicate that both species of bugs prefer to prey on mites and some other small animals, especially having soft-exoskeleton and sluggish movement, such as Dipterous and Lepidopterous larvae. And the bugs avoid to prey on insects with hard body-wall.

As former workers mentioned, some individuals of *L. beneficus* were attracted to the lamp during the summer season (Oho 1955c, Hiura 1957). According to the author's work in the summer season, the nocturnal activities of

Table 25. List of dominant animals in relation to the straw conditions.

	Condition of straw			
	1	2	3	4
<i>Acarus siro</i> Linné (Acaridae)	+++	+	—	++
<i>Armandillidium vulgare</i> Latreille (Armandillidiidae)	—	—	++	—
<i>Onychiurus pseudarmatus</i> Miyoshi (Onychiuridae) j	—	++	+++	—
<i>Ctenolepisma villosa</i> Escherrich (Lepismatidae)	+	+	—	+
<i>Teleogryllus emma</i> (Ohmachi et Matsuura) (Gryllidae)	—	+	+	—
<i>Gryllus minor</i> Shiraki (")	—	—	+	—
<i>Anisolabis maritima</i> Borelli (Psalididae)	—	—	+	—
<i>Trogium pulsatorium</i> Linné (Trogidae)	++	—	—	+
<i>Stenopirates japonicus</i> Esaki (Enicocephalidae)	—	—	—	++
<i>Physopleurella armata</i> Poppius (Anthocoridae)	—	—	—	+
<i>Cardiastethus fulvescens</i> (Walker) (") j	—	—	—	+
<i>Pyralis farinalis</i> Linné (Pyralidae)	—	—	—	+
<i>Pterastichus Zonginqus</i> Bates (Harpalidae)	—	—	+	+
<i>Philonthus macies</i> Sharp (Staphylinidae)	—	—	+	—
<i>Philonthus sericans</i> Sharp (")	—	+	++	—
<i>Falagria fovea</i> Sharp (")	—	+	+	—
<i>Achenomorpus Zithocharoides</i> Sharp (")	—	+	++	—
<i>Platyola paradoxa</i> Bernhauer (")	—	—	+	—
<i>Opetiopalpus obesus</i> Westwood (Cleridae)	+	++	—	+
<i>Monotoma picipes</i> Herbst (Rhizophagidae)	++	++	—	+
<i>Silvanus lewisi</i> Reitter (Silvanidae)	+	+	+	—
<i>Silvanus necticollis</i> Reitter (")	+	+	—	+
<i>Silanoprus scuticollis</i> Walker (")	+	—	—	—
<i>Oryzaephilus surinamensis</i> Linné (")	+	—	—	+
<i>Cryptophagus cellaris</i> Scopoli (Cryptophagidae)	+	—	—	—
<i>Cryptophilus oblitteratus</i> Reitter (Erotylidae)	—	+	++	—
<i>Holoparamerus signata</i> Wollaston (Corylophidae)	+	+	—	+
<i>Stephostethus chinensis</i> Reitter (Lathridiidae)	+	+	—	—
<i>Lathridius transversus</i> Olivier (")	++	+	—	+
<i>Typhaea stercorea</i> Linné (Mycetophagidae)	++	+	—	+
<i>Tribolium castaneum</i> (Herbst) (Tenebrionidae)	—	—	—	+
<i>Anthicus confucii</i> Marseul (Anthicidae)	+	++	—	+
<i>Anthicus floralis</i> Linné (") j	—	++	—	+
<i>Anthicus tanakai</i> Nomura (")	+	+	—	—
<i>Sitophilus oryzae</i> Linné (Rhynchophoridae)	+	—	—	+
<i>Monomorium</i> sp. (Formicidae)	—	+	++	—
<i>Chlorops oryzae</i> Matsumura (Chloropidae)	—	+	+	—

1 : dry paddy straw.

3 : wet paddy straw.

3: moderately dry paddy straw.

4 : dry wheat straw.

Table 26. Prey-predator relation between the two Anthocorid bugs (*Lyctocoris beneficus* and *Xylocoris galactinus*) and other insects found in the straw.

	<i>L. beneficus</i>	<i>X. galactinus</i>
<i>Acarus siro</i>	++	++
<i>Armandillium vulgare</i>	0	0
<i>Ctenolepisma villosa</i>	t-	-t
<i>Anisolabis maritima</i>	0	0
<i>Stenopirates japonicus</i>	<u>t</u>	<u>t</u>
<i>Pyralis farinalis</i>	++	++
<i>Philonthus sericans</i>	0	0
<i>Opetiopalpus obesus</i>	0	—
<i>Monomorium</i> sp.	—	—
<i>Chlorops oryza</i>	++	++

++: frequently preyed by Anthocorid bugs.

+: preyed by Anthocorid bugs.

0: no prey-predator relation exists.

±: cannibalism was seen between the species.

—: Anthocorid bug was attacked.

both species began just before sunset and reached climax about 1 hour after sunset. As no sufficient specimens of *L. beneficus* were collected, the available data was obtained only on *X. galactinus* (Fig. 5).

The nocturnal activity of *X. galactinus* differs significantly between the sexes. A large number of male adults began to appear on the white organdy cloth which was spread on the ground just before sunset. Many male individuals were also caught at the lower zone, about 0-2 meters in height. Later soon they flew high up to the light which was set at 5-6 m in height, and the peak of the number of attracted bugs was seen about one hour after sunset. The females began their nocturnal flight about half an hour later than that of male. At first, they aggregated on the white organdy cloth and later flew up to higher zone as in the case of males. The majority of collected specimens on the organdy cloth were male, while the female was predominant in the higher zone collection. The peak of the number of attracted females at light was observed between 7-9 PM. After this period, the females suddenly decreased in number and finally very few individuals were collected at the light. The nocturnal activity ceased at the end of September probably due to the cooler temperature. The active flight of the bugs were also observed under the laboratory temperature condition higher than 35°C and under strong illumination.

The nocturnal activity of female adult is seemed to have little correlation to the status of ovarian development, because the dissections of the

Fig. 5. Nocturnal activity of *Xylocoris galactinus* attracted to the light on Aug. 30th, 1967 (Sunset at 18:46 o'clock).

Abscissa: hour in o'clock. Ordinate: No. of bugs collected.

collected female specimens indicate the various stages of ovarian development.

To clear the life history in the field condition, a small straw pile was made in the campus of the university. Its size was about 1.5 m in length, width and height respectively. Five pairs of each species were placed in the 3×5 cm (diameter x length) cylindrical screen cage in which the mite infested straws and the rice stem borer (*Chilo suppressalis* (Walker)) were provided as food. Then, the screen cage was inserted in the central part of the pile. Observations and supply of food were carried out once a week. When the bugs became too much crowded in a cage, a part of them were separated to another cage, and the observations were continued (Tables 27-28).

As seen in Tables 27 and 28, *L. beneficus* and *X. galactinus* might have 3 to 5 generations per annum. It seems that the first adult of each generation of *L. beneficus* appears in May, July, August, September and October. In *X. galactinus*, such adult may emerge in May, June, July, August, September and November. As the oviposition period of the adults are long during the warmer seasons, each stage of bugs is observed or co-existed in the cage at the same time. According to the author's observation the last

Table 27. Life cycle of *Lyctocoris beneficus* under the field condition.

May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
Λ	Λ	A A A					
	E E E	E E E	E				
	N N N	N N N	N N N	N			
		A A A	A A A	A A A	A		
			E E E	E E E	E E		
		N	N N N	N N N	N N N	N N	
			A A	A A A	A A A	A A A	A
				E E E	E E E	N N	
				N N	N N	A A A	Λ Λ
					A A		—

Table 28. Life cycle of *Xylocoris galactinus* under the field condition.

May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
Λ	Λ	A A					
	E E	E E					
	N N	N N N	N				
		A A A	A A A	A			
		E E E	E E E	E			
		N N N	N N N	N N N	N		
		A A	A A A	A A A	A A A	A	
		E	E E E	E E E	E E E		
			N N N	N N N	N N N	Λ Λ Λ	Λ
			A	E	A A A	Λ Λ Λ	—
				N N	N N N	N N N	
					E E	Λ Λ Λ	Λ Λ
					N	N N N	N
						A	A A
							—

adult *L. beneficus* died on 16th, the lowest temperature being 1.2°C, and *X. galactinus* on 9th of December, the minimum temperature being 2.8°C. The death was caused partly by the cold temperature of the season. But in the giant straw pile or in the stock yard, the adult can survive until the next spring in the inner warmer part of the pile.

Preying habit

I. Preying record.

Anthocorid bugs are generally known as polyphagous predators. For instance, *Orius insidiosus* (Say) has been recorded as a predator of 53 species of insects and mites (Marshall 1930). According to the prey list of *Anthocoris nemorum* (Linné) compiled by Hill (1957) and Collyer (1967), the species prey upon spiders, mites, aphids, midges, scale insects, *Lygus* bugs, etc. Although

the preying range of *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) have not investigated thoroughly, several insects were previously recorded as their prey under the laboratory or natural condition (Table 29).

Table 29. A list of previously known prey of *Lyctocoris beneficus* and *Xylocoris galactinus*.

L. beneficus

Brachmia triannulella H.-S. (Gelechiidae, Lep.)¹⁾

Aedia leucomelas Linné (Noctuidae, Lep.)¹⁾

Sesamia inferens Walker (Noctuidae, Lep.)¹⁾

Chilo suppressalis (Walker) (Pyralidae, Lep.)^{1,2)}

Arga pagana Panzer (Argidae, Hym.)¹⁾

Pachygrontha antennata Uhler (Lygaeidae, Heter.)¹⁾

Calaphis magnolicolens Takahashi (Aphididae, Homo.)²⁾

Hyalopterus arundinis Fabricius (Aphididae, Homo.)¹⁾

Dactynitus gobonis Matsumura (Aphididae, Homo.)¹⁾

Aranea¹⁾

X. galactinus

Laemophloeus ferrugineus (Stephen) (Cucujidae, Col.)³⁾

Haemolaelaps casalis (Berlese) (Laelaptidae, Acarina)³⁾

Tyrophagus casei (Oudemans) (Acaridae, Acarina)³⁾

Leidionychus krameri (G. & R. Canestrini) (Uropodidae, Acarina)³⁾

1). Oho 1955c. 2). Hiura 1957. 3). Hall 1951.

As pointed out by Thompson (1951) that "An understanding of the behaviour of predacious insects and more particularly of the specificity of their host relation is therefore very necessary in any attempt to evaluate the importance of predators as controlling agents of injurious insect", a thorough investigation on the food range is indispensable on the studies of predators. In the present work, the author tested more than 60 species of insects and other small animals which are composed of paddy plant insect pests, citrus insect pests, straw dwelling insects and stored grain insect pests. The inclusion of the last group of insects was intended mainly for the purpose of finding some preferable prey for the mass culture of bugs under the laboratory condition. The feeding responses of *L. beneficus* and *X. galactinus* against these animals are given in Table 30.

As seen in Table 30, both *L. beneficus* and *X. galactinus* have a wide food range, including 45 species of 23 families and 43 species of 22 families as prey respectively. Though some variations were observed between the two species on the intensity of response to prey, this fact may indicate a close correlation between the body size of prey and predator. In another word,

Table 30. The feeding response of *Lyctocoris beneficus* and *Xylocoris galactinus* to various prey.

Prey	Stage of prey	Response of <i>L. beneficus</i>	Response of <i>X. galactinus</i>
<i>Ctenolepisma villosa</i> Escherich (Lepismatidae Thysa.)	A - N	+	+
<i>Thermobia domestica</i> Packard (Lepismatidae Thysa.)	A - N	+	+
<i>Periplaneta fuliginosa</i> Serville (Blattidae Blatt.)	N	-	-
<i>Coptotermes formosanus</i> Shiraki (Rhinotermitidae Iso.)	A - N	-	+
<i>Liposcelis entomophilus</i> Enderlein (Liposcelidae Psoco.)	A - N	++	++
<i>Trogium pulsatorium</i> Linné (Trogidae Psoco.)	A - N	++	++
<i>Scotinophara lurida</i> Burmeister (Pentatomidae Heter.)	N	—	—
<i>Lagynotomus elongatus</i> Dallas (Pentatomidae Heter.)	N	—	—
<i>Nezara antennata</i> Scott (Pentatomidae Heter.)	N	—	—
<i>Nephotettix cincticeps</i> (Uhler) (Cicadellidae Homo.)	A - N	+	+
<i>Laodelphax striatellus</i> (Fallén) (Delphacidae Homo.)	A - N	+	+
<i>Nilaparvata lugens</i> (Stål) (Delphacidae Homo.)	A - N	+	+
<i>Sogatella furcifera</i> Horváth (Delphacidae Homo.)	A - N	+	+
<i>Cinara piniformosa</i> Takahashi (Adelgidae Homo.)	A - N	+	+
<i>Acyrtosiphon pisum</i> Harris (Aphididae Homo.)	A - N	++	++
<i>Aphis gossypii</i> Glover (Aphididae Homo.)	A - N	++	++
<i>Brevicoryne brassicae</i> (Linné) (Aphididae Homo.)	A - N	++	++
<i>Calaphis magnolicolens</i> Takahashi (Aphididae Homo.)	A - N	++	++

Prey	Stage of prey	Response of <i>L. beneficus</i>	Response of <i>X. galactinus</i>
<i>Hyalopterus arundinis</i> Fabricius (Aphididae Homo.)	A - N	+	++
<i>Macrosiphum avenae</i> Shinji (Aphididae Homo.)	A - N	++	++
<i>Melanaphis bambusae</i> (Fullaway) (Aphididae Homo.)	A - N	++	++
<i>Myzus persicae</i> (Sulzer) (Aphididae Homo. j)	A - N	t t	++
<i>Shivaphis celti</i> Das (Aphididae Homo.)	A - N	++	++
<i>Toxoptera aurantii</i> (Boyer) (Aphididae Homo.)	A - N	t t	++
<i>Trichosiphoniella sasakii</i> Matsumura (Aphididae Homo.)	A - N	++	++
<i>Dialeurodes citri</i> Ashmead (Aleyrodidae Homo.)	N	—	—
<i>Icerya seyschellarum</i> Westwood (Margarodidae Homo.)	A - N	—	—
<i>Phenacoccus azalea</i> Kuwana (Pseudococcidae Homo.)	A - N	—	—
<i>Ceroplastis pseudoceriferus</i> Green (Coccidae Homo.)	A - N	—	—
<i>Ceroplastis rubens</i> Maskell (Coccidae Homo.)	A - N	—	—
<i>Aonidiella aurantii</i> (Maskell) (Diaspidae Homo.)	A - N	—	—
<i>Chrysomphalus bifasciculatus</i> Ferris (Diaspidae Homo.)	A - N	—	—
<i>Hemiberlesia rapax</i> (Cornstock) (Diaspidae Homo.)	A - N	—	—
<i>Pseudaulacaspis pentagona</i> (Targioni) (Diaspidae Homo.)	A - N	—	—
<i>Unaspis yanonensis</i> Kuwana (Diaspidae Homo.)	A - N	—	—
<i>Nemapogon granella</i> Linné (Tineidae Lep.)	L	—	—
<i>Canephora asiatica</i> (Staudinger) (Psychidae Lep.)	L	—	—

Prey	Stage of prey	Response of <i>L. beneficus</i>	Response of <i>X. galactinus</i>
<i>Mahasena minuscula</i> (Butler) (Psychidae Lep.)	L	--	—
<i>Phyllocnistis citrella</i> Stainton (Gracilariidae Lep.)	L	+	+
<i>Adoxophyes orana</i> Fischer (Tortricidae Lep.)	L	-t-	+
<i>Brachmia triannulella</i> H.-S. (Gelechiidae Lep.)	L	ii-	+
<i>Sitotroga cerealella</i> (Olivier) (Gelechiidae Lep.)	L	—	—
<i>Aphomia gularis</i> (Zeller) (Pyrilidae Lep.)	L	—	—
<i>Chilo suppressalis</i> (Walker) (Pyrilidae Lep.)	L	it	+
<i>Cnaphalocrocis medinalis</i> Guénee (Pyrilidae Lep.)	L	++	+
<i>Cadra cautella</i> (Walker) (Pyrilidae Lep.)	L	—	—
<i>Galleria mellonella</i> (Linné) (Pyrilidae Lep.)	L	ii-	ii-
<i>Plodia interpunctella</i> (Hiibner) (Pyrilidae Lep.)	L	—	—
<i>Pyalis farinalis</i> Linné (Pyrilidae Lep.)	L	++	++
<i>Susumia exigua</i> Butler (Pyrilidae Lep.)	L	++	++
<i>Tryporyza incertulas</i> (Walker) (Pyrilidae Lep.)	L	++	+
<i>Naranga aenescens</i> Moore (Noctuidae Lep.)	L	++	+
<i>Prodenia litura</i> (Fabricius) (Noctuidae Lep.)	L	+	+
<i>Sesamia inferens</i> Walker (Noctuidae Lep.)	L	++	+
<i>Pamara guttata</i> Bremer (Hesperiidae Lep.)	L	+	—
<i>Cryptolestes minutus</i> Olivier (Silvanidae Col.)	L	++	++

Prey	Stage of prey	Response of <i>L. beneficus</i>	Response of <i>X. galactinus</i>
<i>Oryzaephilus surinamensis</i> Linné (Silvanidae Col.)	L		
<i>Korynetes coeruleus</i> Degeer (Cleridae Col.)	L		+
<i>Tennochila japonica</i> Reitter (Trogostidae Col.)	L	---	---
<i>Teneboides mauritanicus</i> Linné (Trogositidae Col.)	L	---	---
<i>Rhizopertha dominica</i> Fabricius (Bostrychidae Col.)	L		
<i>Tenebrio molitor</i> Linné (Tenebrionidae Col.)	L		
<i>Tribolium castaneum</i> (Herbst) (Tenebrionidae Col.)	L		
<i>Sitophilus oryzae</i> Linné (Rhynchophoridae Col.)	L		
<i>Musca domestica</i> Linné (Muscidae Dip.)	L	---	---
<i>Chlorops oryzae</i> Matsumura (Chloropidae Dip.)	L		
<i>Sarcophaga</i> sp. (Sarcophagidae Dip.)	L	---	---
<i>Chironomus</i> sp. (Chironomidae Dip.)	L	---	---
DD-136 (Nematodes)	A-L	---	---
<i>Armadillidium vulgare</i> Latreille (Armadillidiidae Isopoda)	A	---	---
<i>Acarus siro</i> Linné (Acaridae Acarina)	A-N		
<i>Carpoglyphus lactis</i> Linné (Carpoglyphidae Acarina)	A-N		
<i>Panonychus citri</i> McGregor (Tetranychidae Acarina)	A-N		

smaller sized *X. galactinus* is not fitted to attack the larger-sized prey. Further, it is said that both species of bugs prey on the comparatively less active larvae of Lepidoptera, Coleoptera, Aphis and mites, etc. The negative response shown in the case of some Pentatomid bugs may probably due to its well-developed exoskeleton. The Muscid and Sarcophagid larvae were also unsuitable for food probably due to its elastic cuticle and vigorous

movement responded against the first attack by the predator. Silver fishes and leafhoppers were not preferable food for the bugs, because of their alertness. It is rather worthy of mention that the larvae of stored product Lepidopterous pests, namely *Plodia interpunctella* (Hübner), *Sitotroga cerealella* (Olivier), *Aphomia gularis* (Zeller) and *Cadra cautella* (Walker) were not suitable food of bugs, and bugs afforded with these larvae as food starved to death finally, though the other Lepidopterous larvae were served as good prey for them. In addition to this phenomenon, high mortality of bugs was observed in such insects infesting corn media. This is partly due to the tangling of bug's legs by the secreted thread of Lepidopterous larvae.

II. Feeding preference.

The food preference experiments of *L. beneficus* and *X. galactinus* were carried out using the larvae of *Galleria mellonella* (Linné), *Chilo suppressalis* (Walker) and *Celaphis magnolicolens* Takahashi. Five individuals of Lepidopterous larvae, 20 of *Tribolium* larvae and *C. magnolicolens* were provided in a 9x8 cm petri-dish which contained 10 adults or 3rd instar nymphs of predator. Among the prey provided, aphids were afforded with a leaf of *Magnolia obovata* Thunberg, and the other prey were placed in the petri-dish without any shelter or so. The number of dead or damaged prey was counted after 24 hours. Table 31 shows the average of 10 replications.

Table 31. Feeding preference of *Lyctocoris beneficus* and *Xylocoris galactinus* against four kinds of prey (10 bugs in each test, average of 10 tests).

		No. of prey consumed per day			
		<i>Galleria mellonella</i>	<i>Chilo suppressalis</i>	<i>Tribolium castaneum</i>	<i>Celaphis magnolicolens</i>
<i>L. beneficus</i>	Adult	0.2	1.5	12.4	17.8
	3rd instar	0.0	0.1	9.3	16.4
<i>X. galactinus</i>	Adult	0.1	0.3	6.5	11.6
	3rd instar	0.0	0.0	3.2	8.1

As indicated in Table 31, it is evident that both adult and nymph of *L. beneficus* feed more prey than *X. galactinus* do. It is undoubtedly due to the larger size of the former species. The effect of body size of the predator on the food amount consumed was more significantly observed in the number of aphids preyed. The preference order was *C. magnolicolens*, *T. castaneum*, *C. suppressalis* and *G. mellonella*, and this tendency was similarly observed in each species. This test also indicated that the Lepidopterous larvae were not preferred prey. If the predators were not afforded with aphids or flour beetles, the bugs attacked the Lepidopterous larvae very frequently especially when the bugs were put into the dish in large numbers or gregariously.

The prey preference of bugs among the larvae of *C. suppressah*, *Tryporyza incertulas* Walker and *G. mellonella* was also tested, but no significant differences of preference were observed among the provided materials (Table 32).

Table 32. Feeding preference of *Lyctocoris beneficus* and *Xylocoris galactinus* against three kinds of Lepidopterous larvae (10 predators and 5 prey of each species).

		No. of prey consumed per day		
		<i>Chilo suppressalis</i>	<i>Tryporyza incertulas</i>	<i>Galleria mellonella</i>
<i>L. beneficus</i>	Adult	2.4	2.2	1.8
	3rd instar	2.2	2.1	1.1
<i>X. galactinus</i>	Adult	1.3	0.9	0.8
	3rd instar	0.2	0.4	0.2

The number of attacked insects indicated in this table was apparently smaller than the result shown in Table 34. This may be attributed to the fact that the bugs prefer rather inactive larvae or damaged individuals to healthy and active ones. Therefore, preyed larvae were frequently faced to death with a number of pierced scars. However, the bugs avoid to attack the dying or dead larvae. As the food preference of bugs seemed to be greatly influenced by the body size of prey, larvae of *G. mellonella* were divided into three groups by their body weight, namely 1) more than 150 mg, 2) between 150 mg and 70 mg 3) less than 70 mg and each group was used for experiments to compared the response of bugs against the various sized prey (Table 33).

Table 33. Number of punctures found on *Galleria mellonella* larvae of different weight by *Lyctocoris beneficus* and *Xylocoris galactinus* (24 hour exposure against 1.0 predators).

		Range of body weight (mg)		
		>150	150-70	<70
<i>L. beneficus</i>	Adult	4.5	8.6	13.3
	3rd instar	2.2	7.4	11.5
<i>X. galactinus</i>	Adult	0.3	8.1	12.4
	3rd instar	0.0	6.0	9.2

The larva of *G. mellonella* is rich enough with blood. According to Richardson et al. (1931) it is about 4 % of body weight. On the other hand, the maximum feeding amount per one meal by the adult *L. beneficus* does not exceed 0.5 mg. Therefore, the feeding amount of the nymph of *L. beneficus*, adult and nymph of *X. galactinus* is seemed, of course, to be less than that

of the adult *L. beneficus*. Consequently, even the smallest-sized *Galleria* larvae may provide adequate amount of food to satisfy the predator's appetite. From this reason, the increase of the number of punctures on the body surface of smaller larvae may not explain the insufficiency of blood supply at one meal even in the adult *L. beneficus*, but it is rather demonstrating the accessibility of the smaller larvae by the predators.

III. Feeding behaviour.

Several steps are normally observed in the feeding behaviour of *L. beneficus* and *X. galactinus*: 1) The antennae, which are usually held in a bended position in the resting insect, are stretched forward straightly towards the prey in front of the head; 2) the bug approaches straightly the prey; 3) The bug depresses the anterior part of body, rises its fore legs and extends its proboscis forwards; 4) Then, the bug pounces the prey and almost at the same time inserts its proboscis. The prey becomes inactive immediately after the insertion. It seems probable to the author that an anaesthetizing fluid might be injected into the wound simultaneously with the insertion of its proboscis. One meal of *L. beneficus* took 2 to 4 minutes and that of *X. galactinus* was a little shorter. In the case of *Tribolium* larvae, one attack by the bug is fatal to the larvae which will be mummified soon after the attack. In the relatively matured larvae of the rice stem borer or the wax moth, no individuals died immediately after the attack. But, the blood sucked prey are weakened gradually and easily accept additional attacks by the other bugs. The wound on the body wall of prey is noticed as a small black speck which enlarges later and changes to a black necrosis area. Even if the damaged larvae were fortunately free from the second attack or intentionally separated from predators, majority of them died within 48 hours. In a very few cases some individuals succeeded to pupate or to make cocoons, but all the pupae failed to accomplish their pupal development. No particular parts of the prey's body seemed to be favoured by the predators, because in the damaged larvae the punctures occur either on the segment or on the intra-segmental membrane from prothorax to telson.

IV. Food requirement and daily consumption.

To evaluate the bugs as natural enemies or to establish a method of mass production of the bugs, it is necessary to make further detailed observations on the feeding habit of the bugs. To acquire more precise data on the food requirement by a single bug, an adult of both species was reared separately in a 50 ml beaker which is provided with 3 individuals of mature larvae of *C. suppressalis*, or 3rd instar larvae of *G. mellonella* or 3 mature larvae of *T. castaneum*. And the number of preyed larvae was counted daily (Table 34). The adults provided in this test were fed with the larvae of *T. castaneum* during their nymphal stages.

Table 34. Number of daily consumed prey during the adult stage of *Lyctocoris beneficus* and *Xylocoris galactinus* (average of 10 days test, with 10 predators).

	<i>Chilo suppressalis</i> ¹⁾	<i>Galleria mellonella</i> ²⁾	<i>Tribolium castaneum</i> ¹⁾
<i>L. beneficus</i>	0.7	1.3	2.1
<i>X. galactinus</i>	0.2	0.5	0.8

1) : mature larvae.

2) : pre-mature larvae.

The results of tests almost coincided with the preceding one. Namely, the frequency of attack was proportional to the body size of prey. Though it is difficult to discuss the importance of the Anthocorid bugs as the predators against the rice stem borer merely from this data, *L. beneficus* seemed to be an excellent predator, so far as its attack of the borer mainly in the field condition was concerned. *X. galactinus* is surely less efficient than *L. beneficus* in the solitary condition, and often the former species is crushed down by the borer when it pounced upon. Nevertheless, it is an efficient predator too when they occur gregariously.

The daily food requirement of the nymphs was estimated with the larvae of *T. castaneum* only, because the large-sized Lepidopterous larvae were unsuitable for the food of nymphs of both species in a solitary condition (Tables 35 & 36).

Table 35. Number of consumed *Tribolium castaneum* larvae in the nymphal stage of *Lyctocoris beneficus*.

Instar of bug	A	B	C
1st	3.4	1.8	0.53
2nd	4.6	3.4	0.74
3rd	5.2	5.2	1.00
4th	4.0	6.6	1.65
5th	8.2	14.0	1.70
Total	25.4	33.0	1.30

A : duration of instar in days.

B : number of larvae preyed during the instar.

C : $\frac{A}{B}$ daily consumption of prey.

Table 35 indicates that the nymph of *L. beneficus* accomplished its nymphal development after consuming 33 *Tribolium* larvae and about 44% of the prey was consumed during the last nymphal stage. The daily con-

Table 36. Number of consumed *Tribolium castaneum* larvae in the nymphal stage of *Xylocoris galactinus*.

Instar of bug	A*	B*	C*
1st	2.2	1.1	0.50
2nd	2.6	1.4	0.54
3rd	4.4	2.6	0.59
4th	3.6	2.4	0.66
5th	4.8	3.4	0.71
Total	17.6	10.9	0.61

* abbreviation same as that of Table 35.

sumption was seemed to be suddenly increased from the fourth instar. In *X. galactinus* (Table 36), it consumed only about 11 larvae during its nymphal stage. Contrary to *L. beneficus*, the consumption gradually increased from 0.50 to 0.71 prey per day through the whole instars. Difference in daily consumption between adult and nymph was less pronounced in *X. galactinus* than in *L. beneficus*, 0.8 : 0.7 and 2.1 : 1.7 respectively. In the actual amount of daily consumption of *Tribolium* larvae by the mature nymph of *L. beneficus* was still prevalent to that of the adult of *X. galactinus*.

The nutritional value of *Tribolium* larvae against each instar of bugs was expressed by the predatory value and conversion ratio which were introduced by Fewks (1960) and Evans (1962) respectively. Also the nutritional value of larvae of *C. suppressalis*, *G. mellonella* and *T. castaneum* against the adult stage was evaluated with the same formula.

The value of conversion ratio and predatory value were calculated using the following expressions.

$$\text{Conversion ratio} = \frac{\text{Increase in weight}}{\text{Weight consumed}} \times 100$$

$$\text{Predatory value} = \frac{\text{Weight consumed during instar}}{\text{Duration of instar}}$$

where all weights are fresh weights.

In this test, the predator was kept singly and provided with weight-known prey. The predator and prey were inspected and weighed daily and fresh prey was provided if necessary. The weighing was done on a single pan automatic balance weighing to 0.1 mg. From the daily inspection it was possible to obtain a record of the weight of food consumed by the predators during their development, and the results were given in Tables 37 and 38. As the first instar was too light to be weighed, the works were started from the second instar.

Table 37. Conversion ratio and predatory value of *Lyctocoris beneficus* nymph fed on *Tribolium castaneum*.

	Instar of nymph			
	2nd	3rd	4th	5th
Initial weight (mg)	0.2	0.6	1.1	1.4
Initial weight of the succeeding instar (mg)	0.6	1.1	1.4	2.2
Increase in weight (mg) (A)	0.4	0.5	0.3	0.8
Weight of food consumed (mg) (Bj)	0.7	0.9	1.3	2.2
Conversion ratio (%) (A/B)	57.1	55.6	44.3	36.4
Duration of instar (in 'days) (C)	7.1	7.3	7.8	11.8
Predatory value (mg/day) (B/C j)	0.10	0.12	0.17	0.19

Table 38. Conversion ratio and predatory value of *Xylocoris galactinus* nymph fed on *Tribolium castaneum*.

	Instar of nymph			
	2nd	3rd	4th	5th
Initial weight (mg)	0.2	0.3	0.5	0.8
Initial weight of the succeeding instar (mg)	0.3	0.5	0.8	1.3
Increase in weight (mg) (A)	0.1	0.2	0.3	0.5
Weight of food consumed (mg) (Bj)	0.2	0.5	0.8	1.5
Conversion ratio (%) (A/B)	50.0	40.0	37.5	33.3
Duration of instar (in days) (C)	4.3	7.2	8.4	9.2
Predatory value (mg/day) (B/C)	0.05	0.07	0.09	0.16

The data indicated in Tables 37 and 38 have some variations from the results of previous experiments both in the duration of nymphal stage and the amount of food consumed. This was probably due to the excessive handling of materials by daily weighing, and high mortality of nymphs was also observed. Although the bugs were too small to make accurate estimation, the data showed higher conversion ratio in *L. beneficus* than in *X. galactinus*. And earlier instars showed a higher conversion ratio than the later ones as observed in *Cimex lectularius* Linné and *Phonoctonus nigrofasciatus* Stål by Johnson (1960) and Evans (1962) respectively. Compared with *X. galactinus*, the conversion ratio of *L. beneficus* was higher through 4 nymphal stages. This may suggest that *L. beneficus* nymph has more efficient metabolic mechanism than *X. galactinus*. The predatory value was lower in earlier instars in both species. The lower conversion ratio and higher predatory value of the late nymphal instars may prove the active predatory action of mature nymphs.

The nutritional value of 3 kinds of foods against *L. beneficus* and *X. galactinus* was estimated with the newly emerged female adults for ten days. The weight change and feeding record were illustrated in Figs. 6 and 7, and the result was also summarized in Table 11 with the value of conversion ratio and predatory value. The materials used in this test were fed to the adult with *T. castaneum* larvae.

As seen from Figures, the body weight of *L. beneficus* increased almost stably irrespective of kinds of prey provided. While in *X. galactinus*, the stability of body weight was only observed in individuals afforded with *Tribolium*. And even the decrease of body weight was observed in *Chilo*-fed individuals. This is apparently indicating the unsuitability of those larvae as food of *X. galactinus* at least in the solitary rearing condition. The increase of body weight was apparently depending upon the development of ovaries, and this was ascertained by the swelling of abdomen as well as the anatomical dissection.

Fig. 6. Feeding record and body weight change of the adult female *Lyctocoris beneficus*; **Abcissa**, body weight of bug (mg); **ordinate**, days from emergence.

A: fed on larvae of *Tribolium castaneum*.

B: fed on larvae of *Chilo suppressalis*.

C: fed on larvae of *Galleria mellonella*.

O: shows the day not fed. X: shows the day fed.

Fig. 7. Feeding record and body weight change of the adult female *Xylocoris galactinus* ; *Abscissa*, body weight of bug (mg) ; *ordinate*, days from emergence.
A : fed on larvae of *Tribolium castaneum*,
B: fed on larvae of *Chilo suppressalis*.
C: fed on larvae of *Galleria mellonella*.
O: shows the day not fed, X: shows the day fed.

Table 39. Conversion ratio and predatory value of the adult female *Lyctocoris beneficus* and *Xylocoris galactinus* against different food.

Prey-predator	A		B		C	
	L.	X.	L.	X.	L.	x.
Initial weight (mg)	2.3	1.5	2.3	1.5	2.2	1.4
Final weight (mg) (after 10 days)	2.6	1.4	2.8	1.6	2.7	1.7
Increase in weight (mg)	0.3	-0.1	0.5	0.1	0.5	0.3
Weight food consumed (mg)	0.8	0.3	2.1	0.9	2.4	1.2
Conversion ratio (%)	25	-3	24	11	21	25
Predatory value mg/day	0.12	0.03	0.19	0.09	0.24	0.12
Number of meals	7	2	13	5	18	8

A : mature larvae of *Chilo suppressalis*.
B: premature larvae of *Galleria mellonella*.
C : mature larvae of *Tribolium castaneum*.
L : *Lyctocoris beneficus*.
X: *Xylocoris galactinus*.

The conversion ratio in the adult stage was surely lower than that of nymphal stage in *Tribolium*-fed groups in both species. Except the *Chilo*-fed *X. galactinus*, the conversion ratio was negatively proportional to the numbers of meals. It coincided with Kasting's opinion that less time for feeding consume less food but made more efficient use of it (Kasting and McGinnis 1959).

V. Effect of density on feeding habit.

It is a general conception that population density affects the growth and development of insects, and moderately dense population with abundance of food promotes the growth and food intakes in insects (Pettit 1940, Willis et al. 1958, Feir 1963, McFarlane 1962, etc.). In the present work, to estimate the variation of food intake under different population density conditions, various numbers of adults and 3rd instar nymphs of *L. beneficus* and *X. galactinus* were reared in a 100 ml beaker and the daily consumption of food was counted (Tables 40-43).

As Tables 40-43 illustrate, the population density of Anthocorids scarcely influenced the number of *Tribolium* larvae consumption. Because the body size of *Tribolium* larvae is so small that one larva is enough to be preyed by one Anthocorid individual at one meal. While the daily food amount consumed in denser population decreased in the groups provided with *Chilo* and *Galleria* larvae. But in rough estimation, in gregarious conditions, bugs surely showed better growth and survival than the bug reared in the solitary condition. This conflict may be simply explained by the fact that the bugs prefer to attack inactive prey which was already damaged by another predator. Therefore, many blood sucked larvae died with a number of punctures though other larvae were quite safe. In fact, several

Table 40. Number of prey consumed daily by the adult *Lyctocoris beneficus* under different population density conditions (in 50 ml beaker).

No. of bugs per beaker	Prey	Total of consumed prey per day	No. of daily consumed prey per predator
20	T (<i>Tribolium</i>)	42.6	2.13
	G (<i>Galleria</i>)	1.08	0.54
	C (<i>Chilo</i>)	5.6	0.28
10	T	20.8	2.08
	G	6.2	0.62
	C	3.2	0.32
5	T	9.8	1.96
	G	3.4	0.68
	c	1.8	0.36
1	T	2.0	2.00
	G	0.8	0.80
	C	0.6	0.60

Table 41. Number of prey consumed daily by the 3rd instar nymph of *Lyctocoris* *beneficus* under different population density conditions (in 50 ml beaker).

No. of bug per beaker	Prey	Total of consumed prey per day	No. of daily consumed prey per predator
20	T	27.2	1.36
	G	8.8	0.44
	C	4.4	0.22
10	T	12.8	1.28
	G	4.8	0.48
	C	2.4	0.24
5	T	5.6	1.12
	G	2.6	0.52
	C	1.2	0.28
1	T	1.2	1.20
	G	0.6	0.60
	C	0.2	0.20

Table 42. Number of prey consumed daily by the adult *Xylocorisgalactinus* under different population density conditions (in 50 ml beaker).

No. of bugs per beaker	Prey	Total of consumed prey per day	No. of daily consumed prey per predator
20	T	16.6	0.83
	G	4.8	0.24
	C	4.0	0.22
10	T	8.4	0.92
	G	2.4	0.24
	C	2.2	0.22
5	T	4.4	0.88
	G	1.8	0.36
	C	1.0	0.20
1	T	0.8	0.80
	G	0.4	0.40
	C	0.2	0.20

bugs were sometimes piercing the body of a single prey at the same time. The active attack by the adult and nymph of *X. galactinus* on the rice stem borer and *Galleria* larvae was almost proportional to their gregarious condition. Actually 1 or 2 individuals of bugs were certainly difficult to attack the vulnerable rice stem borer larvae, and they were sometimes smashed to death by the vigorous movement of prey when the bug's stylet pierced the larvae. It is apparent that the adult of *L. beneficus* which preys approximately upon one borer every two days, is a superior predator against

Table 43. Number of prey consumed daily by the 3rd instar nymph of *Xylocoris galactinus* under different population density conditions (in 50 ml beaker).

No. of bugs per beaker	Prey	Total of consumed prey per day	No. of daily consumed prey per predator
20	T	10.2	0.51
	G	3.6	0.18
	C	2.8	0.14
10	T	4.8	0.48
	G	1.8	0.18
	c	1.4	0.14
5	T	2.6	0.52
	G	1.0	0.20
	C	0.6	0.12
1	T	0.4	0.40
	G	0.2	0.02
	C	0.0	0.00

T : larvae of *Tribolium castaneum*.

G : larvae of *Galleria mellonella*.

C : larvae of *Chilo suppressalis*.

the rice stem borer, and the third instar nymph of the bug may also become a borer predator to a certain extent. This, of course, indicates the high efficiency of the fourth and fifth instar nymphs as borer predators. The rearing of the first and second instar nymphs of *L. beneficus* with mature rice stem borers was scarcely successful. In the laboratory condition, these young nymphs were easily reared with larvae of *T. castaneum*, and in the field condition it is presumably feeding on small sized insects of other creatures. In *X. galactinus*, the rearing of young instar nymphs with rice stem borer larvae was quite impossible. Only from the fourth instar the nymph can feed on the mature borer larvae, but in this case, high mortality of nymph was observed. In any case, through all the developmental stages, *X. galactinus* was not an excellent predator in the case of lower population density, while it sometimes showed extremely voracious appetite to rice stem borer larvae when this bug was in gregarious condition.

VT. Phytophagous habit.

Anthocorid bugs are generally known as predatory insects and including some important natural enemies. However, review of the literature reveals that numbers of observations have been recorded on the phytophagous habit of Anthocorid bugs during the past 90 years. The phytophagous data on the Anthocorid bugs are summarized below in chronological order (Table 44):

Although many phytophagous data were recorded, it is a general con-

Table 44. Phytophagous data of Anthocorid bugs.

Author and year of publication	Anthocorid bug	Host plant	Note
McLachlan 1879	<i>Anthocoris nemorum</i>	hop	
McLachlan 1880	<i>A. nemorum</i>	hop	
Theobald 1895	<i>A. nemorum</i>	hop	
	<i>A. confusus</i>	"	
Reuter 3.908	<i>Lyctocoris campestris</i>	vegetable	
Garman et al. 1914	<i>Orius insidiosus</i>	corn	
Matsumura 1920	<i>O. sauteri</i>	mulberry bud	
Bodenheimer 1921.	<i>O. majusculus</i>	chrysanthemum	The adults suck the leaves, shoot and buds, leaving black scale.
Prohaska 1923	<i>A. nemorum</i>	willow, catkins and alder bud	common and very harmful.
van Poetran 1930	<i>O. majusculus</i>	chrysanthemum	
Barber 1936	<i>O. insidiosus</i>	cotton pollen	
Doeksen 1944	<i>O. majusculus</i>	maize cobs	feed on the seeds through innermost bract.
Iglinsky 3.950	<i>O. insidiosus</i>	cotton pollen	male prefer it, female feed on spider, mites or insects.
Sands 1.951	<i>A. nemorum</i>	salix	feed on plant to some extent.
Carayon et al. 1959	<i>O. pallidicornis</i>	flower pollen	
Hill 1961.	<i>A. sarothamni</i>	sap of broom	
Dicke et al. 1962	<i>O. insidiosus</i>	corn pollen	principal food of nymph as well as adult.

ception that animal diets were indispensable for the bugs to continue their activity and survival, and especially to keep their reproductive activity, and Hill (1965) pointed out the indispensability of animal diet on the egg production of *Anthocoris confusus* (Reuter). Additionally, the importance of animal diet on the egg production was also verified on *Orius insidiosus* (Say) by another worker (Barber 1936). To emphasize this conception, all the rearing data of Anthocorid bugs previously published by many workers are concerned exclusively with animal diet. In Japan, only one phytophagous record of the Anthocorid bug was made by Matsumura (1920) on *Orius sauteri* (Poppius) as an insect pest of mulberry buds. But this record was almost opposed by Esaki and Hashimoto (1955) and predation of leafhopper was demonstrated by the latter workers.

Before the trial of making liberation of *L. beneficus* and *X. galactinus* in

the field as the rice stem borer controlling agents, it is rather important to discuss their influence on the plants if any. Additionally, for the purpose of maintaining the cost for mass production of these bugs under economic level, the study on the reliability of the bugs on the vegetable substance is also indispensable. Because the vegetable substances are in almost all cases less expensive and easier to be handled than animal substances.

At first, the first and third instars and newly emerged adults of both species were reared individually in 3 × 8 cm flat bottomed glass tube, providing various substances such as ; 1) dry filter paper, 2) moist filter paper, 3) water soaked urethane foam, 4) sugar solution soaked urethane foam, 5) moist paddy straw, 6) cherry twig, 7) molded cracked corn, and 8) corpse of *G. mellonella* larvae. The last material was added in this series of test for the purpose of testing the necrophagous habit of the bugs. These substances except the cracked corn were renewed daily and survival of bugs was checked. The observation was conducted with 5-7 individuals in each afforded materials and average longevity is given in Table 45.

Table 45. Longevity of solitary reared *Lyctocoris beneficus* and *Xylocoris galactinus* on vegetable diet of different conditions.

	Stage of bug set	Longevity (in days)	
		<i>L. beneficus</i>	<i>X. galactinus</i>
Dry filter paper	1st instar	1.2	1.3
	3rd instar	1.5	1.4
	Adult	2.1	1.0
Wet filter paper	1st instar	3.2	2.4
	3rd instar	3.1	2.8
	Adult	3.8	5.1
Water soaked urethane foam	1st instar	5.2	3.7
	3rd instar	12.6(1)	14.3(1)
	Adult	27.5	22.2
Sugar solution soaked urethane foam	1st instar	7.4	5.5
	3rd instar	14.0(2)	13.2(1)
	Adult	24.7	25.8
Moist paddy straw	1st instar	5.1	4.2
	3rd instar	11.4(1)	12.0(2)
	Adult	14.2	17.9
Cherry twig	1st instar	2.1	2.4
	3rd instar	4.1	4.4
	Adult	5.2	7.8
Molded cracked corn	1st instar	43.2(3)	41.4(3)
	3rd instar	48.5(3)	44.6(3)
	Adult	46.1	48.3
Corpse of larvae of <i>Galleria mellonella</i>	1st instar	25.4(3)	23.1(3)
	3rd instar	32.2(3)	33.5(3)
	Adult	20.9	22.8

Figure in parenthesis shows the frequency of moulting in maximum value.

As Table 45 shows, water or moisture was an indispensable factor to maintain their lives. In both species, the longer longevity was shown in water soaked urethane foam than moist filter paper. It is probably due to the easier sucking of water in the former materials. Sugar solution showed certain effects to prolong the longevity of both bugs. Although Anderson (1962a) indicated that *Anthocoris nemoralis* (Fabricius) adult reared with sugar solution had 3 times longer longevity than the individuals reared water only, the present experiment showed that sugar solution did not exhibit such a high nutritional value. Cherry twig was only suited for a oviposition substratum, and the phytophagosity on this plant was scarcely recognized in both species. The bugs reared on moist straw had shorter longevity than the individuals reared on sugar solution. Both molded cracked corn and corpse of *G. mellonella* larvae had an excellent capacity to maintain the life and restricted developments were observed in these groups. However, the development of the nymph was always much inferior to that of the nymph afforded with animal substances.

To observe the cannibalistic behaviour of bugs, 5 individuals were put together in a tube, and fed with sugar solution, molded cracked corn or corpse of *G. mellonella* larvae. In this case, cannibalism was observed frequently in both species, and generally only a single individual could survive finally. Cannibalism was frequently occurred at the moulting time, and the individuals just after ecdysis and still keeping soft skin were very often devoured by the other individuals. And the result of the experiment is illustrated in Table 46.

Compared the result of Table 45 with 46, the longevity given in Table 46 was shortened, but the growth was rather accelerated by the improvement of nutritional condition by cannibalism in the majority of cases. Moulting

Table 46. Longevity of gregariously reared *Lyctocoris beneficus* and *Xylocoris galactinus* on vegetable diet of different conditions (in maximum value).

	Stage of bug set	Longevity (in days j	
		<i>L. beneficus</i>	<i>X. galactinus</i>
Sugar solution	1st instar	12(1)	12(2)
	3rd instar	25(1)	23(1)
	Adult	32	18
Molded cracked corn	1st instar	32(5)	38(4)
	3rd instar	31(3)	42(3)
	Adult	28	32
Corpse of larvae of <i>Galleria mellonella</i>	1st instar	23(4)	31(5)
	3rd instar	27(3)	28(3)
	Adult	34	27

Figure in parenthesis shows the frequency of moulting.

was also observed more frequently in this test. A part of the adult female *X. galactinus* reared on the corpse from their third instar laid a few eggs from which the newly hatched nymphs have developed to normal nymphs. The oviposition under an abnormal nutritional condition seemed to exhaust the vitality of the mother, the oviposited female died 1-2 days after the oviposition. It revealed that the Anthocorid bugs were essentially carnivorous insects, and cannibalism between nymphs and adults are often occurred under the animal diet defficient condition, and in the animal material supplying condition, cannibalism seldom occurred. Plant substances were of little value as food sources for both species of bugs. Fungus and the corpse of animal were nutritional but not adequate to accomplish the nymphal growth from the first instar, and its maximum limitation of the development of bugs was only attained to the fourth instar in both species. In cannibalistic bugs a part of the first nymphs developed to the adult. It is nothing but indicating that cannibalism is nutritionally superior to necrophagosity and fungiphagosity. Though in the majority of tested cases the third instar nymphs succeeded to become adults, animal diet was exactly an indispensable material for the fecundity of females. The importance of fungus or corpse as food was only observed in the case of scarce prey. This fact was also proved by the combination tests of copulated males and females which were reared with two different food sources from their fifth instar. In this test males and females were reared singly and a male was supplied to a vial containing one virgin adult female of two day old for one day so as to make a desired couple. (Table 47).

Table 47. Comparison of nutritional effect of fungi and animal diet on the fecundities of *Lyctocoris beneficus* and *Xylocoris galactinus*.

	Combination of pairs	Oviposition period	No. of eggs oviposited
<i>L. beneficus</i>	P ♂ x P ♀	—	0
	C ♂ x P ♀	—	0
	P ♂ x C ♀	25.3	31.7
	C ♂ x C ♀	27.1	32.4
<i>X. galactinus</i>	P ♂ x P ♀	—	0
	C ♂ x P ♀	1.5	3.5
	P ♂ x C ♀	26.5	31.9
	C ♂ x C ♀	24.2	38.4

C : fed with larvae of *Tribolium castaneum*.

P : fed with molded cracked corn.

It seems that the lack of animal prey does not influence the reproductive potential of male adult. On the other hand, animal food has a decided influence on the fecundity of female. The animal diet seems to play as another ovipositional stimulant. Namely, the mature female in the oviposition

stage immediately ceased to oviposit when she was reared in the absence of animal food, even if her ovaries were containing mature eggs. Although Sinha (1961) listed *X. galactinus* as a fungivorous insect, fungus is a secondary important food for this bug. In conclusion, the feeding habits of *L. beneficus* and *X. galactinus* are almost similar and their frequency of occurrence or preference may be arranged in the following order : Carnivorous > Cannibalistic > Necrophagous > Fungivorous > Phytophagous.

Among these feeding habits, only the carnivorous habit can maintain the species from generation to generation.

Effects of food on the fecundity

It is a well known fact that the fecundity of various insects may be greatly affected by environmental conditions, and Klomp (1964) made a detailed review on this problem. For the factors influencing the fecundity, nutritional condition is also very important, and voluminous studies have been published on this point (Orr 1964, Bracken 1965, Turner et al. 1967, etc.). The works on this problem with carnivorous insects were reviewed by House (1958). Besides, some remarkable works were carried out on *Chrysopa carnea* Stephen (Hagen 1950, Sundby 1967), Coccinellid beetles (Smith 1965), Anthocorid bugs (Anderson 1962b), etc. In Anderson's work, the effect of prey on the fecundity was illustrated by the length of pre-oviposition stage. He indicates the occurrence of reproductive diapause when the newly emerged adults of some *Anthocoris* spp. are fed on various prey. For example, *Aulacorthum circum-flexum* (Buckt.)-fed *Anthocoris confusus* (Reuter) began oviposition one week after emergence, whereas the pre-oviposition stage prolonged to three weeks when *Psylla mali* Schid. was offered as food. The effect of food on the egg-production and egg vitality on Coccinellid beetle is presented by Smith (1965). He indicated the indispensability of animal prey for the oviposition of the adult beetles. Also Sundby (1967) pointed out that the artificial media are not enough to make full fecundity on *C. carnea*, and the addition of the fresh flower (*Matricaria indorsa* L.) raised the number of produced eggs about three times.

In the present experiment, the fifth instar *Lyctocoris beneficus* (Hiura) and *Xylocoris glactinus* (Fieber) were reared with larvae of *Chilo suppressalis* (Walker), *Galleria mellonella* (Linné) and *Tribolium castaneum* (Herbst). The pre-oviposition stage, number of produced egg and the rate of egg hatching were estimated as the indices of fecundity of female adults which were gregariously reared in a 50 ml beaker (Table 48). Then the pre-oviposition stage was recorded by the day when the first egg was produced on the corresponding group.

As seen in Table 48, the influence of *C. suppressalis*, *G. mellonella* and *T. castaneum* as diet for the bugs showed the same tendency in the fecundity of both *L. beneficus* and *X. galactinus*. Namely, the *Tribolium*-fed bug showed

Table 48. A comparison of the influence of food on the fecundity and longevity of the adult *Lyctocoris beneficus* and *Xylocoris galactinus*.

	Food	Pre-oviposition period(in days)	No. of eggs per female	Percent of egg hatching(%)	Longevity (in days)
<i>L. beneficus</i>	A	9	18.2	55.0	19.4
	B	28	6.1	34.3	31.1
	C	7	26.8	78.6	28.4
<i>X. galactinus</i>	A	9	21.6	63.3	21.2
	B	26	8.3	21.7	29.0
	C	5	32.1	83.5	22.7

A : larvae of *Chilo suppressalis*.

B : larvae of *Galleria mellonella*.

C : larvae of *Tribolium castaneum*.

the highest value both in the egg-production and percent of egg hatching and indicated the shorter pre-oviposition period. The result of *Chilo*-fed bug was evidently inferior to that of the *Tribolium*-fed individuals, while the worst result was obtained in the *Galleria*-fed group. The longer reproductive diapause was observed in the latter two groups. The longevity of adults were in reciprocal proportion to the fecundity.

As to the cause of inferior fecundity in the *Chilo*- and *Galleria*-fed bugs, the intake of small amount of food was first considered but the counting of the numbers of pierced scars or punctures on the skins of attacked prey indicated the sufficient feeding by tested predators. Therefore, it is rather conceivable to attribute the cause of inferior fecundity shown by the *Chilo*- and *Galleria*-fed individuals to the nutritional quality or quantity of food, especially of the component of body -fluid.

The review of previous works on the relation between insect fecundity and nutrition shows that the essentiality of some nutritional components are variable with the species of insects. For instance, inorganic salts are required for egg-production in Tryptid flies (Hagen 1958) and *Drosophila melanogaster* Meigen (Sang and King 1961) but apparently not for *Aedes aegypti* (Linné) (Sing and Brown 1957, Dimond et al. 1956). Whereas protein and amino acids are unexceptionally essential for the reproduction of all tested insects, and Lea et al. (1956) indicated the importance of total amino acids on the egg-production of mosquitoes.

To evaluate the nutritional value of *Galleria* and *Tribolium* larvae on the fecundity of bugs, amounts of free amino acids were quantitatively analyzed. Also to estimate the effect of food on the predators, the amino acids of *L. beneficus* fed on *Galleria* or *Tribolium* larvae were analyzed by the same method. The samples for the amino acid analysis were prepared by the following procedure ; about 200 mg of precisely weighed row

material were immersed in 60°C warm water for 1 minute for the purpose of preventing melanosis, then it was homogenized thoroughly with 80 % ethanol. After the filtration, precipitate was washed for several times with 80 % ethanol. Ethanol was at last extracted by chloroform and water solution of amino acid was condensed in low pressure condition to 0.5 ml. The ninhydrinpositive compounds were analyzed quantitatively with the automatic amino acid analyzer employing a slight modification of the procedure of Piez and Morris (1960). The results of analysis are given in Table 49.

Table 49. Change of free amino acids contents of *Lyctocoris beneficus* among various foods ($\mu\text{mol/g}$).

Amino acid	T	G	T/G	TL	GL	T-TL	G-GL
Unknown 1*	0.26	0.03	8.67	0.37	0.08	-0.11	-0.05
Unknown 2*	0.14	-	-	0.20	0.09	-0.06	-0.09
Unknown 3*	0.04	0.17	0.24	trace	0.10	-0.04	-0.07
Unknown 4*	-	0.35	-	-	-	-	0.35
Taurin?*	0.51	0.35	1.46	2.91	0.89	-2.40	-0.54
Urea?*	0.13	0.23	0.57	0.11	0.08	0.02	0.15
Methionine suloxide?*	0.40	1.86	0.22	0.50	0.71	-0.10	1.15
Aspartic acid	0.34	0.10	3.40	0.13	0.13	0.21	-0.03
Threonine**	4.45	2.92	1.51	0.88	1.33	3.57	1.59
Serine	4.91	1.24	3.96	0.81	0.60	4.10	0.64
Glutamic acid	7.41	3.67	2.02	3.72	1.43	3.69	2.24
Proline	9.94	4.62	2.15	0.28	1.79	9.66	2.83
Citrulline*	-	1.60	-	-	0.32	-	1.28
Glycine	3.96	2.38	1.66	1.03	0.91	2.93	1.47
Alanine	17.92	5.72	3.13	2.30	2.18	15.62	3.54
Unknown 5*	-	0.14	-	-	trace	-	0.14
Valine**	4.56	1.10	4.15	trace 0.44	0.47	4.12	0.63
Methionine**	0.51	0.17	3.00	0.18	0.08	0.33	0.09
Isoleucine**	1.89	0.41	4.61	0.13	0.17	1.76	0.24
Leucine**	2.87	0.78	3.68	0.26	0.28	2.61	0.50
Unknown 6*	trace	-	-	trace	-	-	-
Tyrosine	5.44	1.04	5.23	0.23	0.54	5.21	0.50
Phenylalanine**	1.32	0.34	3.88	0.12	0.15	1.20	0.19
Unknown 7*	1.01	0.32	3.16	trace	trace	1.01	0.32
Lysine**	2.53	1.20	2.11	0.38	0.15	2.15	1.05
Histidine**	2.87	2.41	1.19	0.49	0.86	2.38	1.55
Ammonia	1.60	0.31	5.16	0.34	0.30	1.26	0.01
Arginine**	2.58	0.93	2.77	1.05	0.51	1.53	0.42
Total of essential amino acids	23.58	8.10	2.91	3.93	4.00	19.65	4.10
Total of amino acids	73.50	30.63	2.40	12.43	11.95	61.07	18.68
Grand total	77.59	34.39	2.26	16.86	14.35	60.73	20.04

T : larvae of *Tribolium castaneum*.

G : larvae of *Galleria mellonella*.

TL : *Tribolium-f* ed *L. beneficus*.

GL : *Galleria-f* ed *L. beneficus*.

* : calculated by the coefficient of aspartic acid.

** : essential amino acid.

The interpretation of the results shows that the ninhydrin-positive substance of *Tribolium* larvae is much more abundant than *Galleria* larvae, and the ratio of *Tribolium*/*Galleria* (T/G) is 2.26. Further, the ratio becomes more significant when identified amino acids are compared, namely that is 2.40. Although no data are available to discuss the essential amino acids for the Anthocorid bugs, ten following amino acids required by the rat are also essential for the majority of insects (House 1965). They are: arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophane and valine. On the other hand, amino acids that are non-essential for rat have been shown to be essential for other insects. According to the literature, the essentiality of the kinds of amino acids differs with the species. Cystine and perhaps glycine are essential for *Aedes aegypti* (Linné) (Golberg and DeMeillon 1948, Sing and Brown 1957). Alanine, glycine, serine and tyrosine are presumably essential for *Agrius affinis* (Fallen) (House 1954). The essential amino acids on a Coccinellid beetle (*Coleomegilla maculata* (DeGeer)) were investigated by Atallah and Killebrew (1967) and they recorded 8 essential amino acids. In this case lysine, methionine and tryptophane were not included among the above mentioned 10 essential amino acids but added alanine. The essential amino acids of *Rhodnius prolixus* (Stål), a Reduviid bug, was studied by Pickett and Friend (1965) and the result is almost coincided with the requirement of rat, except that they added glycine to the essential group. The ratio of T/G in these 10 essential amino acids shows 2.91, and the actual amount is $15.48 \mu\text{mol/g}$ (23.58–8.10). Accordingly the *Tribolium* larva contains about 3 times of essential amino acids than the *Galleria* larva, and the amount of each essential amino acid except histidine in the *Tribolium* larva exceeds twice that of the *Galleria* larva, especially in valine and isoleucine the ratio was 4.15 and 4.16 respectively. Although the physiological role of these amino acids on insect reproduction still remain unknown, such quantitative difference in the essential amino acids contents may show some suggestions to solve this problem.

Apparent quantitative differences are also obtained in some non-essential amino acids and other ninhydrin positive substances. The superior amount of these substances in the *Galleria* larvae to the *Tribolium* larvae is only observed in urea (?), methionine sulfoxide (?), citrulline (?) or some unknown substances, while the adverse relation are shown in many kinds of material which include all members of identified non-essential amino acids such as : aspartic acid, serine, glutamic acid, proline, glycine, alanine and tyrosine. The amount of these dispensable amino acids in the *Tribolium* larva is $49.92 \mu\text{mol/g}$ and that of *Galleria* larva is only 18.77. Among these amino acids, alanine which is recognized to be a very important amino acid in *Blattella* and *Agria* sp. by Hilchey (1953) and House (1954) respectively is significantly abundant in the *Tribolium* larva, being 17.92μ

mol/g and it occupies about 23.2 % of the total amino acids, and that of *Galleria* larva is only 5.72 μ mol/g and 18.7 %. T/G value of these amino acids always exceeds 2 except glycine, but 1.66 even in glycine. The importance of amount of dispensable amino acids as the supplement to the essential ones for normal growth, development or egg-production is postulated by several workers (Ishii and Hirano 1955, Lea et al. 1956). For instance, alanine, glycine, serine and tyrosine in *A. affinis*, and hydroxyproline, proline and serine in *A. aegypti* are not essential for the growth and development, but a significant retardation occurs when any one of them is lacking (House 1954, Sing et al. 1957).

Some qualitative differences are also observed between these 2 samples which contain such unknown substances 2, 4, 5, 6 and citrulline (?). But the discussion of these substances upon the fecundity is of course impossible unless the nature of these substances are not identified. The presence of citrulline in the body fluid was previously reported by Chu (1962a, b) in the *Papilio* larvae, but the physiological function of this amino acid is still unknown. However, Hinton et al. (1951) and DeGroot (1953) reported the partly replacement to arginine, an essential amino acid in *Drosophila melanogaster* Meigen and *Apis mellifera* Linné respectively.

The amount of ninhydrin-positive substances, total amino acids and essential amino acids of *L. beneficus* are surely lower than those of the prey. There are little qualitative variations on the ninhydrin-positive substances of *L. beneficus* reared on different prey. The only significant qualitative difference is the presence of citrulline (?) in the *Galleria*-fed bug and this amino acid is only detected in the *Galleria* larva but not in *T. castaneum*. The apparent quantitative variations of amino acids are rather observed in some components, for instance ; amounts of taurin (?), glutamic acid, methionine, lysine and arginine. And such components are surely prevalent in the *Tribolium*-fed individuals, but a reverse tendency is seen in proline, tyrosine and histidine. Generally speaking, the composition of amino acids in predators is not decidedly influenced by food both in qualitatively and quantitatively. While the discrimination of amino acid amount between the prey and predator is seemed to be an important key to solve this problem. That is, for example, in the *Tribolium*-fed bug can convert 15.62 μ mol. of alanine but only 3.54 μ mol. in the *Galleria*-fed individuals. A similar tendency is observed in all identified amino acids including 9 essential amino acids. The value of T-TL/G-GL are evidently significant both in the amount of ninhydrin-positive substances and total amino acids which are about 3.0 and 3.3 respectively. While the ratio, 4.8, is more significant when the amount of essential amino acids is compared. This may indicate that the *Tribolium*-fed bugs are afforded with about 3 times of amino nitrogen than the *Galleria*-fed individuals, and the value of T-TL/G-GL raises up to 4.8 times when the essential amino acids are only compared. There-

fore, in the *Tribolium*-fed bugs, they are able to convert more essential nitrogen source for reproductive use. And it is worth to notice that arginine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophane and valine were already proved as the essential amino acids for the egg-production of certain insects (Greenberg 1951, Dimond. et al. 1956, Lea et al. 1956).

At the moment we have few available data relating to the function of individual amino acid on the insect fecundity. It is difficult to continue the present discussion. To make a more accurate study on this point, it is at least indispensable to rear bugs with certain amino acid deficient diet and compare its fecundity. Moreover, the nutritional factor affecting the insect fecundity is not restricted to amino acid only. To clear this problem, further studies should be conducted with other nutrients and by another biochemical method.

For the next step, some works on the lipid and its component should be referred. Because it is a general conception that the *Galleria* larva is more abundant in lipid content than the *Tribolium* larva. In addition, some evidences of the importance of certain fatty acids in the nutrition of some insect species have been reported on *Argyrotaenia velutinana* (Walker), *Carpocapsa pomonella* (Linné), and *Anthonomus grandis* Boheman (Rock et al. 1965, Rock 1967, Earle et al. 1967). Also a review on the nutritional requirement by insects is compiled by Fast (1964). On the other hand, the nutritional invalidity of fatty acids on the larvae of *Chilo suppressalis* is also reported by Hirano (1963), and he also indicated the growth inhibiting activity of these compounds at higher dosages.

The total lipid content of the larvae of *G. mellonella* and *T. castaneum* were extracted by the Soxhlet apparatus, almost following the initial steps of the procedure of Ascher and Neri (1961) : The larvae were ground in a mortar with twice their weight of anhydrous Na_2SO_4 , the mixture was left for one hour in a covered mortar and was then transferred quantitatively into a filter paper cylinder for Soxhlet extraction. The larvae were first extracted with ether for 15 hours, after which the vessel with ether was exchanged for one with acetone. Re-extraction of the package with acetone was then continued for another 10 hours. The greater part of the solvent was distilled off on the water bath, then dried in vacuum exsiccators over CaCl_2 and further continued to dry until the solid paraffin wax weight became constant.

The result of extraction shows significant abundance of lipid in the *Galleria* larvae, being 0.22 g per 1 g of wet material and that of the *Tribolium* larvae was only 0.04 g. The nature of the extracts are also very different, the sample from *Galleria* is bright yellow coloured and that of *Tribolium* is dark brown green.

Paper chromatographic separation was employed to separate some components on the hexane solution of these ether-acetone extracts. The separa-

tion was carried out with Tōyō No. 51 filter paper in ascending method. The mixture of white kerosene and xylene (4 : 1 v/v) was used as a solvent. When the solvent had travelled a distant of 35 cm from the spotted point, the chromatogram was removed from the chamber. The colour development procedure was followed by Viswanathan and Meera (1962) : Thus, the chromatogram was at first washed twice in tap water to remove the solvent. It was then kept immersed for 15 min. in a 0.1 % mercuric acetate solution containing 0.5 ml of acetic acid per litre. Excess mercuric acetate was removed by washing the chromatogram in removing tap water for 45 min. The chromatogram was air dried, then sprayed with 0.2 % solution of S-diphenyl carbazide in 95 % ethanol.

Fig. 8. Diagram showing chromatographic patterns produced by paper chromatography of lipid content of ether-acetone extract of *Tribolium* larvae (T) and *Galleria* larvae (G).

Although the separation was almost failed, the long purple coloured bands were observed in both samples. Then the top of the band in the *Tribolium* larvae was extended to the point of R_f 0.86 and that of *Galleria* was ceased at 0.78. In the *Tribolium* sample, 4 significant purple colour concentrations were recognized and they were placed at the points of 0.92, 0.75, 0.55 and 0.42 of R_f value. In another sample, only 3 spots appeared and their R_f value were 0.73, 0.62 and 0.42 (Fig. 8). As the inferiority of chromatographic separation, no further work on the identification was made on these samples. But it is apparent that the composition of ether acetone extracts is different from each other.

The studies on the lipid and fatty acid in the *Galleria* larvae was conducted by Young (1964, 1967). And his latter paper identified 6 fatty acids from the chloroform-methanol extract of the larvae (Table 50). The result of his analysis indicates the abundance of 2 fatty acids of 16 carbons and 18 carbons with 1 double bond. For the former compound,

Table 50. Fatty acids of larvae of *Galleria mellonella* (After Young 1967)

No. of carbon in fatty acid	14	16	16 ¹⁼	18	18 ¹⁼	18 ²⁼
percent by weight	trace	39.6	2.9	3.1	47.2	6.5

palmitic acid is corresponded. And oleic and elaidic acids etc. belong to the latter group (Table 49).

It is a noticeable fact that the growth of *Grylodes sigillatus* (Walker) is inhibited by the existence of some fatty acids and their methyl ester, including the methyl ester of oleic acid and palmitic acid. Above all, inhibiting action is very significant in methyl palmitate (McFarlane and Henneberry 1965). Although the inhibiting action was tested with the use of a compound absorbed filter paper in a cricket rearing cage in their experiment, and not by feeding test, it is possibly conceivable that these fatty acids also show some inhibiting actions when they are taken into the digestive organs of insects. However, a question still exists that whether these fatty acids show any growth inhibition activity on the growth of Anthocorid bugs as in the case of *G. sigillatus*. In addition, Hirano (1963) indicates the inhibiting action for the growth of rice stem borer of oleic acid and linoleic acid ($C^{2=}$), in the presence of higher dosages in the diet of the borer. Because of the lack of an available datum, the author can not make any discussion on the fatty acid of the *Tribolium* larvae.

Rearing with artificial media

The review of previous works reveals that the rearing of suctorial insects with artificial diets was already established in some aphids, leafhoppers and Heteropterous bugs. The first work on the Heteropterous bug was attempted by Scheel et al. (1957) on *Oncopeltus fasciatus* (Dallas) and *Euschistus variolarius* (Palisot de Beauvois). Since then, the works on these bugs were also made by Beck et al. (1958) and Scheel et al. (1958). The studies on another Bug, *Lygus hesperus* Knight, have been published by Lands and Strong (1965), Auclair and Raulston (1966) and Vanderzant (1967), and the last author succeeded to rear both *L. hesperus* and *L. lineolaris* (Palisot de Beauvois) with artificial diet up to the stage of oviposition. Anyway, such works have been confined chiefly to the phytophagous bugs. On the other hands, Bronnimann (1964) succeeded the rearing of *Ranthocoris nemorum* (Linné) and *A. nemoralis* (Fabricius) from the first instar to the adult stage with sweetened condensed milk. However, these Anthocorids failed to produce eggs unless some animal diets were not supplied in their adult stage. This may be the only work on the rearing of predacious bug with artificial media. The same trials on the other groups of predacious insects have been conducted on *Chrysopa carnea* Stephen (Hagen and Tassen 1965, 1966, Sundby 1967) and Coccinellid beetles (Smirnoff 1958, Smith 1960, Tanaka and Maeta 1965, Atallah and Newson 1966). Although the oviposition of *Chrysopa* female which was reared with artificial diet was observed without exception, the addition of animal diets was almost an indispensable procedure to the majority of Coccinellid beetles. Otherwise, the beetles did not lay their eggs.

Presumably the only successful trial on artificial diets was obtained with *Coleomegilla maculata* DeGeer by Atallah and Newson (1966) who added some liver fractions and the extract of cotton leaves into the media. But in some other species of lady beetles the oviposition did not occur when fed with this medium.

The study on the rearing of *Lyctocoris beneficus* (Hiura) and *Xylocoris galatinus* (Fieber) with artificial media was at first begun with the selection of suitable rearing methods and the principal chemical components of media solution. As to the first point, 4 following methods were preliminary compared.

1) *Bronnimann's straw method* (Bronnimann 1964) : That is, a thin twist of

Fig. 9. Rearing apparatus for paraffin film method.

A : glass vial (media container).

B : stretched parafilm.

C : polyvinyl tube.

D : glass tube (bug adaptor).

E : organdy cloth.

cotton wool dipped in the media was drawn through a plastic drinking straw by means of a fine wire. The straw containing the media was then cut into pieces of certain length after the refrigeration to solid.

2) *Hagen's paraffin-covered droplet method* (Hagen and Tassen 1965) : At first 40 ml. of media solution was mixed with 0.2 g paraffin (mp. 45°C) in a 50 ml beaker and the mixture was heated to about 60°C, then the paraffin forms thin layer on the surface of the solution. Droplets were made by dipping into the warm mixture with a glass applicator rod which was lifted out and gently touched to a piece of parafilm.

3) *Ouchi's paraffin method* (Ouchi 1967) : The structure of the apparatus in this method is illustrated in Fig. 9, that is mainly composed of a glass tube and a glass vial both of which are 1 × 3 cm in diameter and height respectively. The tube was used for the bug adaptor, and

its bottom was covered with a sheet of organdy cloth. The cloth side of vial was used for the container of media solution and the opening was covered with a stretched parafilm after the solution was provided. Then the parafilm covered opening of the vial and the uncovered end of the tube were connected with a piece of polyvinyl tube. Thus the bugs in the adaptor can take the media solution by piercing the film with proboscis.

4) *Uretane foam method*: This is simply prepared by the soaking of a piece of urethane foam which was cut to about 1 cm³ with a media solution.

For the media solution, 4 following formulations were preliminary compared.

Diet A. (after Bronnimann 1964)

sweetened condensed milk+H₂O (2 : 1 v/v)

Diet B. (after Hag-en and Tassen 1965)

enzymatic protein hydrolysate of yeast	10.0 g
enzymatic protein hydrolysate of casein	10.0 g
fructose	17.5 g
ascorbic acid	2.0 g
choline chloride	25.0 mg
water	100 ml

Diet C. (modified Beck et al. 1958)

soluble starch	2.0 g
dextrose	2.0 g
sodium caseinate	2.0 g
beef extract	2.0 g
Brewer's yeast powder	4.0 g
cholesterol acetate	0.1 g
Wassen's salt	0.1 g
water	100 ml

Diet D. (modified Scheel et al. 1957)

dextrose	2.0 g
soluble starch	2.0 g
sodium caseinate	2.0 g
Brewer's yeast powder	2.0 g
beef extract	0.5 g
Wassen's salt	0.2 g
cholesterol	80.0 mg
water	100 ml

To select the suitable methods and formulations, 10 first instar *X. galactinus* were reared separately by the above mentioned methods and with the above cited formulations, and the longevity of nymphs was recorded for 10 days (Table 51). In the rearing experiments 1, 2 and 4, all nymphs were reared in a flat-bottomed glass tube, sized 2.5 × 8 cm (diameter × height), and the tube was covered with cork stopper. The medium was renewed every day.

As seen in Table 51, it is apparent that the fourth method, namely the urethane foam method, was the best and the parafilm method followed. In the latter method, the death of majority of nymphs was caused by the drowning in the solution flown from the leakage of parafilm which occurred most frequently when the diluted milk medium was provided.

Table 51. Longevity of the first instar *Xylocoris galactinus* reared with 4 kinds of artificial diets by 4 kinds of rearing methods (started with 10 individuals).

Rearing method	Diet No.	No. of nymphs survived in days									
		1	2	3	4	5	6	7	8	9	10
1	A	6	4	3	2	0	-	-	-	-	-
	B	7	2	2	0	-	-	-	-	-	-
	C	5	3	2	2	0	-	-	-	-	-
	D	6	4	3	1	0	-	-	-	-	-
2	A	5	3	2	0	-	-	-	-	-	-
	B	6	4	3	3	1	0	-	-	-	-
	C	7	2	1	0	-	-	-	-	-	-
	D	5	1	1	1	1	0	-	-	-	-
3	A	5	4	3	3	3	3	3	3	3	2
	B	7	6	6	6	6	5	5	5	5	5
	C	6	6	6	5	5	5	5	5	4	4
	D	8	7	7	6	6	6	5	5	5	5
4	A	10	10	9	9	7	7	7	7	7	7
	B	9	9	9	8	8	8	8	8	8	8
	C	10	10	8	8	7	5	4	4	4	3
	D	10	10	10	9	9	8	8	7	7	7

Although parafilm method is surely a time-consuming work, it may become one of the ideal technique to evaluate the nutritional requirement of bugs when membrane is improved to a more tenacious material. The lowest mortality obtained in the urethane foam method was partly due to the contamination of mold on the wall of glass tube. Because, as previously mentioned, the bugs are facultatively fungivorous, and undoubtedly the fungus may play some nutritional role to some extents on the development of bugs. However, it is not a suitable method to study nutritional requirement of bugs because of the heavy fungi-infection, but may be used as the most effective method for the practical rearing. Both the first and second rearing methods were not advisable for this bug. In the first method, the cotton thread entangled the legs of bugs and this caused the death of all the nymphs. In the second method, high mortality is assumed to be caused by the difficulty of sucking solution through a paraffin membrane. As to the formulation of media, diet C was somewhat inferior, at least on the survival of the nymph, and no significant superiority was observed among 3 tested formulations. Therefore, the further work was concentrated to the urethane foam method with diets A, B and D on both species of bugs.

Rearing results of newly hatched nymphs and the third instars of *L. beneficus* and *X. galactinus* with the formulations A, B and D were presented in Table 52. In this experiment, the nymphs were reared separately in a flat-bottomed glass tube as in the case of the preceding experiment.

Table 52. Growth of *Lyctocoris beneficus* and *Xylocoris galactinus* fed with artificial diets (25 individuals per each test).

Diet No.	Instar set	No. of nymphs to adult		Avg days to adult	
		<i>L. beneficus</i>	<i>X. galactinus</i>	<i>L. beneficus</i>	<i>X. galactinus</i>
A	1st	9	5	32.4	22.4
	3rd	21	19	23.7	13.4
B	1st	7	7	33.8	24.8
	3rd	19	19	24.1	14.7
D	1st	12	15	30.9	19.3
	3rd	23	22	21.2	14.2
CK*	1st	18	21	26.3	16.2
	3rd	24	25	19.1	12.4

* reared with larvae of *Tribolium castaneum*.

It is evident that the diet D was the best one tested so far as the longevity and developmental rate of both species were concerned, and diet B showed the worst result. The rearing of the first instar nymphs was much inferior both in longevity and developmental rate to that of the rearing from the third instar. In the rearing of first instar nymphs, almost all the tested individuals died in the fourth or fifth instar. If comparison was made on the development between the nymphs which were reared with artificial media and fed with animal diets, the prolongation of nymphal stage in the former group was exclusively pronounced during the fifth instar. The rearing of nymphs with artificial media from the third instar was almost successful on all diets. However, some delay of development was observed. Although no detailed data were available, the addition of animal prey to the third instar nymphs which had been reared with artificial media from the first instar was very helpful to accomplish the normal development in the later nymphal stage. The copulation was normally observed in both species when they were reared under gregarious condition, but no eggs were deposited at all.

To examine the nutritional value of artificial media on adult bugs, 2 day old fertile females which were fed with animal prey during their nymphal stages were reared with diets A, B and D. And the pre-oviposition period and the number of oviposited eggs were recorded (Table 53). However, some individuals did not deposit any eggs at all. The minimum value of pre-oviposition period was given in the same table.

As seen in Table 53, the adult bugs laid eggs except the bug group of *L. beneficus* which was fed with diet B. But the egg production was evidently inferior to groups which were fed with animal diets. Although diet D showed the best result in both species among the 3 formulations tested, the egg production ability was less than one-third of the bug fed with animal

Table 53. Number of eggs laid by a female *L. beneficus* or *X. galactinus* reared with artificial media in her adult stage.

Bug	Diet	Pre-oviposition period(in days)*(min-aver-max)	No. of eggs/female (min-aver-max)
<i>L. beneficus</i>	A	8	0-4.2-10
	B	—	0
	D	9	0-12.3-24
	CK	8	1.2-38.7-52
<i>X. galactinus</i>	A	4	0-3.4-8
	B	5	0-0.3-2
	D	5	8-14.4-25
	c K	4	12-48.3-67

* minimum value.

diets. Although no difference was observed on the length of pre-oviposition period between the individuals fed with animal diets and artificial media, the oviposition period was much shorter in the individual groups fed with artificial media, and even the maximum oviposition period lasted for 3 days. On the contrary, in the individual groups fed with animal food, the oviposition period lasted for more than 20 days in both species.

As Tables 52 and 53 indicate, diet D is the most promising formulation among the 4 tested media. As the next step, the improvement of dietal component was tried with this medium. In Table 54, the modification of diet D was given. In this case, the addition of enzymatic protein hydrolysate of casein, ascorbic acid, choline chloride and cholesterol acetate to the diet seemed prospective, but the addition of these materials to the formulation caused precipitation.

The comparison of rearing experiments with these diets was conducted by using the first and third instar *L. beneficus* and *X. galactinus* (Table 55).

Table 54. Improved diets of D for *Lyctocoris beneficus* and *Xylocoris galactinus* (in g/100 ml water).

	1*		Diet No.			
	1	2	3	4	5	6
Dextrose	2	2	2	2	2	2
Soluble starch	2	2	2	2	2	2
Sodium caseinate	2	2	2	2	2	2
Beef extract	0.5	1	1.5	3	1	1
Cholesterol	0.08	0.08	0.08	0.08	0.08	0.08
Wassen's salt	0.2	0.2	0.2	0.2	0.2	0.2
Brewer's yeast powder	2	2	2	2.5	3.0	3.5

* original formulation of diet D.

Table 55. Growth of *Lyctocoris beneficus* and *Xylocoris galactinus* fed with improved diet D.

Diet No.	Instar set	Percent emergence of adult(%)		Avg no. of days to adult	
		<i>L. beneficus</i>	<i>X. galactinus</i>	<i>L. beneficus</i>	<i>X. galactinus</i>
D-1	1st	28	24	31.5	21.8
	3rd	64	64	21.2	14.0
D-2	1st	56	36	28.4	17.8
	3rd	72	72	20.4	12.8
D-3	1st	24	16	32.4	19.2
	3rd	60	64	22.8	13.2
D-4	1st	36	24	33.6	21.8
	3rd	64	56	21.3	13.0
D-5	1st	44	28	29.4	21.3
	3rd	56	60	22.6	14.8
D-6	1st	28	20	31.0	22.4
	3rd	58	68	22.8	15.6

Although the results showed no noticeable difference among the diets, a slight improvement was observed in diet D-2. But all of these diets were not effective to promote the reproductive potential of adults.

The analysis of amino acids in the larvae of *Tribolium castaneum* (Herbst) and *Galleria mellonella* (Linné) suggests the importance of amino acid amount in food for the egg production of bugs. In the experiments shown above, the protein emphasized media, namely diets D-2 and D-3, were scarcely effective in the promotion of both growth and fecundity, and even the rearing result of D-4 was inferior to that of D-1. Presumably it is partly ascribed to the difficulty in sucking the media solution by nymphs due to the high viscosity of media which occurred by the excess amounts of beef extracts.

The prevention of cannibalism is another important factor to decide the practical value of artificial media especially in the mass rearing of predatory insects. Although cannibalistic behaviour was rarely seen in *L. beneficus* and *X. galactinus* in the presence of sufficient animal food, the occurrence of cannibalism was already demonstrated only in the case of deficiency of animal diets (cf. phytophagous habit). To examine the cannibalism in the rearing condition of artificial media, 10 of the third instar nymphs were reared gregariously in a 50 ml beaker provided with artificial media and the number of survived bugs was recorded (Table 56).

The results of experiments showed very violent cannibalism between the individuals, and in most cases, about half of tested individuals were annihilated within 2 days. After 10 days, generally only a single bug survived. And in almost all cases the cannibalistic phenomenon occurred during the moulting process or just after the ecdysis when the integument

Table 56. Survival of *Lytocoris beneficus* and *Xylocoris galactinus* reared gregariously with artificial media (started with 10 3rd instar nymphs, average of 5 tests).

	Diet No.	No. of survived bugs after days				
		2	4	6	8	10
<i>L. beneficus</i>	D-1	3.6	2.2	1.6	1.4	1.2
	D-2	5.0	3.2	2.6	2.2	1.4
	D-6	3.4	2.8	2.0	2.0	0.8
<i>X. galactinus</i>	D-1	5.0	3.8	1.8	1.6	1.2
	D-2	5.2	4.4	3.0	2.2	1.6
	D-6	4.8	4.4	2.2	1.8	1.0

of body was not completely sclerotized. Although the diet D-2 was very slightly preferable, it was still not practical for the mass rearing of bugs.

The addition of vitamin E (alpha-tocopherol) into the diet of some insect artificial media was attempted by several workers (Fraenkel and Blewett 1946, Beck et al. 1949, Vanderzant 1957, etc.). But according to several workers, its nutritional requirement in insects has not yet been recognized (Trager 1953, House 1962, Gilmour 1961, Dadd 1963, etc.). On the other hand, the improvement of the growth, especially of reproductive potential, has been proved in some insects such as : *Cryptolaemus montrouzieri* Mulsant, *Achetadomesticus* (Linné), *Agria affinis* (Fall&), *Exeristes comstockii* (Cresson), etc. (Chumakova 1962, Meikle and McFarlane 1965, House 1966, Bracken 1966 respectively). As seen in the above works, the rearing of bugs with an artificial medium is not superior to that with animal diets both in growth and survival, and this defect is especially significant because of the decrease of reproductive potential. To compensate this defect the addition of vitamin E was attempted to the artificial media. Thus, 0.5, 1.2 and 4 mg of vitamin E were dissolved in small amount of ethyl ether and added to the 100 ml of D-2 artificial medium. The rearing test was started with 10 third instar nymphs of both bugs which were kept in a 50 ml beaker, and the result of survival is presented in Table 57. The experiments were repeated 5 times.

As illustrated in Table 57, an increase in survived individuals was obtained in the media which contained 0.5 to 2.0 mg per 100 ml media. Accordingly, in the medium devoid of vitamin, only a single bug generally survived on the 10th day of rearing, while in the media added with vitamin, in most cases, 2 to 3 bugs still co-existed for a certain period of time. The excess amount of vitamin gave lower survival rate than the vitamin omitted media. The longevity was also prolonged in the bugs which were fed with media added with vitamin moderately. The vitamin content was seemingly more suitable in the concentration of 1.0 and 0.5mg per 100 ml of media solution in *L.*

Table 57. Survival of *Lyctocoris beneficus* and *Xylocoris galactinus* reared gregariously with vitamin E added artificial media (started with 10 3rd instar nymphs, average of 5 tests).

	Content of vitamin E (mg/100 ml)	No. of survived bugs after days					Longevity (in days)*
		2	4	6	8	10	
<i>L. beneficus</i>	0.0	4.2	3.2	2.4	2.0	1.2	15.0
	0.5	7.2	6.4	4.8	3.6	2.6	23.4
	1.0	6.6	5.4	3.8	3.2	2.8	24.6
	2.0	6.4	5.8	4.2	2.0	1.4	20.6
	4.0	5.4	3.4	2.6	1.8	1.2	13.8
<i>X. galactinus</i>	0.0	5.2	3.6	2.2	1.4	1.4	12.8
	0.5	7.4	6.8	6.4	4.4	3.2	22.0
	1.0	7.8	5.6	5.0	3.6	2.8	20.8
	2.0	7.4	6.2	4.8	2.8	2.2	17.6
	4.0	6.4	4.4	3.2	1.8	1.2	14.2

* average of maximum value from each test group.

beneficus and *X. galactinus* respectively. Although the copulation was observed between the emerged adults but no egg was oviposited at all. From this fact it is apparent that vitamin E is not responsible for the egg production of *L. beneficus*. and *X. galactinus*. But a significant decrease in cannibalism was observed.

The result of the present experiments shows that some artificial media permit the restricted growth of bugs and the media of D-2 containing 0.5-1.0 mg/100 ml are most prospective. But compared with the rearing result with animal diet, the rearing with artificial media is certainly inferior both in nutritional value and taste to the animal. diet.

Ovarian development under certain nutritional conditions

It is well known that insect reproduction is affected by many factors. Among them, nutrition is one of the most influential factors. And both qualitative and quantitative nutritional conditions give decided influences on the fecundity (Rasso and Fraenkel 1954, Harlow 1956, Wigglesworth 1960, Orr 1964, etc.). In *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber), higher fecundities were obtained when these bugs were reared with the larvae of *Tribolium castaneum* (Herbst). While the bugs which were fed with the larvae of *Chilo suppressalis* (Walker) or *Galleria mellonella* (Linné) as food showed lower fecundities (cf. Chap. 9). Moreover, the bugs reared under the animal diet difficient condition scarcely laid eggs (cf. Chap. 8). From these facts, it is suggested that the quality of foods supplied influences the ovarian development of the bugs. In the individuals of both species which were reared with the larvae of *T. castaneum*, the adult female frequently

has a well-swollen abdomen. And sometimes, the abdomen becomes enlarged that the wings are covering only the four basal abdominal segments. However, in the newly emerged female or mature male the wings are covering the abdomen entirely. On the contrary, in the individuals which were fed with plant substances, the abdomen is always dwindled, and its length has few differentiations from that of male.

In the present work, the newly emerged adults which were fed from their 1st nymphal stadium with the larvae of *T. castaneum* were reared with the larvae of *T. castaneum*, *C. suppressalis*, *G. mellonella* or molded cracked corn. And the ovarian development was observed by dissecting the abdomen of 1, 3 and 5 day old adult females. About 5 adults were dissected for each test and the number of eggs was counted and classified into three ranks according to their size (Tables 58 & 59).

The development of ovarioles are also illustrated in Figures 10 & 12. In this experiment all the bugs were reared under 30°C condition. The pre-oviposition period of *L. beneficus* and *X. galactinus* under this condition was about 7 and 5 days respectively.

The reproductive organs of both species consist of one common oviduct, a pair of oviducts and ovaries. A oviduct is connected with 7 ovarioles. In the pre-mature adult, the terminal filament is extended to the mesothorax and the ends of the filaments are aggregated to one another. When the ovary is matured, the filaments bend posteriorly and their ends separate from each other. The filament is apparent in the pre-mature individuals, but shortened in the mature ones. The development of each ovariole in an

Table 58. Number and size of eggs in the ovariole of *Lyctocoris beneficus* reared with various foods.

Days after emergence	Food	Size of egg in ovariole (mm)		
		1.00<	1.00-0.70	(0.70
1	ml	0.0	1.4	4.0
	C	0.0	0.6	2.8
	G	0.0	0.4	2.4
	P	0.0	1.0	3.2
3	ml	0.0	6.8	14.4
	c	0.0	4.6	7.8
	G	0.0	3.2	4.4
	P	0.0	0.2	1.2
5	T	8.6	17.9	11.8
	C	3.8	3.2	5.4
	G	2.2	4.0	4.6
	P	0.0	0.0	0.8

T: larvae of *Tribolium castaneum*.

C: larvae of *Chilo suppressalis*.

G: larvae of *Galleria mellonella*.

P: molded cracked corn.

Table 59. Number and size of eggs in the ovariole of *Xylocoris galactinus* reared with various foods.

Days after emergence	Food	Size of egg in ovariole (mm)		
		0.6<	0.60~0.40	<0.40
1	T	0.0	12.4	8.8
	C	0.0	8.8	6.0
	G	0.0	7.2	12.4
	P	0.0	5.4	8.2
3	T	13.2	12.0	9.6
	C	1.2	8.6	7.4
	G	0.0	7.8	6.4
	P	0.0	4.6	2.0
5	T	13.8	18.2	12.6
	c	9.8	4.2	7.4
	G	2.4	4.8	6.0
	P	0.0	4.2	4.2

T: larvae of *Tribolium castaneum*.

C: larvae of *Chilo suppressalis*.

G: larvae of *Galleria mellonella*.

P: molded cracked corn.

individual is almost uniform, but the individual variation is certainly significant. The ovariole is generally transparent, but partly whitish opaque. The egg in the ovariole is white in colour, and the shape is globular at first and becomes oblong later. When the egg is fully matured, it becomes kidney-shaped and provides with a marked operculum which is easily detectable by a slight brown ring. The size of ovary is very variable with the age, nutritional condition etc., in the fully matured individuals it is about 3.2 mm in *L. beneficus* and 2.4 mm in *X. galactinus*.

The newly emerged adults of both species have small ovaries with slightly developed germarium and the abdominal coelum is filled with fat body. In *Tribolium*-fed *X. galactinus*, the significant ovarian development is recognized within one day rearing. On the third day, several mature eggs each with a distinct operculum appeared. While through the author's experiment, no example was obtained as to the production of an egg after 3 day pre-oviposition period. On the fifth day slight numerical increment on mature eggs was observed, however, a marked change was characterized rather by the development of pre-mature eggs. In the individuals reared with the larvae of *C. suppressalis* and *G. mellonella*, the appearance of mature eggs were surely retarded than *T. castaneum* reared individuals, and mature eggs were scarcely recognized in 3 day old adults. The number of pre-mature eggs was also less than that of *Tribolium* reared bugs. In the individuals which were fed with only plant-substances after emergence, some retardation on the ovarian development was already observable within 24 hours. While the differentiation became more significant in the later stage. The especially noticeable fact

Fig. 10. Ovarial development of *Lyctocoris beneficus* reared with various foods.

T: reared with larvae of *Tribolium castaneum*. C: reared with larvae of *Chilo suppressalis*. G: reared with larvae of *Galleria mellonella*. P: reared with molded cracked corn. Figure after hyphen indicates the day after emergence.

was that till the third day slight ovarian development was still seen, but on the fifth day the ovary rather shrinkled. And in this group, of course, no maturation of eggs was observed at all. This phenomenon postulates the indispensability of the animal diet on the ovarian development of these Anthocorid bugs. The shrinkage of the ovary in the individuals fed with plant-substances was also observed in *L. beneficus*.

In *L. beneficus*, the ovarian development was apparently slow than that of *X. galactinus*, and in the *Tribolium*-fed bugs the mature egg appeared on the fifth day. The effect of other foods on the ovarian development was very similar to that of *X. galactinus*.

In the fully matured female the number of mature eggs in the ovary was unexpectedly high, and the maximum figure was 42 in *L. beneficus* and 35 in *X. galactinus*. In such individuals, of course, the oviposition takes place, but they do not deposit all of the mature eggs at once. The plumpness of abdomen continues almost through her life, and the female frequently dies with many mature eggs left in the ovary.

Fig. 11. Ovarial development of *Xylocoris galactinus* reared with various foods.

Effects of population density on the survival, growth and fecundity

The effects of population density on the survival, growth and fecundity of many species of animals greatly depend on species and developmental stage (Allee 1934). When the population density increases, food, available space, sanitation, cannibalism and certain other factors increase their importance. Among insects, population growth has been shown to be influenced by the contracting influences of mating and cannibalism or predatism (especially of the eggs). Further the cannibalism or predatism is more prevalent in the denser population (Park 1932, Chapman 1928, Chapman and Whang 1934, Maclagen 1932, Boyce 1946, Park et al. 1965). Generally, crowding is recognized to be unfavourable to the growth of some cockroaches, *Tenebrio* beetle, etc., and the bad effects caused by crowding on animals are attributed predominantly to physical rather than chemical factors (Allee 1934, Chauvin 1946), and to chemical influences especially of specific pheromones by some workers (Butler 1954, Norris 1954, Lüscher 1956, Loher 1961). Prior to start their practical mass production, it is an important problem to decide the most suitable density on the growth process and fecundity of *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber). The author reared various

developmental stages of both bugs under various population conditions and observed the effects on their development and offspring production.

In this experiment the laboratory reared individuals were used, and they were bred in a plastic container which was 12 cm in length, width and height each, or some different sized containers and always supplied with sufficient *Tribolium* larvae as food. Although the density of the prey also may influence the growth of predator, this factor was not taken into consideration in the present work. The population counting was carried out every three days, and the detailed experimental procedures are described in each experiment.

I. Effect of population density on the survival of adults.

Various numbers of newly emerged *L. beneficus* and *X. galactinus* adults were kept in a plastic container which was provided with molded or dry cracked corn about 1 cm thick and sufficient numbers of *Tribolium* larvae. The survival of adult bugs was counted every 3 days (Table 60).

Table 60. Survival of *Lyctocoris beneficus* and *Xylocoris galactinus* adults under various population densities.

No. of adults set per container(A)		Condi- tion of corn	No. of survived adults after 9 days								$\frac{B+C}{A} \times 100(\%)$	
			--		in media(B)		on cover(C)		B+C			
			L.b.	X.g.	L.b.	X.g.	L.b.	X.g.	L.b.	X.g.		
50	dry	44	45	1	0	45	45	90.0	90.0			
	mold	48	46	0	0	48	46	96.0	98.0			
100	dry	71	77	4	5	75	82	75.0	82.0			
	mold	80	83	3	3	82	86	82.0	86.0			
150	dry	71	114	3	6	74	120	49.3	80.0			
	mold	101	120	21	12	123	132	82.0	88.0			
200	dry	66	119	5	6	71	125	35.5	62.5			
	mold	112	153	44	18	156	171	78.0	85.5			

During this test, there was observed not a few individuals of *L. beneficus* on the cover of the container especially in denser groups. The result of marking test with diluted white poster colour on their pronotum indicated that the majority of individuals found on the cover were inactive for a long time. Therefore, these adults were supposed to be expelled individuals from the media by the density pressure. Consequently, they are counted in the other column in the Table. It is worth to notice that when the medium was over moistened, more individuals climbed up to the cover, while in such a medium the survival was, on the contrary, higher in both species. This is considered as the results of broadening of their territories in the

moist condition. The higher rate of survival was obtained in the moist corn medium, while in the excessively moist medium the mortality was apparently higher than that of the dry medium. The cover staying individuals were scarcely observed in *X. galactinus* which has much smaller body size and was generally living aggressively under more moist condition in the field. Though the survival data shown in Table 60 are the result of 9th day, the significant decrease of population occurred during the first 3 days and further decrease of population was almost negligibile in the later six days. If the bugs staying on the cover are considered as individuals which are not participate in the reproduction of the species, the maximum supporting capacity of *L. beneficus* and *X. galactinus* in the dry corn medium was about 70 and 120 respectively, and that on the molded corn medium was 100 and 150 respectively. This fact is seemingly illustrating a conflict that the moist medium has larger capacity than the dried one. But this is rather explained by the less occurrence of cannibalism or predatism in the moderately molded medium. The bug supporting capacity of medium has such a limit that even the addition of more medium material does not increase the supporting capacity of the given container.

II. Effects of population density on the egg production and hatching rate.

The fecundity, or oviposition rate of many animals shows a sensitivity to the population density (Norris 1950, 1952, 1959, Albrecht et al. 1958, Zaher 1959, Watt 1960, Green 1954). Copulation occurs most frequently in the moderately dense population which permits chance to mate and produces more fertile eggs (Pearl et al. 1922, 1927, Pearl 1932). To examine the density effect on the fecundity of the bugs, 1, 2, 5, 10 and 20 pairs of adults of 8 to 10 day old *L. beneficus* and 6-8 day old *X. galactinus* were reared separately in a 6×9 cm (height x diameter) petri-dish which was provided with 1 cm depth of molded corn, namely about 64 cm^3 in volume that was approximately half the volume of corn medium of the preceding test, namely 144 cm^3 . Therefore, so far as the corn medium volume was concerned, 20 pairs per dish corresponded 80 adults in the plastic container. In this test, the adults deposited all the eggs in the corn medium, because no moist filter paper was provided. As it was difficult to count the number of eggs laid in corn, the corn was renewed every three days, and the number of newly hatched nymphs was counted as an index of fecundity, and the average of 5 replications are shown in Table 61. In another series of test, the fertility of eggs was investigated by counting of the rate of hatching of eggs which were laid on the moist filter paper. Of course in such a test a part of eggs were laid in the corn medium. Therefore, only the ratio of fertilized and unfertilized eggs was obtained.

Though the percentage of fertile eggs and the number of eggs pierced by the bug in the corn medium are very difficult to count, the most ideal

case of the density was 10 pairs in *L. beneficus* and 20 pairs in *X. galactinus* to get the maximum offspring production, and bad-crowding effect was observed in the denser group of *L. beneficus*. The egg-fertility was not affected by the density. However, no detailed data were available. The predation of eggs by the adults was significantly more active in the denser groups, especially when the filter paper harbouring eggs were exposed to the adults for about a day.

Table 61. Number of hatched nymphs and fertility of eggs of *Lyctocoris beneficus* and *Xylocoris galactinus* under various population densities (average of 5 replicates).

No. of adults per dish (pairs)	Total nymphs hatched per day		No. of nymphs per female		Percent fertility(%)	
	L.b.	X.g.	L.b.	X.g.	L.b.	X.g.
1	2.6	3.2	2.6	3.2	88.1	90.3
2						
5	17.0 6.4	19.4 7.0	3.4 3.1	3.9 3.5	94.3 92.2	88.7 89.6
10	34.8	38.6	3.5	3.9	91.7	93.1
20	55.8	86.2	2.8	4.3	87.6	95.6

III. Effect of population density on the growth and survival of nymph.

Although crowding condition is unfavourable to the nymphal growth of many insects, the retardation of growth by isolated rearing is also observed (Michal 1931, Landowski 1937, Petit 1940, Willis et al. 1958, Feir 1963, Kunnel 1966). The density effect on the growth and survival of nymphs was investigated with a similarly provided petri-dish. Six petri-dishes each of which was provided with 1, 10, 20, 40, 60 and 80 newly hatched nymphs respectively were observed to detect the effect of crowding on the body weight and mortality (Table 62). To keep the young nymphs

Table 62. Effect of density on the nymphal growth of *Lyctocoris beneficus* and *Xylocoris galactinus* (average of 2-5 replicates),

Nymphs per dish	Avg. days to adult		Avg. weight of adult (mg)		Percent emergence(%)	
	L.b.	X.g.	L.b.	X.g.	L.b.	X.g.
1	25.2	15.8	2.3	1.4	77.4	87.8
10	24.4	15.5	2.3	1.4	78.6	91.4
20	24.6	15.4	2.4	1.5	72.4	87.4
40	23.9	15.1	2.4	1.5	74.6	78.0
60	23.3	14.8	2.3	1.4	51.2	76.2
80	23.4	14.6	2.3	1.4	48.7	75.6

from the mechanical damage to a minimum level, the dishes have never been disturbed until the emergence of new adults since the release of newly hatched nymphs. Of course, consideration was made in keeping the humidity and supply of food if necessary.

Primarily, on the basis of percent of survival, it is concluded that any number of nymphs of *L. beneficus* from 1 to 40 reared aggressively in a dish, and that of *X. galactinus* from 1 to 60 showed constantly high emergence ratio. When the density exceeded this range, the increase in nymphal mortality was observed in both species, and the offspring were smaller. But this was not due to the shortage of food, as the sufficient food was always supplied in the dish. The results would indicate that moderate gregarious rearing condition or moderate population density is preferable for the normal nymphal development of both species.

IV. Competition between adults and nymphs under the conditions of different densities.

During the process of practical mass production of the species, an adult-nymph complex may occur in the rearing medium. Therefore, the effect of adult population on the nymphal development seems to be an important factor to determine the efficiency of bug production.

In this test, the groups of 1, 5, 10 and 20 adults were placed separately in a 6 x 9 cm petri-dish which was previously provided with 25 newly hatched nymphs and 1 cm depth of molded corn. Then the survival and nymphal growth were observed. As the distinction between the old and newly emerged adult was difficult, the observation was stopped when about half of the tested nymphs became the fifth instar (Table 63).

Table 63. The effect of adult numbers on the nymphal growth of *Lyctocoris beneficus* and *Xylocoris galactinus*.

Initial no. of bugs per dish		Avg days to 5th instar		Nymphal survival — (%) —	
Adult	Nymph	Lb.	X.g.	Lb.	x.g.
0	25	17.3	10.4	78	82
1	25	17.3	10.5	78	80
5	25	17.2	10.8	82	78
10	25	17.5	10.7	72	84
20	25	16.9	11.2	46	76

The data in Table 63 indicate that the bad effect of mixed-rearing of both adults and nymphs on the nymphal growth occurs when 20 adult *L. beneficus* are used, but this is not the case in *X. galactinus*. The shortening of the duration of nymphal stages is observed in denser groups of both species.

In conclusion, to rear the bugs in a 6x 9 cm petri-dish efficiently, the mother strain is advisable to start with less than 20 or 40 adults of *L. beneficus* and *X. galactinus* respectively. The results of rearing experiments in the petri-dishes may be applicable to the rearing in the large plastic container in which it can accomodate about twice the number of individuals.

Overwintering and cold-hardiness

The Anthocorid bugs are generally known to overwinter in the adult stage, especially as fertilized females. They hibernate beneath tree barks, leaves, or litters on the ground. Though there are no detailed observations on the hibernation of *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber), the latter species is generally known from the warmer stock yard where usually few seasonal variations occurred (Hall 1951, Sinha 1961). Moreover, Hall pointed out that the bugs develop continuously throughout the year. Besides, some adult specimens of both species which were collected during the winter months were examined by Hiura (1957, 1966) in his taxonomic work. From this fact, it is evident that in the cold season, these species overwinter at least in the adult stage. Fortunately, the author could collect several adults of *L. beneficus* at the stock yard of a farm in Kyushu in January. During early spring, he also collected many specimens of the adult *X. galactinus* from the straw piles at Saga in Kyushu, and it may be mentioned that in both collections the author could not obtain any nymphs.

In the first experiment on the studies of the overwintering, the work on the cold-hardiness of these bugs was carried out. Various stages of bugs previously kept in 25°C condition were used in this experiment. During the time-temperature exposure, the bugs were kept in organdy clothed 100 ml beaker which was provided with a sheet of moist filter paper and enough prey. Then, they were exposed to the temperatures of -3°, +5° and +7°C for a certain period. The beakers were taken out from the cold chamber and placed under the room temperature condition. The mortality of bugs was estimated 24 hours after the exposure of the room temperature condition and the bugs which did not respond to the warmer temperature were counted as killed. All time-temperature exposure experiments were conducted with the use of bugs and repeated 5 times.

In the test under -3°C condition, all the stages of both species stopped their activities within 3 minutes, and exposure with such coldness for more than 30 minutes was extremely fatal to them, and 1 hour's treatment obtained 100 % mortality in both species. Under 5°C condition, they became inactive within 10 minutes. And within 1 hour almost all the individuals drew their legs and turned over, though few of them wriggled their legs feebly. The period needed to inactivate *L. beneficus* was seemed a little longer than in *X. galactinus*. In a short period treatment, these inacti-

vated individuals were readily recovered when they were removed to the room temperature condition. While in the longer duration of temperature treatment, the time necessary to recover was prolonged proportionally. When the duration of treatment exceeded a certain threshold, the bugs did not recover at all. The mortality after certain periods of exposure is indicated in Table 64.

Table 64. The mortality of *Lyctocoris beneficus* and *Xylocoris galactinus* under 5°C condition (%).

Anthocorid bug and stage	Exposure period (in days) and mortality (in numbers)					
	1	2	3	4	5	6
<i>L. beneficus</i>	1st instar	76	90	100	-	-
	3rd instar	32	18	45	92	100
	5th instar	38	24	24	32	86
	adult ♂	14	14	30	38	84
	♀	6	10	20	18	78
<i>X. galactinus</i>	1st instar	86	100	-	-	-
	3rd instar	14	26	88	100	-
	5th instar	10	24	46	94	100
	adult ♂	14	18	28	82	100
	♀	10	10	22	64	100

It is evident that the first instar nymphs of both species are extremely sensitive against the coldness, and the third instar nymphs are more tolerable than the first instar ones. While the cold-hardiness of these nymphs are certainly inferior to that of the older nymphs and adults. In the mature nymphs and adults, the death caused by coldness began to start from the fifth day and the fourth day in *L. beneficus* and *X. galactinus* respectively. And this fact suggests the higher tolerance of *L. beneficus* than *X. galactinus* against the low temperature.

The temperature condition of 7°C was much more favourable for the survival of bugs, especially in mature nymphs and adults. Compared with the results of rearing at room temperature, no significant increase in the mortality of these individuals was observed under this condition. However, all the tested nymphs of the first instar of *X. galactinus* died within 5 days and none of them developed to the second instar. In *L. beneficus*, a part of the first instar nymphs moulted once after 7-10 days, but all of them died during the second stadium. In the third instar nymphs of both species, some individuals were able to become adults and the percent emergence was higher in *L. beneficus* than in *X. galactinus*. Though the copulation was observed in these adults, the oviposition did not occur at all. The longevity of adults differed significantly between the two species, viz. in the adult

female of *X. galactinus*, it was less than 40 days but more than 90 days in *L. beneficus*. These figures are of course the result obtained under the temperature condition of 7°C. But if reared under a mild temperature condition, no remarkable difference in the longevity was seen. And the results of this test strongly suggests the fact that *L. beneficus* is more cold-adapted than *X. galactinus*. On the other hand, it is also worthy of mention that in a usual environment, there exists only a slight difference in longevity between the female and male individuals, while in such an oviposition restrained condition the longevity of adult female surpassed that of male significantly. And this fact further suggests that the female is consequently more adapted to the survival in the colder season than male.

Table 65. Growth of the 3rd instar nymphs of *Lyctocoris beneficus* and *Xylocoris galactinus* under 7°C condition.

	Duration of stage in days				Percent emergence(%)
	4th	5th	♂ (adult)	♀ (adult)	
<i>L. beneficus</i>	18.6 (5.2)	23.4 (5.4)	38.4 (32.2)	92.1 (32.0)	58 (90)
<i>X. galactinus</i>	12.7 (2.0)	19.5 (4.8 j)	24.3 (30.3)	37.8 (26.8)	32 (95)

The figure in parenthesis shows the data obtained at room temperature.

In autumn, the field collected adults of both species have low oviposition capacity than the spring or summer collected individuals. Even the autumn collected adults were kept under 30°C and 25°C conditions in the laboratory, they could not resume their fecundity. Although the individuals which were reared under 7°C condition in the laboratory readily resumed their fecundity when they were removed to the warmer environment. Therefore, low temperature in the autumn season is not a decisive factor to reduce the fecundity of autumn individuals. The short day-time seems not to influence this phenomenon, because the laboratory strain which are kept in dark condition always show satisfactory results both in growth and reproduction. One of the presumable reason in the low fecundity may be ascribed to the shortage of food, as in this season, compared with summer or spring, the prey fauna in the straw pile is actually poor. Moreover, the results of laboratory experiments indicate that the effect of mal-nutritional condition lasts for a certain long period and is not easily compensated with the additional food supply in the later period. The process of low fecundity in this season seems to be partly explained by the natural regulation in the species so as to keep the population in the cold season as high as possible, namely because the young nymphs cannot withstand the coldness, while the females can tolerate against this if they stop oviposition.

Although the bug cannot be collected on the surface of straw pile in the

winter, the coldness does not seem to play an essential part in reducing the absolute population of bugs. Because, according to the data from the Saga Paper Board Co., the temperature of giant straw pile in winter is unexpectedly high, namely that of the innermost part is recorded as 70-80°C in the fermenting straw pile and sometimes the temperature is so high as to cause the carbonization of straw. Even in a well-dried straw pile, it was more than 20°C. Although this is lower than the optimum temperature for the growth of the bugs, the data in the laboratory rearing experiment pointed out that they can easily keep such a condition for a longer period. Of course, the temperature of the outer part of the straw pile nearly coincides with the natural temperature that is 4°C in mid-winter on an average in northern part of Kyushu, and is often going down to below 0°C for a short period. In the fermenting pile, there exists a gradient of temperature gradually decreasing from 80°C to 4°C from the central part to the outer surface of the pile, and of course it is producing a mild temperature area between the surface and the central part. Therefore, taking in consideration the temperature only, the disappearance of bugs from the surface of straw pile during the winter season and the emigration of bugs into the central part of thatch so as to escape the death of bugs caused by the coldness are the reasonable phenomena. The emigrated individuals will stay in a suitable temperature area, and in the dried thatch the bugs emigrate far deeply. Actually the author collected a number of adult specimens from a considerably inner part of the straw pile in early spring, and these seemed undoubtedly to be the over-wintered individuals in this area. Though Hall (1951) indicates that *X. galactinus* develops through the year in stock yard condition, the author could not collect any nymphs of this species in the field during the winter season. But the bugs of laboratory reared stock continuously grew and produced offspring in the vermiculite or molded corn culture where the condition was more or less similar to that of the grain bins.

Relative toxicity of pesticides

The objective of blending chemical and biological control in an integrated control or modified program for pest suppression has been widely discussed recently (Stern et al. 1959, van den Bosch and Stern 1962, Bartlett 1964, Beirne 1967, etc.). On the other hand, the previous worker-s demonstrated both in laboratory and field tests that some organic pesticides have differential toxicity to numerous entomophagous arthropods (Bartlett 1963, Herne and Putman 1966, Lingren and Ridgway 1967 etc.). Therefore, prior to the establishment of the harmonization of natural enemies and pesticidal applications, the effects of pesticides on the survival and behaviour of natural enemies should be thoroughly studied.

The predation of the rice stem borer (*Chilo suppressalis* (Walker)) by *Lycotocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) was already discussed in the former part of this work. When these Anthocorid bugs are practically liberated in the paddy fields or the other borer infested areas, the influence of pesticidal applications on the bugs would become an unnegligible factor to determine the practical value of these bugs as rice stem borer controlling agents.

The review of the literature reveals that Anthocorid bugs are also destroyed by many pesticides as the other entomophagous insects. The first report on the toxicity of pesticides against Anthocorid bugs was introduced by Hill (1945) who reported the reduction of population of *Orius* spp. by the spray of DDT. Since then, voluminous works have been done on the reduction of Anthocorid bug population in the natural condition after the application of pesticides. The relevant experiments of the laboratory works on the relative toxicity of pesticides against the Anthocorid bugs were done by Burke (1959), Gratwick (1965) and Lingren and Ridgway (1967). Namely, Gratwick evaluated the toxicity of 11 insecticides which were usually used for orchard insect pests control against *Anthocoris nemorum* (Linné) with direct spray and residual toxicity methods. The toxicity of some insecticides against *Orius insidiosus* (Say) was studied by two other authors.

Concerning the insecticidal toxicity against *L. beneficus*, the first work was done in Saga Agricultural Experimental Station in Japan, and the extremely high toxicity of BHC, DDT, dieldrin, aldrin, malathion and parathion dusts was reported (Anonym 1954). The toxicity of BHC was also proved by Oho (1955a, b). But the similar kind of study on *X. galactinus* has not yet been carried out.

In the present work, 8 insecticides and 2 acaricides were tested. Among them, first 5 chemicals are commonly used in the paddy field and latter 5 in orchard in this country.

The names of insecticides and acaricides used are listed as follow :*

Dipterex 50 % E. C. 2, 2, 2-trichloro-1-hydroxyethyl dimethyl phosphonate.

Methyl parathion 40 % E. C. 0, 0-dimethyl 0-p-nitrophenyl phosphorothioate.

Sumithion 50 % E. C. 0, 0-dimethyl-0-3-methyl-4-nitro phenyl phosphorothioate.

Sevin 15 % E. C. 1-naphthyl N-methylcarbamate.

Elsun 50 % E. C. ethyl 0, 0-dimethyl dithiophosphoryl-1-phenylacetate.

Formothion 45 % E. C. 0, 0-dimethyl-S-(N-methyl-N-formoylcarbamoylmethyl) phosphorodithioate.

Dimethoate 43 % E. C. 0, 0-dimethyl S-(N-methylcarbamoylmethyl)-dithiophosphate.

Diazinone 34 % W. P. 0, 0-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate.

* All the pesticides used in this test were kindly furnished by Mikasa Chemical and Industrial Co., Ltd.

Akar (=chlorobenzilate) 21 % E. C. 4,4'- dichlorobenzoic acid ethyl ester.
 Tedion 8 % E. C. 2, 4, 5, 4-tetrachlorodiphenylsulphon.

For the testing method, residual-film method was applied throughout this work. **Thus**, 1.0 ml of pesticidal preparation was uniformly soaked into a set of filter paper which was composed of one rectangular sheet (9x6.3 cm) and two discs (2 cm in diameter). After the immersion to the solution, the filter paper was dried completely under the room temperature condition. The desiccated filter paper was fixed to the interior surface of a glass vial which was 9 cm and 2 cm in length and diameter respectively. Then 10 bugs were released in each vial. After 1 hour's contact with the treated filter paper, the bugs were removed to a clean glass vial of the same size supplied with larvae of *Tribolium castaneum* (Herbst) and a piece of moist filter paper. The survival of tested bugs was estimated 12, 24 and 48 hours after the treatment, then moribound individuals were counted as killed. As a check of test, water soaked filter paper was used instead of chemical soaked paper. The test was replicated 5 times in each test, and all the works were conducted under 25°C condition.

For the preliminary test, laboratory reared adults were used and 5-6 day old *L. beneficus* and 3-4 day old *X. galactinus* were selected for experiments. These adults have been fed with sufficient diet of the larvae of *T. castaneum* from their 1st nymphal stadium. The effect of pesticides on the mortality of *L. beneficus* and *X. galactinus* are given in Tables 66 and 67 respectively.

Among the pesticides tested, the concentration of sevin and two acaricides are apparently lower than the other chemicals. This is merely due to the low concentration of the effective ingredient in the original formulation. Accordingly the other pesticides contain about 40 % of ingredient, and those of sevin, akar and tedion are only 15, 21 and 8 percent respectively. Therefore, it is surely hard to prepare entirely homogenized emulsion in higher concentrated level with those formulation. Nevertheless, in such low concentrations, those evidently surpass the conventional concentration in the field application, namely those are about 0.03 and 0.01 % in sevin and two acaricides respectively. On the other pesticides, the conventional concentrations are about 0.05 %.

Tables 66 and 67 show that *L. beneficus* is slightly more sensitive than *X. galactinus* against all the tested pesticides. For both species of bugs, only parathion shows very high toxicity in the conventional dosage and on this level no other pesticides show significant toxicity. Although sumithion is the next toxic pesticide, in both species more than 80 % mortality is obtained in 0.2 % solution that contains about 4 times as many dosage as that of the conventional preparation. Tables 66 and 67 also indicate that the higher toxicity is demonstrated in organic phosphate compounds than in carbamate compounds. The high toxicity of the former compounds than

Table 66. Effect of pesticides upon the mortality of 5-6 day old adult of *Lyctocoris beneficus* reared in the laboratory.

Pesticides	Concentration of effective ingredient(%)	Percent mortality		
		12 hr	24 hr	48 hr
Dipterex	0.20	4	6	8
	0.50	22	26	26
Dimethoate	0.10	12	14	14
	0.20	38	40	44
	0.50	66	74	74
Diazinone	0.05	10	10	12
	0.10	38	38	42
	0.20	78	80	80
Sumithion	0.05	0	4	4
	0.10	10	10	12
	0.20	84	86	86
Methyl parathion	0.012	94	96	96
	0.025	100	100	100
Formothion	0.20	0	8	8
	0.50	14	18	18
Elsun	0.20	8	10	3.0
	0.50	16	16	16
Sevin	0.05	18	20	20
	0.15	22	22	24
Akar	0.20	2	4	4
	0.40	4	4	4
Tedion	0.10	0	2	2
	0.15	0	0	0
CK		0	2	2

the other series of pesticides against the entomophagous insects are also indicated by some workers (Bartlett 1963, Yun and Ruppel 1964, etc.). Both akar and tedion are very slightly harmful against the bugs even in 10 times as many dosage as the conventional concentration.

The relative toxicity of these pesticides against the field collected adult bugs was also tested with the same method. Hereon, the author used *L. beneficus* of the autumn generation and *X. galactinus* of the spring generation. All the individuals were used to experimental purposes within 12 hours after collection (Tables 68 & 69).

As seen in Tables, field collected individuals were significantly sensitive than laboratory reared ones in both species. Among the highly toxic chemicals, however, only parathion is ranked to most dangerous against the laboratory strain. While diazinone and sumithion belong to the group of dangerous compounds against field collected bugs. Besides, dipterex, dimethoate and formothion showed moderate toxicity. Although other pesticides were less toxic, the mortality of field collected material against these chemicals were apparently higher than that of laboratory reared

Table 67. Effect of pesticides upon the mortality of 5-6 day old adult of *Xylocoris galactinus* reared in the laboratory.

Pesticides	Concentration of effective ingredient(%)	Percent mortality		
		12 hr	24 hr	48 hr
Dipterex	0.02	0	2	2
	0.50	6	8	3.0
Dimethoate	0.10	4	4	4
	0.20	20	22	22
	0.50	58	58	60
Diazinone	0.05	0	2	4
	0.10	20	22	26
	0.20	30	36	36
	0.50	96	100	100
Sumithion	0.05	0	2	2
	0.10	2	4	8
	0.20	70	78	82
	0.50	92	100	100
Methyl parathion	0.012	76	78	78
	0.025	80	82	86
	0.050	98	100	100
Formothion	0.20	0	2	2
	0.50	4	6	8
Elsun	0.20	2	4	4
	0.50	2	2	2
Sevin	0.05	4	4	4
	0.15	2	2	6
Akar	0.20	4	4	4
	0.40	2	2	2
Tedion	0.10	0	0	2
	0.15	0	0	2
CK	-	0	0	0

material. Another noticeable fact proved in this experiment is that the laboratory reared *L. beneficus* was more sensitive to pesticides than *X. galactinus*, while the field collected *L. beneficus* was more tolerable than *X. galactinus*.

For the cause of such higher sensitivity of field collected individuals than laboratory reared ones, the age structure of field collected population may be at first taken into consideration. Namely, field collected adults are composed of various aged individuals. The variations of sensitivity against 7 insecticides in relation to the age were tested with the laboratory reared adults of both species (Table 70). Also the sensitivity against 0.05 % sumithion in the various aged adults of *X. galactinus* is illustrated in Fig. 12.

Table 70 shows the significant increase of sensitivity to insecticides in older adults of both species. And the figure also demonstrates the linear

Table 68. Effect of pesticides upon the mortality of field collected
Lyctocoris beneficus.

Pesticides	Concentration of effective ingredient(%)	Percent mortality		
		12 hr	24 hr	48 hr
Dipterex	0.05	24	26	26
	0.10	48	48	48
Dimethoate	0.05	18	24	24
	0.10	78	80	80
	0.20	94	94	94
Diazinone	0.05	62	62	64
	0.10	88	90	92
Sumithion	0.025	52	56	56
	0.050	96	96	96
	0.100	100	100	100
Methyl parathion	0.012	100	100	100
	0.025	100	100	100
Formothion	0.05	12	14	14
	0.10	24	24	26
Elsun	0.05	2	2	4
	0.10	12	14	14
	0.20	20	24	24
Sevin	0.025	2	4	4
	0.050	24	26	26
Akar	0.20	0	0	2
	0.40	2	2	2
Tedion	0.10	0	0	2
	0.15	2	2	2
CK		0	0	0

increment of sensitivity in accordance with the age.

The nutritional condition of insects also gives certain influences on the tolerance against pesticides. As to this problem, a review was compiled by Gordon (1961). And it has become the general conception that well-fed insects have higher tolerance to pesticides. Although no further study concerning this aspect was conducted in the present work, the superiority of nutritional condition in the laboratory reared material to the field ones may be easily suggested by the presence of larger and heavier individuals in the former material. And this tendency is much more evidently pronounced in *X. galactinus*. Furthermore, the fact that the difference of sensitivity between the laboratory reared and field collected populations of *L. beneficus* is smaller than *X. galactinus* may be explained by the food condition in the field which seems more favourable to *L. beneficus* than *X. galactinus*. The cause of such difference in the susceptibility of pesticides is presumably due to the season of their appearance too. Because, in many insects, the autumn generation is more nutritionally satisfied and have higher tolerance against the detrimental condition than the spring

Table 69. Effect of pesticides upon the mortality of field collected *Xylocoris galactinus*.

Pesticides	Concentration of effective ingredient(s)	Percent mortality		
		12 hr	24 hr	48 hr
Dipterex	0.05	28	28	32
	0.10	54	56	56
	0.20	78	82	82
Dimethoate	0.05	36	38	38
	0.10	100	100	100
	0.20	100	100	100
Di azinone	0.05	76	78	78
	0.10	100	100	100
Sumithion	0.025	62	68	68
	0.050	100	100	100
	0.100	100	100	100
Methyl parathion	0.012	100	100	100
	0.025	100	100	100
Formothion	0.05	26	34	34
	0.10	40	44	46
Elsun	0.05	4	4	8
	0.10	18	18	18
	0.20	36	38	40
Sevin	0.025	12	14	14
	0.050	56	58	58
Akar	0.20	4	4	6
	0.40	8	8	8
Tedion	0.10	2	4	4
	0.15	4	6	6
CK	—	0	0	2

generation (Rainwater 1951, Keiser et al. 1953, etc.).

The sensitivity of nymphs against sumithion was also examined using various stages of both species of bugs (Fig. 13). The result shows the increment of tolerance in accordance with the process of the development of nymphal stages. Sumithion of 0.05 % solution was scarcely harmful to the 5th instar nymphs of both species, while the same dosage of the chemicals was certainly toxic to the young nymphs. In earlier nymphal stage, *L. beneficus* is slightly more sensitive than *X. galactinus*. But the difference in mortality between the two species was not observed in the 4th and 5th instars.

Tedion is widely used for the control of mites on various fruit plants and vegetables. Its acaricidal action concerns the prevention of their offspring. Therefore, the low mortality effect of this acaricide against Anthocorid bug is rather anticipated. Further, the toxicity of tedion against the eggs and young nymphs of the Anthocorid bug is unnegligible. To estimate the ovicidal action of this acaricide, tedion preparation of

Table 70. Effects of pesticides upon the motallity of 20-25 day old adults of *Lyctocoris beneficus* and *Xylocoris galactinus*, both of which are reared in the laboratory.

Pesticides	Concentration of effective ingredient(%)	Mortality after 48 hr (%)	
		<i>L. beneficus</i>	<i>X. galactinus</i>
Dipterex	0.05	28	26
	0.10	44	28
Dimethoate	0.05	14	8
	0.10	26	12
	0.20	68	50
Sumithion	0.05	60	52
	0.10	100	100
Methyl parathion	0.012	100	100
	0.025	160	100
Formothion	0.20	24	12
	0.50	60	48
Elsun	0.20	34	22
	0.50	100	100
Sevin	0.025	28	24
	0.050	42	26
CK	-	2	4

conventional concentration, namely 0.01 %, was sprayed on the eggs which were laid on a filter paper. Then the rate of hatching of eggs and survivality of newly hatched nymphs were observed. The newly hatched nymphs remained on the treated filter paper for about 24 hours, then were placed to a clean petri-dish and reared with the larvae of *T.castaneum*.

The result of experiment reveals the uneffectiveness of tedion on the hatching rate of the eggs of both species. A slight detrimental effect was only observed on the eggs which were treated with tedion on the date of oviposition. But in this case, the rate of hatching still showed more than 70 %. This acaricide has no contact poisonous activity against the newly hatched nymphs, and the majority of young nymphs complete their growth. The acaricidal action of tedion is also known to produce invalid eggs by the treated female. The female treated with 0.1 and 0.15 % solution of tedion laid many eggs which showed normal development.

In the present experiment, only parathion is extremely harmful against laboratory reared individuals, but parathion, sumithion, dipterex and diazinone against field collected individuals. But from the results of these experiments it is too early to conclude the safety of the other pesticides on the Anthocorid bugs. Actually under the field condition the bugs have few opportunities to make contact with such concentrated pesticidal preparations as used in this work. The exposure of Anthocorids for a fairly long period with the treated circumstance is a very important factor of

Fig. 12. Variation of sensitivity to 0.05 % sumithion solution against the age of adult *Xylocoris galactinus*.

increasing the mortality. For example, 3 hour exposure to the filter paper which was treated with 0.1 % sumithion solution gave more than 90 % mortality against the young laboratory reared adult of *X. galactinus*. Whereas, the exposure of such solution for 1 hour gave only 2 % mortality to the adult of the same condition. In another instance, 0.05 % elsun solution which is scarcely harmful to the laboratory reared *X. galactinus* adult showed 100 % mortality to the adult after 6 hour exposure. The bugs are generally acting as a rice stem borer controlling agent in a straw pile which is badly ventilated. When the pesticides is applied in such surroundings, the bugs are easily exposed to the pesticides for a comparatively long period, causing the higher mortality of the bugs. Of course the pesticide may induce the reduction of bug population and at the same time decrease the predatory effect against the rice stem borer.

Fig. 13. Mortality of various stage nymphs of *Lyctocoris beneficus* and *Xylocoris galactinus* exposed to 0.05 % sumithion.
 ■ *L. beneficus*.
 □ *X. galactinus*.

Inter-specific competition between *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber)

Competition between the species under the artificial and natural conditions has been investigated by numerous authors (Andrewartha and Birch 1960, DeBach 1966, etc.). Especially in predacious insects, the intra- and inter-specific cannibalism have been reported by many workers. For example, Wilde (1962) reported the cannibalism among the Anthocorid nymphs and Neuropterous larvae under the laboratory condition. In another example, McMullen and Jong (1967) suggested the competition among *Anthocoris* spp., *A. antevolens* (White) and *A. melanocerus* (Reuter) and the other predacious insects. As they indicated clearly, the population of *Anthocoris* spp. increased in the DDT treated pear orchard. And they attributed the cause of the increase of the *Anthocoris* population to the reduction in numbers of the other predacious species such as *Deraeocoris brevis piceatus* Knight, *D. fasciolus* Knight, *Diaphnocoris provancheri* (Burque), *Chrysopa carnea* Stephane, *C. oculata* Say, etc. The intraspecific cannibalism in the Anthocorid bugs was also pointed out by Barber (1936b) as "*Orius insidiosus* (Say) is cannibalistic, both as a nymph and as an adult. Adults fed upon nymphs, and females frequently feed upon males. The larger nymphs also devour smaller nymphs." A similar phenomenon in *Anthocoris nemorum* (Linné) was observed by Hill (1957). According to the author's preceding experiments, when *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) are reared separately, the intraspecific cannibalism occurred only on the adults that devour their eggs. The cannibalism among adults and nymphs was only observed on the animal diet deficient or artificial diets provided condition. And it is an evident fact that the intraspecific cannibalism exists scarcely among the adults and nymphs of both species under the optimum condition.

The present work was made in an attempt to determine the occurrence of inter-specific competition between *L. beneficus* and *X. galactinus*. All the works were conducted under 30°C condition and the bugs tested were confined to a cylindrical petri-dish (8 x 8 cm in diameter and height) and provided with water soaked filter paper and dry cracked corn. Besides, to avoid the cannibalism caused by the starvation, sufficient prey, namely larvae of *Tribolium castaneum* (Herbst), were supplied.

To investigate the effect of one species on the young nymphs of another species, 10 of the first instar nymphs of one species were reared together with 10 individuals of variable developmental stages of another species. And the number of survivals was counted after 2 and 4 days (Table 71).

As seen in Table 71, the first instar nymphs of *X. galactinus* were not

Table 71. Number of survival of *Lyctocoris beneficus* and *Xylocoris galactinus* reared under an artificial condition (each petri dish contains 20 individuals, avg. of 5 replicates).

Combination	No. of survival			
	After 2 days		After 4 days	
	L. b.	x. g.	L. b.	x. g.
L. b. (1st instar) + X. g. (1st instar)	7.6	8.6	6.8	8.4
L. b. (3rd instar) + X. g. (1st instar)	9.8	8.8	9.8	8.2
L. b. (adult) + X. g. (1st instar)	10.0	8.4	9.4	7.8
L. b. (1st instar) + X. g. (3rd instar)	5.4	10.0	3.6	10.0
L. b. (1st instar) + X. g. (adult)	3.4	10.0	1.8	10.0
L. b. (1st instar)	9.4	-	9.2	-
X. g. (1st instar)		10.0		10.0

L. b. : *L. beneficus*.

x. g. : *X. galactinus*.

influenced by the presence of either old nymphs or adults of *L. beneficus*. But the survival of young nymphs of *L. beneficus* was severely affected by the presence of individuals of *X. galactinus*. When the first instar nymphs of both species were reared together comparing with the result from the pure culture. The survival of *X. galactinus* was scarcely influenced, while *L. beneficus* significantly decreased in numbers of survived individuals as compared with the former.

In the second experiment, 20 first instar nymphs of both species were reared gregariously in a dish until all the tested nymphs became adults, counting the number of survived individuals continuously every other day, and the result was illustrated in Fig. 14.

Among the 20 first instar nymphs of *X. galactinus*, 14 individuals completed their nymphal development, while in *L. beneficus* only 5 nymphs became adults. For the cause of death of nymphs, the cannibalistic behaviour between the two species must be considered. But, no practical preying behaviour of *X. galactinus* on *L. beneficus* was observed at all. During the period of 22 days 12 individuals of *L. beneficus* were consumed for the first 6 days. This is equivalent to four-fifths the whole sacrificed nymphs. This fact also indicates that first and second instar nymphs of *L. beneficus*

Fig. 14. Survival progression of *Lyctocoris beneficus* and *Xylocoris galactinus* (started with 20 1st instar nymphs): **Abscissa, No.** of bugs survived, **ordinate, days.**
 --o-o- *L. beneficus*, ----- *X. galactinus*.
 △ shows the date when all individuals became adults.

are especially sensitive against the mixture of *X. galactinus*. The duration of nymphal stages given in Fig. 14 was 20 and 14 days in *L. beneficus* and *X. galactinus* respectively. These values show little difference from that of the pure culture. The body weights of newly emerged adults were 1.4 and 2.1 mg in *X. galactinus* and *L. beneficus* respectively. Compared these figures with that of the pure culture, no difference was seen in *X. galactinus*, but the value was evidently lighter in the case of *L. beneficus*.

The effect of mixed culture on the egg production was investigated using 10 pairs of both species which were emerged on the same day. In this test, the bugs were kept in a plastic tapper sized 13 x 10 x 7 cm (length x width x height) and provided with moist filter paper, dry cracked corn and sufficient prey (Table 72).

The duration of pre-oviposition period was scarcely influenced in both species, being 7 and 5 days in minimum value in *L. beneficus* and *X. galactinus* respectively. The egg production was studied by counting the numbers of eggs deposited on the filter paper and the results are given in Table 72 and Fig. 15.

Although the mixed culture did not influence the duration of pre-oviposition period in both species, the oviposition period of *L. beneficus* was seriously affected by the presence of *X. galactinus*. And in the former species,

Table 72. Effect of mixed culture on the egg production of *Lyctocoris* *beneficus* and *Xylocoris galactinus* (data from 10 pairs of adults).

		Pre-oviposition period (minimum)	Oviposition period (maximum)	Eggs per female	No. of adults 10 days after emergence	
					♂	♀
<i>L. beneficus</i>	pure	7	28	34.6	7	9
	mixed	7	10	12.3	5	6
<i>X. galactinus</i>	pure	4	23	39.2	5	8
	mixed	5	24	30.5	7	7

Fig. 15. Egg production in the mixed culture of *Lyctocoris beneficus* and *Xylocoris galactinus*: **Abscissa**, No. of eggs per female, **ordinate**, days after emergence. ■ *L. beneficus*. □ *X. galactinus*.

the egg deposition almost stopped within a week and the number of deposited eggs decreased to about one-third of the pure culture. On the contrary, egg production of *X. galactinus* in the mixed culture was scarcely affected by *L. beneficus*. The eggs of both species showed some high degree of hatching ability and this fact may indicate that the copulation was not affected by the mixed culture. In the mixed culture, the male *X. galactinus* sometimes mounted on the female *L. beneficus* and tried to make copulation. However, the apparent difference in body size always prevented their copulation. On the other hand, the author has never observed the male *L. beneficus*

mounting on the female *X. galactinus*. Though the accurate longevity of tested adults was not ascertained, the number of adults survived 10 days after the emergence indicates the shortened longevity of *L. **beneficus*** in the mixed culture.

In conclusion, when *L. **beneficus*** and *X. galactinus* are reared together, *L. **beneficus*** is overwhelmed by *X. galactinus* in every aspect of life activities, especially either the survival of young nymphs or egg production. In fact, under the insectarium condition, when *X. **galactinus*** invades into *L. **beneficus*** culture, the offspring production of *L. **beneficus*** is heavily influenced and only a few new progeny of *L. **beneficus*** are obtained from such a contaminated culture, and within certain periods, the culture is entirely replaced by the invader. In the field condition, although the habitats of both species are very similar to each other, the author has never seen the cases in which the two species co-existed in larger numbers in the same habitat.

Co-existence with *Apanteles chilonis* [Munakata]

A Braconid wasp, *Apanteles chilonis* [Munakata] is one of the most important larval parasites of the rice stem borer in Japan. The availability of this parasite as the controlling agent against the rice stem borer is already pointed out by several workers (Gyotoku 1960, Tateishi 1962, Yasumatsu 1964, 1967, Deloach and Miyatake 1966, etc.). As both ***Lyctocoris **beneficus***** (Hiura) and ***Xylocoris **galactinus***** (Fieber) also devour the larvae of rice stem borer, the study on the inter-relationship between these natural enemies is very important in deciding the effect of natural enemies on the control of rice stem borer in the field condition.

For the rearing method of the parasites, about 10 of newly emerged *A. chilonis* adults were confined to a large test tube which was 20 × 3 cm (length and diameter) in size, providing with water soaked cotton and bee honey. Besides, 3 mature larvae of *Chilo suppressalis* (Walker) were afforded for the oviposition of the parasite. The rice stem borers were readily received the oviposition by the parasites. The borer was changed to new ones every day. The parasitized borers were kept in another test tube in which the paddy straw was afforded under 25°C condition. After 15-16 days, about 20 mature larvae of parasites were escaped from the body of borer and make white small cocoons. Three to four days after pupation the adults emerged. The percent of parasitization was certainly high, being about 80-90 percent.

To test the inter-relationship with the Anthocorid predator, some parasitized borers were placed in a 100 ml beaker into which 10 or 20 adults of *L. **beneficus*** or *X. **galactinus*** were confined. Parts of parasitized larvae were attacked by the predators soon, while some of the borers were attacked after several days. The investigation on the attacked larvae was conduct-

ed twice a day and the borers bearing a pierced puncture were removed from the beaker and observed the further development. Usually, the attacked borers died within 48 hours, and finally became black dwindled corpse. No larvae of parasites were of course escaped from the attacked borers even from the individuals which were parasitized **15** days ago. In the bug container, some mature larvae of parasites came out from the borers which were fortunately escaped the attack of predators. But majority of such mature larvae were readily preyed by the bugs. The predators preferred to prey the larvae of parasites to the borers. Although some mature larvae were seen to succeed to make cocoons, the bugs sucked the blood of larvae through the wall of cocoons, and thus the adult parasites seldom emerged in the bug container.

On the other hand, if the bug-attacked borers were provided for oviposition of *A. chilonis*, then the ovipositional behaviour of the wasp was observed. But all the borers died within 48 hours after the attack by predators, and of course no parasite was developed from those pierced borers. In this case, the parasitization does not seem to accelerate the death of the pierced borers.

Next, the unparasitized borers and those parasitized 3, 6, 9, 12 and 15 days before were marked with magic ink and 5 individuals of each group were put together in a plastic tapper (14 x 10 x 6 cm in long, wide and high) in which about 50 or 100 of *L. beneficus* or *X. galactinus* were supplied. Then the preference of attack by the predators was checked at 2 day interval (Fig. 16).

The experiment indicates that more than 70% of the attacked larvae belonged to the individuals of the earlier parasitized borers. The parasitized larvae in this stage were apparently more sluggish than the normal or newly parasitized individuals. Therefore, they were more easily accepted by the predators.

In conclusion, both *L. beneficus* and *X. galactinus* are directly or indirectly fatal agents to *A. chilonis*, and the Anthocorid bugs always eliminate ***Apanteles*** wasps in the laboratory condition. Although no field survey is conducted as yet, the similar tendency may be conceivable when and where the Anthocorid bugs are abundant in the environment.

Fig. 16. Number of attacked borers by *Lyctocoris beneficus* and *Xylocoris galactinus*. The borers were containing various stage larvae of *Apanteles chilonis*; Abscissa, No. of borers attacked, ordinate, days after parasitization.
 ■ *L. beneficus*.
 □ *X. galactinus*.

Approach to the mass production

The availability of the two bugs as the predators against the rice stem borer was proved in the present work. Therefore, the next necessary step of study is the mass production procedure of these bugs.

I. Culture media and rearing method.

Two kinds of material are advisable for the culture media of the Antho-corid bugs. One is the moist vermiculite and another is the molded cracked corn. The moist vermiculite is prepared to add 15 ml of water into 100g of vermiculite - rough vermiculite is preferable - the growth of bugs is better on the molded vermiculite. The preparation of the molded vermiculite is not difficult. It is easily obtained by adding small amount of wheat flour into the moist vermiculite, then the *Tribolium* larvae can survive for a longer period so long as the food is available.

The molded cracked corn is generally superior to the vermiculite medium both for the offspring production and development of bugs, especially in the case of *L. beneficus*. But it is rather a difficult problem for this medium to prevent its excess fermentation without making any injurious effects on the development of mold and bugs. Such Lepidopterous insect pests as meal moth (*Plodia interpunctella* (Hübner)), grain moth (*Aphomia gularis* (Zeller)), almond moth (*Cadra cautella* (Walker)) etc. are harmful to the bugs. The infestation of these pests is easily prevented by the autoclaving of corn before the mold cultivation. Another difficulty in the corn medium is the outbreak of mites, for example, grain mite (*Acarus siro* Linné) and *Tyrophagus dimidiatus* Herman. Though the mites are excellent food for the bugs especially for young nymphs, the mites also damage bug's eggs, and usually very few nymphs of bugs hatch from the mite contaminated media. In the rearing either with vermiculite or molded corn medium, the bugs oviposit eggs in the media and newly hatched nymphs grow on it. The co-existence of the adults gives very little influences on the growth and survival of nymphs. The maintenance of water content in culture media is very important both for the existence and oviposition of bugs, especially for the latter activity. Further, the drying condition of media substrata is fatal to the hatching ability of eggs on it. The humidity of media should be inspected at least every 2--3 days, and supply some water if necessary.

Although the following method is more time-consuming than the above mentioned methods, higher efficiency of off-spring production is obtained by this method especially in the case of *L. beneficus*. Namely, the bugs are reared in the dry cracked corn or dry vermiculite and the wet filter paper is provided in the rearing container as an oviposition substratum.

Then, the majority of eggs are deposited on the provided filter paper. However, in this case attention should be paid to the prevention of egg-predatism by the adult bugs. Thus, papers should be exchanged to new ones at least twice a day. The presence of eggs is easily visible by the naked eyes, and the papers containing deposited eggs are placed on wet vermiculite. The filter paper should keep certain humidity during the incubation period of eggs, but too much water supply is harmful to eggs. When the eggs change in colour to pinkish and the time of hatching becomes near, it is necessary to provide some amount of *Tribolium* culture medium and premature larvae of *T. castaneum*. The occasional addition of medium is very important both for the survival of *T. castaneum* larvae and the keeping of mold on the medium. The mortality of newly hatched nymph is always kept low in moderately molded media. After hatching, the food and water should be supplied occasionally to the rearing media. When the nymphs grow to the last instar, the nymphs are better to be removed from the vermiculite medium to the dry cracked corn medium. Of course, the vermiculite medium is also suited for adult breeding. Always the rearing process is preferable to be carried out under 30°C condition. The illumination of more than 12 hours per day gives bad influence on the growth of bugs, and the rearing in the dark chamber of incubator showed satisfactory result. The three methods mentioned above are recommendable for the rearing of both species, and especially the mass production of *X. galactinus* was found to be very successful.

For the mass-culture of the bugs, the author mainly used 12x 12x 12 cm (in length, width and height each) plastic tapper which is containing 1 cm depth of culture medium. About 10 or 20 pairs of *L. beneficus* or *X. galactinus* adults may be kept in it respectively.

II. Food.

According to the author's work, it was proved that the larvae of red flour beetle (*Tribolium castaneum* (Herbst)) were the most suitable food for both species of bugs. As it is a cosmopolitan stored-grain insect pest and able to be reared with any of the wide varieties of foodstuffs. Therefore, voluminous studies have been done by many workers on this beetle. Its systematic and morphological studies were conducted by Hinton(1948), Mertz(1961), Ho (1967) etc., and the studies of its biology were accomplished by Good (1933, 1936), Howe (1956, 1960), Khalifa and Badawy (1955a, b), Sinha and Mookhewee (1967), etc. The method of mass culture was developed by Harein and Soderstrom (1966) and Qureshi (1966). The method of the former authors was established at Savannah laboratory in U. S. A., using white flour and corn meal (47.5 % each) and brewer's yeast (5.0 %) for culture medium, while the latter author used a mixture of whole wheat flour, powdered milk and dried yeast.

The present author collected the adults of *T. castaneum* at the stock yard of Kyushu University. Though he did not use "Triboliotrap" which was designed by Stanley (1962), it may be available for the collection of the *Tribolium* beetles. The rearing and propagation of the beetles were carried out by the following method: About 500 individuals of both male and female were released in a 15×20 cm (in diameter and height) glass cylinder which contained about 250 cm³ of powdered rat food and wheat short flour (2 : 1 v/v). Thus, the food was occupying the bottom of the cylinder about 4 cm thick. After 5 day rearing, the adults were separated from the medium by sieving. The adults were transferred to a new medium for further oviposition. The sieved medium which contained eggs and young nymphs of beetle was placed in a 6×22×40 cm (in height, width and length) carton box. Then 400-500 mature larvae were obtained 20-23 days after sieving. The results of rearing with the powdered rat food plus wheat short flour medium were quite excellent and superior to the results obtained from some other media (Table 73).

Table 73. Growth of *Tribolium castaneum* in various media under 30°C (average of about 50 individuals).

Media	Larval stage (in days)	Weight of mature larvae (in mg)
Rat food+wheat short flour (1 : 1v/v)	25.3	3.5
Wheat short flour	30.4	3.4
Refined flour+yeast (95 : 5 w/w)	28.8	3.1
Refined flour	92.5	3.2

The mature larvae were readily separated from media by sieving with a 25 mesh sieve. For the separation and inspection of larval growth and population, Saunder's method was also advisable (Saunders and Kruger 1957). All the breeding works were performed under 30°C condition, while the temperature of 33°C was more preferable.

The renewal of oviposition and feeding media of adults every several days was very important for the mass-production of the beetle, because adults have a predacious habit against their own eggs (Boyce 1946, Sonleithner 1961). The adult beetle is also predacious on the eggs of Anthrenid bugs laid in the bug-culture media, but less harmful to the eggs which are laid on the filter paper. The presence of adult beetles in the bug's media gives no bad effect on the growth of adult and nymphal bugs. Therefore, the supply of numerous mature larvae of the beetle to the bug-culture media at one time should be avoided, and it is a rather preferable method to provide premature larvae instead of mature ones. The removal of the adult beetles from the bug's media is an important pro-

cedure especially to protect bug's eggs which were laid in the culture media.

ITT. Collection of bugs from the culture media.

This work is easily conducted by pouring the whole medium on the white-coloured plate, and shaking the plate gently. Then the adults and mature nymphs leave the medium readily and will be collected by an aspirator. This work is preferably to be done under the warmer condition, higher than 25°C, as under the cooler condition the bugs become too inactive to be collected. The percentage of collection of bugs by this method was tested by using 100 individuals which were composed of both adults and various stages of nymphs. And the number of collected bugs during 5 minutes from 288 cm² of culture medium that corresponds to the contents of two rearing plastic tappers is given in the following Table 74.

Table 74. Percent collection of bugs from the media by pouring method within 5 min work.

	Adult and old nymph		Young nymph	
	<i>L. beneficus</i>	<i>X. galactinus</i>	<i>L. beneficus</i>	<i>X. galactinus</i>
Vermiculite	96.3	94.1	64.3	55.2
Molded corn	85.6	82.8	47.5	36.7
Dry corn	98.0	96.5	71.6	68.2

As seen in the table given above, more than 80 percent of adults and mature nymphs of both species were collected by a short time handling. While the collection of young individuals of both species was always low in any kinds of media. The separation of young individuals or eggs from the culture media is time consuming or almost impossible. Therefore, it is better to keep the media in the rearing condition for additional several days after the first separation and conduct the second separation when the remained nymphs become the matured nymphs or adults. This separation work is more easily conducted if the culture media are spread on the cloth on the bottom of the cabinet, especially in the treatment of the more active species, *L. beneficus*.

Attack of bugs to man

As in the case of the other groups of Heteropterous bugs, the Anthocorid bugs sometimes attack man and suck blood. The biting records of the Anthocorid bugs were previously made by some workers, and the bugs are also dealt as injurious insects of minor medical importance in some

text-books (Riley and Johannsen 1936, Martini 1946, Matheson 1950, Shiraki 1958, etc.). In spite of the blood sucking habit, the bugs are normally recognized as beneficial predators against injurious insects. Up to the present, the following bugs have been reported as blood sucking agents.

Anthocoris congolensis Brumpt (Shiraki 1958)

A. kingi Brumpt (Lewis 1958)

A. musculus Say (Torre-Bueno 1931)

A. nemorum (Linné) (Matheson 1950)

Lyctocoris campestris (Fabricius) (Woodward 1951, Stys and Daniel 1957)

Orius insidiosus (Say) (Tucker 1911, Malloch 1916, Myers 1926j)

O. tantilus (Motschulsky) (Misra 1924, Myers 1926)

Xylocoris afer (Reuter) (Lewis 1958)

The case of attacking by *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber) has not been recorded. During the author's field work with these bugs, he has experienced their attack very rarely. The attack is slightly more painful than that by some leafhoppers such as *Nephotettix cincticeps* (Uhler). The sucked spot becomes red and swollen but without itching as mosquitos-bite and the sore does not remain for a long period. The sucking is encountered in the hot summer season. Anyway, the bugs are nothing but facultative or accidental blood suckers as the other Anthocorid bugs. Therefore, considering their usefulness as the predators of rice stem borers and the other insect pests, such attack against man may be regarded as a permissible fault to the side of the bugs.

Conclusion

Evaluation of bugs as the predators against the rice stem borer

The possibility of utilizing *Lyctocoris beneficus* (Hiura) as a controlling agent against the rice stem borer (*Chilo suppressalis* (Walker)) is already pointed out by some workers (Oho 1955a, b, c, Bess 1967, Yasumatsu 1967, etc.). And the predatory effect of *L. beneficus* and *Xylocoris galactinus* (Fieber) is also proved by the present work. Although the number of prey attacked by a single bug is very much variable with some factors including the population density of the predator itself or the host, etc., the daily attacked rice stem borer numbers are approximately 0.32 and 0.24 for the adult *L. beneficus* and *X. galactinus* respectively. But these data were obtained under the environment where no shelter was provided for the prey and predator, and the predator was directly in contact with the prey. Therefore, such condition may differ in many respects from the natural circumstances where the bugs are playing an important role as the controlling agents against the borers. As the preying amount and feeding be-

haviour of bugs are easily influenced by some environmental factors, the preying behaviour of the two Anthocorids against the borers was investigated under the various temperature, illumination and straw conditions for the purpose of making further evaluation of the effect of bugs as the borer predators. Dry and moist paddy straws were provided in 50 ml beakers separately, and to each beaker were released 5 borers, and 10 or 20 adults of *L. beneficus* or *X. galactinus* respectively. Besides, for the sake of control, some beakers were only provided with the borers and bugs without putting any straw. Then parts of beakers were placed in a dark chamber of an incubator, the temperature being kept 30°C constantly, and the other groups of beakers were placed under 25°C condition. In the 25°C group, half of beakers were placed under the dark condition and the remaining half placed under the illumination of 12 hours per day. Each tested group was consisted of 5 beakers and the number of survived bugs and borers were counted on the fifth day from the start of the experiment (Tables 75 & 76).

Table 75. Number of rice stem borers consumed by *Lyctocoris beneficus* under various conditions during 5 days.

Temp.	Light condition	Condition of straw	No. of individuals per beaker	
			Borers preyed	Bugs survived
25°C	12	wet	2.2	8.2
		dry	2.6	8.6
		no straw	3.0	6.8
25°C	dark	wet	2.0	7.4
		dry	1.6	8.2
		no straw	2.4	6.0
30°C	dark	wet	2.0	7.4
		dry	2.2	7.2
		no straw	3.6	4.6

Table 76. Number of rice stem borers consumed by *Xylocoris galactinus* under various conditions during 5 days.

Temp.	Light condition	Condition of straw	No. of individuals per beaker	
			Borers preyed	Bugs survived
25°C	12 hrs	wet	1.2	12.4
		dry	1.4	17.0
		no straw	1.8	4.4
25°C	dark	wet	1.2	14.6
		dry	1.4	15.8
		no straw	2.0	4.8
30°C	dark	wet	2.4	13.6
		dry	2.8	12.0
		no straw	3.4	2.4

As seen in Tables 75 and 76, the highest preying records were obtained in the no straw group in both species. This is probably due to the easier accessibility of predators to prey. But, in this group the mortality of bugs was also apparently higher than the straw provided group. This tendency was more pronounced in *X. galactinus* and it is perhaps partly attributed to the crush of bugs by the movements of borers when the prey are attacked by the bugs, and also partly due to the cannibalism between the bugs. But no actual cannibalism was observed. Although the number of preyed borers decreased in the straw provided group, the survival of bugs increased. The killed borers were observed both at the outside and inside of the provided straw. The borers found at the outside of the straw might have been attacked when they were at the outside, or might have escaped from the straw after receiving the piercing of the bugs. Under the normal condition all the borers were in the straws provided. Therefore, the latter case is seemingly more possible. In the straw provided beakers, bugs were almost found under the outermost leaf sheath, and some individuals were also found in the pith of straw. Moreover, a part of bugs were invaded into the pith of straw, the opening of which was covered with the layer of borer's excreta. Only few bugs were found on the surface of straw. In a part of attacked borers, pierced punctures were observed on the posterior part of the trunk. This is rather indicating that these borers were attacked in the straw. The low mortality of bugs in the straw provided group is due to the restriction of borer's activity by the narrowness of the inside of straw, and from this reason, the prey are easily attacked by the bugs without receiving any resistance from the borers. In some straw provided beakers no borers were preyed, but the survival of bugs was certainly high. This fact may suggest that the straw seems to contain some nutritional components which can maintain the life of bugs for a certain period. The higher mortality of bugs was obtained in the moist straw than in dry ones, but this may be caused by the drowning of bugs by the excess of water drops on the inner surface of straw and beaker. On the other hand, moderate moist of straw is indispensable for the bugs to keep high survival rate. The preying amount was slightly higher in the dry straw than in the moist one. The illumination condition seems to stimulate slightly the preying activity of *L. beneficus*, but as the majority of bugs are dwelling in the straw, the light condition seems to be ineffective on the activity of bugs. The light condition seems to have no connection to the feeding activity of *X. galactinus*. Concerning the temperature, feeding amount increased in warmer conditions and this tendency was especially pronounced in *X. galactinus*. For the further observation on these tested straw, some young nymphs were hatched from the tested moist straw, but not from dry straw.

The preying activity of bug nymphs against the borers was also investigated using 2 day old individuals. The experiments were divided into

three: 1) 50 nymphs, 2) 50 nymphs+20 borers and 3) 50 nymphs+20 adult bugs + 20 borers. The plastic tapper of 13 x 10 x 6 cm in size (length x width x height) was provided with slightly moist straw, and in these tappers three kinds of experiments mentioned above were performed. The survival of tested bugs was counted after 10 days rearing (Table 77).

Table 77. Effect of predation of 2 day old bug nymphs on the rice stem borer (data obtained after 10 days).

	No. of survival of <i>L. beneficus</i>			No. of survival of <i>X. galactinus</i>		
	Nymph	Adult	Borer	Nymph	Adult	Borer
50 nymphs + 20 adults + 20 borers	34	20	12	24	19	14
50 nymphs + 20 borers	35	-	7	36	-	10
50 nymphs	17	-	-	14	-	-

As seen in Table 77, the preying activity of both species against the borers was apparently high in the group without adult. Although the young instar bug individuals hardly attack the freely crawling borers, the present work showed the predation of borers by the young instar nymphs under the straw provided condition. In the group with adult bugs, however, the number of bugs in a tapper increased by the addition of **20** adults. But, on the contrary, the preying amount of the borers decreased. In this case, the survival of adults was however extremely high, and high nymphal mortality was also observed. The decrease of nymphal population may presumably be attributed to the adult feeding upon the nymphs instead of borers. The predation of nymphs by the adults was apparently less in *L. beneficus* than in *X. galactinus*. In the adult added group, some newly hatched young nymphs which were evidently produced from the eggs deposited by the provided adults were observed. In the group provided with straw only, the survival of nymphs was the lowest of the three tested groups and this is undoubtedly due to the cannibalism between the nymphs induced by the shortage of animal diets. Although the survival of nymphs was not high in the adult added group, part of nymphs survived and developed to adults. Under such a condition the bugs could survive generation after generation, and actually the population of bugs increased gradually.

The preying activity against the overwintering borers in the paddy stubbles was investigated with the stubbles harbouring borers. Therefore, **10** or **20** adults of *L. beneficus* and *X. galactinus* were liberated into a paddy

stubble which was previously provided with 10 mature borers. The tested stubbles were entirely covered with a glass cylinder and organdy cloth to prevent the escape of predators and borers. These stubbles were placed under 25°C condition with 12 hour illumination a day, and the number of survived individuals was counted by the dissection of 5 stubbles every 2 days (Table 78).

Table 78. Preying effect of *Lyctocoris beneficus* and *Xylocoris galactinus* against the rice stem borers in the paddy stubbles.

Days from inoculation	No. of survival of		No. of survival of	
	Borer	<i>L. beneficus</i>	Borer	<i>X. galactinus</i>
0	10,0	10.0	10,0	20.0
2	8.8	8.2	9.2	17.2
4	8.6	8.4	9.0	16.4
6	7.8	7.0	8.4	14.4
8	7.6	5.4	8.8	12.0
10	7.4	6.4	8.4	13.2

Comparison of the results in Tables 75 and 76 shows that the largest difference is characterized by the decrease of preying amount of the borers. For the cause of the decrease of food consumption, it should be taken into consideration that, in the preceding experiment, the environment of predators is only consisted of straw and borer. But in this work, there were found some small-sized Dipterous, Coleopterous and Collembolous insects amidst the tillers and on the soil, and these small creatures were undoubtedly regarded as preferable foods. Although some bugs were found amidst the tillers or inside of the straw, about half of the liberated bugs were recovered from the surface of the soil. And the latter individuals were roving on the soil surface presumably seeking for small insects. On the contrary, all the provided borers entered into the stubbles. Therefore these surface staying bugs did not contribute to control the borers. Some newly hatched nymphs were observed inside the stubbles 8 days in *L. beneficus* and 6 days in *X. galactinus* after the liberation of their adults. Another difference from straw provided test was that majority of the attacked borers died at the outside of the stubbles.

From the above results, *L. beneficus* and *X. galactinus* may be considered as effective natural enemies to check the overwintering borers in the straw and stubbles, and the control is more effective against the straw dwelling borers. As these predators are polyphagous, the preying effect against the rice stem borer may be greatly influenced by the fauna of the environment, especially by the presence of some small-sized creatures. Al-

though Oho (1955a, b) reported the 90 % mortality of rice stem borers by *L. beneficus*, this data might be obtained from somewhat specially conditioned straw piles. As these predators do not attack exclusively the rice stem borers and their attacks are frequently concentrated to some other small-sized and soft-bodied animals. Therefore, the occurrence of these animals may greatly reduce the possibility of attacking the borers.

Anyway, it is apparent that under certain conditions these predators may prey on the rice stem borers effectively, and the attacked borers may almost absolutely receive fatal damage. For this reason, both *L. beneficus* and *X. galactinus* are considered to be effective natural enemies in checking the rice stem borer population.

Summary

The present work was undertaken to make clear the bionomics of two Anthocorid bugs, *Lyctocoris beneficus* (Hiura) and *Xylocoris galactinus* (Fieber), with special reference to the possibility of utilizing these species as the controlling agents against the rice stem borer.

Both species are commonly found in the giant straw piles, warehouses and also in the field.

The rearing of two species was conducted in the laboratory under the constant temperature conditions of 25°C, 30°C and 33°C. The lengths of egg, nymphal and adult stages at 30°C are 4, 20 and 31 days in *L. beneficus* and 3, 13 and 26 days in *X. galactinus*. Both species have 5 nymphal stadia. The egg production varies a great deal with the individuals, the number of egg per female in *L. beneficus* being 10-36.5-65 and 24-44.7-63 (min-avg-max) in *X. galactinus*. The eggs of both species are oviposited on or in several material, such as cherry twig, corn, filter paper, paddy straw, vermiculite, etc. The eggs laid by the young adults show high rate of hatching. The fecundity suddenly decreases from 4 week old female, and the vitality of off-spring produced from such an old female is also inferior to that from the younger one.

In the giant straw pile at Saga Paper Board Co., the spring emergence of *L. beneficus* and *X. galactinus* begins at the middle of May and the end of April respectively. The population of the bugs reaches a peak at the middle of June, and then it decreases toward July. The second emergence occurs at the end of August. Generally, *X. galactinus* is predominant in marshy places, while *L. beneficus* is found in arid places. They scarcely occur together in the same habitat. The sex ratio of field collected adults of both species is about 1: 1 throughout the year.

During the summer season, the nocturnal flight of the male adults begins just before the sunset. They, at first, fly over the ground and later fly up to a

higher range. The peak of appearance at the lower range is at 6.30 pm and the peak at the higher place is 7.00 pm. The flying activity of females follows the males about 30 minutes later. They stop to fly at about 9.00 pm.

In the field, *L. beneficus* has 3 generations per year. New adults emerge in May, July, August, September, and X. *galactinus* has 5 generations, the adults appearing in June, July, August, September and November. But, due to the longer oviposition period, various growing stages are observed throughout the warmer seasons. The emergence of new nymphs stops in October in both species.

The feeding habit of two bugs was investigated with more than 60 species of prey. So far as the author's observation goes, *L. beneficus* is predacious upon 45 species of 23 families and X. *galactinus* on 43 species of 22 families. The prey have such common characters as sluggish movement and soft-body, etc. The feeding preference was studied among *Ceraphis magnolicolens* Takahashi, larvae of wax moth (*Galleria mellonella* Linné), rice stem borer (*Chilo suppressalis* (Walker)) and red flour beetle (*Tribolium castaneum* (Herbst)). The aphid is the most preferred and *Tribolium* larva comes the next. Difference in preference is seldom observed between the two Lepidopterous larvae. The preference experiment was also conducted with various body-sized *Galleria* larvae, and it appears that the small-sized larvae are more suitable as food.

The feeding amount of bug per day is very variable depending upon many factors. The feeding amount against the *Tribolium* larvae was 2.00 and 1.20 in the adult of *L. beneficus* and X. *galactinus* respectively. The daily consumption of the mature *Chilo* larvae by one predator was 0.60 and 0.20 in the separate and gregarious rearing of *L. beneficus* and 0.32 and 0.24 in X. *galactinus*. The reduction of food consumption under a gregarious condition is due to the concentrated attack of many bugs on one prey.

In the earlier nymphal stage, the conversion ratio of food is higher than that of the later nymphal stage in both species. The predatory value in the adult stage is lower than that of the nymphal stage. The value is lower in the earlier nymphal stage in both species.

The nutritional reliance on plant substances seems to be restricted and the sucking of plant juice keeps the life of nymphs and adults for a very short period. No oviposition is observed under the deficiency of animal diet. The order of suitability of feeding habit in both species is as follows : carnivorous>cannibalistic>necrophagous>fungivorous>phytophagous.

The fecundity of bugs is greatly influenced by the kind of foods. The bugs reared with the *Tribolium* larvae show the highest fecundity and the *Chilo*-fed bugs come the next. The *Galleria*-fed bugs show the worst result that is characterized by the longest pre-oviposition period and the lowest egg-production.

For the purpose of studying this phenomenon in detail, the amino acid composition of *Tribolium* and *Galleria* larvae was analyzed and compared. As the result of analysis, the content of free amino acids in the former food was 73.50 $\mu\text{mol/g}$ and that of the latter food was 30.63. The contents of essential amino acids were 23.58 and 8.10 respectively. Among the amino acids, the contents of aspartic acid, serine, alanine, valine, isoleucine, methionine, leucine, tyrosine and phenylalanine in the *Tribolium* larvae exceed 3 times that of the other foods. The amino acid analysis was also conducted with the adults of *L. beneficus* reared with the *Tribolium* and *Galleria* larvae. No significant quantitative variation in the composition of amino acid was observed between the bugs reared with different foods. It becomes evident that the *Tribolium*-fed bug can take more amino acid nitrogen for the reproductive activity than the bug fed with another kind of diets. The contents of ether extract are much more abundant in the *Galleria* larvae than in the *Tribolium* larvae, being 0.22 and 0.04 g/g respectively. The paper chromatogram of these extracts was markedly different, although no identification was conducted.

For the rearing of bugs with artificial media, four methods were used. Among them, the use of the medium soaked uretank foam is most preferable. The most favourable result was obtained with the medium composed of 2 g dextrose, 2 g soluble starch, 2 g sodium caseinate, 2 g yeast extract, 1 g beef extract, 0.05 g cholesterol, 0.2 g Wassen's salt per 100 ml water. But the rearing result with this medium is much inferior to that with the animal diets either on the survival or growth rate. The frequent cannibalism was observed when several bugs were reared gregariously with the medium. No egg was produced by the female reared with the medium. The addition of 1.0 or 0.5 mg α -tocopherol to 100 ml medium prolonged the longevity and prevented the cannibalism for a certain extent.

The ovarian development relating to the nutritional condition was studied anatomically with the bugs reared with various foods. The ovarian development was the best in the *Tribolium* reared bugs, and the mature eggs were recognized on the 5th and 3rd day after the emergence of *L. beneficus* and *X. galactinus* respectively. The ovarian development of *Chilo*- and *Galleria*-fed bugs was inferior to that of the *Tribolium*-fed individuals and no mature egg was observed during the same period. The number of immature eggs was also abundant in the *Tribolium*-fed bugs. The adult fed with plant juices showed a slight ovarian development till the 3rd day, but the ovary rather degenerated on the 5th day.

The survival, egg-production and the rate of egg hatching of bugs under the various population densities were investigated. In the petri-dish of 6 $\times$ 9 cm in height and diameter, the best result was obtained to rear 10 and 20 adult pairs of *L. beneficus* and *X. galactinus* respectively. The nymphal growth and the rate of survival were superior in 40 and 60 individuals

in the same petri-dish.

In both species, the adult was more tolerable against the coldness than nymphs. When adult was exposed to -3°C , the activity stopped within 3 minutes and the exposure of 30 minutes was fatal. Under the temperature of 5°C condition, the paralysis occurred within 30 minutes, the lethal time at this temperature being 6 and 5 days in *L. beneficus* and *X. galactinus* respectively. Although the young nymphs did not survive under the temperature condition of 4°C , the older nymphs reached very slowly to the adults while no oviposition was seen. In the field condition, both species seemed to overwinter in the adult stage under the shelter of warmer places.

The relative toxicities of 10 pesticides against both species were studied with the residual film method. Against the laboratory reared young adults, 0.025 and 0.05 % parathion showed 100 % mortality to *L. beneficus* and *X. galactinus* respectively. Sumithion was the next toxic pesticide, the mortality being 86 and 84 % in 0.2 % solution against *L. beneficus* and *X. galactinus* respectively. The other pesticides were scarcely harmful in their conventional concentrations. To the field collected individuals, the tolerance was more inferior to the laboratory reared ones, and parathion, sumithion, dipterex and diazinone gave high mortality. The high sensitivity in the field collected individuals was partly attributed to the age structure of the field population. And this consideration was verified by the linear increase of the sensitivity to sumithion with the process of age in the laboratory reared population. The pesticidal sensitivity relating to the nutritional condition was also discussed. The pesticidal tolerance in the nymphal stage was higher in the last instar, exceeding that of the laboratory reared young adults.

When the 1st stage nymphs of both species were reared in the same cage, the nymphs of *X. galactinus* were always superior in growth and survival. The presence of *X. galactinus* adults gave serious effect on the survival of younger nymphs of *L. beneficus*. The mixed rearing of the adult *L. beneficus* and the nymphs of *X. galactinus* gave little effects on the nymphal growth. The mixed rearing of the adults of both species gave bad effect on the oviposition period and egg-production of *L. beneficus*, but little influence on *X. galactinus*.

A Braconid parasite of rice stem borer, *Apanteles chilonis* [Munakata], was always influenced by the presence of bugs. The piercing of the borer body by the bugs gave fatal effect on the survival of the wasp which was parasitizing the borer owing to the withering of the host insect. The bug preferred to attack the mature larvae of the wasp just crawled out of the borer body. Therefore the emergence of *Apanteles* wasps under the presence of bugs was very few.

The mass-production procedure of bugs was designed with the use of vermiculite and cracked corn. For the most efficient method, the bugs were reared with cracked corn medium provided with a piece of moist filter paper

for an oviposition substratum. The larvae of *T. castaneum* were the most suitable food, and the method of mass culture of *T. castaneum* was also discussed. The separation of bugs from the culture media was designed too.

The preying effect of bugs on the rice stem borer is evident when the bugs are in contact with the borer directly. Though the amount of prey consumed by the bugs decreased under the straw condition, the survival rate of the bugs rather increased. The preying activity was higher at 30°C than at 25°C. The illumination has little influence on the preying activity. Dry straw condition stimulated the activity than the moist condition of material. But, to keep the moderate moisture condition in the straw was indispensable to maintain the rate of high survival of the bugs. Nymphs were almost impossible to attack the freely crawling borers, but accessible to the borer when the latter was confined to the straw. When the adults and nymphs were present in the same container, the adult bugs attacked their nymphs instead of the borers. The preying activity of bugs was not significant against the borers in the paddy stubbles. Under such an environment the bugs preyed on some small creatures which were found in the stubbles or crawling or walking on the soil surface.

Acknowledgement

The author wishes to express his cordial thanks to Prof. K. Yasumatsu under whose direction and encouragement this work was performed. Many interesting ideas in connection with the planning of this work were suggested by him. In addition, the author is much obliged to him for the advice and constructive criticism he so willingly gave and for the careful help in the preparation of this manuscript.

The author has no words in expressing his hearty appreciation to Prof. S. Miyamoto, for many valuable advice and criticism especially in the field of systematics of the family Anthecoridae.

The author would like to express his thanks to Dr. Y. Hirashima and Dr. K. Yano for their constant warm hospitality in every possible way to perform this study. He is also greatly obliged to Dr. G. Funatsu of Biochemical Laboratory, Kyushu University for his help in the analysis of amino acids. For the field observation, the co-operation of Mr. M. Hata of Saga Paper Board Co. is greatly appreciated.

Further, the author extends his sincere thanks to many colleagues of the Entomological Laboratory and the Institute of Biological Control, Kyushu University for their helpful suggestion and valuable discussion during the course of this study.

Literature cited

- Albrecht, F. O. et al. 1958. Determination de la fertilité par l'effect de groups chez le criquet migrateur. Biol. Bull. 92 : 349-427. (in Watt 1960)
- Allee, W. C. 1934. Recent studies in mass physiology. Biol. Rev. 9 (1): 1-48.
- Anderson, N. H. 1962a. Bionomics of six species of *Anthocoris* in England. Trans. R. ent. Soc. Lond. 114 (3): 67-95.
- Anderson, N. H. 1962b. Growth and fecundity of *Anthocoris* spp. reared on various prey. Ent. exp. & appl. 5 (1) : 40-52.
- Andrewartha, H. G. and Birch, L. C. 1960. Some recent contributions to the study of the distribution and abundance of insects. Ann. Rev. Ent. 5: 219-42.
- Annand, P. N. 1942a. Report of the Chief of the Bureau of Entomology and Plant Quarantine. U.S. D. A. 19 (41): 120 pp.
- Annand, P. N. 1942b. Report of the Chief of the Bureau of Entomology and Plant Quarantine. U.S. D. A. 19 (41-42) : 60 pp.
- Anonym, 1942. Memoria de la Estación experimental agricola de La Molina correspondiente al año 1941. 6: 276 pp. Lima Minist. Peru. (in Rev. Appl. Ent. 1944, 32 (12): 428).
- Anonym. 1953. Proc. Rep. Saga Agri. Exp. Sta. in 1953 pp. 189-91.
- Anonym. 1954. Proc. Rep. Saga Agri. Exp. Sta. in 1954 pp. 182-5.
- Ascher, K. R.S. and Neri, I. 1961. Lipoid content and resistance in the housefly. Ent. exp. & appl. 4 (1): 7-19.
- Atallah, Y.H. and Newson, L. D. 1966. Ecological and nutritional studies on *Coleomegilla maculata* DeGeer. I. The development of an artificial diet and a laboratory rearing technique. Jour. Econ. Ent. 59 (5): 1173-9.
- Atallah, Y. H. and Killebrew, R. 1967. Ecological and nutritional studies on *Coleomegilla maculata* DeGeer. IV. Amino acid requirements of the adults determined by the use of C¹⁴-labeled acetate. Ann. Ent. Soc. Am. 60 (1): 186-8.
- Atkins, E. L., Jr. 1958. The western tussock moth, *Hemerocampa vetusta* (Bdr.) on citrus in southern California. Jour. Econ. Ent. 51 (6) : 762-5.
- Atkins, E. L., Jr. et al. 1957. The "omnivorous leaf roller" *Platynota stultana* on cotton in California; Nomenclature, life history and bionomics (Tortricidae). Ann. Ent. Soc. Am. 50 (3) : 251-9.
- Auclair, J. L. and Raulston, J. R. 1966. Feeding of *Lygus hesperus* on a chemically defined diet. Ann. Ent. Soc. Am. 59 (5) : 1016-7.
- Ballard, E. 1921. A preliminary note on *Triphleps tantilus* Mosch., an enemy of the pink worm. Agric. Jour. India 16 (5) : 571-3.
- Barber, G. W. 1936a. The corn earworm in southern Georgia. Bull. Geor. Exp. Sta. No. 192 (18) : 771.
- Barber, G. W. 1936b. *Orius insidiosus*, an important natural enemy of the corn earworm. U. S. D. A. Tech. Bull. No. 504, 24 pp.
- Barber, G. W. 1943. Oviposition habits of the earworm moth in relation to infestation in the ears and to control. Jour. Econ. Ent. 34 (4j): 611-8.
- Bartholomai, C. W. 1954. Predatism of European corn borer eggs by Arthropods. Jour. Econ. Ent. 47 (2): 295-9.
- Bartlett, B. R. 1963. The contact toxicity of some pesticide residues to hymenopterous parasites and coccinellid predators. Jour. Econ. Ent. 56 (5): 694-8.
- Bartlett, B. R. 1964. Integration of chemical and biological control. In DeBach (ed.),

- Biological control of insect pests and weeds. Pp. 489-511, Chapman and Halls, London.
- Recks, S. D. et al. 1949. Nutrition of the European corn borer, *Pyrausta nubilalis* (Hbn.). I. Development of a satisfactory purified diet for larval growth. Ann. Ent. Soc. Am. **42** (4) : 483-96.
- Beck, S. D. et al. 1958. Feeding and nutrition of the milkweed bug, *Oncopeltus fasciatus* (Dallas). Ann. Ent. Soc. Am. 51 (3) : 283-8.
- Beirne, B. P. 1967. The future of integrated controls. Mushi 39, Suppl. pp. 127-30.
- Bell, K. D. and Whitecomb, W. H. 1964. Field studies on egg predators of the bollworm, *Heliothis zea* (Boddie). Florida Ent. 47 (3): 171-80.
- Bess, H. A. 1967. Feasibility and problems of chemical control and biological control of rice stem borers. Mushi 39, Suppl. pp. 45-50.
- Blatchley, W. S. 1926. Heteroptera of true bugs of Eastern North America. Pp. 621-46. Nature Publ. Co., Indianapolis.
- Bodenheimer, F. 1921. Zur Kenntnis der Chrysanthemum-Wanzen sowie der durch sie hervorgerufenen Gallbildung. Zeitschr. Pflanzenkr. 31: 3-4.
- Bondar, G. 1924. A ferrugem na folha do algodoeiro. Correio-Agricola. 2 (6) :172-4. (in Rev. Appl. Ent. 1924, 12 (9) : 413).
- Boyce, J. M. 1946. The influence of fecundity and egg mortality on the population growth of *Tribolium confusum* Duval. Ecology 27 (4) : 290-302.
- Bracken, G. K. 1965. Effects of dietary components on fecundity of the parasitoid *Exeristes comstockii*. Canad. Ent. 97 (10): 1037-41.
- Bracken, G. K. 1966. Role of ten dietary vitamins on fecundity of the parasitoid *Exeristes comstockii*. Canad. Ent. 98 (9) : 918-22.
- Brönnimann, H. 1964. Rearing Anthocorids on an artificial medium. Tech. Bull. Commw. Inst. Biol. Contr. No. 4, pp. 147-50.
- Burke, H. R. 1959. Toxicity of several insecticides to two species of beneficial insects on cotton. Jour. Econ. Ent. 52 (4): 616-8.
- Busvine, J. R. 1951. Insects and Hygiene. P. 341. Methuen Co., London.
- Butler, E. A. 1923. Biology of the British Hemiptera-Heteroptera. Pp. 324-6. Witherby Co., Holborn, W. C.
- Butler, C. C. 1954. The method and importance of the recognition by a colony of honey bees of the presence of its queen. Trans. R. ent. Soc. Lond. 103 (1) : 11-29.
- Butler, G. D. 1966. Development of several predacious Hemiptera in relation to temperature. Jour. Econ. Ent. 59 (5): 1306-7.
- Carayon, J. 1953. Organ de Ribaga et fécondation hemocoelienne chez les *Xylocoris* du groupe *galactinus*. Comp. Rend. Acad. Sci. Paris 236: 1099-101.
- Carayon, J. et Steffan, J. R. 1959. Observations sur le regime alimentaire des *Orius* et particulièrement d'*Orius pallidicornis* (Reuterj. Cah. Nat. Bull. N. P. (n. s.) 15: 53-64.
- Carayon, J. 1961. Quelques remarques sur les Hémiptères-Hétéroptères: Leur importance comme insectes auxiliaires et les possibilites de leur utilisation dans la lute biologique. Entomophaga 6 (2): 133-41.
- Chalfant, R. B. and Mitchell, E. R. 1967. Some effects on food and substrate on oviposition of the spotted cucumber beetle. Jour. Econ. Ent. 60 (4) : 1010-2.
- Chapman, N. R. 1928. The quantitative analysis of environmental factors. Ecology 9(2) : 698-711.
- Chapman, R. N. and Whang, W. Y. 1934. An experimental analysis of the cause of population fluctuations. Science, N. Y. 8 (2074) : 297-8.

- Chauvin, R. 1946. Notes sur la physiologie comparée des Orthoptères. V. L'effect de groups et la croissance larvaire des Blattes, des Grillons et du Phaneroptère. Bull. Soc. 2001. Fr. 71 (1) : 39-48.
- China, W. E. 1926. A new species of *Triphleps* preying on the eggs of *Heliothis obsoleta* H.-S. in Queensland. Bull. Ent. Res. 16 (4j) : 361-2.
- China, W. E. and Miller, N. C. E. 1955. Check list of family and subfamily names in Hemiptera-Heteroptera. Ann. Mag. Nat. Hist. Lond. 12 (8) : 257-67.
- China, W. E. and Miller, N. C. E. 1959. Check list and key to the families and subfamilies of the Hemiptera-Heteroptera. Bull. Brit. Mus. Ent. 8 (1) : 1-45.
- Chu, Y. I. 1962a. Identification of sugar and amino acids on the haemolymph of lindane treated *Papilio*. Mem. Coll. Agr. Nation. Taiwan Univ. 6 (4) : 38-41.
- Chu, Y. I. 1962b. Quantitative determination of amino acid and sugars on the haemolymph of lindane treated *Papilio*. Mem. Coll. Agr. Nation. Taiwan Univ. 6 (4) : 42-6.
- Chumakova, B. M. 1962. Significance of individual food components for the vital activity of mature predatory and parasitic insects. Vop. Ekol. Kievsk. 8 : 133-4. (in Biol. Abstr, 1962, 45 (44j) : 502).
- Collyers, E. 1967. On the ecology of *Anthocoris nemorum* (L.). Proc. R. ent. Soc. Lond. Ser. (Aj), 42 (7-9) : 107-18.
- Corti, A. 1921. *Lyctocoris campestris* F. un Memico ignorato del Baco da Seta. Atti. Soc. Ital. Sci. Nat. in Milano 110 (1): 1-10. (in Rev. Appl. Ent. 1921, 9 (8): 404).
- Dadd, R. H. 1963. Feeding behavior and nutrition in grasshopper and locust. Adv. Insect Physiol. 1: 47-109.
- Davis, C. J. and Krauss, N. L. H. 1963. Recent introduction for biological control in Hawaii VIII. Proc. Hawaii. Ent. Soc. for 1962, 18 (2): 245-9.
- Davis, C. J. and Krauss, N. L. H. 1964. Recent introduction for biological control in Hawaii IX. Proc. Hawaii. Ent. Soc. for 1963, 18 (3): 391-7.
- Davis, C. J. and 'Krauss, N. L. H. 1965a. Recent introduction for biological control in Hawaii X. Proc. Hawaii. Ent. Soc. for 1964, 19 (1) : 87-90.
- Davis, C. J. and Krauss, N. L. H. 1965b. Recent introduction for biological control in Hawaii XI. Proc. Hawaii. Ent. Soc. for 1964, 19 (2): 201-7.
- DeBach, P. 1966. The comparative displacement and coexistence principles. Ann. Rev. Ent. 11: 183-212.
- DeGroot, A. P. 1953. Protein and amino acid requirements of the honeybee. Physiol. Comparata et Oecol. 3: 197-285.
- DeLoach, C. J., Jr. and Miyatake, Y. 1966. Seasonal abundance and degree of parasitism of the Asiatic rice borer in Fukuoka Prefecture, Japan. Mushi 39 (3): 31-46.
- Dempster, J. P. 1960. A quantitative study of the predators on the eggs and larvae of the broom beetle, *Phytodectet olivacea*, using the precipitin test. Jour. Anim. Ecol. 29 : 149-67.
- Dempster, J. P. 1963. The natural prey of three species of *Anthocoris* living on broom. Ent. exp. & appl. 6 (2): 149-55.
- Diakonoff, A. 1940. Remarks on a predatory bug. Ent. Med. Ned.-Ind. 35 (3): 55-6.
- Dicke, F. F. and Jarvis, J. L. 1962. The habits and seasonal abundance of *Orius insidiosus* (Say) on corn. Jour. Kansas Ent. Soc. 35 (3) : 339-44.
- Dimond, J. B. et al. 1956. The amino acids required for egg production in *Aedes aegypti*. Canad. Ent. 88 (1): 107-8.
- Doeksen, J. 1944. Een nieuwe beschadiging van maiskolven. Tij. Pl. Ziekt. 50 (4) : 107 to 8.

- Douglas, J. W. and Scott, J. 1865. The British Hemiptera-Heteroptera. Pp. 495-6. Ray Soc., London.
- DuPort, E. M. 1917. The eye-spotted bud moth (*Eucosma ocellana* Schiff.), Ninth Ann. Rept. Quebec Soc. Protec. Plants from Insect & Fungus Dis. 1916-7. Pp. 118-37.
- Durrant, A. 1955. Effects of time embryo formation on quantitative characters in *Drosophila*. Nature 175 (4456j) : 560-1.
- Earle, N. W. et al. 1967. Essential fatty acids in the diets of the boll weevil (*Anthonomus grandis* Boheman). Jour. Ins. Physiol. 13 (2): 187-200.
- Esaki, T. and Hashimoto, S. 1935. Report on the leafhoppers injurious to the rice plant and their natural enemies. No. 7, p. 23. Ent. Lab. Dept. Agr. Kyushu Imp. Univ. Fukuoka, Japan.
- Evans, D. E. 1962. The food requirements of *Phonoctonus nigrofasciatus* Stål (Hemiptera, Reduviidae j. Ent. exp. & appl. 5 (1) : 33-9.
- Ewing, K. and Ivy, E. E. 1943. Some factors influencing ball worm population and damage. Jour. Econ. Ent. 36 (4) : 602-6.
- Fast, P. G. 1964. Insect lipids: A review. Mem. Ent. Soc. Can. No. 37, 50 pp.
- Feir, D. 1963. Effects of rearing alone and in groups on the growth of the milkweed bug, *Oncopeltus fasciatus* (Lygaeidae). Ann. Ent. Soc. Am. 56 (3j) : 406-7.
- Fewks, D. W. 1960. The food requirements by weight of some British Nabidae. Ent. exp. & appl. 3 (3) : 231-7.
- Fieber, F. X. 1837. Weit. Beitr. 1: 107. (in Reuter 1884).
- Fieber, F. X. 1860. Die europäischen Hemiptera, Halbflügler, nach der analytischen Methode bearbeitet. 1860. P. 61, Wien. (in Butler 1923).
- Fletcher, R. K. and Thomas, F. L. 1943. Natural control of eggs and first instar larvae of *Heliothis armigera*. Jour. Econ. Ent. 36 (4j) : 557-60.
- Flint, W. P. 1918. Insect enemies of the chinch bug. Jour. Econ. Ent. 11 (5) : 415-9.
- Foster, S. W. and Jones, P. R. 1915. The life history and habits of the pear thrips in California. U. S. D. A. Bull. No. 173, 52 pp.
- Fraenkel, G. and Blewett, M. 1946. Linoleic acid, vitamin E and other fat-soluble substances in the nutrition of certain insects (*Ephestia kuehniella*, *E. elutella*, *E. cautella* and *Plodia interpunctella*). Jour. Exp. Biol. 22 (3-4) : 172-90.
- Fronk, W. D. 1947. The southern pine beetle, its life history. Tech. Bull. Va. Agr. Exp. Sta. No. 108. (in Rev. Appl. Ent. 1950, 38 (10): 415).
- Funasaki, G. Y. 1966. Studies on the life cycle and propagation technique of *Montanodoniola moraguesi* (Puton). Proc. Hawaii. Ent. Soc. 19 (2): 209-11.
- Garman, H. and Jewett, H. H. 1914. The life-history and habits of the corn-ear worm (*Chlorida obsoleta*). Ky. Agr. Exp. Sta. Bull. No. 187, pp. 511-91. (in Dicke 1962).
- Garman, P. and Townsend J. F. 1938. The European red mite and its control. Bull. Conn. Agr. Exp. Sta. No. 416, 6 pp.
- Gilliatt, F. 1937. Natural control of the grey-banded leaf roller *Eulia mariana*, in Nova Scotia orchards. Canad. Ent. 69 (7) : 145-6.
- Gilmour, D. 1961. The biochemistry of insects. Academic Press, New York.
- Golberg, L. and DeMeillon, B. 1948. The nutrition of the larvae of *Aedes aegypti* L.-IV. Protein and amino acid requirements. Biochem. Jour. 43 (3) : 379-87.
- Good, N. E. 1933. Biology of the flour beetles, *Tribolium confusum* Duval and *T. ferrugineum* Fabr. Jour. Agr. Res. 46 (4): 327-31.
- Good, N. E. 1936. The flour beetles of the genus *Tribolium*. Tech. Bull. U.S. D.A. No. 498, 57 pp.
- Gordon, H. T. 1961. Nutritional factors in insect resistance to chemicals. Ann. Rev.

Ent. 6. 27-54.

- Gratiwick, M. 1965. Laboratory studies of the relative toxicities of orchard insecticides to predatory insects. 52nd Rep. E. Malling Res. Sta. 1964, pp. 171-6.
- Green, C. D. 1964. The effect of crowding upon the fecundity of *Folsomia candida* (William) var. **distincta** (Bagnall). Ent. exp. & appl. 7 (1) : 62-70.
- Greenberg, J. 1951. Some nutritional requirements of adult mosquitoes (*Aedes aegypti*) for oviposition. Jour. Nutr. 43 (1): 27-35.
- Gyotoku, N. 1960. [On the natural enemies of *Chilo suppressalis* Walker]. Proc. Assoc. Pl. Prot. Kyushu 6: 1-3.
- Hagen, K. S. 1950. Fecundity of **Chrysopa californica** as affected by synthetic foods. Jour. Econ. Ent. 43 (1) : 101-4.
- Hagen, K. S. 1958. Honeydew as an adult fruit fly diet affecting reproduction. Proc. 10th Intern. Congr. Ent. 3: 25-30.
- Hagen, K. S. and Tassen, R. L. 1965. A method of providing artificial diets to **Chrysopa** larvae. Jour. Econ. Ent. 58 (5): 999-1000.
- Hagen, K. S. and Tassen R. L. 1966. A method of coating droplets of artificial diet with paraffin for feeding **Chrysopa** larvae. (in Ecology of aphidophagous insects, pp. 89-90. Academic Pub. House. Prague).
- Hahn, C. W. 1835. Die Wanzenartigen Insecten 3: 19-20.
- Hall, D. W. 1951. Observations on the distribution, habits and life history of the bug, **Piezostethus galactinus**. Ent. mon. Mag. 86 (14th Ser. No. 121) : 45-52.
- Hambleton, E. J. 1944. **Heliothis virescens** as a pest of cotton, with notes on host plants in Peru. Jour. Econ. Ent. 37 (5): 660-6.
- Hanison, F. P. 1960. Corn earworm oviposition and the effects of DDT on the egg predator complex in corn silk. Jour. Econ. Ent. 53 (6): 1088-94.
- Harein, P. K. and Soderstrom, E. L. 1966. Coleoptera infesting stored products. (in Smith (ed.), Insect colonization and mass production, pp. 247-53. Acad. Press, N. Y. and London).
- Harlow, P. M. 1956. A study of ovarian development and its relation to adult nutrition in the blowfly, **Protophormia terrae-novae**. Jour. Exp. Biol. 33 (4) : 777-97.
- Hassan, A. A. G. et al. 1960. Analysis of the relative abundance of some beneficial insects on cotton plants treated by certain insecticides. Bull. Soc. Ent. Egypte 44 : 179-84.
- Herbst, J. F. W. 1797. Natursystem aller bekannten in- und ausländischen Insecten, als eine Fortsetzung der vom buff onschen Naturgeschichte. Der Käfer, pts. 7, p. 282.
- Herne, D. H. C. and Putman, W. L. 1966. Toxicity of some pesticides to predacious arthropods in Ontario peach orchard. Canad. Ent. 98 (9): 936-42.
- Herring, J. L. 1967. Heteroptera, Anthocoridae. Insect of Micronesia 7 (8) :391-414.
- Hilchey, J. D. 1953. Studies on the quantitative requirement of **Blattella germanica** (L.) for amino acids under aseptic condition. Contri. Boyce Thompson Inst. 17 (3) : 203-19.
- Hill, R. E. 1945. Effect of DDT and other insecticides on several species of potato insects. Res. Bull. Nebr. Agr. Exp. Sta. No. 138. (in Rev. Appl. Ent. 1946, 34 (7) :223).
- Hill, A. R. 1957. The biology of **Anthocoris nemorum** (L.) in Scotland. Trans. R. ent. Soc. Lond. 113 (3) : 41-54.
- Hill, A. R. 1965. The bionomics and ecology of **Anthocoris confusus** Reuter in Scotland. I. The adult and egg production. Trans. Soc. Brit. Ent. 16 (9): 245-56.
- Hinton, H. E. 1948. A synopsis of the genus **Tribolium** Macleay with some remarks on the evolution of its species groups. Bull. Ent. Res. 39 (1) : 13-56.
- Hinton, T. et al. 1951. Amino acid and growth factors in a chemically defined medium

- for *Drosophila*. *Physiol.* 2001. 24 (4) : 335-53.
- Hirano, C. 1963. Effect of dietary unsaturated fatty acids on the growth of larvae of *Chilo suppressalis*. *Japanese Jour. Appl. Ent.* 2001. 7 (1) : 59-62.
- Hiura, I. 1957. Description of a new species of Japanese Anthocoridae, and its biology. *Sci. Bull. Fac. Agr. Kyushu Univ.* 16 (1): 31-9.
- Hiura, I. 1966. Contribution to the knowledge of Anthocoridae from Japan and its adjacent territories (3). *Bull. Osaka Mus. Nat. Hist.* 19: 29-37,
- Ho, F. K. 1967. Identification of *Tribolium* larvae by their setal characteristics. *Ann. Ent. Soc. Am.* 60 (4) : 729-32.
- House, H. L. 1954. Nutritional studies with *Pseudosarcophaga affinis* (Fall.), a Dipterous parasite of the spruce bud worm, *Choristoneura fumiferana* (Clem.). III. Effects of nineteen amino acids on growth. *Canad. Jour.* 2001. 32 (4) : 351-9.
- House, H. L. 1958. Nutritional requirements of insects associated with animal parasitism. *Exp. Parasitol.* 7: 555-609,
- House, H. L. 1962. Insects nutrition. *Ann. Rev. Biochem.* 31: 653-72.
- House, H. L. 1965. Insect Nutrition. (in *The physiology of Insects*, 2: 769-814. Acad. Press., N. Y. and London).
- House, H. L. 1966. Effects of vitamins E and A on growth and development, and the necessity of vitamin E for reproduction in the parasitoid *Agris affinis* (Fallen). *Jour. Ins. Physiol.* 12 (4) : 409-17.
- Howe, R. W. 1952. Entomological problems of food storage in northern Nigeria. *Bull. Ent. Res.* 43 (1): 111-44.
- Howe, R. W. 1956. The effect of temperature and humidity on the rate of development and mortality of *Tribolium castaneum*. *Ann. Appl. Biol.* 44 (2): 356-68.
- Howe, R. W. 1960. The effects of temperature and humidity on the rate of development and the mortality of *Tribolium confusum*. *Ann. Appl. Biol.* 48 (2): 363-76.
- Iglinsky, W. and Rainwater, C. F. 1950. *Orius insidiosus*, as enemy of a spider mite on cotton. *Jour. Econ. Ent.* 43 (4): 567-8.
- Ishihara, T. 1941. Hemiptera-Heteroptera as the natural enemies for controlling the pests. *Botany and Zoology.* 9 (9): 223-9.
- Ishii, S. and Hirano, C. 1955. Qualitative studies on the essential amino acids for the growth of the larvae of the rice stem borer, under aseptic conditions. *Bull. Nation. Inst. Agr. Sci. Japan* C5: 35-48.
- Ito, Y. et al. 1960. Natural mortality of the common cabbage butterfly *Pieris rapae crucivora* Boisduval with consideration on the factors affecting it. *Japanese Jour. Appl. Ent.* 2001. 4 (1): 1-10.
- Johansen, C. and Brekey, E. P. 1949. Insecticides tested against the willamette mites on red raps-berries. *Jour. Econ. Ent.* 42 (3): 562-3.
- Johnson, C. G. 1960. The relation of weight of food ingested to increase in body weight during growth in the bed bug, *Cimex lecturarius*. *Ent. exp. & appl.* 3 (3): 238-40.
- Kamel, M. 1951. The biological control of the cotton leaf-worm (*Prodenia litura* F.) in Egypt. *Bull. Soc. Fouad* 1^{er} Ent. 35^e: 221-70.
- Kasting, R. and McGinnis, A. J. 1959. Nutrition of the pale western cutworm, *Agrotis orthogonia* Morr. *Canad. Jour.* 2001. 37 (3): 259-66.
- Kelly, E. O. A. 1915. A new wheat thrips. *Jour. Agr. Res.* 4 (3): 219-24.
- Kelton, L. A. 1967. Synopsis of the genus *Lyctocoris* in North America and description of a new species from Quebec. *Canad. Ent.* 99 (8) : 807-14.
- Khalifa, A. and Badawy, A. 1955a. The effect of nutrition on the biology of *Tribolium*

- confusum** Duv., **Tribolium castaneum** Hbst. and **Latheticus oryzae** Waterh. Bull. Soc. Ent. Egypte 39: 337-50.
- Khalifa, A. and Badawy, A. 1955b. Biological studies on **Tribolium confusum** Duv., **Tribolium castaneum** Hbst. and **Latheticus oryzae** Waterh. Bull. Soc. Ent. Egypte 39: 351-73.
- Kirkaldy, G. W. 1906. List of the genera of the **Pagiopodous** Hemiptera-Heteroptera, with their type species, from 1758 to 1904 (and also of the aquatic and semi-aquatic **Trochalopoda**). Trans. Am. Ent. Soc. 32: 117-56.
- Lewis, D. J. 1958. Hemiptera of medical interest in the Sudan Republic. Proc. R. ent. Soc. Lond. 33 (1-3) : 43-7.
- Lingren, P. D. and Ridgway, R. L. 1967. Toxicity of five insecticides to several insect predators. Jour. Econ. Ent. 60 (6): 1939-41.
- Loher, W. 1961. The chemical acceleration of the maturation process and its hormonal control in the male of the desert locust. Proc. R. Soc. (B) 153: 380-97.
- Lord, F. T. 1949. The influence of sprays programs on the fauna of apple orchards in Nova Scotia, III. Mites and their predators. Canad. Ent. 81 (8-9) :202-14.
- Luginbill, P. 1928. The fall armyworm. Tech. Bull. U. S. D. A. No. 34, 91 pp.
- Lüscher, M. 1956. Hemmende und fördernde Faktoren bei der Entstehung von Ersatzgeschlechtstieren bei der Termite, **Kaloterme flavicollis** Fabr. Rev. Suisse Zool. 63: 261-7.
- MacLagen, D. S. 1932. An ecological study of the lucerne flea (**Smynturus viridisj**. Bull. Ent. Res. 23 (1-2) : 101-45.
- Malenotti, E. 1926. Impressioni sull'andamento della decorsa campagna bacologica. L'Italia Agricola (in Rev. Appl. Ent. 1926. 14 (1): 36).
- Malloch, J. R. 1916. **Triphleps insidiosus** sucking blood. Ent. News 27 (5) : 200.
- Marcovitch S. and Stanley, W. W. 1941. The tomato fruit worm in Tennessee. Bull, Tenn. Agr. Exp. Sta. No. 174. 18pp.
- Marshall, G. E. 1930. Some observation on **Orius insidiosus** (Say). Jour. Kansas Ent. Soc. 3 (2) : 29-32.
- Martini, E. 1946. Lehrbuch der medizinischen Entomologie. Pp. 145-6. Jena.
- Massee, A. H. and Steer, W. 1926. Tar distillate washes and red spiders. Jour. Minist. Agric. 36 : 253-7.
- Matheson, R. 1950. Medical entomology. 2nd Ed. P. 188. Comstock Publ. Co., Ithaca, New York.
- Matsumura, S. 1920. [Applied Entomology.] p. 427. Keiseisha Co., Tokyo.
- McFarlane, J. E. 1962. A comparison of the growth of the house cricket (Orthoptera : Gryllidae) reared singly and in groups. Canad. Jour. 2001. 40 (4): 559-60.
- McFarlane, J. E. and Henneberry, G. O. 1965. Inhibition of the growth of an insect by fatty acids. Jour. Ins. Physiol. 11 (9): 1247-52.
- McGregor, E. A. 1913. The red spider on cotton. U. S. Bureau of Ent. Circ. No. 172, 22 PP.
- McGregor, W. S. 1942. **Orius insidiosus**, a predator on cotton insects in Western Texas. Jour. Econ. Ent. 35 (3): 454-5.
- McLachlan, R. 1879. P. E. Soc. pp. 93-4. (in 2001. Rec. 1879. 16: 237).
- McLachlan, R. 1880. P. E. Soc. pp. 29-30. (in 2001. Rec. 1880. 17: 230).
- McMullen, R. D. and Jong, C. 1967. The influence of three insecticides on predation of the pear Psylla, **Psylla pyricola**. Canad. Ent. 99 (12) : 1292-7.
- Meikle, J. E.S. and McFarlane, J. E. 1965. The role of lipid in the nutrition of the house cricket, **Acheta domesticus** L. Canad. Jour. 2001. 43 (1): 87-98.

- Mertz, D. B. 1961. Identification of *Tribolium* species by pupal characteristics. Ecology 42 (4) : 811-8.
- Metcalf, R. L. and Newell, I. M. 1962. Investigation of the biochromes of mites. Ann. Ent. Soc. Am. 55 (3) : 350-3.
- Michal, K. 1931. Die Beziehung der Populationsdichte zum Lebensoptimum aus der Zahlenverhältnis der Geschlechter bei Mehworm und Stubenfliege. Biologia Gen. 7 (3) : 631-46.
- Michelbacher, A. E. et al. 1952. Mites on melons in northern California. Jour. Econ. Ent. 45 (3) : 365-70.
- Misra, C. S. 1924. Observations on some Rhynchota from Northern India. Rep. Proc. 5th Ent. Meeting, Pusa(1923), p. 300.
- Mitchell, R. G. and Wright, K. H. 1967. Foreign predator introductions for control of the balsam woolly aphid in the Pacific Northwest. Jour. Econ. Ent. 60 (1): 140-7.
- Miyamoto, S. 1957. List of ovariole numbers in Japanese Heteroptera. Sieboldia 2 (1) : 69-82.
- Miyamoto, S. 1961. Comparative morphology of alimentary organs of Heteroptera, with the phylogenic consideration. Sieboldia 2 (4) : 198-259.
- Miyamoto, S. and Lee, C. E. 1966. Heteroptera of Quelpart Island. Sieboldia 3 (4): 313-411.
- Mizukami, T. and Oho, N. 1955. *Euspudaeus* sp. as a predator of *Chilo suppressalis*. Plant Prot. 9 (1): 10 figs.
- Morkzecki, Z. 1923. Sprawozdanie z. dzialalnosci Zakładów ochrony Lasu i Entomologii w Skierniewicach. Ecole. Sup. Agr. à Varsovie L(1922-23): 32. (in Rev. Appl. Ent. 1924, 12 (3): 105).
- Morley, C. 1905. The Hemiptera of Suffolk. (in Butler 1923).
- Myers, J. G. 1929. Facultative blood-sucking in phytophagous Hemiptera. Parasitol. 21 (4) : 472-80.
- Nickel, J. L. 1964. Biological control of rice stem borer: a feasibility study. Tech. Bull. Intern. Rice Res. Inst. 2 : i-iii, 1-111.
- Norris, M. J. 1950. Reproduction in the African migratory locust (*Locusta migratoria migratorioides* R. & F.) in relation to density and phase. Anti-locust Bull. No. 6, 47 pp.
- Norris, M. J. 1952. Reproduction in the desert locust (*Schistocerca gregaria* Forsk.) in relation to density and phase. Anti-locust Bull. No. 13, 49 pp.
- Norris, M. J. 1954. Sexual maturation in the desert locust (*Schistocerca gregaria* Forsk.) with special reference to the effects of grouping. Anti-locust Bull. No. 18, 44 pp.
- Norris, M. J. 1959. Reproduction in the red locust (*Nomadacris septemfasciana* Serville) in the laboratory. Anti-locust Bull. No. 38, 46 pp.
- Nutting, W. L. and Gerhardt, P. D. 1964. A study of the Khapra beetle, *Trogoderma granarium*, in commercial grain storage in Southern Arizona. Jour. Econ. Ent. 57 (3) : 305-14.
- Ogilvie, L. 1927. Report of the plant pathologist for the year 1926. Rep. Dept. Agr. Bermuda 1926, pp. 35-41. (in Rev. Appl. Ent. 1928, 16 (1): 17).
- Oho, N. 1955a. On the *Euspudaeus* sp., as a parasite of the rice stem borer (*Chilo suppressalis* Walker). Bull. Saga Agr. Exp. Sta. 1 : 45-6.
- Oho, N. 1955b. On the *Euspudaeus* sp. as a predator of the rice stem borer. Proc. Assoc. Pl. Prot. Kyushu 34: 222-4.
- Oho, N. 1955c. On the *Euspudaeus* sp. as a parasite of the rice stem borer, I. The food habits and life history. Proc. Assoc. Pl. Prot. Kyushu 16: 59-61.

- Oku, T. and Kobayashi, T. 1966. Influence of the predation of *Orius* sp. on the aphid population in a soy-bean field. Japanese Jour. Appl. Zool. 10 (2) : 89-94.
- Orr, C. W. H. 1964. The influence of nutritional and hormonal factors on **egg** development in the blow fly. Jour. Ins. Physiol. 10 (1) : 53-64.
- Ouchi, Y. 1967. [Rearing of *Nephotettix cincticeps* by artificial diets] .Rep. Symp. 1967 Meeting, Japanese Soc. Appl. Ent. Zool. p. 19.
- Pant, N. C. and Uberoi, N. K. 1958. On the carbohydrate utilization by the larvae of *Trogoderma granarium*. Experientia 14 (2) : 71-2.
- Park, T. 1932. Studies in population physiology. The relation of numbers to initial population growth in the flour beetle *Tribolium confusum* Duval. Ecology 13 (2): 172-81.
- Park, T. et al. 1965. Cannibalistic predation in populations of flour beetle. Physiol. Zool. 38 (3) : 289-321.
- Parson, E. S. and Ullyett, G. C. 1934. Investigations on the control of the American red boll worms of cotton in South Africa. Bull. Ent. Res. 25 (3): 349-81.
- Pearl, R. and Parker, S. L. 1922. On the influences of density of population upon the rate of reproduction in *Drosophila*. Proc. Nat. Acad. Sci. U.S. A. 8: 212-9.
- Pearl, R. et al. 1927. Experimental studies on the duration of life. XI. Density of population and life duration in *Drosophila*. Amer. Nat. 61 (675) : 289-318.
- Pearl, R. 1932. The influence of density of population upon egg production in *Drosophila melanogaster*. Jour. Exp. Zool. 63 (1) : 57-84.
- Pettit, L. C. 1940. The effect of isolation on growth in the cockroach *Blattella germanica*. Ent. News 51 (10): 293.
- Phillips, W. J. and King, K. M. 1923. The corn ear-worm: its ravages on field corn and suggestion for control. U.S. D. A. Farmer's Bull. No. 1310, 17 pp.
- Phillips, W. J. and Barber, G. W. 1933. Egg laying habits and fate of eggs of the corn ear-worm moth and factors affecting them. Tech. Bull. Va. Agr. Exp. Sta. No. 47, 34 pp.
- Pickett, C. and Friend, W. G. 1965. The nutritionally essential amino acids of *Rhodnius prolixus* (Stål) determined with glucose-U-¹⁴C. Jour. Ins. Physiol. 11 (12) : 1617-23.
- Piez, K. A. and Morris, L. 1960. A modified procedure for the automatic analysis of amino acids. Anal. Biochem. 1 (1) : 187-201.
- Poppius, B. 1909. Beiträge zur Kenntnis der Anthocoriden. Act. Soc. Sci. Fenn. 37 (9) : 1-43.
- Prohaska, K. 1923. Beiträge zur Kenntnis der Hemiptera Kärtens. Carinthia 2: 32-101 (in Rev. Appl. Ent. 1925, 13 (7): 311).
- Putman, W. L. 1964. Occurrence and food of some Coccinellids in Ontario peach orchards. Canad. Ent. 96 (9) : 1149-55.
- Putman, W. L. 1965. Paper chromatography to detect predation on mites. Canad. Ent. 97 (4): 435-41.
- Qureshi, A. H. 1966?, A simple technique for rearing larvae of *Tribolium castaneum* in the laboratory. Rep. Niger. Stored Prod. Res. Inst. 1965: 105-6. (in Rev. Appl. Ent. 1967, 55 (7): 393).
- Rainwater, C. F. 1951. Seasonal decline in the effectiveness of certain insecticides against boll weevil. Jour. Econ. Ent. 44 (6): 971-4.
- Rao, B. R. S. and Chawla, S. S. 1964. A catalogue of Hymenopterous parasites of rice stem borer. Indian Jour. Ent. 26 (3) : 332-44.
- Rasso, S. C. and Fraenkel, G. 1954. The food requirements of the adult female blow-fly,

- Phormia regina*, in relation to ovarian development. Ann. Ent. Soc. Am. 47 (4) : 636-45.
- Reiser, R. et al. 1953. Variations in lipid content of the boll weevil and seasonal variation in its resistance to insecticides. Jour. Econ. Ent. 46 (2): 337-40.
- Reuter, O. M. 1871a. Skandnaviens och Finlands Acanthiider beskrifne. Öfvers, K. Vetensk-Akad. Förh. 28 : 403-29.
- Reuter, O. M. 1871b. Acanthiidae americanae, descriptae. Öfvers, K. Vetensk-Akad. Förh. 28: 557-67.
- Reuter, O. M. 1884. Monographia Anthocoridarum orbis terrestris. Act Soc. Sci. Fenni. 14 : 555-758.
- Reuter, O. M. 1908. Charakteristik und Entwicklungsgeschichte der Hemiptera Fauna der palaearktischen Coniferen. (in Butler 1923 j .
- Richards, G. and Kolderie, M. Q. 1957. Variation in weight, developmental rate and hatchability of *Oncopeltus* eggs as a function of the mother's age. Ent. News 68 (3): 57-64.
- Richardson, C. H. et. al. 1931. The determination of the blood volume of insect larvae. Ann. Ent. Soc. Am. 24 (3) : 503-7.
- Richardson, H. H. and Nelson, R. H. 1933. Field control of the gladiolus thrips (*Taeniothrips gladioli*). Jour. Econ. Ent. 26 (3): 546-54.
- Riley, W. A. and Johannsen, O. A. 1938. Medical Entomology. P. 146. McGraw-Hill Book Co., New York.
- Rock, G. C. et al. 1965. Studies of the fatty acid requirements of *Argyrotaenia velutinana* (Walker). Jour. Ins. Physiol. 11 (1): 91-101.
- Rock, G. C. 1967. Aseptic rearing of the codling moth on synthetic diets: Ascorbic acid and fatty acid requirements. Jour. Econ. Ent. 60 (4): 1002-5.
- Sands, J. H. 1951. The biology of some British Anthocoridae. (M. Sc. thesis, London Univ.), (in Anderson 1962a).
- Sang, J. H. 1956. The quantitative nutritional requirements of *Drosophila melanogaster*. Jour. Exp. Biol. 33 (1): 45-72.
- Sang, J. H. and King, R. C. 1961. Nutritional requirements of axenically-cultured *Drosophila melanogaster*. Jour. Exp. Biol. 38 (4) : 793-809.
- Saunders, J. P. and Kruger, H. 1957. A rapid method for counting larvae of the confused flour beetle. Jour. Econ. Ent. 50 (5): 693.
- Scheel, C. A. et al. 1957. Nutrition of plant sucking Hemiptera. Science 125 (3245) : 444-5.
- Scheel, C. A. et al. 1958. Feeding and nutrition of certain Hemiptera. Proc. 10th Intern. Congr. Entomol., Montreal 1956, 2: 303-8.
- Shiraki, T. 1958. Sanitary insects. Pp. 186-7. Hokuryukan, Tokyo.
- Simón, F. J. E. 1954. Algunas experiencias eu el control de las plagas del algodónero: más intercalado e insecticidas sistémicos. Int. Ectac. Exp. Agr. La Molinar No. 90 (22): 7. (in Rev. Appl. Ent. 1956, 44(3) : 97).
- Sing, K.R.P. and Brown, A. W. A. 1957. Nutritional requirement of *Aedes aegypti* L. Jour. Ins. Physiol. 1 (3): 199-220.
- Sinha, R. N. 1961. Insects and mites associated with hot spots in farm stored grains. Canad. Ent. 93 (8.): 609-21.
- Sinha, B. P. and Mookhewee, P. B. 1967. Studies on the effect of airtightness on the development of *Tribolium castaneum* in wheat and wheat flour. Indian Jour. Ent. 29 (1j) : 79-83.
- Slama, K. 1962. The juvenile hormone-like effects of fatty acids, fatty alcohols and

- some other components in insect metamorphosis. *Acta Soc. Ent. Cechoslov*, 59 (4) : 323-40.
- Smirnoff, W. A. 1958. An artificial diet for rearing Coccinellid beetles. *Canad. Ent.* 90 (9) : 563-5.
- Smith, B. C. 1960. A technique for rearing Coccinellid beetles on dry foods and influence of various pollens on the development of *Coleomegilla maculata*. *Canad. Jour. Zool.* 38 (6): 1047-9.
- Smith, B. C. 1965. Effects of food on the longevity, fecundity and development of adult Coccinellids. *Canad. Ent.* 97 (9) : 910-9.
- Sonleither, F. J. 1961. Factors affecting egg cannibalism and fecundity in populations of adult *Tribolium castaneum*. *Physiol. Zool.* 34 (3) : 233-55.
- Southwood, T. R. E. and Leston, D. 1959. Land & water bug of British Island. Pp. 185-6. New York, Fredk. Warne & Co. Ltd., London.
- Stanley, J. 1962. The Tribolium-trap. *Canad. Jour. Zool.* 40 (8): 1313.
- Stern, V. M. et al. 1959. The integration of chemical and biological control of the spotted alfalfa aphid. I. The integrated control concept. *Hilgardia* 29 (2): 81-101.
- Strong, L. C. 1954. Parental age and characteristics of the offspring. *Ann. N. Y. Acad. Sci.* 57: 451-614.
- Stultz, H. T. 1955. The influence of spray programs on the fauna of apple orchards in Nova Scotia. VIII. Natural enemies of the eye-spotted bud moth, *Spilonota ocellana* (D.&S.). *Canad. Ent.* 87 (Zj) : 79-85.
- Stys, P. and Daniel, M. 1957. *Lycocoris campestris* (F.), as a human facultative ectoparasite. *Acta Soc. Ent. Cechoslov.* 54 (1): 88-97.
- Sundby, R. A. 1967. Influence of food on the fecundity of *Chrysopa carnea* Stephens (Neuroptera, Chrysopidae). *Entomophaga* 12 (5) : 475-9.
- Sweetman, H. L. 1958. Successful biological Control against animals. *Proc. 10th Intern. Congr. Ent. Montreal, 1956*, 4: 449-56.
- Tanaka, M. and Maeda, Y. 1965. Rearing of some predacious Coccinellid beetles by the artificial diets. *Bull. Hort. Res. Sta. Japan, Ser. D.* 3: 17-35.
- Tateishi, I. 1962. [On the parasitization and sex ratio of *Apanteles flavipes* Cameron]. *Proc. Assoc. Pl. Prot. Kyushu* 8: 26-9.
- Tawfik, M. F. S. and Nagui, A. 1965. The biology of *Montandoniola moraguesi* Puton, a predator of *Gynaikothrips ficorum* Marchal in Egypt. *Bull. Soc. Ent. Egypte* 49 : 181-200.
- Theobald, F. V. 1895. Notes on the needle nosed hop bugs. *J. S.-E. Agr. Coll. Wye* 2 : 11-6. (in Hill 1957).
- Thompson, W. R. 1951. The specificity of host relations in predacious insects. *Canad. Ent.* 83 (10) : 262-9.
- Torre-Bueno, J. R. 1931. Biting bugs. *Bull. Brookl. Ent. Soc.* 26 (4) : 176.
- Trägårdh, I. 1914. Skogsentomogiska bidrag, 1-5. *Ent. Tidsk.* 3-4: 188-209.
- Trager, W. 1953. Nutrition. *In* Roeder (ed.), *Insect Physiology*, pp. 350-86. Wiley, New York.
- Tucker E. S. 1911. Random notes on entomological field work. *Canad. Ent.* 43 (1) : 22-32.
- Turner, E. C. 1967. Effect of diet on longevity and fecundity of laboratory-reared face flies. *Jour. Econ. Ent.* 60 (3): 857-60.
- Tuttle, D. M. et al. 1961. The cotton leaf perforator and its control in Arizona. *Jour. Econ. Ent.* 54 (1) : 67-70.

- Underhill, G. W. 1926. Studies on the potato tuber moth during the winter of 1925-6, Bull. Va. Agr. Exp. Sta. No. 251, 21 pp.
- van den Bosch, R. and Stern, V. M. 1962. The integration of chemical and biological control of arthropod pests. Ann. Rev. Ent. 7: 367-86.
- van Duzee, E. P. 1916. Check list of the Hemiptera (excluding the Aphididae, Aleurodidae and Coccidae) of America North of Mexico. Pp. 288-97. New York.
- van Poetran, N. 1930. Verslag over de Werkzaamheden van den Plantenziekten kundigen Dienst in het Jaar 1926. Versl. Plantenziektenk. Dienst in het Jaar 1929. No. 62, 342 pp. (in Rev. Appl. Ent. 1931, 19 (5) : 242j.
- Vanderzant, E. S. 1957. Growth and reproduction of the pink bollworm on an amino acid medium. Jour. Econ. Ent. 50 (2) : 219-21.
- Vanderzant, E. S. 1958. The amino acid requirements of the pink bollworm. Jour. Econ. Ent. 51 (3) : 309-11.
- Vanderzant, E. S. 1967. Rearing *Lygus* bugs on artificial diets. Jour. Econ. Ent. 60 (3) : 813-6.
- Veitch, R. 1927. The corn ear-worm on tomatoes. Queensland Agr. Jour. 28 (6) : 585-8.
- Viswanathan, C. V. and Meera Bai, B. 1962. Application of reversed phase circular paper chromatography to the analysis of higher fatty acid. Jour. Chromatog. 7 (4) : 507-15.
- Wagner, E. 1955. *Lyctocoris nidicola* n.sp., eine neue Anthocoriden Art aus Finnland (Hem., Het.). Notul. Ent. 35 (2) : 60-Z.
- Watson, T. K. and Widde, W. H. A. 1963. Laboratory and field observations on two predators of the pear psylla in British Columbia. Canad. Ent. 95 (4) : 435-8.
- Watt, K. E. F. 1960. The effect of population density on fecundity of insects. Canad. Ent. 92 (9) : 674-95.
- Weber, P. W. 1953. Recent liberation of beneficial insects in Hawaii. II. Proc. Hawaii. Ent. Soc. for 1952, 15 (1) : 127-30.
- Wene, G. P. and Sheets, L. W. 1962. Relationship of predatory and injurious insects in cotton field in the Salt River area of Arizona. Jour. Econ. Ent. 55 (3) : 395-8.
- Wharton, D. R. A. et al. 1967. Population density, survival, growth and development of the American cockroach. Jour. Ins. Physiol. 13 (5) : 699-716.
- Wigglesworth, V. B. 1960. Nutrition and reproduction in insect. Proc. Nutrition Soc. 19 : 18-23.
- Wilde, W. H. A. 1962. The pear psylla in British Columbia. Canad. Dept. Agr. Publ. (in Watson and Widde 1963).
- Willcocks, F. C. 1916. The pink bollworm, the insect and related pests of Egypte. Sul-tanic Agr. Soc. 1(1): 339 pp.
- Wille, J. E. 1951. Biological control of certain cotton insects and the application of new insecticides in Peru. Jour. Econ. Ent. 44 (1):13-8.
- Willis, E. R. et al. 1958. Observation on reproduction and development in cockroaches. Ann. Ent. Soc. Am. 51 (1) : 53-69.
- Wilson, H. F. and Moznette, G. F. 1915. The bud moth *Tomotocera ocellana* Schiff. Second biennial crop pest and horticultural report 1913-1914. Oregon Agr. Exp. Sta, pp. 102-8.
- Winburn, 'I'. F. and Painter, R. H. 1932. Insect enemies of the corn-worm. Jour. Kansas Ent. Soc. 5 (1): 1-28.
- Woodward, T. E. 1951. A case of persistent attacks on a human by *Lyctocoris campestris* (F.). Ent. mon. Mag. 86 (14th Ser. No. 121): 44.

- Yasumatsu, K. 1964. The possible control of rice stem borers by the use of natural enemies. **In:** The maior insect pests of the rice plant. Pp. 431-442. The Johns Hopkins Press, Baltimore.
- Yasumatsu, K. and Watanabe, C. 1964. A tentative catalogue of insect natural enemies of injurious insect in Japan. 1 : 6-7. Ent. Lab, Kyushu Univ., Japan.
- Yasumatsu, K. 1967. Distribution and bionomics of natural enemies of rice stem borers. **Mushi** 39, Suppl. : 33-44.
- Yasumatsu, K. and Torii, T. 1968. Impact of parasites, predators and diseases on rice pests. **Ann. Rev. Ent.** 13 : 295-324.
- Young, R. G. 1964. Lipids of the larvae of the greater wax moth, *Galleriamellonella*. **Ann. Ent. Soc. Am.** 57 (3) : 321-4.
- Young, R. G. 1967. Fatty acids of some Arthropods. **Mem. Cornell Univ. Agr. Exp. Sta.** No. 401, 14 pp.
- Yun, Y. M. and Ruppel, R. F. 1964. Toxicity of insecticides to a coccinellid predator of the cereal leaf beetle. **Jour. Econ. Ent.** 57 (6): 853-7.
- Zaher, M. A. and Long, D. B. 1959. Some effects of larval population density on the biology of *Pieris brassicae* L. and *Plusia gamma* L. **Proc. R. ent. Soc. Lond. (A)** 34 (7-9) : 97-109.