Effect of Virogenic Hydroxylamine-Metabolites on Deoxyribonucleic Acid and RNA-Polymerase

Omura, Hirohisa

Food Chemistry Institute, Department of Food Science and Technology, Faculty of Agriculture,
Kyushu University

Iiyama, Satoru

Food Chemistry Institute, Department of Food Science and Technology, Faculty of Agriculture,
Kyushu University

Fujii, Takato

Food Chemistry Institute, Department of Food Science and Technology, Faculty of Agriculture,
Kyushu University

Yamafuji, Kazuo

Food Chemistry Institute, Department of Food Science and Technology, Faculty of Agriculture,
Kyushu University

<https://doi.org/10.5109/22829>

出版情報：九州大学大学院農学研究院紀要. 17 (2), pp.181-186, 1973-03. Kyushu University
バージョン：
権利関係：

Effect of Virogenic Hydroxylamine-Metabolites on Deoxyribonucleic Acid and RNA-Polymerase

Hirohisa Omura, Satoru Iiyama, Takato Fujii
and
Kazuo Yamafuji

Food Chemistry Institute, Department of Food Science and Technology,
Faculty of Agriculture, Kyushu University, Fukuoka

(Received September 5, 1972)

Single strand scission in duplex DNA is caused by hydroxylamine, Na-nitrite or oximes of glucose and galactose of proper low molarities. The suitable nicking in the DNA-molecule brings about the elevation of its template nature. Excessive breakage of DNA, however, reduces its template activity. The results have been discussed in relation to an effect causing cyto-differentiation and -anomalization.

INTRODUCTION

Hydroxylamine is the first reagent which was successfully used for virus induction (Yamafuji *et al.*, 1944a; 1944b). Enzymatic studies on this amine have led us to the establishment of a cycle producing metabolic mutagens and virogens (Yamafuji, 1963 ; 1964 ; Yamafuji and Watanabe, 1964). The cycle includes nitrite and oxime. The present paper describes the effect of these three nitro-genous substances on the breakage of DNA-thread and the synthesis of mRNA.

MATERIALS AND METHODS

Cleavage of DNA-strand

DNA was prepared from calf thymus according to the method stated before (Yamafuji *et al.*, 1966 ; 1971b). To prove the single strand scission (nick) in the double helix-molecule, a mixture of 40 $\mu\text{g/ml}$ DNA in SSC (0.15 M NaCl + 0.015 M Na-citrate, pH = 7) and hydroxylamine-metabolites of appropriate molarity was first incubated at 37° for 4 hrs., and a sucrose gradient centrifugation was then applied to the solution of pH=12.8 for 4 hrs. at 30,000 r.p.m. In the control, DNA was incubated without metabolites and centrifuged in the same way.

Alteration of RNA-polymerase

The preparation of enzyme from *E. coli* was carried out with the method of Ishihama and Kameyama (1967). In order to measure the enzymatic activity, 0.25 $\mu\text{Ci } ^{14}\text{C-4-UTP}$ was mixed with 0.08 $\mu\text{moles ATP}$, 0.08 $\mu\text{moles GTP}$, 0.08 $\mu\text{moles CTP}$, 1.0 $\mu\text{moles MgSO}_4$, 0.4 $\mu\text{moles MnSO}_4$, 1.0 pmoles mercaptoethanol, 2.4 μmoles

Tris-HCl buffer (pH=7.8), 20 μ g DNA and 150 μ g polymerase ; the total volume was 200 μ l. At regular intervals at 37°, a portion of the mixture was mixed with 10 % trichloroacetic acid. The precipitate was dissolved in 2 M NH_4OH and counted in Beckman LS 250-fluid scintillation counter. The polymerase action was thus expressed in cpm. To determine the template activity, the nicked DNA prepared by 30 min.-pretreatment with metabolic virogens was used in the test after dialyzing against water for 16 hrs. at 5°. In the control, the DNA incubated in the SSC-solution without virogens was employed after the dialysis in the same manner.

RESULTS

Cleavage of DNA-strand

In the preceding investigation (Yamafuji *et al.*, 1971b), it was demonstrated that hydroxylamine or KNO_2 in co-operation with CuSO_4 breaks duplex DNA. We now investigated the effect in NH_2OH or NaNO_2 of much lower concentration on the DNA-breakage. As shown in Fig. 1, nicks produced in a 5×10^{-5} M amine or 5×10^{-5} M nitrite solution without copper.

It was also observed previously (Murakami and Yamafuji, 1970) that glucosyl- or galactosoxime + Cu cleaves native DNA. As illustrated in Fig. 2, we found here that single strand cleavages can be caused by 5×10^{-6} M oximes in the absence of Cu-salt.

The nick formation was further examined using DNA which had been pre-

Fig. 1. Sedimentation pattern of DNA at pH=12.8 after treating with NH_2OH or NaNO_2 .

Fig. 2. Sedimentation pattern of DNA at pH = 12.8 after treating with glucos- or galactosoxime.

treated with hydroxylamine-metabolites for 30 min. The patterns obtained were the same as those in the above figures, but their peaks were pretty low.

Alteration of RNA-polymerase

In the previous paper (Yamafuji *et al.*, 1972), it was reported that the activity of RNA-polymerase is affected by the combined action of noradrenaline and CuSO_4 . We now estimated the template potentiality of DNA pretreated with virogenic metabolites of relatively low and rather high molality without co-operation of Cu. The experiment disclosed that, as is evident from Fig. 3, an activation of enzyme occurs in the solution containing DNA treated with 5×10^{-5} M hydroxylamine. When the DNA was pretreated with 5×10^{-2} M amine, however, the polymerase was inactivated remarkably. This may be due to double strand-breaks of considerable amount. Taylor *et al.* (1970) proved that treatment of T7 DNA with 0.5 M NH_2OH inhibits transcription of this template by *coli*-RNA polymerase.

In the tests of DNA-splitting by hydroxylamine and NaNO_2 , it was found that the nick-producing ability of the former is higher than that of the latter. As is obvious from Fig. 4, the template potency of DNA pretreated with diluted nitrite solution was lower than that of the one treated with the amine of the same concentration. The observation suggests that the RNA-polymerase activity is proportional to the nicking grade in DNA-molecules. It was further observed

Fig. 3. RNA-polymerase activity in solution containing DNA pretreated with NH_2OH .

Fig. 4. RNA-polymerase activity in solutions containing DNA pretreated with NaNO_2 .

Fig. 5. RNA-polymerase activity in solutions containing DNA pretreated with glucosoxime.

that NO, of high normality also hinders the transcriptional capacity of DNA.

Glucosoxime has been regarded as one of the best polyhedral virus inducers and its toxic action is weak. As depicted in Fig. 5, the template capability of DNA was elevated by treating it with 5×10^{-6} M glucosoxime. The treatment of DNA with 5×10^{-2} M oxime, however, weakened its activity.

DISCUSSION

Cytoanomalizations including virogenesis would be primarily introduced by deviation of DNA-duplication and mRNA-synthesis. For instance, production of viral polyhedrosis necessitates intensive replication of special fragment of chromosomal DNA and rapid formation of specific protein by newly synthesized RNA. The activation of template nature of DNA by metabolic virogens should mean the synthesis of new RNA by polymerase. The formation of abnormal protein, therefore, can be the result of appropriate nicking in DNA-molecules. The present investigation may thus give a support for the formulation (Yamafuji et al., 1971a) of an effect causing cellular differentiation and anomalization.

REFERENCES

- Ishihama, A. and T. Kameyama 1967 The molecular mechanism of enzymic reaction in RNA synthesis. *Biochim. Biophys. Acta*, 138, 480
- Murakami, H. and K. Yamafuji 1970 Mode of action of some catecholamines and sugar oximes on deoxyribonucleic acid. *Enzymol.*, **38**, 337

- Taylor, A. T., S. B. Crist and O. W. Jones 1970 Alteration of native DNA transcription by the mutagen hydroxylamine. *Cancer Research*, **30**, 95
- Yamafuji, K. 1963 A metabolic cycle producing virogens and mutagens. *Agric. Biochem. Kyushu Univ.*, **23**, 13
- Yamafuji, K. 1964 Metabolic virogens having mutagenic action and chromosomal previruses. *Enzymol.*, **27**, 217
- Yamafuji, K., F. Hashinaga and T. Fujii 1966 Isolation and identification of polyhedral pre-viral deoxyribonucleic acid from healthy silkworm cells. *Enzymol.*, **31**, 92
- Yamafuji, K., S. Iiyama and H. Murakami 1972 Action of noradrenalin on brain nucleic acids and RNA-polymerase. *Enzymol.*, **42**, 439.
- Yamafuji, K., S. Iiyama and K. Shinohara 1971a Mode of action of steroid hormones on deoxyribonucleic acid. *Enzymol.*, **40**, 259
- Yamafuji, K., K. Shinohara, F. Yoshihara, M. Iio and Y. Torikai 1971b Breakage of nucleic acids as initiative process of cytodifferentiation and cytoanomalization. *Enzymol.*, **40**, 107
- Yamafuji, K. and Y. Shirozu 1944a Die Abhängigkeit der Neubildung des Virusproteins von Erniedrigung der Katalase. *Biochem. Zs.*, **317**, 94
- Yamafuji, K., S. Takeoka and T. Fujiki 1944b On the production of virus in living organisms. *J. Agric. Chem. Soc. Japan.*, **19**, 225
- Yamafuji, K. and K. Watanabe 1964 Mutagenic action of hyponitrous acid and glucose oxime. *Nature*, **203**, 409