Base-catalyzed Intramolecular Rearrangement of Trimethylamine N-oxide

Harada, Katsuhiko

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

Aso, Yoichi

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

Hayashi, Katsuya

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/22878>

出版情報：九州大学大学院農学研究院紀要. 19 (4), pp.159-168, 1975-06. Kyushu University
バージョン：
権利関係：

Base-catalyzed Intramolecular Rearrangement of Trimethylamine N-oxide

Katsuhiko Harada*, Yoichi Aso and Katsuya Hayashi

Laboratory of Sericultural Chemistry, Faculty of
Agriculture, Kyushu University, Fukuoka

(Received March 3, 1975)

The catalytic role of acid and base in the rearrangement and the following decomposition of trimethylamine N-oxide to formaldehyde and dimethylamine has been investigated. The catalyzer which decomposed the N-oxide most extensively was tris (hydroxymethyl) aminomethane-malonate system, while several other tris (hydroxymethyl) aminomethane-dibasic carboxylic acid systems exhibited appreciable catalytic activities. Specific acid and base, hydrogen ion and hydroxide ion, scarcely showed the catalytic activity in the pH region from 0 to 14. The optimum pH of tris (hydroxymethyl) aminomethane-malonate system for the decomposition of trimethylamine N-oxide was 11 and the amounts of the products, formaldehyde and dimethylamine, bore an equimolar relation to each other. On the other hand, several assumed schemes were simulated to elucidate the catalytic mechanism of tris (hydroxymethyl) aminomethane-malonate system.

Based upon the reaction mechanism presumed from the experimental results, it has been concluded that the catalyzed rearrangement of trimethylamine N-oxide may occur through an intermediate to yield dimethylaminomethylol which gives spontaneously formaldehyde and dimethylamine as final products. In addition, the mechanism of rearrangement step was discussed from the standpoint of role of acid-base catalysis with emphasizing the action of base or hydride ion.

INTRODUCTION

It has been well known that trimethylamine N-oxide (TMO) is distributed widely in tissues of many species of fish and shells (Yamada, 1967). The oxide is cleaved by the action of a certain enzyme (Yamada, 1968) or non-enzymatic catalysis (Vaisey, 1956 ; Soudan, 1959, 1961; Craig *et al.*, 1961; Ferris *et al.*, 1967; Hayashi *et al.*, 1964; Otsuka, personal communication) into formaldehyde (FA) and dimethylamine (DMA). The reaction mechanism of formations of FA and DMA from TMO in enzymatic reaction has been preliminarily investigated by Yamada (1968). Contrarily, the mechanism has been fairly well examined in the non-enzymatic reactions (Vaisey, 1956; Craig *et al.*, 1961; Ferris *et al.*, 1967). The majority of presented mechanisms is containing the reaction step of the formation of an intermediate dimethylaminomethylol (Fish *et al.*, 1956; Sweely *et al.*, 1957; Frisell *et al.*, 1959).

* Visiting Research Associate from Department of Food Science and Technology, The Shimonoseki University of Fisheries, Shimonoseki.

Harada (unpublished data) observed that the participation of reduced methylene blue or reduced flavin nucleotide (co-factor) was required in the enzymatic reaction. The decomposition of TMO to FA and DMA is purely unimolecular reaction and there is any exchange of matter with surrounding medium. Therefore, the role of the reduced co-factors may not be the entire donation of electrons or hydride ion (reduction), but the circulation of electrons through the TMO molecule to generate the polarization of electron distribution in the molecule. If this is the case, acid-base species may exhibit also the catalytic activity to some extents for the decomposition of TMO, because one of the catalytic roles of acid-base species has been known to polarize the electron distribution of a substrate molecule. Thus, the present study was focused to the observation of action of acid-base catalyzer toward the decomposition of TMO in connection with the elucidation of the mechanism of catalysis in the enzymatic reaction.

EXPERIMENTAL

Materials

Substrate : Trimethylamine N-oxide·2H₂O was of guaranteed reagent grade, purchased from Wako Pure Chemical Industries, Ltd.

Catalyzer: Lewis acids were aluminum trichloride and boron trifluoride etherate, purchased from Wako Pure Chemical Industries, Ltd. Specific acid and base were hydrochloric acid and sodium hydroxide. General acid-bases were several buffer solutions. Tris-carboxylate buffers were most frequently used.

Methods

Assay of catalytic activity: A 50 μ mole of TMO was added to 5 ml of chloroform containing 1 mmole of aluminum trichloride or 5 ml of boron trifluoride etherate, while for acid-base catalysis 1 ml of 0.2 M TMO solution (200 μ mole) was added to 10 ml of buffer solution. The mixture solution was then incubated at 80°C for desired periods, and an aliquot was subjected to the determinations of FA and DMA.

Determination of FA and DMA : The amounts of FA and DMA were determined by the methods of Nash (1953) and Dyer (1945), respectively.

Computer simulation : Computer simulations were performed with FACOM 230-75 using Graphic Display GDP F6233A in the Computer Center, Kyushu University.

RESULTS

Catalytic activity of Lewis acids

The catalytic formations of FA and DMA from TMO with Lewis acids are shown in Fig. 1. As seen in the figure, the formations of FA and DMA are observed only in the solution of boron trifluoride, but not in the solution of aluminum trichloride.

Abbreviations are ;TMO: trimethylamine N-oxide, FA : formaldehyde, DMA : dimethylamine, Tris : tris (hydroxymethyl) aminomethane.

Fig. 1. Catalytic formations of FA and DMA from TMO by Lewis acids. ○ : FA, ● : DMA.

Catalytic activity of specific acid-base

As shown in Fig. 2, the formations of FA and DMA in the solution of hydrochloric acid and sodium hydroxide were found to be very faint even below pH 1 and above pH 13. This may mean that TMO is not in the methylol form which has been believed to decompose easily to FA and DMA in acid or alkaline medium.

Fig. 2. Catalytic formations of FA and DMA from TMO in specific acid and base solutions. Reaction mixture was incubated at 80°C for 40 hrs. ○ : FA, ● : DMA.

Catalytic activity of general acid-base

The formations of FA and DMA in many kinds of buffer solutions were, first of all, generally surveyed. The results are shown in Fig. 3. The catalytic formations of FA and DMA were observed in several buffer solutions, though the extents were not exceeded 1 % of initial amount of TMO. However, it was expected that the formations of FA and DMA in the Tris-maleate system may increase in high pH region. To confirm the expectation, the pH-dependence of catalytic action of Tris-maleate system was examined in pH region from 5 to 12. The result obtained is shown in Fig. 4. It was noted that the formations of FA and DMA occurred considerably at pH 11. In connection with the catalytic activity of Tris-maleate system, the formations of FA and DMA in the medium containing other dibasic acid used instead of maleate were surveyed. The results are presented in Fig. 5. The formations were observed most strongly in pH

Fig. 3. Catalytic formations of FA and DMA from TMO in buffer solutions. (1) Glycine-NaCl-HCl (pH 1.1-3.0), (2) Sodium citrate-HCl (pH 1.1-4.0), (3) Sodium phosphate dibasic-potassium phosphate monobasic (pH 5.0-8.0), (4) Glycine-NaCl-NaOH (pH 9.0-12.0), (5) HCl-sodium acetate (pH 1.0-5.0), (6) Citric acid-sodium phosphate dibasic (pH 2.2-8.0), (7) Tris-WCl (pH 7.0-9.0), (8) Tris-maleic acid-NaCl (pH 5.0-8.0), (9) NaOH-sodium phosphate dibasic (pH 11.0-12.0). Reaction mixture was incubated at 80°C for 40 hrs. ○: FA, ●: DMA.

Fig. 4. Catalytic formations of FA and DMA in Tris buffer. Solid line: Tris-maleic acid-NaOH, Dotted line: Tris-NaOH. ○: FA, ●: DMA.

Fig. 5. Effect of dibasic acids on the catalytic formations of FA and DMA in Tris buffer. (1) Malonic acid, (2) Azelaic acid, (3) Succinic acid, (4) Oxalic acid. ○: FA, ●: DMA.

region from 10 to 11.5 regardless of kind of dibasic acids. The formations of FA and DMA in Tris-malonate system was superior to those in other Tris-dibasic acid systems.

Catalytic activity in Tris-malonate system

In Tris-malonate system of which the pH values were adjusted by the addi-

tion of sodium hydroxide, the formations of FA and DMA were estimated with changing the concentrations of the catalyzer and the substrate. The results are shown in Figs. 6 and 7. In the early stages of the reaction, it was found that the time-course of the formations exhibited a clear induction periods; at low concentration of the catalyzer, induction periods (lag-time) extended to 20 hrs of incubation. The amounts of product held an equimolar relation to each other. In the late stages of the reaction, such the relation was broken due to the loss of formed FA.

Fig. 6. Formations of FA and DMA in Tris-malonate system. Concentrations of Tris-malonate (catalyzer) were I : 0.152 M, II : 0.091 M, III : 0.03 M. Incubated at 80°C. ○ : FA, ● : DMA.

Fig. 7. Formations of FA and DMA in 0.152 M Tris-malonate system. Concentrations of TMO of I, II, III, IV, V and VI were 0.182, 0.091, 0.046, 0.0182, 0.0091 and 0.0046 M, respectively. ○ : FA, ● : DMA.

Computer simulation

To elucidate the reaction pathway of TMO decomposition by the base (Tris-malonate system) catalysis, the several schemes were assumed and simulated. As a result of simulation, several schemes which can principally explain the characteristic features of time-course in the base-catalyzed decomposition of TMO were selected. Typical schemes thus selected are shown in Scheme 1. The results of the simulation on the selected schemes are shown in Figs. 8 and 9. The rate constants used for the computation of time-courses shown in the figures are

Table 1. Rate constants used for computation.

Scheme	Catalyzer		TMO	
	(1)	(2)	(1)	(2)
k_1 (sec ⁻¹)	0.333×10^{-4}		0.333×10^{-4}	—
k_2 (M ⁻¹ sec ⁻¹)	0.333×10^{-3}	0.333×10^{-5}	0.333×10^{-3}	0.167×10^{-3}
k_3 (sec ⁻¹)	0.333×10^{-2}	0.333×10^{-3}	0.333×10^{-2}	0.167×10^{-3}
k_4 (M ⁻¹ sec ⁻¹)	0.333×10^{-3}		0.333×10^{-3}	
k_5 (sec ⁻¹)	0.333×10^{-2}	—	0.333×10^{-2}	
k_6 (M ⁻¹ sec ⁻¹)	0.333×10^{-3}	0.333×10^{-4}	0.333×10^{-3}	0.167×10^{-3}

DISCUSSION

Vaisey (1956), Craig *et al.* (1961) and Ferris *et al.* (1967) studied the catalytic actions of iron compounds and Otsuka (personal communication) investigated the decomposition of TMO in the presence of acetic anhydride. Catalytic cleavage of TMO by ferrous or ferric compounds gives the oxido-reduction products, trimethylamine and formic acid, in addition to the normal products, FA and DMA. Since the enzymatic decomposition of TMO gives stoichiometrically equimolar amounts of FA and DMA, but not trimethylamine (Harada, unpublished data), it is not believed that the enzyme provides for a similar catalytic mechanism to that by which iron compound exerts its catalytic activity.

In the enzymatic reaction, it was not observed that the change of co-factor reduced flavin nucleotide or reduced methylene blue to their oxidation forms took place incidentally to the decomposition of the substrate TMO. Since the enzymatic reaction was conducted in anaerobic milieu, there might not be autooxidation of reduced co-factor during the enzymatic reaction. Furthermore, it is clear that the decomposition TMO to FA and DMA is purely unimolecular reaction.

Thus, the catalytic action of enzyme and co-factors toward the substrate may be assumed to occur *via* (a) the enzyme protein moiety acts as acid-base catalyzer and the co-factor is not participating directly in the reaction steps, or (b) both enzyme protein and co-factor act as acid-base catalyzer cooperatively. To approve the above assumptions or to justify the possibility of the above assumptions, catalytic actions of non-enzymatic acid-base species were examined.

Specific acid-base, hydrogen and hydroxide ion, exhibited very low activity in pH region from 0 to 13 as shown in Fig. 2. It had been supposed that an amine N-oxide has partly the methylol structure in aqueous solution. If this is the case, trimethylamine N-oxide must be rapidly decomposed to FA and DMA in the presence of moderate concentration of hydrogen ion or hydroxide ion. The experimental results exclude the methylol structure of TMO in aqueous solution.

It should be emphasized that the some buffer solutions exhibited more catalytic activity in various pa-regions than specific acid-base did. Especially,

(A) Base

(B) Hydride ion

duction periods and the consumption of formed FA in the late stages. In order to explain such the specific features, several reaction schemes were assumed and their behaviors were simulated using digital computer. As a result, it is concluded that the induction periods were caused by the sequence of several reaction steps in the whole pathway, and that the formed FA was consumed in a slow rate by malonate in the catalyzer system,

It is, however, not likely that the reaction which is consisted of several reaction steps exhibits always the induction periods. The problems on the factors which cause the induction periods in reaction time-course will be discussed in the succeeding paper.

ACKNOWLEDGEMENTS

The authors wish to express their thanks to Dr. K. Yamada of The Shimomoseki University of Fisheries, for his valuable advices and discussions.

REFERENCES

- Craig, J. C., F. B. Dwyer, A. N. Glazer and C. Horning 1961 Tertiary Amine Oxide Rearrangements. I. Mechanism. *J. Am. Chem. Soc.*, 83: 1871-1878

- Dyer, W. J. and Y. A. Mounsey 1945 Amines in Fish Muscle II. Development of Trimethylamine and other Amines. *J. Fish. Res. Bd. Canada*, 6: 359-367
- Fish, M. S., N. M. Johnson and E. C. Horning 1956 t-Amine Oxide Rearrangements. N, N-Dimethyltryptamines Oxide. *J. Am. Chem. Soc.*, 78: 3668-3671
- Frisell, W. R., C. W. Chung and C. G. Mackenzie 1959 Catalysis of Oxidation of Nitrogen Compounds by Flavin Coenzymes in the Presence of Light. *J. Biol. Chem.*, 234: 1297-1302
- Ferris, J. P., R. D. Gerwe and G. R. Gapski 1967 Detoxication Mechanism II. The Iron-Catalyzed Dealkylation of Trimethylamine Oxide. *J. Am. Chem. Soc.*, 89: 5270-5275
- Hayashi, Y., Y. Nagano, S. Hongyo and K. Teramura 1974 Trapping an Intermediate of the Polonovski Reaction. *Tetrahedron Letters*, 14 : 1299-1302
- Nash, T. 1953 The Colorimetric Estimation of Formaldehyde by Means of the Hantzsch Reaction. *Biochem. J.*, 55: 416-421
- Sweely, C. C. and E. C. Horning 1957 Rearrangement and Decarboxylation Reactions of N, N-Dimethylglycine Oxide. *J. Am. Chem. Soc.*, 79: 2620-2625
- Soudan, F. 1959 Le Microdosage du Formol dans les Produits Marins. *Rev. Trav. Inst. Pêches Marit.*, 23: 203-210
- Soudan, F. 1961 Sobre la Presencia de Formol Natural en los Productos Marinos. Presented Paper at FAO International Conference on Fish in Nutrition, Washington, D. C., Sept., 19
- Vaisey, E. B. 1956 The Non-Enzymic Reduction of Trimethylamine Oxide to Trimethylamine, Dimethylamine and Formaldehyde. *Can. J. Biochem. Physiol.*, 34: 1085-1090
- Yamada, K. 1967 Occurrence and Origin of Trimethylamine Oxide in Fishes and Marine Invertebrates. *Bull. Jap. Soc. Sci. Fish.*, 33: 591-603
- Yamada, K. 1968 Post-mortem Breakdown of Trimethylamine Oxide in Fishes and Marine Invertebrates. *Bull. Jap. Soc. Sci. Fish.*, 34: 541-551