Developmental ‘Changes in Activities of Several Enzymes Involved in Glycolysis and TCA Cycle during the Pupal Period of Bombyx mori

Miake, Fumio

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

Torikata, Takao

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

Hanada, Shuichi

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

Soeda, Eiichi

Laboratory of Sericultural Chemistry, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/22896>

出版情報：九州大学大学院農学研究院紀要. 20 (3/4), pp.143-149, 1976-10. Kyushu University
バージョン：
権利関係：

Developmental 'Changes in Activities of Several Enzymes Involved in Glycolysis and TCA Cycle during the Pupal Period of *Bombyx mori*

Fumio Miake, Takao Torikata, Shuichi Hanada*, Eiichi Soeda,
Katsumi Koga and Katsuya Hayashi**

Laboratory of Sericultural Chemistry, Faculty of Agriculture,
Kyushu University, Fukuoka

(Received May 14, 1976)

In order to elucidate metabolic regulation with regard to carbohydrates and lipids during metamorphosis of *Bombyx mori*, eight enzymes related to glycolysis and TCA cycle were measured for their levels of activity during the period from the day of pupal ecdysis to 2 days after adult emergence, by use of crude extract from the whole bodies of the insect. The resulting patterns could be grossly divided into two groups. Glucose-B-phosphate dehydrogenase, NADP-specific isocitrate dehydrogenase, malic enzyme, hexokinase and pyruvate kinase were characterized by their apparent high activity levels during larval-pupal development followed by low levels at emergence. On the contrary, aconitase, NAD-specific isocitrate dehydrogenase and malate dehydrogenase were characterized by their apparent low levels during pupal period and by an abrupt increase in activity before emergence. The roles of these enzymes in metamorphosis are briefly discussed.

INTRODUCTION

Recently, attention has been focused on metabolic conversions underlying the remarkable, morphological changes during the development of insects (Morishima, 1975; Walker and Bailey, 1970; Yamashita and Hasegawa, 1974). In metamorphosing endopterygote insects like the silkworm, certain larval tissues are degraded and regenerated into a new form adaptable to the adult life. Such a process must be controlled by highly ordered, elaborate systems of decomposition and reconstruction which are closely correlated with each other. The regulatory mechanisms concerning the system may be approached by an enzymological study. The enzymes involved in the glycolytic pathway and TCA cycle will be of primary importance. The present paper deals with the changes in activity levels for the several of these enzymes in the crude extracts from the developing pupae of *Bombyx mori*.

Present address: * Central Research Laboratories of Green Cross Corporation, Miyakojima, Osaka 534.

** National Institute of Genetics, Mishima 411.

MATERIALS AND METHODS

Materials

NAD*, NADP, glucose-6-phosphate, phosphoenolpyruvate, ADP and ATP were obtained commercially (Boehringer, Mannheim). Malate dehydrogenase, glucose-6-phosphate dehydrogenase and lactate dehydrogenase used for coupling reactions were the products of Boehringer, Mannheim. Other reagents were of analytical grade.

Preparation of crude extract

Hybrid strains of the silkworm were used. Ten individuals were taken at intervals during the developmental stage from the day of larval-pupal ecdysis to 2 days after adult emergence. Male and female animals were separately processed. The pooled silkworms were weighed and homogenized with a Waring blender in three volumes per weight of a solution composed of 0.1 M KCl, 0.01 M sodium phosphate buffer (pH 7.0) and 1 mM 2-mercaptoethanol. The homogenate was passed through double layers of gauze and centrifuged at 8000 x g for 25 min. The supernatant was subjected to spectrophotometric assay of enzymatic activity as described below.

Measurement of enzymatic activity

Activities of the enzymes were routinely assayed by use of a spectrophotometer in a cell with a light path of 1-cm, by recording an increase or a decrease in the absorbance of a UV-absorbing substrate, product, or coenzyme. A Hitachi 124 spectrophotometer equipped with a recorder and a temperature controlling cell-holder was used. As to all the enzymes concerned the changes in the absorbance were linear during the first several minutes of reaction: the initial velocities of the enzymatic reaction could be calculated by the gradient of the recorded curves. One unit of an enzyme was defined as its amount that caused the change of 1 μ mole of the substrate or so (see above) used for the spectrophotometry per min per ml at 30°C.

Glucose-6-phosphate dehydrogenase

Enzymatic activity for this was measured in 0.1 M Tris-HCl buffer (pH 7.5) containing 0.5 μ moles of glucose-6-phosphate, 1.0 μ mole of $MgCl_2$ and 1.0 μ mole of NADP in a total volume of 3.0 ml. The reaction was initiated by the addition of 10 or 25 μ l of the extract. Formation of NADPH was estimated by the absorbance at 340nm.

NADP-specific isocitrate dehydrogenase

Activity of the enzyme was measured by an increase in the absorbance at 340 nm of NADPH. The mixture used was composed of 0.1 M Tris-HCl buffer

* Abbreviations used are: NAD, nicotinamide-adenine dinucleotide; NADH, reduced form of NAD; NADP, nicotinamide-adenine dinucleotide phosphate; NADPH, reduced form of NADP; ADP, adenosine-5'-diphosphate; ATP, adenosine-5'-triphosphate; G6P-DH, glucose-6-phosphate dehydrogenase; NAD-ICDH, isocitrate dehydrogenase specific for NAD; NADP-ICDH, do. specific for NADP; ME, malic enzyme; HK, hexokinase; PK, pyruvate kinase; MDH, **malate** dehydrogenase.

(pH 7.8) containing 0.5 μ moles of DL-isocitrate, 1.0 μ mole of MnCl_2 and 0.1 μ moles of NADP in a total volume of 3.0 ml. The reaction was initiated by the addition of 10 or 25 μ l of the extract.

Malic enzyme

Activity of the enzyme was measured in 0.05 M Tris-HCl buffer (pH 7.5) containing 10 μ moles of malate, 0.5 μ moles of NADP and 5 μ moles of MnCl_2 in a total volume of 3.0 ml. The reaction was initiated by the addition of 10 to 50 μ l of the extract. Production of NADPH was measured by the absorbance at 340 nm.

Hexokinase

Production of glucose-6-phosphate from glucose by the enzyme was coupled with the reaction of added glucose-6-phosphate dehydrogenase and was measured by an increase in the absorbance of NADPH at 340nm. The reaction mixture used was made up of 0.1 M Tris-HCl buffer (pH 7.5), 666 μ moles of glucose, 20 μ moles of MgCl_2 , 8.1 μ moles of ATP, 2.2 μ moles of NADP and 1.5 units of glucose-6-phosphate dehydrogenase in a total volume of 3.0 ml. The reaction was initiated by the addition of 100 μ l of the extract.

Pyruvate kinase

Production of pyruvate from phosphoenolpyruvate by the enzyme was coupled with the reaction of added lactate dehydrogenase and was spectrophotometrically followed by a decrease in the absorbance of NADH at 340nm. The reaction mixture used for this consisted of 0.1 M Tris-HCl buffer (pH 7.6), 1.6 μ moles of phosphoenolpyruvate, 7.5 μ moles of MgSO_4 , 30 μ moles of KCl, 14.1 μ moles of ADP, 0.6 μ moles of NADH and 45 units of lactate dehydrogenase in a total volume of 3.0 ml. The reaction was initiated by the addition of 20 μ l of the crude enzyme solution.

Aconitase

The reaction was started by the addition of less than 50 μ l of the extract to 3 ml of a solution containing 10 μ moles of DL-isocitrate in 0.05 M potassium phosphate buffer (pH 7.5). The initial increase in the absorbance at 240nm that was caused by the formation of aconitate from isocitrate was measured according to the method previously reported (Soeda *et al.*, 1973).

NAD-specific isocitrate dehydrogenase

The reaction mixture used for the enzyme was composed of 0.05 M Tris-HCl buffer (pH 7.0), 20 μ moles of DL-isocitrate, 3 μ moles of NAD and 15 μ moles of MgCl_2 in a total volume of 3.0 ml. The reaction was initiated by the addition of 10 or 50 μ l of the extract. An increasing amount of NADH was measured in the absorbance at 340nm.

Malate dehydrogenase

Activity for the enzyme was measured in 0.05 M Tris-HCl buffer (pH 7.5) containing 10 μ moles of oxaloacetate and 1.3 μ moles of NADH in a total volume of 3.0 ml. The reaction was initiated with 10 or 20 μ l of the extract, and a decrease in the absorbance of NADH was measured at 340 nm.

RESULTS AND DISCUSSION

Many different patterns of activity were observed for different enzymes during developmental stage, they could be, however, grossly divided into two groups. A group of enzymes, GGP-DH, NADP-ICDH, ME, HK and PK, was characterized by their apparent high levels during larval-pupal development and low levels at emergence. The activities of GGP-DH (Fig. 1-A), NADP-ICDH (Fig. 1-B) and ME (Fig. 1-C) gradually decreased just after pupation and increased gradually on 3rd to 6th days of the pupal period. At 6 days of age,

Fig. 1. Changes of GGP-DH (A), NADP-ICDH (B) and ME (C) activity during pupal period. • male, Δ female.

the activities took their maximum values and gradually decreased thereafter. No significant differences between males and females occurred with the three enzymes. The HK activity (Fig. 2-A) decreased till 3 to 4 days of age and later increased gradually, reaching its maximum level at day 7 of development. Then the activity decreased gradually till emergence. The activity level was slightly higher in females than that of males. The activity of PK (Fig. 2-B)

Fig. 2. Changes of HK (A) and PK (B) activity during pupal period. • male, Δ female.

gradually decreased during the period from pupation to 6 or 7 days of development and attained to a maximum value at 9 days of age, then it decreased steeply. The enzyme showed higher activity in females than in males.

Another group of enzymes, aconitase (Fig. 3-A), NAD-ICDH (Fig. 3-B) and

Fig. 3. Changes of aconitase (A), NAD-ICDH (B) and MDH (C) activity during pupal period. ● male, △ female.

MDH (Fig. 3-C), was characterized by their apparent low levels during pupal period and by a remarkable rise in the activity shortly before emergence. The activities of NAD-ICDH and MDH showed their maximum values just at the time of emergence and then decreased gradually, while the aconitase activity remained high after emergence. No significant differences between males and females were observed.

It has been shown that a number of metabolites, including cyclic nucleotides (Kafatos, 1975), carbohydrates (Wyatt, 1967; Yamashita and Hasegawa, 1974), lipids and fatty acids (Ichimasa, 1975; Ichimasa and Hasegawa, 1973), attain interesting fluctuation patterns in accordance with the pupal-adult development of certain silkmoths and *Bombyx mori*. Only a few investigators, however, have pointed out the correlations of these changes to their related enzymes, as shown in studies on the glycogen-trehalose system (Wyatt, 1967; Yamashita and Hasegawa, 1974; Yamashita *et al.*, 1974). It is well known that absorption of oxygen by many insects changes in a U-shaped curve during pupal age. As this holds also for *Bombyx mori* (Nittono and Nakasone, 1955; Kobayashi, 1957), it is easily presumed that in the same species the activities of the enzymes that are involved in TCA cycle are low during the middle stage of the pupa, and increase at the end of the pupal age or just after emergence. The present experiments showed that the activities of aconitase, NAD-ICDH and MDH abruptly increased at emergence, although they slightly increased 2 to 4 days before emergence. At the time of emergence, the thoracic muscles attain rapid development of sarcosomes, which correspond to mitochondria in biochemical actions. Moreover, there was evidence that the aconitase was of a sarcosomal origin (Soeda *et al.*, 1973). These facts support the concept that aforementioned enzymes may play their parts in respiration and that activation or biosynthesis of these enzymes may be a critical step for the differentiation of the muscles.

Contrary to above, the activities of G6P-DH, NADP-ICDH and ME were maximum at 6 days of age and that of HK was maximum at day 7 in *Bombyx mori* (Figs. 1-A, B, C and Z-A). During these periods, aerobic respiration systems in this insect may be low in activity (Nittono and Nakasone, 1955; Kobayashi, 1957), but histolysis and histogenesis with concomitant degradation and regeneration of carbohydrates (Yamashita and Hasegawa, 1974), lipids and

fatty acids (Ichimasa, 1975 ; Ichimasa and Hasegawa, 1973), nucleic acids (Chinzei and Tojo, 1972) and proteins (Tojo, 1971) are actively taking place. It is also probable that the cuticle chitin of the larva degrades during larval-pupal ecdysis and some part of it may be stored in the form of glycogen or so for the reconstruction of the adult cuticles as was shown for *Hyalophora cecropia* (Bade and Wyatt, 1962). A part of the energy for these processes may be supplied by NADPH, which is conceivably produced by the action of GGP-DH, ME and NADP-ICDH.

The activity of PK was maximum in pupae of 9 days old, and it is possible that the enzyme may supply pyruvate which can serve as a starting point for synthetic pathways such as those for fatty acids and amino acids. In fact, it was shown that fatty acid composition in the glycerides changes significantly during pupal age of *Bombyx mori* (Ichimasa, 1975; Ichimasa and Hasegawa, 1973; Nakasone and Ito, 1967), and this is taken to indicate that fatty acids are synthesized at least to some extent in the pupal tissues, although these substances accumulate most abundantly during the last larval instar (Shigematsu and Takeshita, 1959).

During pupal-adult development, the fat body and the ovaries exhibit glycolysis and glycogenesis through a pathway in which the level of trehalose, the specific blood-sugar of insects, is homeostatically kept constant in *Bombyx mori* (Yamashita and Hasegawa, 1974) and in *Hyalophora cecropia* (Wyatt, 1967). It is worth mentioning here that, according to Suzuki *et al.* (1973), significant activity of fructose-1,6-diphosphatase (FDPase) was detected in the ovaries of *Bombyx mori* at the middle pupal age, suggesting that the organ is capable of generating glucose also from metabolites other than trehalose. The results of our investigation (Figs. 1 and 2) showed that GGP-DH, NADP-ICDH, ME, HK and PK varied in somewhat similar ways to that for FDPase, permitting the idea that the pupal tissues exhibit wide-ranging interconversions of metabolites. It should be noted here that the whole body extracts were used in the present study, and that, as a matter of course, solely the present data on the levels of enzymatic activities do not reveal how the metabolic pathways are functioning. Distribution of the enzymes and behavior of metabolites among tissues remain to be studied. The kinetic properties of the purified enzymes will also be of value from the viewpoint of metabolic regulation. We, therefore, have tried to purify and characterize some of the enzymes from the pupae of *Bombyx mori*; the findings have been published separately (Miake *et al.*, 1976; Soeda *et al.*, 1973 ; Torikata, 1974).

REFERENCES

- Bade, M. L. and G. R. Wyatt 1962 Metabolic conversions during pupation of the cecropia silkworm. *Biochem. J.*, 83: 470-478
Chinzei, Y. and S. Tojo 1972 Nucleic acid changes in the whole body and several organs of the silkworm, *Bombyx mori*, during metamorphosis. *J. Insect Physiol.*, 18: 1683-1698
Ichimasa, Y. 1975 Studies on the mode of action of the diapause hormone with special reference to lipid metabolism in the silkworm, *Bombyx mori* L. II. Lipid component and

- fatty acid composition of glyceride in pupal ovary, fat body and haemolymph. *J. Sericult. Sci. Japan*, **44**: 137-145
- Ichimasa, Y. and K. Hasegawa 1973 Studies on the mode of action of the diapause hormone with special reference to lipid metabolism in the silkworm, *Bombyx mori* L. 1. Effect of the hormone on lipid in pupal ovaries and matured eggs. *J. Sericult. Sci. Japan*, **42**: 380-392
- Kafatos, F. C. 1975 Personal communication.
- Kobayashi, M. 1957 Studies on the neurosecretion in the silkworm, *Bombyx mori* L. Bull. *Seric. Exp. Sta. Japan*, **15**: 181-273
- Miake, F., T. Torikata, K. Koga and K. Hayashi 1976 Affinity chromatography of NADP-specific isocitrate dehydrogenase from the pupa of *Bombyx mori*. *Agr. Biol. Chem.*, **40**: 1269-1271
- Morishima, I. 1975 Cyclic nucleotide phosphodiesterase in silkworm. Developmental change of cyclic AMP and cyclic GMP phosphodiesterases. *Biochim. Biophys. Acta*, **403**: 106-112
- Nakasone, Y. and T. Ito 1967 Fatty acid composition of the silkworm *Bombyx mori* L. *J. Insect Physiol.*, **13**: 1237-1246
- Nittono, Y. and Y. Nakasone 1955 Effects of oxygen tensions and carbon monoxide on the respiration of silkworm pupa (*Bombyx mori* L.). *J. Sericult. Sci. Japan*, **25**: 5-10
- Shigematsu, H. and H. Takeshita 1959 On the change in the weight of the fat body and of its chief constituents in the silkworm, *Bombyx mori* L., during metamorphosis. *Jap. J. Appl. Ent. Zool.*, **3**: 123-127
- Soeda, E., T. Torikata and S. Akune 1973 Sarcosomal aconitase of silkworm. *J. Biochem.*, **73**: 529-536
- Suzuki, K., O. Yamashita and K. Hasegawa 1973 On the activity of fructose-1, 6-diphosphatase in pupal ovaries of the silkworm, *Bombyx mori* L. *J. Sericult. Sci. Japan*, **42**: 357-362
- Tojo, S. 1971 Uric acid production in relation to protein metabolism in the silkworm *Bombyx mori*, during pupal-adult development. *Insect Biochem.*, **1**: 249-263
- Torikata, T. 1974 Studies on a malate decarboxylase in the pupa of *Bombyx mori*. "Dissertation, Kyushu Univ."
- Walker, P. and E. Bailey 1970 Changes in enzymes associated with lipogenesis during development of the adult male desert locust. *J. Insect Physiol.*, **16**: 679-690
- Wyatt, G. R. 1967 The biochemistry of sugars and polysaccharides in insects. In "Advances in Insect Physiology," Vol. 4, ed. by J. W. L. Beament, J. E. Treherne and V. B. Wigglesworth, Academic Press Inc., New York, pp. 287-360
- Yamashita, O. and K. Hasegawa 1974 Mobilization of carbohydrates in tissues of female silkworms, *Bombyx mori*, during metamorphosis. *J. Insect Physiol.*, **20**: 1749-1760
- Yamashita, O., M. Sumida and K. Hasegawa 1974 Developmental changes in midgut trehalase activity and its localization in the silkworm, *Bombyx mori*. *J. Insect Physiol.*, **20**: 1079-1085