

THE GENUS ANTISPILA FROM JAPAN, WITH
DESCRIPTIONS OF SEVEN NEW SPECIES (Lepidoptera,
Heliozelidae)

Kuroko, Hiroshi

<https://doi.org/10.5109/2347>

出版情報 : ESAKIA. 3, pp.11-24, 1961-08-05. Hikosan biological laboratory, Faculty of
Agriculture, Kyushu University

バージョン :

権利関係 :



THE GENUS *ANTISPILA* FROM JAPAN, WITH
DESCRIPTIONS OF SEVEN NEW SPECIES*

(Lepidoptera, Heliozelidae)

By

Hiroshi Kuroko

Antispila is a small genus, including only over thirty described species in the world. In Japan not a single species has hitherto been reported. In the present paper the author gives a key to the species, descriptions of seven new species on the basis of specimens which were gained by breeding of the larvae. The type specimens designated in the present paper are deposited in the collection of the Entomological Laboratory, Kyushu University.

Before going further the author wishes to express his hearty thanks to Prof. K. Yasumatsu for his constant direction ; to Dr. K. Sattler, of Zoologische Sammlung des bayerischen States for the gift of valuable specimens for comparison.

Genus *Antispila* Hübner

Antispila Hübner, 1825, Verz. bekannt. Schmett., (27) : 419; Stainton, 1870, Nat. Hist. Tin., 11: [298; Meyrick, 1895, Handb. Brit. Lep. : 683; Spuler, 1910, in Hofmann-Spuler, Schmett. Eur., 2 : 471; Hering, 1932, Tierwert Mitteleurop. 3: 15.

Type species : *Antispila pfeifferella* (Hübner).

Head and face smooth; ocelli absent; tongue developed, naked. Antennae 1/2—2/3 of forewing, thick, simple in both sexes. Labial palpi 3 jointed, short and thick, drooping, terminal joint pointed. Maxillary palpi are seen inside labial palpi and concealed by frontal scales, nearly 1/2 the length of labial palpi. Posterior tibiae smooth, with long hairs beneath. Forewings broad-lanceolate, with silvery or golden-metallic markings, namely, a transverse fascia or two opposite spots (anterior costal and dorsal-spots) at 1/3, and two opposite

* Contribution Ser. 2, No. 11, Hikosan Biological Laboratory, Kyushu University, Hikosan.

Contribution Ser. 2, No. 99, Entomological Laboratory, Kyushu University, Fukuoka.

spots (posterior costal and dorsal spots) at near $2/3$, sometimes with an apical spot. Hindwings $3/4$ in width, narrow-lanceolate, with a long frenulum near base in male; cilia 2.




Fig. 1. *Antispilella purplella* sp. nov.

A. Head, lateral view.

B. Labial palpus and maxillary palpus.

$1/4$ of discoidal cell; sometimes a fine trachea running longitudinally in discoidal cell as a remnant of media. On hindwing, lc not reaching dorsal margin; 2 and 3 stalked, or 2 absent; 6 and 7 stalked, 7 to costa or near apex, sometimes 6 absent; 4 diverging from before middle of wing; 8 highly curved at base.

Male genitalia : Scaphium weakly sclerotized, uncus undeveloped, with a hairy pad; gnathos consists of a pair of hairy tubercles; valva triangular or trapezoidal, with a comb of teeth, ampulla projected like a finger; saccus large and deep, U-shaped; aedoeagus long and slender, with a peculiar shaped, moderately sclerotized phallobase (basal plate of phallus) which is in contact with its ventral side, and apex of aedoeagus with various shaped terminal appendages; anellus spined; transtilla and juxta present.




Fig. 2. Male genitalia of *Antispilella hikosana* sp. nov., lateral view.

Ae: aedoeagus, Am: projection of ampulla, An: anus, Ct: comb of teeth, Gn: gnathos, Jx: juxta, Ph: phallobase, Sa: saccus, SC: scaphium, Tr: transtilla, v: valva.

- All spots not reaching half across wing, reflections of forewing moderate
 *A. corniella* sp. nov.

***Antispila orbiculella* sp. nov.**

(Pl. 7, fig. 31; Pl. 8, fig. 38)

♂♀. 4.0-4.5 mm. Head, antennae, thorax and abdomen shining bronzy-fuscous. Palpi pale fuscous.

Forewings rather broad-lanceolate ; discocellulars slightly curved between vein 2 and 4+5, vein 4-t-5 from middle of discocellulars, vein 9 from the upper apex of discoidal cell, dark bronzy-fuscous with reddish reflections; basal area shining leaden-bronze ; markings silvery-metallic with yellowish or bluish reflections in some lights; a rather broad, slightly curved oblique transverse fascia at 1/3; a rather large quadrilateral spot on costa at 2/3, reaching nearly half across wing, a rather large opposite spot on dorsum, bifurcated in disc; cilia pale fuscous, basal half dark fuscous, its margin rounded. Hindwings, vein 2 absent, vein 6 and 7 stalked; pale fuscous, with feeble purplish lusters; cilia pale fuscous.

Male genitalia (Pl. 4, fig. 11): Edge of scaphium slightly bilobed; valva trapezoidal ; ampulla moderately projected, slightly downward ; sacculus angulated ; saccus broad; apex of aedeagus with some spines and two projections, one rounded at apex and another tapered; phallobase short, its apex broadly swollen.

Female genitalia (Pl. 6, figs. 16 & 24): Ovipositor five pointed, central point indent; enlarged part of ductus bursae with 2 pairs of plates, 1 pair of short bars and 1 pair of spined rows.

Holotype : ♂, Mt. Hikosan, N. Kyushu, 1 July 1955 (H. Kuroko).

Allotype : ♀, the same as holotype, 1 July 1955 (H. Kuroko).

Paratypes : 1 ♀, the same as holotype, 24 June 1955; 1 ♂, 26 June 1955; 1 ♀, 2 July 1955; 1 ♂, 3 July 1955 (H. Kuroko).

Distribution : Japan (Kyushu & Yakushima).

This new species is easily distinguishable from the other species by the rather broad transverse fascia and the bifurcated posterior dorsal spot. The larvae are not rarely found in autumn on Mt. Hikosan.

Biological notes

Host plant : *Ampelopsis brevipedunculata* (Maxim.) Trautvetter (Jap. name : Nobudō).

Mine (Pl. 9, fig. 45): Orthogenous blotch mine, full depth type; pale greenish brown, central area blackish brown, appearing just as disease blotch; 5-6 mm. in diameter. Usually one mine found on a single leaf, but sometimes 2-3 are found, and extending between two ribs of the leaf.

Frass: Blackish brown grains of frass are arranged in a circle around central area and some of frasses are smeared inside of epidermis of central area of the mine.

Case (fig. 3): Discal, central parts of both faces convex; dark brown; 2.5—3.0 mm. in diameter. Thorn-like projections more or less pointed; pale brown, 1—21 in number; 0.09—0.12 mm. in length. The case is cut out from the central part of the mine.

Development: One generation in a year. The larvae may be found feeding from the end of August to September; when they become full grown at the end of September, they cut out circular cases and descend to the ground; in case they hibernate as a non-feeding larval stage, pupation occurs at the end of May of the following year; the adults appear from the middle of June to the beginning of July.




Fig. 3. Case of the larva of *Antispila orbiculella* sp. nov.

***Antispila ampelopsia* sp. nov.**
(Pl. 7, fig. 32; Pl. 8, fig. 40)

♂♀. 4.5-5.5 mm. Head, antennae, thorax and abdomen shining bronzy-fuscos, Palpi whitish.

Forewings, venation the same as in *A. orbiculella* Kuroko; dark bronzy-fuscos with reddish reflections; basal area shining leaden-bronze; markings silvery-metallic, with yellowish or bluish reflections; slightly curved oblique transverse fascia moderately, constricted on fold, broad toward dorsum; posterior costal spot subtriangular or rectangular, and opposite dorsal spot placed slightly before costal spot, triangular and nearly reaching half across wing; cilia pale fuscous, basal half dark fuscous with purplish reflections. Hindwings, veins 2 and 6 absent, vein 7 to near apex; fuscous with purplish lusters feebly; cilia pale fuscous.

Male genitalia (Pl. 5, fig. 14): Scaphium trapezoidal; valva triangular, projection of ampulla well developed and broadened; sacculus swollen; basal edge of saccus prolonged at middle; apex of aedoeagus with two finger-shaped projections and a membranous projection bearing a bifurcated, spined piece; phallobase spear-shaped.

Female genitalia (Pl. 6, figs. 18 & 26): Ovipositor seven pointed, central point deeply indent; enlarged part of ductus bursae with two oblong plates and a curved plate.

Holotype: ♂, Mt. Hikosan, N. Kyushu, 11 Aug. 1956 (H. Kuroko).

Allotype: ♀, the same as holotype, 24 June 1955 (H. Kuroko).

Paratypes: 2 ♀♀, the same as holotype, 14 & 15 Sept. 1954; 1 ♀, 13 Aug. 1956 (H. Kuroko). The following paratypes were bred in 25°C during the winter. 2 ♀♀, Ambo, Yakushima, 5 & 11 Feb. 1960 (H. Kuroko).

Distribution: Japan (Kyushu & Yakushima).

This new species differs from *A. ampelopsiella* Chambers, *A. ampelopsifoliella* Chambers and *A. voraginella* Braun of which the larvae are known as the leaf-

miner of *Ampelopsis* or *Vitis* in N. America, by the absence of the apical spot. This species is similar to *A. isabellella* Clemens from N. America, but being smaller and having the posterior dorsal spot triangular. From *A. viticordifoliella* Clemens this species is distinguishable by the absence of silvery-white joints at the apex of the antennae. The larvae are commonly found in autumn on Mt. Hikosan.

Biological notes

Host plants: *Ampelopsis brevipedunculata* Maxim). Trautvetter (Jap. name : Nobudō), *Vitis flexuosa* Thunberg (Jap. name: Sankakuzuru).

Mine (Pl. 9, fig. 46): Blotch mine, full depth type ; semitransparent, whitish, faintly tinged with yellowish green; feeding area 74-120 mm? Usually extending at the apical area or marginal area of the leaf, and one mine is made on a single leaf.

Frass: Blackish grains of frass are scattered more or less in a zigzag line.




Fig. 4. Case of the larva of
Antispila ampelosia sp. nov.

Case (fig. 4): Elliptical, with a longitudinal ridge at the middle of each face; brown to dark brown; 3.5 x 1.2 mm. -4.0 x 2.0 mm. in size. Thorn-like projections pointed and radiating; brown ; 11-24 in number ; 0.3-0.42 mm. in length. The case is cut out from the end of the mine.

Development: Number of generations unknown, but the larvae are found at the end of July, at the middle of August and from the end of September to the middle of October, and the adults appear at the middle of August, at the middle of September and at the end of June. Therefore, it may be apparent that this new species repeats two or three generations in a year. If it is so, this habit may be exceptional.

Antispila corniella sp. nov.

(Pl. 7, fig. 33 ; Pl. 8, fig. 41)

♂ ♀, 5 - 6 mm. Head, antennae and thorax shining bronzy-fuscos. Palpi whitish. Abdomen bronzy-fuscos above, shining white beneath.

Forewings, venation as in *A. ampelosia* Kuroko, but vein 9 stalked with vein 7+8 at a short distance beyond the upper apex of discoidal cell; dark bronzy-fuscos with brassy reflections, becoming purplish reflections on the apical half; basal area shining leaden-fuscos ; makings silvery-metallic with yellow or bluish reflections ; a subtriangular costal spot at 1/3, and a triangular dorsal spot slightly before the costal spot; a similar posterior costal spot at near 2/3, and dorsal spot slightly before just mentioned costal spot; both costal spots

not reaching half across wing but two dorsal spots reaching half; cilia pale fuscous, basal half dark fuscous. Hindwings, venation the same as in *A. ampelapsia* Kuroko; pale fuscous with feeble purplish lusters; cilia pale fuscous.

Male genitalia (Pl. 5, fig. 15): Edge of scaphium convexed at middle; valva triangular, with a rather large comb of teeth; projection of ampulla short, straightened; saccus prolonged and rather narrow; apex of aedoeagus with two pouch-shaped projections; phallobase spear-shaped, apex not pointed.

Female genitalia (Pl. 6, figs. 22 & 23): Ovipositor five pointed, central point indented; enlarged part of ductus bursae with two conspicuous filaments, bearing a basal 'knot'.

Holotype: ♂, Mt. Hikosan, N. Kyushu, 12 Aug. 1955 (H. Kuroko).

Allotype: ♀, the same as holotype, 3 Aug. 1956 (H. Kuroko).

Paratypes: 1 ♀, the same as holotype, 25 July 1955; 1 ♂, 2 ♀♀, 1-5 Aug. 1956 (H. Kuroko).

Distribution: Japan (Kyushu).

The present new species is allied to *A. treitschkiella* Fischer von Röslerstamm of which the larva is known as a leaf-miner of *Cornus* in Europe, but in the latter species the forewing has a transverse fascia instead of anterior spots, and the projection of ampulla is much longer.

Biological notes

Host plants: *Corms controversa* Hemsley (Jap. name: Mizuki), *C. brachypoda* C. A. Mey. (Jap. name: Kumano-mizuki).

Mine (Pl. 9, fig. 47): Linear-blotch, full depth type; linear mine whitish, semitransparent, extends in a line along the leaf margin or a wavy line and becoming gradually broader; about 35 mm. in length. After the 3rd moult, the mine develops into a whitish green blotch, with greenish grains by feeding tracks; feeding area 156—192 mm². One or two mines are found on a single leaf.

Frass: Blackish grains of frass, in linear mine arranged in a line, but in blotch mine appearing like a large black patch in the middle of the mine.

Case (fig. 5): Elliptical, with a longitudinal ridge at the middle of each face; brown to dark greyish brown; 4.0 x 2.0 mm. — 4.0 x 2.5 mm. in size. Thorn-like projections triangular and radiating; brown; 10—20 in number; 0.18-0.3 mm. in length.

Development: One generation in a year. The larvae may be found feeding from the September to October; the cases are cut out from the end of mine and the larvae descend to the ground; the larvae hibernate as a




Fig. 5. Case of the larva of *Antispila corniella* sp. nov.

non-feeding larval stage, and change to pupae at the beginning of June of the following year; the adults appear from the end of July to August.

***Antispila iviella* sp. nov.**

(Pl. 7, fig. 34 ; Pl. 8, fig. 41)

♀. 5--6 mm. Head, antennae and thorax shining dark bronzy-fuscous. Palpi whitish. Abdomen dark bronzy-fuscous above, silvery-whitish beneath.

Forewings, vein 4+5 from near the upper apex of discoidal cell, vein 9 stalked with vein 7+8 at some distance beyond the upper apex of discoidal cell ; dark bronzy-fuscous with brassy reflections, becoming purplish reflections towards apex and termen; basal area shining dark leaden; markings rather raised and silvery-metallic tinged with faint golden; a subtriangular or oblong costal spot at 1/3, and a triangular dorsal spot slightly before costal spot; a similar posterior costal spot at near 2/3 and a dorsal spot slightly before posterior costal spot; all spots almost reaching half across wing; cilia fuscous, basal half dark fuscous with strong purplish reflections. Hindwings, vein 2 absent, veins 6 and 7 stalked at near apex ; pale fuscous with feeble lusters ; cilia pale fuscous

Female genitalia (Pl. 6, figs. 17 & 25) : Ovipositor similar to that of the preceding species; enlarged part of ductus bursae with a sclerotized loop.

Holotype : ♀, Miyanoura, Yakushima, 14 July 1960 (H. Kuroko).

Paratypes : 1 ♀, the same as holotype, 13 Feb. 1960; 1 ♀, 22 July 1960 (H. Kuroko). All the specimens were bred under 25°C during the winter.

Distribution : Japan (Yakushima).

This new species is closely allied to *A. argostoma* Meyrick, of which the larva is known as a leaf-miner of *Vitis* in India, but separable from it by golden-metallic markings.

Biological notes

Host plant : *Parthenocissus tricuspidata* (Sieb. et Zucc.) Planchon (Jap. name : Tsuta).

Mine (fig. 6) : Linear-blotch, full depth type; the larva at first making whitish brown linear mine, proceeds in an irregular wavy gallery, sometimes spiral at the beginning; about 60 mm. in length. After continuing to feed in this linear track for some distance, the mine develops rather abruptly into a pale brown irregular blotch ; feeding area about 284 mm².

Frass : Blackish brown grains of frass arranged in a line along the center of the linear mine, but in blotch mine being scattered a large pale brown patch in the middle of the mine.

Case : Elliptical, with a longitudinal ridge at the middle of each face; brown to dark brown; 3.5 × 1.8 mm.—4.4 × 2.6 mm. in size. Thorn-like projections, pointed and arranged near both ends; whitish to pale brown ; 9-16 in number; 0.36—0.48 mm. in length. The cases are settled to some suitable places by one or two silken strands.

Development : Number of generations unknown, but the larvae were collected on the leaves of the host plant at the end of October on Yakushima; they

passed the winter as a non-feeding larval stage, and appeared in July of the following year.




Fig. 6. Leaf of *Parthenocissus tricuspidata* Planchon mined by the larva of *Antispila iviella* sp. nov. and the larval case.

***Antispila hikosana* sp. nov.**

(Pl. 7, fig. 36; Pl. 8, fig. 44)

♂♀. 8-9 mm. Head, antennae, thorax and abdomen shining bronzy-fuscos. Palpi pale ochreous.

Forewings, vein 4+5 from near the upper apex of discoidal cell, vein 9 stalked with 7+8 at some distance beyond the upper apex of discoidal cell; bronzy-fuscos with coppery reflections, becoming purplish reflections towards apex; basal area shining dark leaden-fuscos; markings pale golden-metallic; oblique transverse fascia rather narrow, constricted on fold and broadened towards dorsum, rarely interrupted in disc; posterior costal spot triangular, not reaching half across wing, and dorsal spot rather large triangular, slightly before costal spot, reaching half across wing; cilia pale greyish fuscos, basal half bronzy-fuscos with brassy reflections. Hindwings, veins 2 and 6 absent, vein 7 near apex; pale fuscos with feeble purplish lusters; cilia pale fuscos.

Male genitalia (fig. 2; Pl. 4, fig. 10): Edge of scaphium slightly concave at middle; valva with two swellings at posterior margin; projection of ampulla short and slender; apex of aedoeagus with a bifurcated appendage and a pouch-shaped projection; phallobase anchor-shaped.

Female genitalia (Pl. 6, figs. 20 & 28): Ovipositor five pointed, edge of the points straightened, center point indent; enlarged part of ductus bursae with

two intricate patterns and many microscopical spines.

Holotype : ♂, Mt. Hikosan, N. Kyushu, 21 Apr. 1956 (H. Kuroko).

Allotype : ♀, the same as holotype, 21 Apr. 1956 (H. Kuroko).

Paratypes: 2 ♂♂, 2 ♀♀, the same as holotype, 21 Apr. 1956; 1 ♂, 1 ♀, 23 Apr. 1956 (H. Kuroko).

Distribution : Japan (Kyushu).

This new species is most closely related to *A. pfeifferella* Hübner of which the larva is known as a leaf-miner of *Cornus* in Europe, but separable from the latter in having coppery reflections on the forewings and rather shorter projection of ampulla in male.

Biological notes

Host plants : *Cornus controversa* Hemley (Jap. name : Mizuki), *C. brachypoda* C. A. Mey. (Jap. name : Kumano-mizuki).

Mine (fig. 7) : Linear-blotch, full depth, similar to that made by the larva of *A. corniella* Kuroko, but much larger; linear mine extends along the leaf margin, pale greyish fuscous, and the mines formed by the larvae of 1st instar, 2nd instar and 3rd instar, measure about 8 mm., 10 mm. and 12 mm. in length respectively, the width gradually increased. Blotch mine, formed by 4th instar larva, expands along or near the leaf margin, with dark greenish grains figured by feeding tracks; pale fuscous green. Usually one mine is observed in a single leaf.

Frass : In linear mine, blackish grains of frass form a row occupying the whole of the width, but in blotch mine they are scattered irregularly.

Case : Rather broad elliptical, with a longitudinal ridge at the middle of each face; dark greyish brown; 6.0 x 4.5 mm. - 6.5 x 4.7 mm. in size. Thorn-like projections, obtuse and arranged near both ends; whitish; 2-8 in number; 0.18-0.3 mm. in length.

Development : One generation in a year. The larvae feed from the end of May to June; the cut cases descend to the ground at the beginning of July; the larvae hibernate as a non-feeding larval stage in their cases and change to




Fig. 7. Leaf of *Cornus brachypoda* C. A. Mey. mined by the larva of *Antispila hikosana* sp. nov. and the larval case.

pupae in the spring of the following year; the adults appear from the end of April to the beginning of May.

***Antispila hydrangifoliella* sp. nov.**

(Pl. 7, fig. 35 ; Pl. 8, fig. 39)

♂ ♀. 5-6 mm. Head, antennae, thorax and abdomen shining bronzy-fuscous. Palpi pale ochreous.

Forewings, venation as in *A. hikosana* Kuroko; dark bronzy-fuscous, reflections feeble; basal area shining leaden-bronze; markings silvery-white, metallic reflections feeble; transverse fascia moderately to rather narrow and nearly straight; posterior costal spot and dorsal spot moderately to rather small, sometimes becoming nearly to dot; cilia pale grey, basal half dark bronzy-fuscous. Hindwings, veins 2 and 6 absent, vein 7 to near apex of the wing; pale fuscous, with feeble purplish lusters; cilia pale fuscous.

Male genitalia (Pl. 5, fig. 13): Edge of scaphium rounded; ampulla moderately projected, slightly downward; sacculus swollen; saccus V-shaped; apex of aedoeagus with two mushroom-shaped and a clavate appendages; phallobase spear-shaped.

Female genitalia (Pl. 6, figs. 19 & 27): Ovipositor with about 14 blunt serrations; enlarged part of ductus bursae with many wrinkles and winding fillets.

Holotype: ♂, Mt. Hikosan, N. Kyushu. 24 July 1954 (H. Kuroko).

Allotype: ♀, the same as holotype, 25 July 1954 (H. Kuroko).

Paratypes: 2 ♀♀, the same as holotype, 15 & 23 July 1954; 2 ♀♀, 28 & 29 July 1957; 1 ♂, 9 Aug. 1957 (H. Kuroko).

Distribution: Japan (Kyushu).

This species is allied to *A. hydrangaeella* Chambers, of which the larva is known as a leaf-miner of *Hydrangea* in N. America, but the apical spot is absent in the new species. The larvae may be observed in large numbers on Mt. Hikosan at the end of autumn.

Biological notes

Host plants: *Hydrangea macrophylla* (Thunb.) Sering var. *acuminata* (Sieb. et Zucc.) Makino (Jap. name: Yama-ajisai), *H. petiolaris* Siebold et Zuccarini (Jap. name: Gotbuzuru).

Mine (Pl. 9, fig. 48): Linear-blotch, full depth type; linear mine pale yellowish green to pale brown, semitransparent; about 80 mm. in length; proceeds in an irregular wavy gallery, somewhat confined by ribs. After 3rd moult, the mine expands into an irregular, semitransparent, whitish green blotch, often along the leaf margin; feeding area 224-264 mm? Usually one mine is found on a single leaf, sometimes 2 mines are made.

Frass: Blackish grains of frass form a scattered row occupying nearly the whole of the linear mine, but in blotch mine they are thinly scattered, sometimes deposited along the margin of the mine.




Fig. 8. Case of the larva of
Antispila hydrangifoliella
sp. nov.

Case (fig. 8) : Elliptical, broadened at the middle of both sides, with a longitudinal ridge at the middle of each face; light brown ; 4.0 x 2.5 mm. - 4.5 x 2.7 mm. in size. Thorn-like projections rounded and arranged near both ends; whitish; 3-7 in number; 0.3-0.42 mm. in length.

Development : One generation in a year. The larvae feed from the middle to the end of October, pass the winter as a non-feeding larval stage in the cut cases; pupation occurs at the middle of July, and the adults appear from the end July to the beginning of August.

***Antispila purplella* sp. nov.**

(Fig. 1; Pl. 7, fig. 37; Pl. 8, fig. 43)

♂♀. 6.5-7.5 mm. Head, antennae, thorax and abdomen shining bronzy-fuscous. Palpi pale fuscous.

Forewings, vein 4+5 from the upper apex of discoidal cell, vein 9 stalked with 7+8 at some distance beyond the upper apex of discoidal cell; dark bronzy-fuscous with reddish or purplish reflections; basal area shining dark leaden-fuscous ; markings silvery-metallic with somewhat coppery reflections ; transverse fascia slightly curved but not obliqued, its posterior margin obscured and with purplish lusters; posterior costal spot triangular, rather small, and dorsal spot narrow triangular, barely reaching half across wing; cilia fuscous, basal half with reddish or purplish reflections. Hindwings, veins 2 and 3 stalked, veins 6 and 7 stalked; fuscous with feeble purplish lusters; cilia pale fuscous.

Male genitalia (Pl. 4, fig. 12) : Edge of scaphium slightly convexed at middle; valva nearly triangular; ampulla rather prolonged, slightly upward; comb of teeth rather large ; sacculus with a small swelling; basal edge of saccus with a triangular sclerotized plate; apex of aedoeagus with a bended plate, a pouch-shaped appendage and a bifurcated appendage; apex of phallobase nearly equilateral triangle.

Female genitalia (Pl. 6, figs. 21 & 29) : Ovipositor five pointed, edge of points rounded, central point indent ; enlarged part of ductus bursae with two conspicuous filaments and twin intricate patterns.

Holotype : ♂, Mt. Hikosan, N. Kyushu, 3 June 1956 (H. Kuroko).

Allotype : ♀, the same as holotype, 4 June 1956 (H. Kuroko).

Paratypes : 4 ♂♂, 4 ♀♀, the same as holotype, 1-4 June 1956 (H. Kuroko).

Distribution : Japan (Kyushu).

This new species is easily distinguishable from the other species by the transverse fascia not oblique.

Biological notes

Host plants: *Cornus controversa* Hemsley (Jap. name : Mizuki), *C. brachypoda* C. A. Mey. (Jap. name : Kumano-mizuki).

Mine and frass : Similar to those made by the larva of *A. hikosana* Kuroko, but the mine of this species being made in autumn.

Case (fig. 9) : Elliptical, with a longitudinal ridge at the middle of each face; dark greyish brown; 4.2 x 2.3 mm. - 4.8 x 3.0 mm. in size. Thorn-like projections pointed and arranged near both ends ; pale brown ; 8-19 in number; 0.18-0.36 mm. in length.

Development : One generation in a year. The larvae feed from the end of September to the beginning of October; when full fed, the larvae fall to the ground with the cut cases, and change to pupae at the end of May of the following year; the adults appear at the beginning of June.




Fig. 9. Case of the larva of *Antispila purplella* sp. nov.

Explanation of Plates

Plate 4

Male genitalia, showing scaphium, valva, saccus and aedoeagus.

Fig. 10. *Antispila hikosana* sp. nov.

Fig. 11. *Antispila orbiculella* sp. nov.

Fig. 12. *Antispila purplella* sp. nov.

Plate 5

Male genitalia, showing the same parts as Plates 4.

Fig. 13. *Antispila hydrangifoliella* sp. nov.

Fig. 14. *Antispila ampelopsia* sp. nov.

Fig. 15. *Antispila corniella* sp. nov.

Plate 6

Female genitalia, showing enlarged part of ductus bursae and ovipositor.

- Figs. 16 & 24. *Antispila orbiculella* sp. nov.
 Figs. 17 & 25. *Antispila iviella* sp. nov.
 Figs. 18 & 26. *Antispila ampelopsia* sp. nov.
 Figs. 19 & 27. ***AntispiZa*** *hydrangifoliella* sp. nov.
 Figs. 20 & 28. *Antispila hikosana* sp. nov.
 Figs. 21 & 29. *Antispila purplella* sp. nov.
 Figs. 22 & 23. *Antispila corniella* sp. nov.

Plate 7

Venation


- Fig. 30. *Antispila pfeifferella* Hiibner
 Fig. 31. ***AntispiZa*** *orbiculella* sp. nov.
 Fig. 32. ***AntispiZa*** *ampelopsia* sp. nov.
 Fig. 33. ***AntispiZa*** *corniella* sp. nov.
 Fig. 34. *Antispila iviella* sp. nov.
 Fig. 35. *Antispila hydrangifoliella* sp. nov.
 Fig. 36. *Antispila hikosana* sp. nov.
 Fig. 37. *Antispila purplella* sp. nov.


Plate 8


- Fig. 38. *Antispila orbiculella* sp. nov.
 Fig. 39. *Antispila hydrangifoliella* sp. nov.
 Fig. 40. *Antispila ampelopsia* sp. nov.
 Fig. 41. *Antispila corniella* sp. nov.
 Fig. 42. *Antispila iviella* sp. nov.
 Fig. 43. *Antispila purplella* sp. nov.
 Fig. 44. *Antispila hikosana* sp. nov.


Plate 9

- Fig. 45. Leaf of *Ampelopsis brevipedunculata* Trautvetter mined by the larva of *Antispila orbiculella* sp. nov.
 Fig. 46. Leaf of *Ampelopsis brevipedunculata* Trautvetter mined by the larva of ***AntispiZa ampelopsia*** sp. nov.
 Fig. 47. Leaf of *Cornus brachypoda* C. A. Mey. mined by the larva of *Antispila corniella* sp. nov.
 Fig. 48. Leaf of ***Hydrangea macrophylla*** Seringe var. *acuminata* Makino mined by the larva of *Antispila hydrangifoliella* sp. nov.












30




31



32




33



34



35



36



37



