

Comparative Studies on Chloroplast Development and Photosynthetic Activities in C₃- and G-plants : II. Studies on Carbon Dioxide Fixation in Barley and Maize Leaves

Lee, Sun Hi

Laboratory of Plant Nutrition, Faculty of Agriculture, Kyushu University

Ikeda, Motoki

Laboratory of Plant Nutrition, Faculty of Agriculture, Kyushu University

Kang, Young Hee

Laboratory of Plant Nutrition, Faculty of Agriculture, Kyushu University

Yamada, Yoshio

Laboratory of Plant Nutrition, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23691>

出版情報 : 九州大学大学院農学研究院紀要. 24 (1), pp.1-9, 1979-08. Kyushu University
バージョン :
権利関係 :



Comparative Studies on Chloroplast Development and Photosynthetic Activities in C₃- and C₄-plants

II. Studies on Carbon Dioxide Fixation in Barley and Maize Leaves

**Sun Hi Lee*, Motolci Ikeda, Young Hee Kang*
and Yoshio Yamada**

Laboratory of Plant Nutrition, Faculty of Agriculture, Kyushu
University 46-02, Fukuoka 812

(Received April 5, 1979)

This study was undertaken to investigate the fixation patterns of carbon dioxide and chlorophyll accumulation following the chloroplast development by illumination to the etiolated leaves of barley and maize seedlings grown in the dark for a week. CO₂ fixation rate was remarkably increased after 3 hours illumination in barley and maize leaves. These results were very closely correlated with differentiation of grana and chlorophyll accumulation. At an early stage of the chloroplast development in barley leaves, CO₂ fixation patterns were not different from those in leaves cultured in a green house. In maize leaves, however, ¹⁴C was much incorporated into aspartate at an early stage of the chloroplast development in the leaves illuminated for 3 hours to the etiolated plant, and ¹⁴C was much incorporated into malate at later stages of the chloroplast development as in the leaves cultured in a green house.

INTRODUCTION

It is well known that the evolution of the photosynthetic apparatus from its simplest form in the green photosynthetic bacteria to the elaborate system of green plants is accompanied by the evolution of a more complicated pigment structure. Moreover, chloroplast is differentiated from its simplest system to very complicated system even in the same higher green plants. But the chloroplasts of vascular bundle sheath cells are devoid of grana even in mature leaves of maize seedlings. In the previous report (Lee et al., 1977), prolamellar bodies were formed in the etioplast of barley and maize leaves grown at a dark room for a week. These bodies were completely dispersed into lamellae in the chloroplasts of barley leaves illuminated by 1,000 ft-c of light for 3 hours after 6 days of dark germination, while in the chloroplasts within mesophyll cells of maize leaves the bodies still persisted during 18 hours under the same light condition and could not be observed in the chlo-

* Present address: Department of Biology, College of Science, Yonsei University, Seoul, Korea.

roplasts of parenchyma vascular bundle sheath cells illuminated for 6 hours. Grana were formed at the chloroplasts of chlorenchyma cells in barley leaves and of mesophyll cells in maize leaves, but not at the chloroplasts of vascular bundle sheath cells in maize leaves. Ultrastructural aspects of chloroplasts within parenchyma vascular bundle sheath cells of maize leaves were similar to those of chloroplasts within chlorenchyma cells of barley at an early stage of the chloroplast development.

On the other hand, phosphorylated compounds are synthesized as initial photosynthetic products (Quayle *et al.*, 1954; Calvin and Bassham, 1962) and compounds such as glycine, serine, phosphoglycolate and glycolate are much synthesized as photorespiratory products in C_3 -plants (Tolbert, 1971; Kisaki *et al.*, 1972; Zelitch, 1973). While in C_4 -plants dicarboxylic acids are synthesized as initial photosynthetic products (Kortschak *et al.*, 1965; Hatch and Slack, 1966; Slack and Hatch, 1967) and compounds related to the glycolate pathway are not much synthesized because of low photorespiration (Jackson and Volk, 1970; Osmond and Harris, 1971). Moreover, C_4 -plants can be divided into two groups, malate formers such as *Zea mays* and *Sorghum sudanense* in which malate is preferentially labelled on exposure to $^{14}CO_2$, and aspartate formers such as *Amaranthus edulis*, *Atriplex spongiosa* and *Panicum miliaceum* in which aspartate is preferentially labelled on brief exposure to $^{14}CO_2$ (Downton, 1970; Edwards and Gutierrez, 1972).

The aim of the present studies was to investigate the patterns of CO_2 fixation in barley and maize leaves following the chloroplast development.

MATERIALS AND METHODS

Barley (*Hordeum vulgare* L.) and maize (*Zea mays* L., Golden Cross Bantham) were grown by water culture in a dark room at $25^\circ C$ for one week. Leaf samples were prepared as described in the previous paper (Lee *et al.*, 1977).

After exposure to light (ca. 1,000 ft-c), leaves were detached by cutting under water to expose to $^{14}CO_2$ atmosphere in the light or in the dark for 3 minutes. The detached first foliage leaves were placed vertically in a fixation chamber and the bases of leaves were immersed into water to prevent from shortage of water during $^{14}CO_2$ fixation. The light intensity at the leaf surface was about 2,500 ft-c. $^{14}CO_2$ was generated in the CO_2 fixation chamber by pouring 50 % (v/v) lactic acid into $NaH^{14}CO_3$ solution (65 $\mu Ci/ml$) and CO_2 concentration was adjusted to about 0.03 %.

After feeding of $^{14}CO_2$ for 3 minutes, the leaves were transferred to boiling ethanol (70 %, v/v). Then they were extracted in sequence with boiling ethanol (50 %, v/v) and boiling water. The extracts were concentrated under reduced pressure in a rotary evaporator below $40^\circ C$ and passed through a column of Dowex-50 (H+) resin. And the fraction retained on the column was eluted with 2N NH_4OH . This fraction was used for amino acid analysis by paper chromatography (phenol : water=4 : 1, v/v). The effluent from the Dowex-50 (H+) column was passed through a column of Dowex-1 (CH_3COO^-) resin for organic acid analysis by Zelitch's method (Zelitch, 1965). The frac-

tion which was not adsorbed on the column was sugar. Radioactivity was measured by a liquid scintillation counter (Beckman Instrument Ltd., LS-250). Chlorophyll was measured by Comar's method (Comar and Zscheile, 1942).

Barley and maize were grown by an ordinary method in a green house as control to compare the CO₂ fixation pattern.

RESULTS

CO₂ assimilation

The rate of CO₂ assimilation was not increased both in light and dark fixation until 1 hour illumination stage of etiolated first foliage leaves of barley and maize plants. The rate of light and dark fixation in maize leaves was remarkably increased after 3 hours illumination stage, while the rate of dark fixation in barley was not increased throughout the chloroplast development. CO₂ assimilation rate was about 2 times higher at light fixation and 2-9 times higher at dark fixation in maize leaves than in barley leaves. However, the rate of light fixation was a little higher in barley leaves than in maize leaves cultured in a green house (Table 1). The rate of CO₂ assimilation and enzyme activities concerned in photosynthesis are decreased at aged leaves (Khanna and Sinha, 1973; Kennedy, 1976). It is therefore considered that the rate of CO₂ assimilation in the control maize leaves was due to aging of the first foliage leaves at this stage in green house cultivation.

Table 1. Total activity of ¹⁴C fixed during 3 minutes exposure to ¹⁴CO₂ following the chloroplast development (cpm/mg. fr. wt.).

Illumination	Fixation	Barley	Maize	Maize /Barley
0 hour	Dark	4		0.3
	Light	10	2	0.2
1 hour	Dark	6	14	2.3
	Light	15	30	2.0
3 hours	Dark	29	129	4.5
	Light	400	650	1.6
6 hours	Dark	23	106	4.6
	Light	600	1100	1.8
9 hours	Dark	29	125	4.3
	Light	1500	2300	1.5
18 hours	Dark	23	164	7.1
	Light	2500	4800	1.9
24 hours	Dark	38	278	7.3
	Light	3900	6100	1.6
Green house	Dark	51	449	8.8
	Light	5700	4500	0.8

Distribution of ¹⁴C incorporated into sugar, organic acid and amino acid fractions

I) Barley

As in the previous paper (Lee et al., 1977), grana formation initiated at the chloroplasts of the leaves illuminated for 3 hours. In this stage ¹⁴C was incorporated 21.7 % into a sugar, 44% into an organic acid and 33.2 % into

an amino acid fraction, respectively. As successive illumination to the etiolated leaves, the incorporation rate of ^{14}C was increased in a sugar fraction following the chloroplast development, while decreased gradually in organic acid and amino acid fractions. By the illumination for 24 hours, ^{14}C was incorporated 46.3 % into a sugar fraction, 23.2 % into an organic acid fraction and 29.2 % into an amino acid fraction, respectively. This result was the same as in the leaves cultured during the same period in a green house. ^{14}C was incorporated 42.3 % into a sugar, 24.4 % into an organic acid and 32.7 % into an amino acid fraction, respectively in the leaves cultured in a green house (Fig. 1).




Fig. 1. Proportion of total ^{14}C incorporated into sugar, organic acid and amino acid compounds in alcohol extract of barley leaves during 3 minutes photosynthesis following the chloroplast development.

2) Maize

The proportion of total ^{14}C fixed was 4.5 % in a sugar, 35.5% in an organic acid and 59.4 % in an amino acid fraction, respectively in the leaves of 3 hours illumination stage. In this stage of the chloroplast development total ^{14}C fixed was incorporated much more into an amino acid fraction than any other two fractions. But following the chloroplast development, ^{14}C incorporated into a sugar fraction was gradually increased. Generally, ^{14}C was much more incorporated into organic acids than amino acids with the chloroplast development. In the leaves of 24 hours illumination stage, the rate of total ^{14}C fixed was 41.7 % in a sugar, 28.8 % in an organic acid and 27 % in an amino acid fraction, respectively. This was the same tendency as in the




Fig. 2. Proportion of total ^{14}C incorporated into sugar, organic acid and amino acid compounds in alcohol extract of maize leaves during 3 minutes photosynthesis following the chloroplast development.

leaves cultured in a green house (Fig. 2).

Proportion of ^{14}C incorporated into photosynthetic intermediates of each fraction

^{14}C was distributed as a proportion of 6.6 % in serine+glycine, 4.8 % in phosphoglycolate and 9 % in glycolate in barley leaves of 3 hours illumination stage, but the proportion of total ^{14}C incorporated into aspartate and malate which are primary photosynthetic intermediates in C_4 -plants, was 7.1% and 8.5 %, respectively. In maize leaves illuminated for the same period as barley leaves, ^{14}C incorporated into aspartate and malate was 41.3 % and 22.1%. However, ^{14}C was incorporated 2.8 % into serine+glycine, 2.2 % into phosphoglycolate and 3.4% into glycolate. In barley leaves illuminated for 6 hours, ^{14}C was incorporated 17.3 % into serine+glycine, 4.6 % into phosphoglycolate, 10.4 % into glycolate, which were 32.3% of total fixed ^{14}C , but 5 % into aspartate and 12.5 % into malate. The same tendency was observed in leaves illuminated for 18 hours and 24 hours, and in the leaves cultured in a green house (Table 2).

Table 2. Proportion of total ^{14}C incorporated into individual compounds following the chloroplast development in barley leaves (% of total ^{14}C fixed).

Metabolic intermediate	Illumination				Green house
	3 hrs	6 hrs	18 hrs	24 hrs	
Aspartate	7.13	5.00	3.39	4.69	2.68
Glutamate	1.72	1.11	0.70	0.90	0.67
Serine + Glycine	6.60	17.27	12.99	11.05	16.00
Alanine	13.50		12.10	11.39	10.58
Other amino acids	4.26	11.10 2.09	1.62	1.19	2.74
	33.21	36.57	30.80	29.23	32.67
Glycolate	9.03	10.36	9.61	4.66	9.59
Malate	8.48	9.60	5.28	4.74	1.53
Citrate	19.27	12.49	11.10	8.28	8.30
Phosphoglycerate	1.81	0.87	1.50	1.12	0.59
Phosphoglycolate	4.82	4.63	6.12	4.17	4.27
Other organic acids	0.60	0.14	0.15	0.24	0.11
	44.01	38.09	33.76	23.21	24.39
Sugar	21.70	24.40	34.57	46.27	42.26
Residue	1.08	0.94	0.87	1.29	0.68

In maize leaves illuminated for 6 hours, its proportion was 24.8 % in aspartate and 29.7% in malate. Generally, there was no characteristically different pattern of ^{14}C incorporation in each stage of the chloroplast development compared with that in leaves cultured in a green house (Table 3). However, at an early stage of the chloroplast development ^{14}C was much more incorporated into aspartate than into malate in maize leaves (Table 3).


Chlorophyll accumulation

Chlorophyll was accumulated linearly after 3 to 6 hours lag phase in the etiolated leaves of barley and maize seedlings as the chloroplast development.

Table 3. Proportion of total ^{14}C incorporated into individual compounds following the chloroplast development in maize leaves(% of total ^{14}C fixed).

Metabolic intermediate	Illumination				Green house
	3 hrs	6 hrs	18 hrs	24 hrs	
Aspartate	41.34	24.81	12.37	12.10	4.56
Glutamate	4.37	3.60	3.03	1.25	0.78
Serine +Glycine	2.81	4.62	5.98	3.72	2.98
Alanine	8.96	7.90	11.98	8.77	8.24
Other amino acids	1.88	1.54	1.37	1.14	0.76
	59.36	41.93	34.73	26.98	17.32
Glycolate	3.40	3.59	5.86	3.34	5.45
Malate	22.07	29.74	33.58	13.52	21.39
Citrate	5.99	8.28	7.72	4.05	13.37
Phosphoglycerate	1.55	2.27	1.54	4.26	1.44
Phosphoglycolate	2.16	2.58	1.16	3.05	2.01
Other organic acids	0.36	0.31	0.14	0.54	0.11
	35.53	46.77	50.00	28.76	43.77
Sugar	4.52	9.60	14.13	41.66	37.69
Residue	0.59	1.70	1.14	2.60	1.22

The amount of chlorophyll was higher in barley than in maize leaves (Fig. 3). Grana were not well developed and chlorophyll b was deficient in the chloroplasts of vascular bundle sheath cells in maize leaves. But grana were present and chlorophyll b was not deficient in the chloroplasts of mesophyll cells in maize leaves and of chlorenchyma cells in barley leaves (Thorner *et al.*, 1967; Woo *et al.*, 1970; Pylotis *et al.*, 1971; Lee *et al.*, 1977). Therefore, the amount of chlorophyll was considered to be still higher in barley than in maize leaves.

**Fig. 3.** Total chlorophyll content following the chloroplast development.

DISCUSSION

CO_2 fixation rate was not so much increased until 1 hour illumination

stage in barley as well as in maize leaves, but increased afterwards as the chloroplast development. These results were very closely correlated with an increase in total chlorophyll contents and grana formation following the chloroplast development as reported in the previous paper (Lee *et al.*, 1977). CO₂ fixation was about 2 times higher in light fixation and 2-9 times higher in dark fixation in maize leaves than in barley leaves. These suggest that phosphoenolpyruvate carboxylase activity is higher even in the etiolated leaves of maize seedlings than barley seedlings and also is stimulated by light (Stabenau, 1972; Kamiya and Miyachi, 1975).

Rhodes and Yemm (1966) reported that grana formation was promoted by light and coincided with chlorophyll formation and an increase in photosynthetic activity in barley seedlings. Wieckowski (1969), using very young bean seedlings, found that O₂ evolution was first detectable after about 2 hours of illumination, and this corresponded with the appressed lamellae in the plastid. But grana formation is generally correlated with photosystem II, but not with O₂ evolution (Anderson and Boardman, 1964; Gyldenholm and Whatley, 1968; Sane *et al.*, 1970). And also Robertson and Laetsch (1974), using each region of a barley leaf, reported that membrane appression was not strictly correlated with the presence of chlorophyll and photosynthetic activities. In addition, a barley mutant lacking chlorophyll b exhibited high photosystem I and II activities in spite of reduction of appressed lamellae (Highkin and Frenkel, 1962; Goodchild *et al.*, 1966). We also reported that there were many appressed lamellae even in the etioplasts (Lee *et al.*, 1977). In this result, chlorophyll could be detected after about 1 hour illumination. So we agreed that appressed lamellae were not strictly correlated with the presence of chlorophyll. But chlorophyll formation was stimulated by light. Both ATP and NADPH were generated as chlorophyll harvested quanta and also Hill reaction was stimulated (Gyldenholm and Whatley, 1968; Oelge-Karow and Butler, 1971). Such cumulative results led us to conclude that an increase of CO₂ fixation was closely correlated with chlorophyll formation and accumulation as the chloroplast development.

On the other hand, in maize leaves, a malate former, proportion of total ¹⁴C incorporated into aspartate was much higher than into malate in the leaves illuminated for 3 hours. With the chloroplast development, ¹⁴C was much incorporated into malate as in the leaves cultured in a green house. Hatch (1973) and Hatch and Mau (1973) reported that aspartate aminotransferase was about equally distributed between mesophyll and bundle sheath cells. Aspartate aminotransferase in mesophyll cells was associated with chloroplasts or other subcellular organelles, but the major aspartate aminotransferase isoenzyme in bundle sheath cells was associated with mitochondria in aspartate formers such as *Atriplex spongiosa* and *Panicum miliaceum*. However, in maize seedlings, mitochondria were well differentiated even in the mesophyll as well as in the vascular bundle sheath cells after 3 days of dark germination, but chloroplasts were not differentiated until 3 hours illumination to etiolated leaves of 6 days dark-grown seedlings (Lee *et al.*, 1977). And also aspartate aminotransferase activity was not increased following the

chloroplast development (Lee *et al.*, unpublished data). Moreover, the activity of NADP malate dehydrogenase was low in etiolated leaves of maize seedlings grown at dark, but increased 10–20 folds together with chlorophyll when leaves were illuminated (Johnson and Hatch, 1970). Therefore, we considered that oxaloacetate produced by phosphoenolpyruvate carboxylase will be transformed into aspartate rather than malate by aspartate aminotransferase originated from mitochondria and other organelles except chloroplasts at an early stage of the chloroplast development, but by further development of chloroplasts oxaloacetate can be transformed into malate by increased NADP malate dehydrogenase.

REFERENCES

- Anderson, J. M. and N. K. Boardman 1964 Studies on the greening of dark-grown bean plants. II Development of photochemical activity. *Aust. J. Biol. Sci.*, 17: 93-101
- Calvin, M. and J. A. Bassham 1962 *The Photosynthesis of Carbon Compounds*. W. A. Benjamin Inc., New York, pp. 10
- Comar, C. L. and F. P. Zscheile 1942 Analysis of plant extracts for chlorophyll a and b by a photoelectric spectrophotometric method. *Plant Physiol.*, 17: 198-209
- Downton, W. J. S. 1970 Preferential C₄-dicarboxylic acid synthesis, the postillumination CO₂ burst, carboxylic transfer step, and grana configuration in plants with C₄-photosynthesis. *Can. J. Bot.*, 48: 1795-1800
- Edwards, G. E. and M. Gutierrez 1972 Metabolic activities in extract of mesophyll and bundle sheath cells of *Panicum miliaceum* (L.) in relation to the C₄-dicarboxylic acid pathway of photosynthesis. *Plant Physiol.*, 50: 728-732
- Goodchild, D. J., H. R. Highkin and N. K. Boardman 1966 The fine structure of chloroplasts in a barley mutant lacking chlorophyll b. *Exp. Cell Res.*, 43: 684-688
- Gyldenholm, A. O. and F. R. Whatley 1968 The onset of photophosphorylation in chloroplast isolated from developing bean leaves. *New Phytol.*, 67: 461-468
- Hatch, M. D. 1973 Separation and properties of leaf aspartate aminotransferase and alanine aminotransferase isoenzymes operative in C₄ pathway of photosynthesis. *Arch. Biochem. Biophys.*, 156: 207-214
- Hatch, M. D. and S. L. Mau 1973 Activity, location, and role of aspartate aminotransferase and alanine aminotransferase isoenzymes in leaves with C₄ pathway of photosynthesis. *Arch. Biochem. Biophys.*, 156: 196-206
- Hatch, M. D. and C. R. Slack 1966 Photosynthesis by sugarcane leaves. A new carboxylation reaction and the pathway of sugar formation. *Biochem. J.*, 101: 103-113
- Highkin, H. R. and A. W. Frenkel 1962 Studies of growth and metabolism of a barley mutant lacking chlorophyll b. *Plant Physiol.*, 37: 814-820
- Jackson, W. A. and R. J. Volk 1970 Photorespiration. *Ann. Rev. Plant Physiol.*, 21: 385-432
- Johnson, H. S. and M. D. Hatch 1970 Properties and regulation of leaf nicotinamide-adenine dinucleotide phosphate malate dehydrogenase and 'malic' enzyme in plant with the C₄-dicarboxylic acid pathway of photosynthesis. *Biochem. J.*, 119: 273-280
- Kamiya, A. and S. Miyachi 1975 Blue light-induced formation of phosphopyruvate carboxylase in colorless *Chlorella* mutant cells. *Plant Cell Physiol.*, 16: 729-736
- Kennedy, R. A. 1976 Relationship between leaf development, carboxylase enzyme activities and photorespiration in the C₄-plant *Portulaca oleracea* L., *Planta* (Berl.), 128 : 149-154
- Khanna, R. and S. K. Sinha 1973 Change in predominance from C₄ to C₃ pathway following anthesis in Sorghum. *Biochem. Biophys. Res. Commun.*, 52: 121-124

- Kisaki, T., A. Imai and N. E. Tolbert 1971 Intracellular localization of enzymes related to photorespiration in green leaves. *Plant Cell Physiol.*, **12**: 267-273
- Kortschak, H. P., C. E. Hartt and G. O. Burr 1965 Carbon dioxide fixation in sugarcane leaves. *Plant Physiol.*, **40**: 209-213
- Lee, S. H., M. Ikeda and Y. Yamada 1977 Comparative studies on chloroplast development and photosynthetic activities in C₃- and C₄-plants. 1. Studies on ultrastructure of developing chloroplasts within vascular bundle sheath and mesophyll cells of barley and maize leaves. *J. Fac. Agr., Kyushu Univ.*, **22**: 65-74
- Oelge-Karow, H. and W. L. Butler 1971 The development of photophosphorylation and photosynthesis in greening bean leaves. *Plant Physiol.*, **48**: 621-625
- Osmond, C. B. and B. Harris 1971 Photorespiration during C₄ photosynthesis. *Biochim. Biophys. Acta*, **234**: 270-282
- Pylotiot, N. A., K. C. Woo and W. J. S. Downton 1971 Thylakoid aggregation correlated with chlorophyll a/ chlorophyll b ratio in some C₄ species. In "Photosynthesis and Photorespiration," ed. by M. D. Hatch, C. B. Osmond and R. O. Slatyer, John Wiley and Sons, New York, pp. 406-412
- Quayle, J. R., R. C. Fuller, A. A. Benson and M. Calvin 1954 Enzymatic carboxylation of ribulose diphosphate. *J. Amer. Chem. Soc.*, **76**: 3610-3611
- Rhodes, M. J. C. and E. W. Yemm 1966 The development of chloroplasts and photosynthetic activities in young barley leaves. *New Phytol.*, **65**: 331-342
- Robertson, D. and W. M. Laetsch 1974 Structure and function of developing barley plastids. *Plant Physiol.*, **54**: 148-159
- Sane, P. V., D. J. Goodchild and R. B. Park 1970 Characterization of chloroplast photosystems 1 and 2 separated by a non-detergent method. *Biochim. Biophys. Acta*, **216**: 162-178
- Slack, C. R. and M. D. Hatch 1967 Comparative studies on the activity of of carboxylases and other enzymes in relation to the new pathway of photosynthetic carbon dioxide fixation in tropical *grasses*. *Biochem. J.*, **103**: 660-665
- Stabenau, H. 1972 Aktivitätsänderungen von Enzymen bei *Chlorogonium elongatum* unter dem Einfluß von rotem und blauem Licht. *Z. Pflanzenphysiol.*, **67**: 105-112
- Thornber, J. P., R. P. F. Richard, C. A. Smith and J. L. Bailey 1967 Studies on the nature of the chloroplast lamella. I. Preparation and some properties of two chlorophyll-protein complex. *Biochemistry*, **6** : 391-396
- Tolbert, N. E. 1971 Leaf peroxisomes and photorespiration. In "Photosynthesis and Photorespiration," ed. by M. D. Hatch, C. B. Osmond and R. O. Slatyer, John Wiley and Sons, New York, pp. 458-471
- Wieckowski, S. 1969 Studies on the activity and ultrastructure of the photosynthetic apparatus in the earliest stage of primary bean leaves development. *Acta Soc. Bot. Pol.*, **38**: 103-114
- Woo, K. C., J. M. Anderson, N. K. Boardman, W. J. S. Downton, C. B. Osmond and S. W. Thorne 1970 Deficient photosystem II in agranal bundle sheath chloroplast of C₄-plant. *Proc. Nat. Acad. Sci. U. S. A.*, **67**: 1869-1876
- Zelitch, I. 1965 The relation of glycolic acid synthesis to the primary photosynthetic carboxylation reaction in leaves. *J. Biol. Chem.*, **240**: 1869-1876
- Zelitch, I. 1973 Plant productivity and the control of photorespiration. *Proc. Nat. Acad. Sci. U. S. A.*, **70** : 579-584