Effect of Some Hydroxylamine-Metabolites on the Priming Activity for DNA Polymerase

Tomita, Yoshifumi

Department of Public Health, School of Medicine, Kurume University

Fujiki, Hiroaki

Institute of Food Technology, Meiji Seika Co. Ltd.

Shinohara, Kazuki

Food Chemistry Institute Faculty Of Agriculture Kyushu University

Murakami, Hiroki

Food Chemistry Institute Faculty Of Agriculture Kyushu University

他

<https://doi.org/10.5109/23727>

出版情報：九州大学大学院農学研究院紀要. 25 (4), pp.161-166, 1981-04. Kyushu University
バージョン：
権利関係：

Effect of Some Hydroxylamine-Metabolites on the Priming Activity for DNA Polymerase

Yoshifumi Tomita*, Hiroaki Fujiki**, Kazuki Shinohara,
Hiroki Murakami and Hirohisa Omura***

Laboratory of Food Chemistry, Faculty of Agriculture,
Kyushu University 46-09, Fukuoka 812

(Received December 4, 1980)

Mutagenic and virogenic hydroxylamine-metabolites, such as nitrite, hydroxylamine, oxime and hydrogen peroxide, stimulated more or less the priming activity of template DNA for DNA polymerase, although their actions were not so high as those of enediol reductones. Hydroxylamine and hydrogen peroxide were more effective for the stimulation than the others.

INTRODUCTION

It was reported that some hydroxylamine-metabolites, such as nitrite (Yamafuji *et al.*, 1971 a), hydroxylamine (Yamafuji and Hashinaga, 1966, 1967; Yamafuji *et al.*, 1966, 1971 a; Omura and Yamaguchi, 1967; Omura *et al.*, 1973), oxime (Murakami and Yamafuji, 1970; Yamafuji *et al.*, 1971 a, b; Iiyama *et al.*, 1973; Omura *et al.*, 1973) and hydrogen peroxide (Yamafuji and Uchida, 1966, 1968) caused breakage of nucleic acids. These substances were found to be mutagenic and/or virogenic agent. Yamafuji (1970) presented a working hypothesis that some appropriate cleavage introduced in chromosomal nucleic acids might induce mutagenesis, virogenesis, carcinogenesis or carcinostasis, as the initiating stage of these cellular anomalies. Further studies elucidated that cleavage in DNA brought about some alteration of the priming activity for DNA-dependent RNA polymerase, stimulating or repressing according to the degree of the cleavage (Yamafuji *et al.*, 1972; Iiyama *et al.*, 1973; Omura *et al.*, 1973). On the other hand, antitumor reductones induced changes of the priming activity for DNA polymerase as well as RNA polymerase in connection with DNA cleavage by them (Omura *et al.*, 1975; Tomita *et al.*, 1979). It is expected that hydroxylamine-metabolites may affect the priming activity for DNA polymerase, since they have abilities to cleave nucleic acids and to alter the priming activity for RNA polymerase.

* Present address: Department of Public Health, School of Medicine, Kurume University, Kurume 830

** Present address: Institute of Food Technology, Meiji Seika Co. Ltd., Kawasaki 210

*** Reprint request should be addressed to this author.

MATERIALS AND METHODS

Reagents: Hydroxylamine hydrochloride (the special grade), sodium nitrite (the special grade) and hydrogen peroxide (the first grade) were purchased from Wako Chemical Co. Galactose oxime was kindly supplied by Dr. Iio (Kumamoto Women University). ^3H -thymidine triphosphate (^3H -TTP, 20.9 Ci/mmol) was obtained from New England Nuclear.

DNA: DNA from calf thymus was purified according to the method reported in the previous paper (Omura *et al.*, 1975).

DNA polymerase: DNA polymerase was prepared from *E. coli* strain B after Richardson *et al.* (1964) by autolysis, ammonium sulfate fractionation, DEAE cellulose and phosphocellulose chromatographies, as described in the foregoing paper (Tomita *et al.*, 1979).

Treatment of DNA with reagent: DNA (100 $\mu\text{g}/\text{ml}$) was incubated with 1 μM or 1 mM reagent in 25 mM tris-HCl buffer (pH 6.8) at 37°C for 1 hr. After dialysis against $0.1 \times \text{SSC}$ (0.15 M NaCl-0.015 M Na-citrate) at 4°C for 24 hr, the priming activity of the DNA was estimated.

Priming activity assay: The assay mixture (400 μl in total) contained 20 μmoles of tris-HCl buffer (pH 7.4), 2 μmoles of dithiothreitol, 2 μmoles of MgCl_2 , 20 nmoles of dATP, dGTP, dCTP each, 14 pmoles of ^3H -TTP (0.41 nCi), 20 μg of DNA and 12.5 μg of DNA polymerase preparation. An aliquot (50 μl) of the mixture incubated at 37°C was transferred into 2 ml of cold 0.5 N perchloric acid at 5 and 25 min after the beginning of the incubation and allowed to stand for 30 min. Acid-insoluble fraction was collected on glass filter paper (Whatman GF/C) and washed successively with 3 ml of cold 0.5 N perchloric acid, cold water and 95 % cold ethyl alcohol. Radioactivity of the dried filter paper was counted by a Beckman LS-250 liquid scintillation counter with 0.4 % PPO-toluene as scintillant.

RESULTS

(1) Hydroxylamine

The effect of hydroxylamine on the template DNA of DNA-polymerase was assayed. As described in Fig. 1, the priming activity was enhanced with a low concentration of 1 μM hydroxylamine.

(2) Nitrite

DNA was treated with sodium nitrite as with hydroxylamine. Result in Fig. 2 indicates that the effect of 1 μM nitrite is not remarkable. However, a slight enhancement of the priming activity was observed with the treatment of 1 mM nitrite.

(3) Galactose oxime

Galactose oxime showed a similar effect to that of nitrite for the priming activity, as seen in Fig. 3.

(4) Hydrogen peroxide

DNA treatment with hydrogen peroxide was samely performed with hy-

Fig. 1. Priming activity of DNA treated with NH_2OH . DNA ($100 \mu\text{g/ml}$) was incubated with $1 \mu\text{M}$ or 1mM NH_2OH in 25mM tris-HCl buffer (pH 6.8) at 37°C for 1 hr. After dialysis against $0.1 \times \text{SSC}$ (0.15M NaCl- 0.015M Na-citrate) at 4°C for 24 hr, priming activity of the DNA was assayed. x---x: control, o---o: $1 \mu\text{M}$ NH_2OH , ●—●: 1mM NH_2OH .

Fig. 2. Priming activity of DNA treated with NaNO_2 . DNA treatment and the priming assay were described in the legend of Fig. 1, except that NaNO_2 was used instead of NH_2OH . x---x: control, o---o: $1 \mu\text{M}$ NaNO_2 , ●—●: 1mM NaNO_2 .

Fig. 3. Priming activity of DNA treated with galactose oxime. DNA treatment and the priming activity assay were described in the legend of Fig. 1, except that galactose oxime was used instead of NH_2OH . x---x: control, o--o: 1 μM galactose oxime, ●—●: 1 mM galactose oxime.

Fig. 4. Priming activity of DNA treated with H_2O_2 . DNA treatment and the priming activity assay were described in the legend of Fig. 1, except that H_2O_2 was used instead of NH_2OH . x---x: control, o--o: 1 μM H_2O_2 , ●—●: 1 mM H_2O_2 .

droxylamine, although peroxide is not hydroxylamine-metabolite. The effect of hydrogen peroxide on the template DNA is shown in Fig. 4. Hydrogen peroxide caused an enhancement of the priming activity by the treatment even with 1 μ M of the concentration.

DISCUSSION

The bacterial mutagenicity assay and animal experiments suggested the overlapping of mutagens and carcinogens. Yamafuji (1970) demonstrated that the cellular anomalization including mutagenesis, virogenesis, carcinogenesis and carcinostasis might be induced by some proper scissions in the chromosomal DNA. Along this working hypothesis, it was disclosed that the breakage of nucleic acid strands was brought about by some mutagens, virogens, carcinogens and antitumoric substances and that such breakages caused the alteration of RNA polymerase, DNA polymerase and protein synthesis. Hydroxylamine-metabolites are the first ones established as the chemical virogen as well as mutagen (Yamafuji and Shirozu, 1944; Yamafuji and Omura, 1950; Yamafuji and Yoshihara, 1950). As cited above, cleavage of nucleic acids and alteration of the priming activity for RNA polymerase by them were disclosed. In this study, an analogous effect is shown for DNA polymerase too. These results may provide an additional support to the Yamafuji's hypothesis.

All reagents tested increased more or less the incorporation of ^3H -TMP into DNA, as shown in Figs. 1, 2, 3 and 4, although their effects were in general not so remarkable as those of ascorbic acid and its related substances (Omura *et al.*, 1975; Tomita *et al.*, 1975). Treatment with 1 mM reagents is more effective than with 1 μ M. Galactose oxime and nitrite were relatively moderate for the effect compared with hydroxylamine and hydrogen peroxide, as especially seen in 1 μ M treatment. In the case of ascorbic acid and its related substances, the enhancement of the priming ability was observed when single strand scissions were introduced into DNA strand, but double strand scissions which had been provoked in the much severer conditions decreased the ability. Within the range of concentrations applied, higher concentrations of reagents did not reduce the priming activity. Cleavage of nucleic acids by several DNA breaking reagents were accelerated in the coexistence of Cu^{2+} . Concerning with hydroxylamine-metabolites, cooperation of Cu^{2+} should be examined. The precise mechanism for the increase of incorporation of ^3H -TMP into DNA is not clear at present, but a possible interpretation may be that 3'-OH terminal groups, initiation sites for DNA synthesis (Mittra and Kornberg, 1966), increase as the strands are subjected to scissions, as shown by Iiyama *et al.* (1973) for RNA polymerase. Thus, mutagenic and virogenic hydroxylamine-metabolites may modulate the priming ability for DNA synthesis depending on the mode and degree of strand breakages. Such modulation for the replication and transcription may be related to anomalization of cell functions through affecting on protein synthesis.

REFERENCES

- Iiyama, S., H. Murakami, M. Kuraki and H. Omura 1973 Action of DNA breaking reagents on RNA, deoxyribonucleotide and RNA polymerase reaction. *Nippon Nogei Kagaku Kaishi (J. Agr. Chem. Soc. Japan)*, **47**: 455-461
- Mitra, S. and A. Kornberg 1966 Enzymatic mechanism of DNA replication. *J. Gen. Physiol.*, **49**: 59-79
- Murakami, H. and K. Yamafuji 1970 Mode of action of some catecholamines and sugar oximes on deoxyribonucleic acid. *Enzymologia*, **38**: 337-345
- Omura, H., S. Iiyama, T. Fujii and K. Yamafuji 1973 Effect of virogenic hydroxylamine-metabolites on deoxyribonucleic acid and RNA polymerase. *J. Fac. Agr., Kyushu Univ.*, **17**: 181-186
- Omura, H., S. Iiyama, Y. Tomita, Y. Narazaki, K. Shinohara and H. Murakami 1975 Breaking action of ascorbic acid on nucleic acids. *J. Nutr. Sci. Vitaminol.*, **21**: 237-249
- Omura, H. and K. Yamaguchi 1967 Action of hydroxylamine on deoxyribonucleic acid in vitro. *Enzymologia*, **33**: 1-18
- Richardson, C. C., C. L. Schildkraut, H. V. Aposhian and A. Kornberg 1964 Enzymic synthesis of deoxyribonucleic acid XIV. Further purification and properties of deoxyribonucleic acid polymerase of *Escherichia coli*. *J. Biol. Chem.*, **239**: 222-232
- Tomita, Y., K. Shinohara, H. Fujiki, H. Murakami and H. Omura 1979 Effect of some reductones on DNA and RNA polymerases. *J. Fac. Agr., Kyushu Univ.*, **24**: 21-36
- Yamafuji, K. 1970 *Food, Cancer and Cytodifferentiation (40th Anniversary of Prof. K. Yamafuji's Research)*. Shukosha, Fukuoka, Japan
- Yamafuji, K., T. Fujii, Y. Kanegae and Y. Matsuo 1972 Action of carcinogenic tryptophan metabolites on deoxyribonucleic acid and RNA-polymerase. *Enzymologia*, **43**: 73-82
- Yamafuji, K. and F. Hashinaga 1966 Isolation of a previral deoxyribonucleic acid. *Nature*, **210**: 1158
- Yamafuji, K. and F. Hashinaga 1967 Hybridization between polyhedral viral and previral deoxyribonucleic acid. *Enzymologia*, **32**: 364-370
- Yamafuji, K., F. Hashinaga and T. Fujii 1966 Isolation and identification of polyhedral previral deoxyribonucleic acid from healthy silkworm cells. *Enzymologia*, **31**: 92-104
- Yamafuji, K. and H. Omura 1950 The formation of silkworm virus by acetoxime feeding. *Enzymologia*, **14**: 120-123
- Yamafuji, K., K. Shinohara, F. Yoshihara, M. Iio and Y. Torikai 1971a Breakage of nucleic acids as initiative process of cytodifferentiation and cytoanomalization. *Enzymologia*, **40**: 107-119
- Yamafuji, K., K. Shinohara, F. Yoshihara, H. Omura and N. Ogata 1971b Control of genes in silkworms by steroid hormones, catecholamines and hexose oximes. *Enzymologia*, **41**: 183-199
- Yamafuji, K. und Y. Shirozu 1944 Die Abhangigkeit der Neubildung des Virusproteins von der Erniedrigung der Katalase. *Biochem. Zs.*, **317**: 94-98
- Yamafuji, K. and Y. Uchida 1966 Liberation of adenine from deoxyribonucleic acid by hydrogen peroxide. *Nature*, **209**: 301
- Yamafuji, K. and Y. Uchida 1968 Effect of some mutagenic virogens and carcinogens on deoxyribonucleic acid. *Enzymologia*, **35**: 131-138
- Yamafuji, K. and F. Yoshihara 1950 On the virus production and oxime formation in silkworms fed with nitrites. *Enzymologia*, **14**: 124-127