

# Ecological Studies on *Formica yessensis* Forel, with Special Reference to Its Effectiveness as a Biological Control Agent of the Pine Caterpillar Moth in Korea : III. Nest Distribution and Environmental Conditions of Nest Sites

Kim, Chang Hyo

Division of Insect Natural Enemies, Institute of Biological Control, Faculty of Agriculture,  
Kyushu University

Murakami, Yozo

Division of Insect Natural Enemies, Institute of Biological Control, Faculty of Agriculture,  
Kyushu University

<https://doi.org/10.5109/23734>

---

出版情報 : 九州大学大学院農学研究院紀要. 26 (1), pp.9-19, 1981-10. Kyushu University  
バージョン :  
権利関係 :



**Ecological Studies on *Formica yessensis* Forel, with Special  
Reference to Its Effectiveness as a Biological Control  
Agent of the Pine Caterpillar Moth in Korea**

**III. Nest Distribution and Environmental  
Conditions of Nest Sites**

**Chang Hyo Kim\* and Yôzô Murakami**

Division of Insect Natural Enemies, Institute of Biological Control,  
Faculty of Agriculture, Kyushu University 46-13, Fukuoka 812

*(Received May 6, 1981)*

Seasonal change of the nest distribution of *Formicayessensis* in the hill and biotic and abiotic environmental conditions of nest sites were investigated in red pine stands in Gyeongsang-Nam-Do, Korea in 1973-77. The nests scarcely occurred in the upper zone of slopes and concentrated in the lower and middle zones. The distribution of nests changes seasonally but the density per unit area is relatively constant (0.014-0.022 per m<sup>2</sup>). The active nests increase in number from spring to summer by budding, then decrease in autumn by abandonment of nests. The life span of nests was 1-4 months in most cases. The abandonment occurs *more* frequently in newly founded and revived nests. Abandoned nests are re-utilized, the number of times the same nest was utilized being ranged between 1 and 11. The humidity above ground and soil moisture are important factors for establishing nests. The content of organic matters was higher in soils at the lower and middle zones inhabited by the ant than in the upper zone. While the content of the exchangeable Ca was lower in the former than the latter. In the upper zone the density of pine trees is relatively low and that of *Zoisia* is higher than the middle and lower zones. Such conditions are unfavourable for establishing nests. The lower and middle zones, on the contrary, are favourable because of the relatively high density of pine trees, but sufficient light intensity and presence of suitable plants for nest cores such as long graminaceous grasses. *Arundinella hirta*, *Pinus densiflora* (red pine trees) and *Miscanthus sinensis* f. *purpurascens* as well as stones serve as the core for building nests.

**INTRODUCTION**

*Formica (Formica) yessensis* Forel prefers grasslands close to the forest margin (Ito, 1971) or sparse secondary forests of the white birch and other deciduous trees (Morisita, 1945) as nesting sites in Japan. In Korea, however, red pine trees are relatively sparse in forests, and sufficient light intensity

---

\* Present address: College of Agriculture, Gyeongsang National University, Jinju 620, Korea.

provides favourable conditions for nesting of the ant (Kim and Murakami, 1980 a). The ant, however, inhabits only limited stands among red pine forests in Korea. Some physical and biotic conditions may be responsible to establish nests.

The present paper deals with seasonal changes of distribution, density and abundance of *F. yessensis* nests in the hill growing red pine trees in Gyeongsang-Nam-Do, Korea, and various environmental conditions of nest sites such as humidity above ground, chemical properties of the soil, densities of red pine trees and floor-layer plants and suitable nest cores.

## METHODS

In order to clarify the nest distribution in the hill, the slope was conveniently divided into three zones, the lower, the middle and the upper ones, approximately equal in distance, then the number of nests were counted in each zone. The investigation was made at two red pine stands in each locality in Gyeongsang-Nam-Do, Korea from October 20 to November 20, 1973. The localities are Jinyang, Goseung, Hadong, Sacheun, Sancheung and Namhae.

The seasonal census of distribution, density and abundance of nests was made by the quadrat method in a red pine stand in Jinseung (Station A), Jinyang-gun, Gyeongsang-Nam-Do. The home range in the slope of the stand was divided into 195 quadrats of  $10 \times 10$  m area. A precise map of the area was made on April 30, 1973, the day of the first census, and afterwards newly appeared, abandoned and revived nests were checked monthly from April to October in 1973-77.

One of the physical factors which may affect the distribution of nests, the relative humidity above ground was determined in the same study area with intervals of a week from May to November, 1975. Twelve screens were placed in the slope and the relative humidity at 10 cm above ground was recorded twice a day (at 10 : 00 and 15 : 00).

**Soil** chemical analysis was conducted with samples collected from the same slope on April 25, 1973. From each zone of the slope, three sites were selected for sampling soil. Five hundred g of soil sample was collected from each depth of 0-20, 20-40 and 40-60 cm at each site. In order to determine the content of exchangeable Ca, 500 ml of  $\text{CH}_3\text{COONH}_4$  was added to 5 g of each dried soil sample. After shaking for 30 min, it was filtered through a Toyo No. 2 filter paper. Then 20 ml distilled water was added to 5 ml of the filtrate. Five ml of 8 M-NaOH and 1 ml of 10 %  $\text{NH}_2\text{OH} \cdot \text{HCl}$  were added to it. It was then titrated with 0.005 M-EDTA using about 0.1 g of N-N indicator (2-hydroxy-1-(2-hydroxy-4-sulfo-1-naphthylazo)-3-naphthoic acid). For the determination of the content of exchangeable Mg, 5 ml of 1.26 M- $\text{NH}_4\text{Cl}$ ·3.70 M- $\text{NH}_4\text{OH}$  buffer (pH 10.5) was added to 5 ml of the above mentioned filtrate. Then 1 ml of 10%  $\text{NH}_2\text{OH} \cdot \text{HCl}$  was added to it. The solution was titrated with 0.005 M-EDTA using about 0.1 g EBT (1-(1-hydroxy-2-naphthylazo)-6-nitro-2-naphthol-4-sulfonic acid) as indicator. The content of organic materials was determined by the Walkley-Black method.

As for the biotic conditions of nesting sites, densities of red pine trees and floor-layer plants were estimated in the same study slope. For the red pine density, five quadrats of  $10 \times 10$  m area were sampled at random from the lower, the middle and the upper zones of the slope respectively. The density was represented by the number of trees taller than 1m and older than 10 years per  $100 \text{ m}^2$ . For the density of floor-layer plants, 10 quadrats were sampled at random with a frame of  $1 \times 1$  m area from each zone of the slope, then the species and the individual number of plants growing there were examined. These investigations were made on July 10, 1974.

The material utilized as the nest core was investigated in several natural habitats of the ant in Jinyang, Jinseung, Goseung, Sacheun and Namhae in July, 1974.

## RESULTS

### Nest distribution and abundance

#### I) Vertical distribution of nests

Vertical distribution of nests at twelve red pine stands in six localities was shown in Fig. 1. The nests scarcely occurred in the upper zone of slopes but were concentrated in the lower and the middle zones. Of 966 nests examined in these forest stands, 631 nests or 65.3 % were in the lower and 325 nests or 33.6 % were in the middle, but only 10 nests or 1.0 % were in the upper zone of slopes.




Fig. 1. Percentage ratios of nests distributed in the lower, middle and upper zones of hills inhabited by *F. yessensis*.

#### 2) Seasonal change in vertical distribution of nests

Fig. 2 shows vertical distributions of nests in a slope of a hill at Jinseung (Station A) in April, July and October, 1973. In April, nests were mostly distributed in the lower zone. The nests, however, spread over the middle zone by July, then became similar distribution to that of April again in October.




Fig. 2. Diagram of seasonal change of nest distribution in 1973 in a natural habitat (Jinseung, Station A) of *F. yessensis*. A slope of the hill was conveniently divided into three zones, and the home range of the ant was divided into 195 quadrats of  $10 \times 10$  m area.

### 3) Seasonal and annual changes of nest density


Based upon the result obtained from the quadrat census in the home range in a slope of a hill at Jinseung (Station A), seasonal changes of nest densities in 1973-77 were estimated. The density means the number of nests per  $m^2$  in quadrats where one or more nests occurred. The number of nests in the whole census area ( $19,500 m^2$ ) and the density are shown in Table 1. Although the number of whole nests in the census area increased in July, the density per  $m^2$  was rather constant, slightly decreasing in July.

Table 1. Seasonal and annual changes of nest density in a slope of a hill at Jinseung (Station A) in 1973-77.

Year	No. of nests in the whole census area			No. of nests/ $m^2$ in quadrats nests occurred		
	April	July	October	April	July	October
1973	66	124	90	0.017	0.015	0.018
1974	66	110	89	0.017	0.015	0.022
1975	77	122		0.017	0.016	0.022
1976	50	147	112	0.018	0.017	0.018
1977		64	50	0.015	0.014	0.017

### 4) Seasonal change of the nest abundance


*Formica yessensis*, a polydomous ant, exhibits nest proliferation by budding (Ito, 1973; Higashi, 1976; Kim and Murakami, 1980 b). Fig. 3 shows the results of the monthly census of nests from April to October in 1973-77 at the study area (Jinseung, Station A). The initiated nests mean the nests newly appeared during a month after the previous census, the continued nests mean those continued from the previous census, the deserted nests mean those abandoned during a month after the previous census, and the revived nests mean those re-utilized which had been abandoned previously. The number of existing active nests at each census is a total of the initiated, continued and revived




**Fig. 3.** Seasonal trends of the nest number for successive years and percentage of nests deserted in a slope of a hill at Jinseung (Station A) in 1973-77.

ones.

The active nests increased after the first census in April till July, and decreased in August to October. In July the number of the active nests attained 1.6-2.8 times as many as in April. In August the active nests decreased because the deserted nests increased in spite of increasing of newly initiated nests. The seasonal trends of the revived nests were similar to those of the initiated nests.



**Fig. 4.** Frequency distributions of nest longevity in a slope of a hill at Jinseung (Station A) in 1973-75.



**Fig. 5.** Frequency distributions of number of times the same nest was utilized during a year in a slope of a hill at Jinseung (Station A) in 1973-75.

### 5) Longevity of nests

The frequency distributions of the period during which each newly founded nest was being utilized were shown in Fig. 4 for the results obtained in 1973-75. From these results it was concluded that the life span of nests was 1 to 4 months in most cases and 13 months in maximum. Thus, many new nests are abandoned within the year. The abandonment occurs more frequently in the inactive nests such as newly founded and revived nests, but relatively rare in the active nests. The number of times the same nest was utilized during a year is as shown in Fig. 5.


## Environmental conditions in natural habitat of the ant

### 1) Humidity above ground

From the relative humidity determined at 10cm above ground at twelve sites in the natural habitat of the ant in a slope of the red pine stand in Jinseung (Station A), a map of vertical distributions of the averaged relative humidity from May to September and from October to November is shown in Fig. 6. With the relationships between nest distribution and the humidity from May to September when nests of the ant spread over lower to middle zones in the slope, it is supposed that the upper limit of the nest distribution (near the boundary between middle and upper zones) almost coincides with the line of 75 % RH. In the upper zone where the humidity is lower, the nest of the ant never occurred. In pre-hibernation period from October to November, nests are retreated to the lower zone of the slope where the humidity is higher than 74% RH. At that time the humidity becomes lower than in the previous season in the middle zone.

### 2) Chemical properties of the soil

The result of the soil chemical analysis in the natural habitat of the ant in the slope at Jinseung is shown in Table 2. In the lower and middle zones



**Fig. 6.** Contour lines of relative humidities at 10 cm above ground in a slope of a hill at Jinseung (Station A) in 1975. Broken lines show average humidity from May to September, solid lines from October to November. Solid square : points of determination.

inhabited by the ant, the content of exchangeable Ca in soil was extremely low, and organic matters and water contents were considerably higher than in the upper zone, though pH and the content of exchangeable Mg were not significantly different.

**Table 2.** Results of the soil chemical analysis in a slope of a hill at Jinseung (Station A), a natural habitat of *F. yessensis*.

Zone	Depth (cm)	pH	Exchangeable		Organic mat. (%)	Moisture (%)
			Ca	Mg		
			(me /100 g)			
Lower	0-20	5.2	0.71	1.10	2.88	21.23
	20-40	5.1	0.75	1.84	2.75	22.10
	40-60	5.1	0.62	1.9%	2.41	22.47
Middle	0-20	5.1	0.74	0.96	2.86	50.11
	20-40	5.2	0.73	1.00	2.60	21.36
	40-60	5.2	1.79	1.10	2.10	21.45
upper	0-20	5.1	2.31	1.10	1.21	11.56
	20-40	5.1	2.14	1.04	1.54	14.14
	40-60	5.1	2.22	1.26	1.19	13.31

### 3) Density of red pine trees

The density of red pine trees in the natural habitat of the slope at Jinseung (Station A) is shown in Table 3. In the lower and middle zones inhabited by the ant, the density of red pine trees older than 10 years age was higher than in the upper zone.

### 4) Density of floor-layer plants

Fifty-seven species of floor-layer plants were found in the study area of



Table 3. Density of red pine trees taller than 1 m and older than 10 years per 100 m<sup>2</sup> in a slope of a hill at Jinseung (Station A), a natural habitat of *F. yessensis* in July, 1974.

Age (years)	Lower zone	Middle zone	Upper zone
10-14	1.2	5.6	5.8
15-19	16.8	13.6	7.2
20-24	5.2	1.2	0.2
25-29	0	0.4	0
30-34	0	0.4	0
35-39	0	1.6	0
40-44	0	0.8	0
45 <	0	0.4	0
Total	23.2	24.0	13.2

the red pine stand inhabited by the ant. Among them dominant species were the following 10 species; *Andropogon brevifolius* Swartz, *Arundinella hirta* Tanaka, *Cymbopogon tortilis* var. *Goeringii* Handmazz, *Spodiopogon cotulifer* Hackel, *Festuca ovina* L., *Miscanthus sinensis* forma *purpurascens* Makino, *Eulalia speciosa* Kuntze, *Themeda triandra* subsp. *japonica* Tanaka, *Zoisia japonica* Steud and *Rhododendron yedoense* var. *poukhanense* Nakai. The densities and frequencies of these plants are shown in Table 4. In the lower and middle zones inhabited by the ant, it was grown with *A. brevifolius*, *A. hirta*, *C. tortilis* var. *Goeringii*, *S. cotulifer* and *F. ovina* but not with *E. speciosa* nor *Z. japonica*.

#### 5) Suitable nest cores utilized by the ant

Thirty-one kinds of materials were utilized as the nest core in five natural habitats of the ant as shown in Table 5. Among them, two species of long graminaceous grasses, *Arundinella hirta* (33.7 %) and *Miscanthus sinensis* f. *purpurascens* (17.7 %), red pine trees, *Pinus densiflora* Sieb. et Zucc. (23.9 %), oak trees, *Quercus* sp. (5.1 %), wild azalea trees, *Rhododendron yedoense* var. *poukhanense* (3.8 %) were main core plants. Stones (8.0 %) also serve sometimes as the core.

Table 4. Densities and frequencies of the ten dominant species of floor-layer plants of a red pine stand in a slope of a hill at Jinseung (Station A), a natural habitat of *F. yessensis* in July, 1974.

Species	Density (per m <sup>2</sup> )			Frequency (%)		
	Lower zone	Middle zone	Upper zone	Lower zone	Middle zone	Upper zone
<i>Andropogon brevifolius</i>	42.0	120.5	22.6	100	90	50
<i>Cymbopogon tortilis</i> var. <i>Goeringii</i>	38.7	30.2	4.7	100	90	90
<i>Spodiopogon cotulifer</i>	11.4	18.6	19.5	90	80	60
			6.1	70	50	30
<i>Miscanthus sinensis</i> f. <i>purpurascens</i>	16.0	2.4	5.6	20	90	30
<i>Rhododendron yedoense</i> var. <i>poukhanense</i>	4.7	1.1	0	50	30	0
<i>Eulalia speciosa</i>	0	0	14.7	0	0	80
<i>Themeda triandra japonica</i>	0	4.2	1.0	0	30	10
<i>Zoisia japonica</i>	0	0	3.5	0	0	20

Table 5. Number of nests built by utilizing various plants and stones as nest cores in Gyeongsang-Nam-Do, Korea.

Kind of nest cores	Locality					Total
	Jinseung	Namhae	Goseung	Jinyang	Sacheun	
<i>A. hirta</i>	23	73	88	63	30	277
<i>Pinus densijlora</i> (red pine)	48	39	68	32	9	196
<i>M. sinensis</i> f. <i>purpurascens</i>	7	25	79	29	5	145
<i>S. cotulifer</i>	0	0	0	12	0	12
<i>T. triandra japonica</i>	0	0	17	2	0	19
<i>Impera cylindrica</i>	0	0	7	5	0	12
<i>Quercus</i> sp.	7	11	20	0	4	42
<i>R. yedoense</i> var. <i>poukhanense</i>	21	3	0	6	1	31
<i>Juniperus chinensis</i>	0	3	0	0	0	3
<i>Castanea crenata</i>	1	0	0	0	0	1
<i>Indigofera kirilowii</i>	0	0	0	10	0	10
<i>F. ovina</i>	1	0	0	5	1	7
Stones	2	21	15	26	2	66
Total	110	175	294	190	52	821

## DISCUSSION

Nest distribution in slopes of hills naturally inhabited by *F. yessensis* is concentrated in the lower and middle zones but scarcely in the upper zone. The distribution changes with seasonal fluctuation of nest number. From autumn to early spring, the nests are distributed only in the lower zone of the slope, while in summer they spread over the middle zone according to the increase in nest number since the density of nests per unit area is constant seasonally.

Although red pine forests in Korea provide a favourable environment for *F. yessensis*, the upper zone of the hill seems not to be so favourable for establishing nests because of several reasons. In the upper zone the density of pine trees is relatively low and that of *Zoisia* grasses is higher than in the middle and lower zones. Such floral conditions are associated with lower humidity above ground and moisture in soil. The lower moisture of soil is also related to lower content of organic matter in the upper zone. In the lower and middle zones inhabited by the ant the content of exchangeable Ca in soil was extremely lower than in the upper zone, but it is uncertain whether it would be a cause or a result. Petal (1978) stated that ants modified the physical and chemical properties of both the soil used construction of nests and the adjacent soil. The concentration of exchangeable cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^{+}$ ,  $\text{Na}^{+}$ ) in nests, however, is slightly higher than in the surrounding soil in *Myrmica* sp. and *Lasius flavus* (Fabricius) according to the data cited by him.

In Ishikari Shore, Hokkaido, the distribution of nests of *F. yessensis* shows a remarkable correlation with the spatial change of vegetation, mostly confined within the zone occupied by eulalia (*Miscanthus sinensis* Anderss.) and sweet brier (*Rosa rugosa* Thunb.) and that occupied by vines (*Rubus parvifolius* L., *Vitis Coignetiae* Pulliat, etc.) with moderately dense plant matrix, while absent

both in sparse shore side vegetation and thick inland oak forest (Ito, 1971). Lower and middle zones of the hill, a study station at Jinseung, are favourable for the ant owing to relatively high density of red pine trees, but sufficient light intensity and presence of suitable plants for nest cores such as long graminaceous grasses.

Contrasting with European species of *Formica rufa* group, *F. yessensis* can establish nests without appropriate cores. The establishment, however, is facilitated by the presence of certain structures suitable as cores. In Ishikari Shore, more than half of nests were found around or under the root system of *Miscanthus sinensis* (Ito, 1971). Hayashida (1960) also suggested that the best site was around the root of living trees. Thus, the presence of suitable nest cores was regarded as one of factors responsible for the nest abundance. In Korea, various plant including *Arundinella hirta*, *Pinus densiflora* (red pine trees) and *Miscanthus sinensis* forma *purpurascens* as well as stones serve as the core for building nests.

#### ACKNOWLEDGEMENTS

The authors wish to express their hearty thanks to Professor Y. Hirashima for his constant guidance. Thanks are also due to Dr. Jin Sik Choi, Mr. Jong Man Kim, Mr. Sok Hyon Kim and Mr. Bu Gun Jeung of Gyeongsang National University for their kind cooperation in the course of field investigations in Korea, Professor Sam Sik Kim of the University for his kind identification of plants, Mr. Han Saeng Lee of Research Bureau, Office of Rural Development, Gyeongnam Province for his help in soil chemical analysis. The senior author also thanks Dr. Tae Gyu Yoon, the ex-President of Gyeongsang National University, for his help promoting the present study and encouragement.

#### REFERENCES

- Hayashida, K. 1960 Studies on the ecological distribution of ants in Sapporo and vicinity (1 et 2). *Insectes Sociaux*, 7: 125-162
- Higashi, S. 1976 Nest proliferation by budding and nest growth pattern in *Formica (Formica) yessensis* in Ishikari Shore. *J. Fac. Sci., Hokkaido Univ., Ser. VI, Zool.*, 20: 359-389
- Ito, M. 1971 Nest distribution of *Formica yessensis* Forel in Ishikari Shore, in reference to plant zonation. *J. Fac. Sci., Hokkaido Univ., Ser. VI, Zool.*, 18: 144-154
- Ito, M. 1973 Seasonal population trends and nest structure in a polydomous ant, *Formica (Formica) yessensis* Forel. *J. Fac. Sci., Hokkaido Univ., Ser. VI, Zool.*, 19: 270-293
- Kim, C. H. and Y. Murakami 1980a Ecological studies on *Formica yessensis* Forel, with special reference to its effectiveness as a biological control agent of the pine caterpillar moth in Korea I. Geographical and ecological distributions of *Formica yessensis* Forel (Hymenoptera: Formicidae). *J. Fac. Agr., Kyushu Univ.*, 24: 239-245
- Kim, C. H. and Y. Murakami 1980b Ecological studies on *Formica yessensis* Forel, with special reference to its effectiveness as a biological control agent of the pine caterpillar moth in Korea II. Bionomics of *Formica yessensis* Forel (Hymenoptera: Formicidae) in Korea. *J. Fac. Agr., Kyushu Univ.*, 25: 119-133

- Morisita, M. 1945 Ants. *In* "Insects," Vol. 2. ed. by H. Furukawa, Kenkyusha, Tokyo. pp. 1-56 (in Japanese)
- Petal, J. 1978 The role of ants in ecosystems. *In* "Production Ecology of Ants and Termites," ed. by M. V. Brian, Cambridge Univ. Press, Cambridge, pp. 293-325