Properties of Active Substance, BCF, which Synergistically Enhances the Antimicrobial Activity of Hexametaphosphate

Tsutsumi, Masakazu

Laboratory of Food Hygienic Chemistry, Faculty of Agriculture, Kyushu University

Miyamoto, Takahisa

Laboratory of Food Hygienic Chemistry, Faculty of Agriculture, Kyushu University

Suda, Ikuo

Laboratory of Food Hygienic Chemistry, Faculty of Agriculture, Kyushu University

Watanabe, Tadao

Laboratory of Food Hygienic Chemistry, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23740>

出版情報：九州大学大学院農学研究院紀要. 26 (2/3), pp.71-78, 1982-02. Kyushu University
バージョン：
権利関係：

Properties of Active Substance, BCF, which Synergistically Enhances the Antimicrobial Activity of Hexametaphosphate

Masakazu Tsutsumi, Takahisa Miyamoto, Ikuo Suda
and Tadao Watanabe

Laboratory of Food Hygienic Chemistry, Faculty of Agriculture,
Kyushu University 46-09, Fukuoka 812

(Received July 29, 1981)

An active substance named BCF was found in the culture broth of *Bacillus subtilis*. This substance synergistically enhanced the antibacterial activity of hexametaphosphate against gram-negative bacteria such as *Escherichia coli* and markedly weakened the salt tolerance of some bacteria such as *Aerobacter aerogenes* and *Staphylococcus aureus*, and yet no antibacterial activity was shown in itself. BCF was extractable with n-butanol at pH 9.0 and dialyzable against water using cellulose tubing; besides, it was stable at alkaline pH and to heat also.

INTRODUCTION

Polyphosphates such as hexametaphosphate (HP) or ultraphosphate are effective against most gram-positive bacteria, but are inactive against most gram-negative bacteria (Tsutsumi *et al.*, 1976). Polyphosphates, however, show strong growth inhibition against gram-negative bacteria in the presence of sodium cholate or glycerol monocaprates (Tsutsumi *et al.*, 1977).

In a series of investigations we found an active substance named BCF in the culture broth of *Bacillus subtilis*, which synergistically enhanced the antibacterial activity of polyphosphates against *Escherichia coli*. We also found out the fact that the salt tolerance of *Staphylococcus aureus* was markedly weakened by adding BCF. Whereas no antibacterial activity was shown in BCF alone.

Being interested in studying such bacteriologically active substance, we have tried purifying and characterizing BCF. In this paper, we tried elucidating the properties of BCF first though it is not yet isolated in a pure state.

MATERIALS AND METHODS

Strains

Escherichia coli IFO 3301, *Pseudomonas aureofaciens* IFO 3521, *Bacillus cereus* IFO 3060, *Saccharomyces cerevisiae* IFO 2044 and *Torulopsis etchellsii* IFO 1229 were obtained from Institute for Fermentation, Osaka.

Bacillus subtilis and *Aerobacter aerogenes* were generously provided by a certain Institute.

Media and cultural conditions

Nutrient broth containing 1% polypeptone and 1% Ehrlich meat extract was used as a growth medium for bacteria. The broth pH was adjusted to 7.0.

Sabouraud medium (pH 7.0) improved by addition of 0.2% yeast extract was used for yeasts. The culture was carried out at 30°C for 24 hours on a reciprocal shaker.

Determination of antibacterial activity

The antibacterial activity of BCF was routinely determined on *Escherichia coli* in the presence of HP. To 5 ml of 2% broth were added 3 ml of the culture filtrate (BCF-CF) containing BCF, 1 ml of sterile HP, 0.5 ml of sterile water and 0.5 ml of overnight culture of *Escherichia coli* growing in 1% broth, and the inoculated medium was incubated at 30°C for 20 hours with shaking. The same volume of sterile water was added instead of the culture filtrate and HP in control experiment 1, the culture filtrate in control experiment 2 and HP in control experiment 3. The turbidity (OD_{660}) of the overnight culture used was adjusted to 0.3 with sterile water before use. Sodium chloride (NaCl) or HP was added to give the final concentrations indicated.

Growth was routinely monitored by the change in absorbance of the culture with a colorimeter fitted with a red 660 filter and graded from – through ++ as follows: –, no growth ($OD_{660}=0\sim0.04$); ±, very slight growth ($OD_{660}=0.05\sim0.09$); +, moderate growth ($OD_{660}=0.1\sim0.4$); ++, good growth (OD_{660} =above 0.4).

n-Butanol extraction

The culture broth (20 ml) containing BCF was adjusted to pH 5.0, pH 7.0 and pH 9.0 with HCl or NaOH solution, and then the same volume of n-butanol was added. After shaking vigorously, n-butanol extracts were separated from the broth and dried *in vacuo* at 40°C. The residues were dissolved in 10 ml of n-butanol and the insoluble materials were removed by centrifugation. After the supernatant was evaporated to dryness *in vacuo* at 40°C, the residues were dissolved in 20 ml of water. This was named BCF-Buol and those extracted at pH 5.0, pH 7.0 and pH 9.0 were named BCF-Buol-5, BCF-Buol-7 and BCF-Buol-9, respectively.

RESULTS

1. Accumulation of BCF in the culture broth

In the growth curve of *Bacillus subtilis*, the accumulation of BCF began to appear at mid-logarithmic phase and reached the highest level at early stationary phase (data not shown).

After *Bacillus subtilis* was cultured at different temperatures from 20°C to 45°C, the cells were separated from the culture broth by centrifugation and then the activity in the supernatant was determined in the presence of HP.

Though the maximal activities were detected when the bacterium was grown at 30°C and 35°C, no activity was observed in the cells grown at 45°C (data not shown).

2. Antimicrobial activity of BCF

To 5 ml of 2% broth were added 3 ml of the culture filtrate containing BCF, 1.5 ml of sterile water and 0.5 ml of microorganisms suspension ($OD_{660} = 0.3$), and the inoculated cultures were incubated at 30°C for 20 hours.

Table 1 shows that BCF-CF alone was inactive against all microorganisms tested.

3. Synergism between BCF-CF and other chemicals

The synergistic effects between BCF-CF and other chemicals (HP, NaCl, sodium cholate or glycerol monocaprates) were determined using some microorganisms. The effects between BCF-CF and either HP, sodium cholate or glycerol monocaprates against gram-positive microorganisms were not determined because these chemicals alone showed the antimicrobial activity against these microorganisms.

a) In the case of *Escherichia coli*

1) BCF-CF and HP

Growth was always observed in the control experiments 1 and 3, and therefore the rest omitted the data of these results.

2) BCF-CF and NaCl

3) BCF-CF and sodium cholate

4) BCF-CF and glycerol monocaprates

Table 1. Antimicrobial activity of BCF-CF against several microorganisms. Growth was graded from—(no growth), $\pm$ (very slight growth), + (moderate growth) through ++ (good growth).

Microorganisms	Growth
<i>E. coli</i>	++
<i>P. aureofaciens</i>	++
<i>S. aureus</i>	++
<i>B. cereus</i>	++
<i>B. megaterium</i>	++
<i>S. cerevisiae</i>	++
<i>T. etchellsii</i>	++

Table 2. Synergism between BCF-CF and HP against *E. coli*. Symbols are the same as in Table 1.

HP(%)	Growth			
	Control 1	Control 2	Control 3	Test
0	++	++	++	++
0.25	++	++	++	—
0.50	++	++	++	—
0.75	++	++	++	—
1.00	++	++	++	—

Table 3. Synergism between BCF-CF and NaCl against *E. coli*. Symbols are the same as in Table 1.

NaCl (%)	Growth	
	Control 2	Test
0	++	++
1	++	++
2	++	+
3	+	+
4	+	+
5	+	+
6	+	—
7	+	—
8	+	—
9	+	—
10	—	—

Table 4. Synergism between BCF-CF and cholate against *E. coli*. Symbols are the same as in Table 1.

Cholate (%)	Growth	
	Control 2	Test
0	++	++
0.025	++	++
0.050	++	++
0.075	++	++
0.100	++	++

Table 5. Synergism between BCF-CF and glycerol monocaprato against *E. coli*. Symbols are the same as in Table 1.

Glycerol monocaprato (%)	Growth	
	Control 2	Test
0	++	++
0.025	++	++
0.050	++	++
0.075	++	++
0.100	++	++

The synergistic effects between BCF-CF and either HP or NaCl was observed (Tables 2 and 3), but not either sodium cholate or glycerol monocaprato (Tables 4 and 5).

b) In the case of *Aerobacter aerogenes*

1) BCF-CF and NaCl

The synergistic effect between BCF-CF and NaCl against *Aerobacter aerogenes* was apparent. As the effects between BCF-CF and either HP, sodium cholate or glycerol monocaprato were similar to the results of *Escherichia coli*, the data were omitted. In the case of *Pseudomonas aureofaciens*, the data were similar to the results of *Escherichia coli*.

c) In the case of *Staphylococcus aureus*

Table 6. Synergism between BCF-CF and NaCl against *A. aerogenes*. Symbols are the same as in Table 1.

NaCl (%)	Growth	
	Control 2	Test
0	++	++
1	++	+
2	++	—
3	++	—
4	++	—
5	++	—
6	++	—
7	+	—
8	—	—
9	—	—

Table 7. Synergism between BCF-CF and NaCl against *Staph. aureus*. Symbols are the same as in Table 1.

NaCl (%)	Growth	
	Control 2	Test
0	++	++
1	++	++
2	++	++
3	++	++
4	++	+
5	++	—
6	++	—
7	++	—
8	++	—
9	++	—
10	++	—
11	++	—
12	+	—
13	+	—
14	—	—
15	—	—

1) BCF-CF and NaCl

Staphylococcus aureus could grow in a medium containing 13% NaCl. The salt tolerance of the bacterium, however, was markedly weakened by adding BCF-CF, as shown in Table 7.

d) In the case of *Saccharomyces cerevisiae*

1) BCF-CF and NaCl

No synergistic effect was observed (Table 8).

4. Properties of BCF

As described in Method, BCF was extracted with n-butanol. Each BCF-Buol alone was inactive against the test microorganisms (data not shown); however, BCF-Buol-7 and BCF-Buol-9 were effective against *Escherichia coli* in the presence of HP, as shown in Table 9.

The n-butanol extracts (BCF-Buol-9) were stable between pH values 7~12

Table 8. Synergism between BCF-CF and NaCl against *Sacch. cerevisiae*. Symbols are the same as in Table 1.

NaCl (%)	Growth	
	Control 2	Test
0	++	++
1	++	++
2	++	++
3	+	++
4	±	+
5	—	±
6	—	—
7	—	—

Table 9. Synergism between each BCF-Buol and HP against *E. coli*. Symbols are the same as in Table 1.

BCF-Buol	Growth			
	Control 1	Control 2	Control 3	Test
BCF-Buol-5	++	++		
BCF-Buol-7	++	++		
BCF-Buol-9	++	++	++	++

when left for 60 min at room temperatures between pH values 2~12 (data not shown). When kept for 30 min in boiling water, for 20 min at 120°C (in autoclave) or for 72 days at -20°C, the extracts were stable (data not shown).

The n-butanol extracts (BCF-Buol-9) were dialyzed against running water (deionized) using Cellulose tubing (Visking Co.). The dialyzed extracts were withdrawn at intervals and then the activities were determined in the presence of HP. The active substance (BCF) in the extracts completely disappeared by dialysis (data not shown).

DISCUSSION

Polyphosphates such as hexametaphosphate show antimicrobial action; therefore their application as food preservatives has been studied. Polyphosphate, however, were inactive against most gram-negative bacteria.

The fact may be disadvantageous as the preservatives, hence we proposed some attempts to overcome the difficulties by use of the substances enhancing synergistically the activity of HP against gram-negative bacteria (Tsutsumi *et al.*, 1977; Watanabe *et al.*, 1977; Watanabe *et al.*, 1979). In a series of investigation an active substance named BCF which is capable of employing for this purpose was found in the culture broth of *Bacillus subtilis*. As elucidating the properties of BCF may be more useful for isolating it, we tried carrying out this work first.

As a result, this substance (BCF) aroused our interest because it had no antimicrobial activity in itself, but enhanced the activity of HP against gram-

negative bacteria and weakened the salt tolerance of *Staphylococcus aureus*; besides, BCF was unstable at acid pH though it was stable at alkaline pH, and it was stable to heat. These properties were different from those of many antibiotics (Shoji *et al.*, 1975 a, b, c, d; Shoji *et al.*, 1976 a, b, c, d, e, f, g; Shoji *et al.*, 1977; Shoji *et al.*, 1978; Shoji *et al.*, 1980; Umezawa, 1967) which are produced by the genus *Bacillus*.

To our knowledge, therefore, there is no known substance identical with BCF. Thus, BCF may be a new substance which is bacteriologically active.

REFERENCES

- Shoji J., H. Hino, Y. Wakizaka, K. Koizumi and M. Mayama 1976 a Isolation of a new peptide antibiotic complex, B-43. *J. Antibiotics*, **29**: 813-817
- Shoji J., R. Sakazaki, Y. Wakisaka, K. Koizumi and M. Mayama 1975 a Isolation of a new peptide antibiotic complex 61-26. *J. Antibiotics*, **28**: 129-131
- Shoji J., T. Kato, K. Matsumoto, Y. Takahashi and M. Mayama 1976 b Production and isolation of cerexins C and D. *J. Antibiotics*, **29**: 1281-1285
- Shoji J., H. Hino, Y. Wakisaka, K. Koizumi, M. Mayama and Matsuura 1977 Isolation of two new plmyoxyin group antibiotics *J. Antibiotics*, **30**: 1029-1034
- Shoji J., H. Hino, Y. Wakisaka, K. Koizumi, M. Mgyama and S. Matsuura 1975 b Isolation of a new peptide antibiotic TL-119. *J. Antibiotics*, **28**: 126-128
- Shoji J., H. Hino, Y. Wakisaka, K. Koizumi, M. Mayama and S. Matsuura 1976 c Isolation of a new peptide antibiotic, 339-29. *J. Antibiotics*, **29**: 809-812
- Shoji J., R. Sakazaki, Y. Wakisaka, K. Koizumi, M. Mayama and S. Matsuura 1975 c Isolation of galantins 1 and 2, water-soluble basic 3. *J. Antibiotics*, **28**: 122-125
- Shoji J., H. Hino, Y. Wakisaka, K. Koizumi, M. Mayama, S. Matsuura and K. Matsumoto 1975 d Isolation of two new related peptide antibiotics, cerexins A and B. *J. Antibiotics*, **28**: 56-59
- Shoji J., H. Hino, Y. Wakisaka, K. Koizumi, M. Mayama, S. Matsuura and K. Matsumoto 1976 d Isolation of three new antibiotics, thiocillins 1, 2 and 3, related to micrococcin P. *J. Antibiotics*, **29**: 366-374
- Shoji J., H. Hino, Y. Wakisaka, K. Koizumi, M. Mayama, S. Matsuura and K. Matsumoto 1976 e Isolation of a new antibiotic 333-25, related to antibiotic EM 49. *J. Antibiotics*, **29**: 516-520
- Shoji J., R. Sakazaki, Y. Wakisaka, K. Koizumi, H. Miwa, S. Matsuura and M. Mayama 1980 Isolation of octapetin D. *J. Antibiotics*, **33**: 182-185
- Shoji J., R. Sakazaki, Y. Wakisaka, K. Koizumi, M. Mayama, S. Matsuura and K. Matsumoto 1976 f Isolation of brevistin, a new peptide antibiotic. *J. Antibiotics*, **29**: 375-379
- Shoji J., R. Sakazaki, Y. Wakisaka, K. Koizumi, M. Mayama, S. Matsuura and K. Matsumoto 1976 g Isolation of a new antibiotic, laterosporamine. *J. Antibiotics*, **29**: 390-393
- Shoji J., H. Hino, R. Sakazaki, T. Kato, Y. Wakisaka, M. Mayama H. Hiwa and S. Matsuura 1978 Isolation of tridecaptins A, B and C. *J. Antibiotics*, **31**: 646-651
- Tsutsumi M., K. Nishimura, K. Yasui, A. Matsuoka and T. Watanabe 1976 Synergistic action of cholate on some antimicrobial substances. *Shokuhin Eiseigaku Zasshi (J. Food Hyg. Soc. Jpn.)* **17**: 273-275
- Tsutsumi M., K. Yasui, K. Isshiki and T. Watanabe 1977 Preservative effect of polyphosphates (Part 2). Action of hexametaphosphate on *Escherichia coli*. *Shokuhin Eiseigaku Zasshi (J. Food Hyg. Soc. Jpn.)* **18**: 341-345

- Umezawa S. 1967 Khokinseibussitsu, Baifukan, Tokyo (in Japanese)
- Watanabe T., G. Hirata and M. Tsutsumi 1977 Bacterostatic technique by application of the synergism between sodium polyphosphate $1.67 > (\text{Na}_2\text{O} + \text{H}_2\text{O}) / \text{P}_2\text{O}_5 > \text{O}$ and sodium cholate. *Nihon Tokkyo No* 858746 (in Japanese)
- Watanabe T., M. Tsutsumi, K. Isshiki, G. Hirata 1979 New food preservative containing both A (either hexametaphosphate or ultraphosphate) and B (either glycerol monocatecholate or glycerol monocaprylate). *Tokkyo Koho Showa* 54-32057 (in Japanese)