

Effect of Temperature and Fertilizer Level on the Chemical Composition and Nutritive Value of Green Panic (*Panicum maximum* var. *trichoglume* cv. *Petrie*)

Fleischer, Joseph Edmund

Department of Animal Science, Faculty of Agriculture, Kyushu University

Masuda, Yasuhisa

Department of Animal Science, Faculty of Agriculture, Kyushu University

Goto, Ichiro

Department of Animal Science, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23780>

出版情報：九州大学大学院農学研究院紀要. 28 (2/3), pp.83-90, 1983-12. Kyushu University
バージョン：
権利関係：



Effect of Temperature and Fertilizer Level on the Chemical Composition and Nutritive Value of Green Panic (*Panicum maximum* var. *trichoglume* cv. Petrie)

Joseph Edmund Fleischer, Yasuhisa Masuda and Ichiro Goto.

Department of Animal Science, Faculty of Agriculture,
Kyushu University 46-06, Fukuoka 812.

(Received September 5, 1983)

The effect of temperature and fertilizer level on the chemical composition and nutritive value of green panic (*Panicum maximum* var. *trichoglume* cv. Petrie) was studied. There were three temperature treatments i. e., 20, 25 and 30°C; and three fertilizer treatments at the rate of 0.98, 1.96 and 2.94 kg N, P₂O₅ and K₂O per are i.e., F₁, F₂ and F₃ respectively. Plants were raised from seeds and were grown in the open for seven weeks before transferring into the phytotron. The plants were harvested after 10, 20 and 30 days in the temperature rooms. Dry matter yield increased with increased fertilizer level but was little affected by temperature. Nitrogen increased with increased fertilizer level but decreased with advancing growth period and increasing temperature. All the cell wall constituents increased with increasing temperature but was little affected by fertilizer level.

In vitro organic matter digestibility (IVOMD) decreased with increasing temperature ($P < 0.01$) and F₁ was higher than both F₂ and F₃ ($P < 0.05$). IVOMD decreased at the rate of 0.78% unit per degree rise in temperature. Both temperature and nitrogen were negatively correlated with IVOMD but only that of temperature was statistically significant.

INTRODUCTION

Two of the most important factors influencing the growth and development of herbage and hence its nutritive value are growth temperature and soil nitrogen when moisture is not limiting (Wilson, 1982).

With tropical forages, the level of temperature maximum for photosynthesis is reported to be high (Ludlow and Wilson, 1968) and this together with the high insolation to which they are exposed, make them potentially more productive in terms of dry matter production compared to temperate or cool season forages (Cooper and Tainton, 1968; Cooper, 1970).

Even though temperature has been thought to influence the digestibility of pastures through increasing the fibre and lignin content, it is only recently that direct evidence of this has been established. Minson and McLeod (1970) reported a negative correlation ($r = -0.76$) between growth temperature and digestibility. Minson and Wilson (1980) also reported an average decline of 0.60 % unit in digestibility per unit rise in growth temperature. These latter workers noted that temperature accounted for about 60 % of the variations in digestibility.

The nitrogen content of most tropical forages is relatively lower than that of most temperate forages (Minson, 1976) mainly because of the low fertility status of the soils on which they are grown, and also their rapid maturity leading to a dilution of the crude protein content. Tropical forages respond well to fertilizer nitrogen which not only increases the crude protein content but also productivity. However, the effect of nitrogen fertilization on dry matter digestibility has been mixed (Wilson, 1982).

Thus, this experiment was initiated to examine the effect of ambient growth temperature and fertilizer levels on the trend of changes in the chemical composition and nutritive value of tropical forages.

MATERIALS AND METHODS

The species used in this experiment was green panic (*Panicum maximum* var. *trichoglume* cv. Petrie). Three fertilizer levels and three growth temperatures were employed. The three temperatures imposed were 20, 25 and 30°C day and night i. e., T_{20} , T_{25} and T_{30} ; and a compound fertilizer (14 % N, 14% P_2O_5 and 14 % K_2O) was applied at the rate of 0.98, 1.96 and 2.94 kg per are of each element i. e., F_1 , F_2 and F_3 respectively. There were three pots per treatment.

Pots of diameter 12 cm and depth 12cm were loosely filled to the brim with sandy-clayey loam soil which had been sifted through a 3mm sieve and air-dried. The pots were divided into three groups, each of which received one of the three levels of fertilizer in two equal applications at sowing and six weeks thereafter. Thinning and transplanting of plants to 5-6 per pot was done at three weeks after germination such that plants would be of approximately equal size at the beginning of temperature treatment. A week after the second fertilizer application, plants of three pots from each of three fertilizer groups were harvested. The remainder of each of the fertilizer groups were then subdivided into three groups and placed in the constant temperature rooms of the phytotron. The relative humidity of each of the temperature rooms was 70% throughout the experiment. Natural sunlight passing through the glass wall and roof of the phytotron was employed.

Harvesting of plants was done 10, 20 and 30 day i. e., P_1 , P_2 and P_3 respectively, after placement in the phytotron. Plants were harvested at 5cm above soil level. After harvesting, samples were dried at 70°C for more than 24 hours for dry matter determination. The dried samples were bulked for each temperature and fertilizer groups, ground to pass through 1mm sieve and stored until analyses.

Samples were analyzed for total nitrogen by the Kjeldahl method, cell wall constituents by the method of Goering and Van Soest (1970) excluding sodium sulphite and antifoaming agents; and *in vitro* organic matter digestibility (IVOMD) by the method of Minson and McLeod (1972).

RESULTS

Dry matter yield as affected by temperature and fertilizer levels is presented in Table 1. Dry matter yield increased with fertilizer level ($P<0.01$) but differences among temperature groups were not significant.

Table 1. Dry matter yield and cell wall constituents of green panic as affected by fertilizer level and temperature.

Characteristic	Period	T ₂₀			T ₂₅			T ₃₀		
		F ₁	F ₂	F ₃	F ₁	F ₂	F ₃	F ₁	F ₂	F ₃
Dry Matter	1	4.63	10.10	12.37	6.35	13.86	12.27	9.09	15.33	9.63
	2	8.81	16.09	16.86	11.47	19.88	17.98	10.91	18.47	18.02
	3	11.93	24.87	19.42	12.14	22.53	16.23	12.14	17.59	18.57
Neutral Detergent Fibre	1									
	2	74.47	66.96	72.74	66.96	66.18	68.78	69.29	71.01	72.40
	3	74.29	77.53	74.42	75.41	73.71	74.94	76.24	74.65	76.56
Acid Detergent Fibre	1			35.90	37.10	40.58	38.24	43.47	40.12	40.28
	2	34.37	43.50	35.91	44.25	43.31	46.65	43.51	43.98	49.16
	3	44.40	46.56	46.70	46.92	46.03	46.33	48.09	48.93	48.93
Cellulose	1									
	2	40.32	31.53	39.59	32.86	32.57	33.51	36.79	34.65	39.14
	3	39.85	41.25	41.47	38.24	40.15	40.22	41.36	40.56	42.58
Acid Detergent Lignin	1	1.96	2.43	2.21	2.35	3.03	2.91		3.32	3.17
	2	2.83	3.73	3.86	4.20	4.88	4.79	4.96	5.16	5.59
	3	3.59	4.53	4.35	4.81	5.36	5.52	5.68	6.23	6.01

Nitrogen content is shown in Fig. 1. Nitrogen content increased with increasing level of fertilization but decreased with advancing growth period. There was a significant decrease in nitrogen content with increasing temperature ($P<0.01$). Nitrogen content decreased most at T₃₀ by the end of P₁ in the phytotron; however, by the end of P₃ it has decreased to about 32-38 % in T₂₅ and T₃₀ and 35-40 % in T₂₀ of the pre-temperature treatment value.

The results of cell wall constituents analyses are shown in Table 1. In general, significant increase ($P<0.01$) occurred between P₁ and P₂ but no difference was observed between P₂ and P₃.

Neutral detergent fibre (NDF) : There was a consistently large increase in NDF with increasing temperature ($P<0.01$) than with increasing level of fertilization. There was a difference of about 3-5 % units as temperature moved from T₂₀ to T₂₅ or T₃₀ during P₁ but by the end of P₃ these differences have greatly narrowed down. On the contrary, differences due fertilizer level were negligible throughout the temperature treatment period.

Acid detergent fibre (ADF) : The differences in ADF were 2.88, 3.02 and 5.09 for T₂₀ and T₂₅, T₂₅ and T₃₀, and T₂₀ and T₃₀ respectively during P₁. However, by the end of P₂ and P₃ there were no differences between T₂₀ and T₂₅ but these were about 3 % units lower than that at T₃₀ ($P<0.05$). No differences were observed among fertilizer levels.




Fig. 1. Effect of temperature and fertilizer level on the nitrogen content of green panic.

Cellulose: The increase in cellulose occurred mainly between P_1 and P_2 after which it remained virtually unchanged. No differences were observed among fertilizer levels but the differences among temperature groups were significant ($P < 0.01$).

Acid detergent lignin (ADL): ADL increased with both fertilizer levels ($P < 0.01$) and temperature ($P < 0.01$) but the interaction between these two factors was not significant. The lignin contents of F_2 and F_3 were almost the same but higher than that of F_1 .

In vitro organic matter digestibility (IVOMD) is shown in Fig. 2. There was a general decline in IVOMD with advancing growth period. T_{30} was about 6.0 % units lower than T_{20} by the end of P_1 and the difference reached about 9.0 and 7.5 % units at the end of P_2 and P_3 respectively. The differences between T_{20} and T_{25} were 2.9, 5.0 and 6.0 % units, while those between T_{25} and T_{30} were 3.1, 4.0 and 1.5 % units for the three periods respectively. These differences were significant ($P < 0.01$). With regard to fertilizer level, though there were smaller differences compared to those due to temperature, only F_1 was significantly higher ($P < 0.05$) than those of F_2 and F_3 . There were, however, no interactions between temperature and fertilizer level.

In Fig. 3 are shown the response curves of IVOMD to temperature and fertilizer level. The graphs indicate that IVOMD was affected linearly by temperature and quadratically by fertilizer level.




Fig. 2. Effect of temperature and fertilizer level on the in vitro organic matter digestibility of green panic.




Fig. 3. Temperature and fertilizer response curves of *in vitro* organic matter digestibility of green panic.

Table 2. Correlation between *in vitro* organic matter digestibility and temperature or fertilizer level.

Period in Phytotron	Temperature	Fertilizer
All Harvests	-0.4026*	-0.1333
First Harvest	-0.8018**	-0.0741
Second Harvest	-0.8864**	-0.4323
Third Harvest	-0.8567**	-0.2861

* P < 0.05, ** P < 0.01

In Table 2 is shown the coefficient of correlation between IVOMD and temperature or fertilizer level. Fertilizer level was negatively correlated to IVOMD but the values were small and not significant. On the contrary, the values with regard to temperature were also negative but larger and significant. When all the data for the respective periods were taken together the correlation values were relatively small yet significant with regard to temperature.

DISCUSSION

The lower nitrogen contents with increasing temperature have been reported by Wilson and Ford (1971). In the present experiment temperature had little effect on the dry matter yield and so the lower nitrogen content cannot be fully explained by the increased dry matter alone. Increasing temperature has been reported to increase the rate of leaf senescence (Wilson and t'Mannetje, 1978) and stem development (McWilliams, 1978). Since these plant fractions are generally lower in nitrogen content than green leaves, they might have contributed to the lower nitrogen contents under higher temperatures.

The increase in structural components with increasing temperature is in accordance with published results (Wilson and Ford, 1973; Wilson *et al.*, 1976; Moir *et al.*, 1977). The increase in NDF beyond P₂ may be due to the increase in ADF content. However, since no increase in cellulose occurred beyond this period it seems reasonable to think that beyond a certain stage of the plant's development photosynthates are transformed into lignin as has been reported by Deinum and Knoppers (1979) with maize.

A number of workers have reported a decline in digestibility with increasing growth temperature (Wilson and Ford, 1971, 1973; Wilson *et al.*, 1976; Moir *et al.*, 1977). Minson and McLeod (1970) found a very high negative correlation ($r = -0.76$) between growth temperature and digestibility. In the present experiment, the correlations were found to range between -0.89 and -0.80 for the three respective periods. The lower correlation value when all the data for the three periods were taken together might suggest that the effect of temperature on IVOMD is modified by stage of development or some other factors (see Fig. 2).

There was an average decline of 0.78 % unit per degree rise in temperature in all the fertilizer groups, which appears higher than 0.60 % unit reported

by Minson and Wilson (1980).

The decrease in digestibility with fertilizer was unexpected though there are some supporting literatures to this effect (Blaser, 1964; Gomide *et al.*, 1969; Hutton, 1970). One possible reason is the increased and sustained growth brought about by the higher levels of fertilizer (Wood *et al.*, 1972) and the increased lignin content with increased fertilizer (Hutton, 1970; Vicente-Chandler, 1974). Nevertheless, the low negative correlation observed between fertilizer level and digestibility compared to that of temperature suggest that their effects differed in extent, and the non-significant interaction between temperature and fertilizer also suggests that these two factors might be affecting IVOMD differently.

REFERENCES

- Blaser, R. E. 1964 Effects of fertilizer levels and stage of maturity on forage nutritive value. *J. Anim. Sci.*, 23: 246-253
- Cooper, J. P. 1970 Potential production and energy conversion in temperate and tropical grasses. *Herb. Abstr.*, 40: 1-15
- Cooper, J. P. and N. M. Tainton 1968 Light and temperature requirements for growth of tropical and temperate grasses. *Herb. Abstr.*, 38: 167-176
- Deinum, B. and J. Knoppers 1979 The growth of maize in the cool temperate climate of Netherlands : Effect of grain filling on production of dry matter yield, chemical composition and nutritive value. *Neth. J. Agric. Sci.*, 27: 116-130
- Goering, H. K. and P. J. Van Soest 1970 *Forage Fiber Analyses*. United States Department of Agricultural Husbandry. Number 379
- Gomide, J. A., C. H. Holler, G. O. Mott, C. J. H. Conrad and D. L. Hill 1969 Effects of plant age and nitrogen fertilization on the chemical composition and *in vitro* cellulose digestibility of tropical grasses. *Agron. J.*, 61: 116-120
- Hutton, E. M. 1970 Tropical forages. *Adv. in Agron.*, 22: 1-73
- Ludlow, M. M. and G. L. Wilson 1968 Studies on the productivity of tropical plants. 1. Growth analysis, photosynthesis, and respiration of hamil grass and siratro in a controlled environment. *Aust. J. Agric. Res.*, 19: 35-45
- McWilliams, J. R. 1978 Response of pasture plants to temperature. In "Plant Relations in Pastures", ed. by J. R. Wilson, Melbourne, CSIRO, pp. 17-34
- Minson, D. J. 1976 Nutritional significance of protein in temperate and tropical pastures. In "From Plant to Animal Protein" ed. by T. M. Sutherland, J. R. McWilliams and R. A. Leng, Reviews in Rural Science, Number II. pp. 27-30
- Minson, D. J. and M. N. McLeod 1970 The digestibility of temperate and tropical grasses. *Proc. XIth Int. Grassl. Cong.*, Surfers Paradise, Australia. pp. 719-722
- Minson, D. J. and M. N. McLeod 1972 The *in vitro* technique: Its modifications for estimating digestibility of large numbers of tropical pasture plants. *Australia, CSIRO, Div. of Trop. Past. Tech. Pap. No. 8*
- Minson, D. J. and J. R. Wilson 1980 Comparative digestibility of tropical and temperate forages-a contrast between grasses and legumes. *J. Aust. Inst. Agric. Sci.*, 46: 247-249
- Moir, K. W., J. R. Wilson and G. W. Blight 1977 The *in vitro* digested cell wall and fermentation characteristics of grasses as affected by temperature and humidity during growth. *J. Agric. Sci., Camb.*, 88: 217-222
- Vicente-Chandler, J. 1974 Fertilization of humid tropical grasslands. "In Forage Fertilization" ed. by D. A. Mays, A. S. A., C. S. S. A., S. S. S. A. Publication, pp. 277-300
- Wilson, J. R. 1982 Environment and nutritional factors affecting herbage quality. *In*

- "Nutritional Limits to Animal Production from Pastures", ed. by J. B. Hacker, Farnham Royal, U. K., Commonwealth Agricultural Bureaux, pp. 117-130
- Wilson, J. R. and C. W. Ford 1971 Temperature influences on the growth, digestibility and carbohydrate composition of two tropical grasses, *Panicum maximum* var. *trichoglume* and *Setaria sphacelata* and two cultivars of the temperate grass *Lolium perenne*. **Aust. J. Agric. Res.**, **22**: 563-571
- Wilson, J. R. and C. W. Ford 1973 Temperature influences on the in *vitro* digestibility and soluble carbohydrate accumulation of tropical and temperate grasses. **Aust. J. Agric. Res.**, **24**: 187-198
- Wilson, J. R. and L. t'Mannetje 1978 Senescence, digestibility and carbohydrate content of buffel grass and green panic leaves in swards. **Aust. J. Agric. Res.**, **29**: 503-519
- Wilson, J. R., A. O. Taylor and G. R. Dolby 1976 Temperature and atmospheric humidity effects on cell wall content and dry matter digestibility of some tropical and temperate grasses. **N. Z. J. Agric. Res.**, **19**: 41-46
- Wood, J. T., D. J. Greenwood and T. J. Cleaver 1972 Interactions between the beneficial effects of nitrogen, phosphate and potassium on plant growth. **J. Agric. Sci., Camb.**, **78**: 389-391
-