Studies on the Trisomics in Rice Plants (*Oryza sativa* L.) : VII. Some Marker Genes Located on Chromosome 2 and Their Sequence on the Linkage Map

Iwata, Nobuo

Plant Breeding Laboratory, Faculty of Agriculture, Kyushu University

Satoh, Hikaru

Plant Breeding Laboratory, Faculty of Agriculture, Kyushu University

Omura, Takeshi

Plant Breeding Laboratory, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23800>

出版情報 : 九州大学大学院農学研究院紀要. 29 (2/3), pp.139-144, 1984-12. Kyushu University
バージョン :
権利関係 :

**Studies on the Trisomics in Rice Plants (*Oryza sativa* L.)
VII. Some Marker Genes Located on Chromosome 2
and Their Sequence on the Linkage Map**

Nobuo Iwata, Hikaru Satoh and Takeshi Omura

Plant Breeding Laboratory, Faculty of Agriculture,
Kyushu University, Fukuoka 812, Japan

(Received September 28, 1984)

Using trisomic method, some genes located on chromosome 2 were detected, and their gene sequence on the chromosome was inferred.

A single recessive gene for a bright green leaf mutant, *bgl*, was newly described in this report.

INTRODUCTION

Nagao and Takahashi (1963) tentatively proposed twelve linkage groups which corresponded to a haploid chromosome number of rice ($n=12$).

By the use of trisomic method, Iwata and Omura (1976) suggested a possibility that three linkage groups VI, IX and XII were located on the same chromosome, chromosome 2, which is the extra chromosome of trisomic type L.

By the use of reciprocal translocations involving chromosome 2, Sato *et al.* (1982) found the linkage relations between *nl-I*, a marker gene of linkage group IX, and both *d-l* and *gl-I*, marker genes of linkage groups VI and XII, respectively.

Yoshimura *et al.* (1982) also confirmed by the reciprocal translocation method that *d-l*, marker gene of linkage group VI, and *nl-I* and *ri*, marker genes of IX, were located together on chromosome 2.

In the present study, a new marker gene located on chromosome 2 was found by trisomic analysis, and a sequence of genes on chromosome 2 was inferred.

MATERIALS AND METHOD

Marker genes used in this study were *d-l*, *nl-2* and *ops* of linkage group VI, *al-K-2*, *al-K-3*, *nl-I*, *ri*, *spl-7* and *spl-8* of linkage group IX, *gl-1* of linkage group XII, and *bgl* described newly in the study (Table 1). A mutant for bright green leaf was induced by treating fertilized egg cells with N-methyl-N-nitrosourea (MNU) solution and designated as *bgl*. It was single recessive to the normal. Trisomic type L (Iwata *et al.*, 1970) was used for the trisomic analysis to find the genes located on chromosome 2. Recombination values

Table 1. List of marker genes used and their linkage groups.

Linkage ¹⁾ group	Gene ²⁾ symbol	Character	Reference
VI	d-1	daikoku dwarf	Nagao and Takahashi (1963)
	nl-2	neck leaf 2	Iwata and Omura (1977)
	<i>ops</i>	open spikelet sterile	" (1971)
IX	<i>al-K-2</i>	albino Kyushu 2	Iwata and Omura (1978)
	<i>al-K-3</i>	albino Kuysu 3	"
	<i>nl-1</i>	neck leaf 1	Nagao and Takahashi (1963)
	<i>ri</i>	verticillate rachis	"
	<i>spl-7</i>	spotted leaf 7	Iwata <i>et al.</i> (1978)
	<i>spl-8</i>	spotted leaf 8	"
XII	<i>gl-1</i>	glabrous leaf and hull 1	Nagao and Takahashi (1963)
	<i>bgl</i>	bright green leaf	

1) By Nagao and Takahashi (1963)

2) Although the standardization of gene symbols in rice has been discussed and new symbols are recommended for some genes by Japanese Committee on Nomenclature and Linkage Group of Rice, original symbols are used in this study to avoid the confusion, because the symbols recommended are still not authorized.

were calculated in F_2 and F_3 by the method of maximum likelihood.

RESULTS AND DISCUSSION

Segregation of marker genes in F_2 of the crosses with trisomic type L were shown in Table 2. Three marker genes of linkage group VI, d-1, *nl-2* and *ops*, two of IX, *nl-1* and *ri*, and one of XII, *gl-1*, showed the trisomic segregation. Regarding *bgl*, the segregation mode of dominant 63: recessive 1 was observed in the F_2 with type L and it was also considered to be the trisomic segregation. It is reasonably expected, therefore, that these seven genes compose a linkage group located on chromosome 2.

Linkage relations were observed between *bgl* and some marker genes belonging to linkage groups VI and IX (Table 3). From F_2 and F_3 data, the weighted means of recombination values on *bgl-d-1*, *bgl-ops*, *bgl-ri*, *bgl-spl-7* and *bgl-spl-8* were estimated at 34.2%, 12.6%, 2.9%, 16.1% and 15.4%, respectively. These facts show that the *bgl* links closely with *ri* of linkage group IX and also links with both *d-1* and *ops* of linkage group VI. It is obvious, therefore, that both linkage groups VI and IX which have been proposed by Nagao and Takahashi (1963) are combined in one group through the *bgl* gene.

The linkage relations among some marker genes belonging to the linkage groups VI or IX were investigated (Table 4). Recombination values of *nl-2-d-1*, *nl-2-ops* and *d-1-ops* were estimated at 22.3%, 44.0% and 30.7%, respectively, suggesting a sequence of *nl-2-d-1-ops*.

A recombination value of *spl-8-spl-7* was 29.6%. From this linkage intensity and the linkage intensities of *bgl-spl-7* (16.1%) and *bgl-spl-8* (15.4%) men-

Table 2. Trisomic segregation of the marker genes belonging to three linkage groups and of a marker gene described newly, in F_2 of crosses with trisomic type L.

Linkage group	Gene	Observed number			χ^2 for 3: 1
		Dominant	Recessive	Total	
VI	d-1	390	48	438	46.05***
	nl-2	477	32	509	95.06***
	ops	402	36	438	65.78***
IX	nl-1	281	24	305	47.74***
	ri	424	16	440	107.10***
XII	gl-1	410	24	434	87.75***
	bgl	63	1	64	18.75***

***: Significant at 0.1% level.

These data are partially quoted from Iwata and Omura (1976).

Table 3. Linkage relations between **bgl** and five genes belonging to the linkage groups Vi or IX.

Gene pair	Source of data	Phase ¹⁾	Segregation ²⁾	Total	Recombination value (%)	Weighted mean of recombination value (%)
bgl—d-1	F_2					
	F_3 (Ab)	R	326: 130: 115: 15	586	34.7±3.6	
		R	47: 41	88	30.4±4.5	
bgl—ops	F_3 (aB)	R	37: 28	65	39.8±6.0	34.2±2.5
	F_2					
	F_3 (Ab)	R	108: 49: 42: 0	199	≠0	
bgl—ri		R	38: 11	49	12.6±3.8	
	F_2					
	F_3 (Ab)	R	228 : 91: 61: 0	380	≠0	
bgl—spl-7		R	84: 4	88	2.3±1.2	
	F_3 (aB)	R	53: 5	58	4.5±2.0	2.9±1.0
bgl—spl-8	F_2					
	F_3 (Ab)	R	390: 194: 171: 5	760	16.5±3.5	
		R	68: 23	91	14.5±3.0	
bgl—spl-8	F_3 (aB)	R	63: 27	90	17.6±3.3	16.1±1.9
	F_2					
	F_3 (Ab)	R	191: 94: 91: 3	379	17.6±4.9	
bgl—spl-8		R	70: 24	94	14.6±3.0	15.4±2.5

1) R=Repulsion phase

2) F_2 =AB: Ab: aB: ab. F_3 (Ab)=AAbb: Aabb. F_3 (aB)=aaBB: aaBb.

tioned before, the sequence of *spl-8*, *spl-7* and **bgl** is considered as *spl-7—bgl—spl-8*.

Recombination values of *spl-7—nl-1*, *ri—spl-7* and *ri—nl-1* were estimated at 10.4%, 14.7% and 30.7%, respectively. The sequence of the three genes, thus, should be *ri—spl-7—nl-1*. The genes **bgl** and *ri* linked closely (2.9%) as mentioned above, therefore, a gene sequence of *spl-8—bgl—ri—spl-7—nl-1* or *spl-8—ri—bgl—spl-7—nl-1* is considered at this group. Moreover, based on the

Table 4. Linkage relations among the marker genes belonging to the linkage groups VI or IX.

Gene pair	Source of data	Phase ¹⁾	Segregation ²⁾	Total	Recombination value (%)	Weighted mean of recombination value (%)
<i>nl-2-d-1</i>	F ₂	R	398: 163: 194: 9	764	22.3±3.0	
<i>nl-2—ops</i>	F ₂	R	212: 80: 77: 19	338	44.0±4.1	
<i>d-1—ops</i>	F ₂	C	573: 142: 127: 104	946	34.1±2.0	
	F ₃ (Ab)	RR	294: 27: 24 120: 112: 2	528.51	30.8±6.0/14.8±4.2	30.7f1.7
<i>d-1-spl-7</i>	F ₂	R	106: 47: 41: 9	203	40.1f5.8	
<i>spl-8—spl-7</i>	F ₂	C	224: 58: 50: 47	379	33.0±3.1	
	F ₂	R	107: 49: 56: 1	213	13.6±6.7	29.6F2.8
<i>ri-spl-7</i>	F ₂	R	198: 91: 98: 1	388	10.4±5.0	
	F ₃ (Ab)	R	53: 14	67	11.7±3.1	
	F ₃ (aB)	R	43: 28	71	24.6±4.5	14.7±2.3
<i>ri-nl-1</i>	F ₂	c	241: 48: 45: 54	388	27.3±2.7	
	F ₂	R	220: 84: 93: 18	415	40.5±4.1	30.7±2.1
<i>spl-7—nl-1</i>	F ₂	R	194: 102: 92: 0	-388	0	
	F ₃ (Ab)	R	34: 11	45	13.9±4.2	
	F ₃ (aB)	R	56: 11	67	8.9±2.7	10.4±2.3

1) R=Repulsion phase. C=Coupling phase.

2) F₂=AB: Ab: aB: ab. F₃(Ab)=AAbb: Aabb. F₃(aB) =aaBB: aaBb.

Table 5. Linkage relations between two albino genes belonging to the linkage group IX and three marker genes locating on chromosome 2.

Gene pair	Source of data	Phase	Segregation ¹⁾	Total	Recombination value (%)	Weighted mean of recombination value (%)
<i>al-K-2—spl-7</i>	F ₃ (Ab)	R	52: 14	66	11.9±3.1	
	F ₃ (AB)	R	17: 107	124	15.0±3.8	13.1±2.4
<i>al-K-3—nl-2</i>	F ₃ (Ab)	R	46: 51	97	35.7f4.7	
	F ₃ (AB)	R	26: 90	116	26.8±6.0	32.3±3.7
<i>al-K-3-d-1</i>	F ₃ (Ab)	R	91: 38	129	17.3±2.3	
	F ₃ (AB)	R	52: 170	222	28.4±4.6	20.2±2.4

1) F₃(Ab)=AAbb: Aabb. F₃(AB)=AAB_: AaB_.

linkage intensities of *bgl-d-1* (**34.299**, *bgl—ops* (12.6%), *d-1-spl-7* (**40.1%**) and so on in Table 3 or 4, the gene sequence of ***d-1-spl-S-ops-bgl-spl-7*** was tentatively proposed, although a three point analysis was essential to get the exact order of these genes on the map.

Two albino genes, *al-K-2* and *al-K-3*, which linked to *nl-1* with recombination values of 1.1% and 34.6% respectively have been reported by Iwata and

Omura (1978). In the present study, the linkage relations between the albino genes and some marker genes located on chromosome 2 were observed (Table 5). Weighted mean of recombination values of *al-K-2-spl-7* was estimated at 13.1% and a gene sequence of either *spl-7-nl-1-al-K-2* or *spl-7-al-K-2-nl-1* is considered. The *al-K-3* was linked to *nl-2* and *d-1* with the recombination values of at 32.3% and 20.2%, respectively, suggesting the gene sequence of *nl-2-d-1-al-K-3*. The gene arrangement of *al-K-3* and *spl-8* at the map may be expected as *d-1-al-K-3-spl-8-&l*, however, further investigation will be necessary to confirm it.

Based on the results of the present study, the sequence of the ten genes on the chromosome 2 can be drawn tentatively as shown in Fig. 1.

Fig. 1. A linkage map of 10 genes on chromosome 2

Thus, the linkage group corresponded to the chromosome 2 was reconstructed with the marker genes of the two linkage groups which have been proposed tentatively by Nagao and Takahashi (1963).

Sato *et al.* (1982) estimated the gene sequence of *d-1-nl-1-gl-1* on the chromosome 2 from the facts that some interchanged points of reciprocal translocation lines involving chromosome 2 linked with both *d-1* and *nl-1*, and others linked with both *nl-1* and *gl-1*. Although ten genes were found to locate on the chromosome 2 in the present paper, none of the linkage relations between these genes and *gl-1* were found. To find out the genes linked with *gl-1*, trisomic analysis of mutant lines has been conducting by the use of type L.

REFERENCES

- Iwata, N. and T. Omura 1971 Linkage analysis by reciprocal translocation method in rice plants (*Oryza sativa* L.). I. Linkage groups corresponding to the chromosome 1, 2, 3 and 4. *Japan. J. Breed.*, **21**: 19-28 (in Japanese with English summary)
- Iwata, N. and T. Omura 1976 Studies on the trisomics in rice plants (*Oryza sativa* L.). IV. On the possibility of association of three linkage groups with one chromosome. *Japan. J. Genet.*, **51**: 135-137
- Iwata, N. and T. Omura 1977 Linkage studies in rice (*Oryza sativa* L.). On some mutants derived from chronic gamma irradiation. *J. Fac. Agr., Kyushu Univ.*, **21**: 117-127

- Iwata, N. and T. Omura 1978 Linkage studies in rice (*Oryza sativa* L.). Some albino genes and their linkage relations with marker genes. **Sci. Bull. Fac. Agr., Kyushu Univ.**, 33: 11-18 (in Japanese with English summary)
- Iwata, N., T. Omura and M. Nakagahra 1970 Studies on the trisomics in rice plants (*Oryza sativa* L.). I. Morphological classification of trisomics. Japan. **J. Breed.**, 20: 230-236
- Iwata, N., T. Omura and H. Satoh 1978 Linkage studies in rice (*Oryza sativa* L.). On some mutants for physiological leaf spots. **J. Fac. Agr., Kyushu Univ.**, 22: 243-251
- Nagao, S. and M. Takahashi 1963 Trial construction of twelve linkage groups in Japanese rice. Genetical studies on rice plants, XXVII. **J. Fac. Agr., Hokkaido Univ.**, 53: 72-130
- Sato, S., K. Muraoka and Y. Sano 1982 Reconstruction of a linkage group corresponding to the Nishimura's second chromosome in rice, *Oryza sativa* L. **Japan. J. Breed.**, 32: 232-238
- Yoshimura, A., N. Iwata and T. Omura 1982 Linkage analysis by reciprocal translocation method in rice plants (*Oryza sativa* L.). III. Marker genes located on chromosomes 2, 3, 4 and 7. **Japan. J. Breed.**, 32: 323-332