Pollen Tube Growth and Embryo Development in Interspecific Crosses of Cucurbita

Kwack, Soo Nyeon
University Farm, Faculty of Agriculture, Kyushu University

Fujieda, Kunimitsu
University Farm, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23809>

出版情報：九州大学大学院農学研究院紀要. 30 (1), pp.1-8, 1985-09. Kyushu University
バージョン：
権利関係：

Pollen Tube Growth and Embryo Development in Interspecific Crosses of *Cucurbita*

Soo Nyeon Kwack and Kunimitsu Fujieda

University Farm, Faculty of Agriculture, Kyushu University 46-06,
Kasuyamachi, Fukuoka 811-23

(Received March 15, 1985)

Pollen tube growth and seed production were investigated in a diallel involving three cultivated species (*Cucurbita moschata*, *C. maxima*, *C. pepo*) and a *PM* line selected from a cross between *C. pepo* × *C. moschata* in order to study the relationship between pollen tube growth and cross compatibility. All species combinations produced seeds with an embryo except one combination, *C. pepo* × *C. maxima*. The *PM* line, 'PM 143' showed similar results to *C. moschata* cv. 'Seoul-Dadagi' in pollen tube growth, fruit set and seed production in interspecific crosses. There were reciprocal differences in pollen tube growth, fruit set and seed production in *C. moschata* × *C. maxima*, *PM* × *C. maxima* and *C. maxima* × *C. pepo*. The differences in seed production between reciprocal crosses seem a result of the differences in pollen tube growth and in embryo development after fertilization.

INTRODUCTION

Interspecific hybridization in *Cucurbita* has been studied in order to transfer promising characters from one species to other species and to determine taxonomic relationships among taxa. Whitaker and Davis (1962) summarized studies on interspecific crosses in the genus *Cucurbita*. A later striking work was the breeding of *PM* lines, which have the early maturity and high female/male flower ratio of *C. pepo* cv. 'Zucchini' and the good flesh quality of *C. moschata* cv. 'Seoul-Dadagi' (Kim and Seo, 1976). Non-cultivated species have been crossed to cultivated species in order to exploit their disease resistance and to study evolution in this genus (Whitaker and Bemis, 1964; De Vaulx and Pitrat, 1980; Washek and Munger, 1983). Many workers have studied interspecific relationships by assessing fruit and seed set, and fertility of hybrid offspring (Whitaker and Davis, 1962). However, only a few reports focus on pollen tube growth in interspecific crosses (Hayase, 1950). Several techniques have been shown to aid in expediting interspecific crosses in *Cucurbita*, i.e. the use of highly heterozygous parents (Wall and York, 1962), bridging crosses (Rhodes, 1959), and embryo culture (Wall, 1954).

In this study, *in vivo* pollen tube growth and seed production were investigated in a diallel involving a *PM* line and three cultivated species (*C. moschata*, *C. maxima* and *C. pepo*) in order to study the relationship between pollen tube growth and cross compatibility.

MATERIALS AND METHODS

These experiments were carried out at the University Farm, Faculty of Agriculture, Kyushu University in 1984. We used *C. moschata* cv. 'Seoul-Dadagi', *C. maxima* cvs. 'Kuri' and 'Ebisu', *C. pepo* cvs. 'Zucchini' and 'Ford Zucchini' and a **PM** line, 'PM 143', selected from *C. pepo* cv. 'Zucchini' x *C. moschata* cv. 'Seoul-Dadagi', which resembles *C. moschata* in most characteristics. All of these cultivars are highly inbred lines. They were sown on March 20 and transplanted to the field on April 25. For each combination ten to twelve flowers were pollinated. About 50 days after pollination, fruits were harvested and after-ripened. Obtained seeds were sorted into empty seed, seed with a partially developed embryo and seed with a fully developed embryo.

For examining pollen tube growth, the same materials were sown directly in the field on August 24. To compare pollen tube growth in a self with that in an interspecific cross, we used 'Zucchini' selfs and 'Zucchini' x 'Kuri'. The pollinated pistils were harvested at 1, 3, 6, 10, 15, 21, 36, 72, and 96 hours after pollination. In the diallel crosses, more than six flowers were pollinated for each cross, and the pistils were harvested three days after pollination. The lengths of the style above the ovary, the ovary, and the stylar canal in the ovary of selfs were measured. Pollen tube growth was examined by fluorescence microscopy using Kho and Baer's method (1968). We observed pollen tube growth at six cross sections of each flower, i.e. the middle part of the style above the ovary, the point where the style enters the ovary, and 1/4, 2/4, 3/4, and 4/4 the depth of the stylar canal in the ovary. The number of pollen tubes which had penetrated to each depth were counted.

RESULTS

Percent fruit set in intra- and inter-specific crosses among a **PM** line and three cultivated species of *Cucurbita* are presented in Table 1. Fruit set in all intraspecific crosses was at least 80 %. Interspecific crosses showed 0 to 100% fruit set, with only one combination, 'Zucchini' x 'Ebisu', producing no fruit. Percent fruit set of the **PM** line in interspecific crosses was in most cases similar to that of *C. moschata* cv. 'Seoul-Dadagi'. When used as the female par-

Table 1. Percent fruit set in intra- and inter- specific crosses of *Cucurbita*

♀ \ ♂	<i>PM*</i>	<i>C. moschata</i>	<i>C. maxima</i>		<i>C. pepo</i>		
	PM 143	Seoul-Dadagi	Kuri	Ebisu	Zucchini	Ford	Zucchini
PM 143	—	90.9	83.3	33.3	10.0		75.0
Seoul-Dadagi	100	—	75.0	50.0	70.0		100
Kuri	33.3	33.3	—	85.7	10.0		58.3
Ebisu	100	100	80.0		58.3		50.0
Zucchini	75.0	8.3	16.7	0			100
Ford Zucchini	100	83.3	100	100	100		—

* A line selected from *C. pepo* x *C. moschata*.

ent, *C. pepo* cv. 'Ford Zucchini' showed a higher percent fruit set than the other cultivars. When used as the pollen parent, *C. pepo* cv. 'Zucchini' had lower ability to set fruit than any other cultivar.

Table 2 presents seed production per pollinated flower on the basis of embryo development. Embryo development in the cross between the **PM** line, 'PM 143', and *C. moschata* cv. 'Seoul-Dadagi' was similar to that in intra-specific crosses of *C. maxima* and *C. pepo*. Seeds with a fully developed embryo were obtained in crosses between **PM**, *C. moschata* and *C. maxima*, and another cross, 'Zucchini' x 'PM 143'. *C. maxima* x *C. moschata* and *C. maxima* x **PM** produced about ten seeds with a fully developed embryo per pollinated flower. In their reciprocal crosses, 'Ebisu' showed lower ability to produce seeds with a fully developed embryo than 'Kuri'. **PM** x *C. pepo* and *C. moschata* x *C. pepo* produced a few seeds with a partially developed embryo reciprocally. Unilateral incompatibility was found in the cross between *C. maxima* and *C. pepo*. *C. maxima* x *C. pepo* produced 1.5-8.5 seeds with a partially developed embryo, but its reciprocal cross produced no seeds with an embryo.

The lengths of the style above the ovary, the ovary, and the stylar canal

Table 2. Seed production per pollinated flower in intra- and inter- specific crosses of *Cucurbita*

♀	♂	<i>PM</i>	<i>C. moschata</i>	<i>C. maxima</i>		<i>C. pepo</i>	
		PM 143	Seoul-Dadagi	Kuri	Ebisu	Zucchini	Ford Zucchini
PM 143	A*	—	0	13.5	0.8	1.0	19.3
	B	—	2.5	2.9	2.9	0.1	3.8
	C	—	224.2	9.2	0	0	0
Seoul-Dadagi	A	0	—	15.4	2.3	14.6	44.0
	B	1.1	—	3.7	1.9	2.3	8.0
	C	198.5	—	9.4	0.8	0	0
Kuri	A	13.8	8.3	—	0	5.6	21.3
	B	8.8	3.7	—	4.1	1.5	1.9
	C	7.5	11.4	—	140.3	0	0
Ebisu	A	8.3	4.0	0	—	22.1	17.9
	B	34.6	52.4	8.9	—	8.5	1.6
	C	11.5	15.7	74.1	—	0	0
Zucchini	A	1.3	4.4	2.9	0	—	0
	B	3.9	1.9	0	0	—	2.4
	C	1.4	0	0	0	—	98.0
Ford Zucchini	A	7.7	5.1	14.9	13.1	0	—
	B	4.6	1.9	0	0	3.3	—
	C	0	0	0	0	117.6	—

* A, empty seed; B, seed with a partially developed embryo; C, seed with a fully developed embryo.

in the ovary of selfs measured at three days after pollination are presented in Table 3. The range in length of the style above the ovary was 2.0-3.2 cm. Total stylar canal length was very different among cultivars. The longest total stylar canal was 12.1 cm in a pistil of 'PM 143', and the shortest was 4.1 cm in a pistil of 'Ebisu'.

Table 3. Lengths of style above the ovary, ovary, and stylar canal in the ovary of selfs measured three days after pollination

Cultivar	Style above the ovary (cm)	Ovary (cm)	Stylar canal in the ovary (cm)	Total stylar canal (cm)
PM 143	3.0 ± 0.2	11.6 ± 1.5	9.1 ± 1.2	12.1 ± 1.4
Seoul-Dadagi	3.2 ± 0.1	11.2 ± 0.6	7.8 ± 0.7	11.0 ± 0.3
Kuri	2.5 ± 0.1	4.0 ± 0.5	2.1 ± 0.3	4.6 ± 0.4
Ebisu	2.2 ± 0.2	3.9 ± 0.4	1.9 ± 0.1	4.1 ± 0.5
Zucchini	2.4 ± 0.1	12.6 ± 2.2	9.2 ± 1.5	11.6 ± 1.4
Ford Zucchini	2.0 ± 0.1	10.3 ± 1.0	6.9 ± 0.6	8.9 ± 0.7

Fig. 1 depicts the pollen tubes at cross sections of a pistil. With the aid of the fluorescence technique the pollen tubes in the stylar canal were clearly discernible,

Fig. 2 shows pollen tube growth in 'Zucchini' selfs and 'Zucchini' × 'Kuri' observed at various hours after pollination. One hour after pollination, pollen had germinated on the stigma in both crosses. Six hours after pollination, pollen tubes of 'Zucchini' had penetrated into the stylar canal in the ovary of 'Zucchini', but those of 'Kuri' had not. By 72 hours after pollination, pollen tubes of 'Zucchini' selfs had grown into about 4/4 the depth of the stylar canal in the ovary. However, pollen tubes of 'Kuri' had grown to about 1/4 the depth of the stylar canal in the ovary of 'Zucchini' and had been arrested there.

Pollen tube growth in intra- and inter-specific crosses is shown in Fig. 3. Selfs and intraspecific crosses resulted in good pollen tube growth. Pollen tube growth in the crosses between 'PM 143' and *C. moschata* cv. 'Seoul-Dadagi' was similar to that in intraspecific crosses. When *C. maxima* was used as the female parent, the interspecific crosses showed good pollen tube growth. In *C. pepo*, the interspecific crosses of 'Zucchini' showed poor pollen tube growth, but those of 'Ford Zucchini' showed good growth.

DISCUSSION

Reports on interspecific hybridization within the genus *Cucurbita* are rather extensive (Whitaker and Davis, 1962). But according to Whitaker and Davis' summary, interspecific hybrids can be obtained only with difficulty from most species combinations of the cultivated *Cucurbita*. Our results showed that all species crosses produced seeds with an embryo except *C. pepo* × *C. maxima* (Table 2). In our study, 'PM 143', selected from *C. pepo* cv. 'Zucchini' × *C. moschata* cv. 'Seoul-Dadagi', was crossed with three cultivated species to investi-

Fig. 1. **Cross** sections showing the penetration of pollen tubes into the styler canal. A, many pollen tubes at the point where the style enters the ovary; B, many pollen tubes in the styler canal in the ovary; C, three pollen tubes in the styler canal in the ovary; D, no pollen tubes in the styler canal in the ovary.

gate the possibility of using 'PM 143' as a bridge plant. Although 'PM 143' has the cytoplasm of *C. pepo*, 'PM 143' x 'Seoul-Dadagi' reciprocally produced as many seeds as intraspecific crosses, and pollen tube growth and seed production in interspecific crosses of 'PM 143' showed similar results to those of 'Seoul-Dadagi'. These results indicate that the interaction between the cytoplasm and the nuclear genes does not play an important role in the success of interspecific crosses of *Cucurbita*. The differences between cultivars of *C. maxima* resulted in the great differences in seed production in *PM* x *C. maxima* and *C.*

Fig. 2. Pollen tube growth in 'Zucchini' selfs and 'Zucchini' x 'Kuri' observed at various hours after pollination. □, more than 30 pollen tubes observed; ◻, 10-30 pollen tubes observed; ◻, 1-9 pollen tubes observed.

moschata x **C. maxima**. Accordingly, the genetic constitution seems important in the success of interspecific crosses of **Cucurbita**.

Total stylar canal length was very different among cultivars (Table 3). **C. maxima**, with a short stylar canal, showed good pollen tube growth in the ovary of 'Ford Zucchini', with a long stylar canal (Fig. 3). So we suggest that the difference between the stylar canal length of the cultivars is not so important in interspecific crosses of **Cucurbita**.

Pollen tubes of **C. maxima** cv. 'Kuri', compared with those of 'Zucchini', grew slowly and were arrested at about 1/4 the depth of the stylar canal in the ovary of **C. pepo** cv. 'Zucchini' (Fig. 2). These results were consistent with Hayase's findings (1950) that pollen tube growth in **C. pepo** x **C. maxima** is inhibited in the upper region of the ovaries. However, pollen tubes in other crosses, **C. pepo** cv. 'Ford Zucchini' x **C. maxima** cv. 'Kuri' or 'Ebisu', grew into 3/4 the depth of the stylar canal in the ovary (Fig. 3). These differences may result from the difference in cultivars involved.

The observations on pollen tube growth in the stylar canal agreed with fruit set as a rule, but did not always agree with seed production. Although 'Ford Zucchini' x **C. maxima** showed good pollen tube penetration into the deep stylar canal in the ovary, they produced no seeds with an embryo. We observed many ovules penetrated by pollen tubes in these crosses. These results indicate that after fertilization the embryo may degenerate in the early developmental stage.

Reciprocal crosses showed great differences in pollen tube growth, fruit set and seed production in **C. moschata** x **C. maxima**, **PM** x **C. maxima** and **C. max-**

Fig. 3. Pollen tube growth in intra- and inter-specific crosses of *Cucurbita*. A, a PM line, 'PM 143'; B, *C. moschata* cv. 'Seoul-Dadagi'; C and D, *C. maxima* cvs. 'Kuri' and 'Ebisu'; E and F, *C. pepo* cvs. 'Zucchini' and 'Ford Zucchini'. □, more than 30 pollen tubes observed; ▨, 10-30 pollen tubes observed; —, 1-9 pollen tubes observed.

ima x *C. pepo*. The differences in pollen tube growth and embryo development may result in the differences in seed production between reciprocal crosses.

ACKNOWLEDGEMENTS

We wish to thank the Hungnong Seed Company, Seoul, Korea for providing seed materials and also Miss G. M. Caddell for help in preparing this manuscript.

REFERENCES

- De Vaulx, R. D. and M. Pitrat 1980 Realization of the interspecific hybridization (F,

- and BC₁) between *Cucuybita pepo* and *C. ecuadoryensis*. *Cucuybit Genetics Cooperative*, 3 : 42
- Hayase, H. 1950 *Cucuybita-crosses*, I. The pollen tube growth in interspecific crosses. *Jap. Jour. Genet.*, 25: 181-190 (in Japanese with English Resume)
- Kho, Y. and J. Baer 1968 Observing pollen tubes by means of fluorescence. *Euphytica*, 17: 298-302
- Kim, B. H. and Y. G. Seo 1976 Studies on the interspecific crosses. (II) Breeding process and characteristics of Bulam Sacheol Aehobak. *Jour. Kor. Soc. Hoyt. Sci.*, 17: 38-46 (in Korean with English abstract)
- Rhodes, A. M. 1959 Species hybridization and interspecific gene transfer in the genus *Cucuybita*. *Proc. Amer. Soc. Hort. Sci.*, 74: 546-551
- Wall, J. R. 1954 Interspecific hybrids of *Cucuybita* obtained by embryo culture. *Proc. Amer. Soc. Hort. Sci.*, 63: 427-430
- Wall, J. R. and T. L. York 1962 Gametic diversity as an aid to interspecific hybridization in *Phaseolus* and *Cucuybita*. *Proc. Amey. Soc. Hor. Sci.*, 75: 419-428
- Washek, R. L. and H. M. Munger 1983 Hybridization of *Cucurbita pepo* with disease resistant *Cucuybita* species. *Cucurbit Genetics Cooperative*. 6: 92
- Whitaker, T. W. and W. P. Bemis 1964 Evolution in the genus *Cucurbita*. *Evolution*, 18: 553-559
- Whitaker, T. W. and G. N. Davis 1962 Cucurbits, botany, cultivation and utilization. Leonard Hill Books Ltd., London (England)