

Synopsis of Andrena (Andrena) of Japan (Hymenoptera, Andrenidae) Part II

Tadauchi, Osamu
Entomological Laboratory, Faculty of Agriculture, Kyushu University

Hirashima, Yoshihiro
Entomological Laboratory, Faculty of Agriculture, Kyushu University

Matsumura, Takeshi
National Grassland Research Institute

<https://doi.org/10.5109/23833>

出版情報：九州大学大学院農学研究院紀要. 31 (1/2), pp.37-54, 1987-02. Kyushu University
バージョン：
権利関係：



Synopsis of *Andrena* (*Andrena*) of Japan
(Hymenoptera, Andrenidae)
Part II ^{1),2)}

Osamu Tadauchi*, Yoshihiro Hirashima* and Takeshi Matsumura †

* Entomological Laboratory, Faculty of Agriculture,
Kyushu University 46-01, Fukuoka 812

† National Grassland Research Institute, Nishinasuno,
Tochigi 329-27

(Received May, 27, 1986)

In this paper (Part II) seven Japanese species of *Andrena* (*Andrena*) are treated. A new species, *Andrena* (*A.*) *babai*, and a new subspecies, *Andrena* (*A.*) *lapponica sumizome* are described. *Andrena* (*A.*) *shirozui* Hirashima is reduced to the subspecies of *A. lapponica* Zetterstedt. The males of *A. (A.) lapponica shirozui* Hirashima and *A. (A.) maukenis* Matsumura are described for the first time. The distribution map, floral records and flight records of each species are presented.

10. *Andrena* (*Andrena*) *hondoica* Hirashima

(Figs. 9, 24, 39, 72, 94, Table 4)

Andrena (*Andrena*) *hondoica* Hirashima, 1962, J. Fac. Agr., Kyushu Univ., 12 : 144, female & male ;
Hirashima, 1966, *ibid.*, 14 : 103.

Distribution : Japan (Hokkaido and Honshu). (Fig. 72)


Floral records : Hirashima (1962) recorded *Veronica caninotesticulata* and *Salix* sp. We examined 19 females and 6 males collected on 7 plants as follows : Salicaceae : *Salix Pet-susu* Kimura (2 ♀) ; *S. gracilistyla* Miq. (8 ♀ 1 ♂) ; *S. sachalinensis* Fr. Schm. (1 ♀) ; *S. gigiana* Seemen (3 ♀) ; *S. sp.* (3 ♀ 2 ♂). Compositae: *Petasites japonicus* subsp. *giganteus* Kitam. (1 ♀ 3 ♂). Scrophulariaceae : *Veronica didyma* var. *lilacina* Yamaz. (1 ♀).

Flight records : Female : mid March to late May (Honshu) ; late April to late May (Hokkaido). Male: mid March to early May (Honshu) ; late April to mid May (Hokkaido).

Specimens examined : We have examined 95 females and 18 males. Some of them are cited as follows: HOKKAIDO: 1 ♀ 3 ♂, Kussharo-ko, Teshikaga, 6. v. 1984 (O. Tadauchi) ; 2 ♀ 1 ♂, Kyushu Univ. Forests, Ashoro, 28. iv. 1984 (O. Tadauchi) ; 1 ♀, Yamabe, Furano, 23. v. 1984 (O. Tadauchi) ; 1 ♂, Jozankei, Sapporo, 28. iv. 1957 (T. Nanbu) ; 1 ♀, Moheji, Kamiiso, Oshima, 19. v. 1984 (O. Tadauchi). HONSHU : 8 ♀, Aomori, Aomori Pref., 18. v. 1984 (O. Tadauchi) ; 3 ♀, Shiga-kogen, Nagano Pref., 7.

¹⁾ Partly supported by a grant (1984) from the Japan Society for the Promotion of Science, Tokyo.

²⁾ Contribution from the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka (Ser. 3, No. 213).



Figs. 72-73. Maps showing the distributions of *Andrena*(*Andrena*) ***hondoica*** Hirashima (72) and *A. (A.) longitibialis* Hirashima (73). A square indicates the type locality.

v. 1971 (S. Hashimoto) ; 1 ♀, Rokuroshi, Ohno, Fukui Pref., 5. iv. 1972 (Y. Haneda) ; 1 ♀, Suwara, Ohno, Fukui Pref., 31. v. 1981 (Y. Haneda) ; 1 ♂, Hikimi, Shimane Pref., 12. iv. 1986 (O. Tadauchi).

Other localities (*new records) : Obihiro, Sapporo, Narukawa, Hakodate, Kikonai (Hokkaido) ; *Takizawa-mura, *Kuriyagawa, *Ashiro (Iwate Pref.) ; Kadota-mura (Fukushima Pref.) ; *Kawasaki-Mukogaoka (Kanagawa Pref.) ; Kurokawa (Niigata Pref.) ; Ina (*type* locality), Ueda, *Utsukushigahara (Nagano Pref.) ; *Masutomi (Yamanashi Pref.) ; Asahi-mura, Minami-mura (Gifu Pref.) ; Tsurugi (Ishikawa Pref.) ; Katsuyama, Izumi (Fukui Pref.) ; Sasayama (Hyogo Pref.) ; *Mt. Daisen (Tottori Pref.) ; *Hiwa (Hiroshima Pref.).

11. *Andrena* (*Andrena*) ***longitibialis*** Hirashima

(Figs. 10, 25, 40, 73, 94, Table 4)

Andrena(*Andrena*) ***longitibialis*** Hirashima, 1962, J. Fac. Agr., Kyushu Univ., 12 : 139, female & male ; Hirashima, 1966, *ibid.*, 14 : 102 and 116.

Distribution : Japan (Hokkaido, Rishiri Is., Honshu, Sado Is., Izu-Oshima Is., Oki Is., Shikoku, and Kyushu). This is the first records of this species from the Izu-Oshima

and Ōki Is. (Fig. 73)

Floral records : Ikudome (1980) recorded 3 plants. We examined 25 females and 36 males collected on 14 plants as follows : Ericaceae : *Rhododendron reticulatum* D. Don (8 ♂); *R. metternichii* Sieb. et Zucc. (17 ♂); *R. kiusianum* Makino (12 ♀); *R. macrosepalum* Maxim. (1 ♀); *R. sp.* (3 ♀); *Enkianthus perulatus* Schneid. (1 ♀ 7 ♂). Salicaceae : *Salix yoshinoi* Koidz. (1 ♀). Rosaceae: *Prunus persica* Batsch. (1 ♂). Cruciferae: *Brassica Napus* L. (2 ♀ 1 ♂); *Capsella Bursa-pastoris* Medic. (1 ♂). Labiatae : *Lamium album* var. *barbatum* Franch. et Savat. (2 ♀); *Ajuga decumbens* Thunb. (1 ♀). Compositae : *Chrysanthemum leucanthemum* Linn. (2 ♀). Saxifragaceae: *Philadelphus satsumi* Sieb. (1 ♀).

Flight records : Female : early May to mid July (Kyushu) ; early May to early June (Shikoku) ; mid March to early July (Honshu) ; mid June (Hokkaido). Male : mid May (Kyushu) ; mid April to late May (Honshu).

Specimens examined : We have examined 70 females and 50 males. Some of them are cited as follows : HONSHU : 3 ♀, Nasu-Yumoto, Tochigi Pref., 22. v. 1975 (O. Tadauchi); 1 ♂, Hodosan, Saitama Pref., 14. iv. 1972 (T. Nanbu); 2 ♀, Kurokawa, Niigata Pref., 17. v. 1985 (K. Baba) ; 1 ♂, Mt. Haku, Ishikawa Pref., 26. v. 1962 (I. Togashi); 1 ♂, Mitsudani, Shiramine-mura, Ishikawa Pref., 29. v. 1974 (I. Togashi); 1 ♀, Mt. Fujikura, Imajo, Fukui Pref., 23. v. 1981 (H. Kurokawa) ; 3 ♂, Hatahoko, Mt. Norikura, Gifu Pref., 8. v. 1976 (O. Tadauchi) ; 1 ♂, Taishaku-kyo, Hiroshima Pref., 30. iv. 1978 (H. Makihara). SADO IS. : 2 ♀, Washizaki, Ryotsu, 28. v. 1985 (K. Baba) ; 2 ♀, Urakawa, Ryotsu, 28. v. 1985 (K. Baba). IZU-OSHIMA IS.: 1 ♂, 30. iii. 1949 (K. Maruyama). OKI IS. : 1 ♀, Omosu, 10. v. 1982 (Y. Maeta). KYUSHU : 3 ♂, Chojabaru, Kuju, Oita Pref., 17-18. v. 1970 (K. Kanmiya) ; 17 ♂, Mt. Kuju, Oita Pref., 11. v. 1972 (Y. Hirashima) ; 14 ♂, Danbaru, Mt. Kuju, Oita Pref., 12. v. 1972 (Y. Hirashima) ; 5 ♂, Mt. Daisen, Mts. Kuju, 12. v. 1978 (Y. Hirashima) ; 5 ♀, Hokkein, Kuju, Oita Pref., 5. vi. 1985 (O. Tadauchi).

Other localities (*new records) : Rishiri Is., Hakodate (Hokkaido) ; Mt. Hakkoda (Aomori Pref.) ; *Morioka (Iwate Pref.) ; Mt. Zao (Miyagi Pref.) ; *Kuroiso (Tochigi Pref.) ; *Hanno (Saitama Pref.) ; *Mt. Mitake (Tokyo) ; Karuizawa (Nagano Pref.) ; *Masutomi (Yamanashi Pref.) ; Senami, Mt. Yahiko ; Mt. Kinpoku-Sado Is. (Niigata Pref.) ; Fukui, Kamishihi, Ohno, Izumi, Ikeda, Takefu, Tsuruga, Mikata (Fukui Pref.) ; Kibi (Wakayama Pref.) ; *Kyoto (Kyoto Pref.) ; *Sasayama (Hyogo Pref.) ; *Nagi (Tottori Pref.) ; *Matsue (Shimane Pref.) ; Mt. Saragamine (Ehime Pref.) ; Tosayama, Godaisan (Kochi Pref.) ; Mt. Hikosan, *Mt. Homan (Fukuoka Pref.) ; *Mt. Aso (Kumamoto Pref.).


12. *Andrena* (*Andrena*) *maukensis* Matsumura

(Figs. 11, 26, 41, 74-79, 94, Table 4)

Andrenamaukensis Matsumura, 1911, J. Coll. Agr., Tohoku Imp. Univ., 4 : 107, female ; Yasumatsu 1941, Peking Nat. Hist. Bull., 15 : 277.

Andrena (*Andrena*) *maukensis* Hirashima, 1962, J. Fac. Agr., Kyushu Univ., 12 : 142, female ; Hirashima, 1966, *ibid.*, 14 : 103.

The description of male, which is new to science is given below.




Figs. 74-78. Genital capsule and associated structures of *Andrena (Andrena) mauken-sis* Matsumura, 74 : dorsal view of genital capsule, 75 : lateral view of the same, 76 : ventral view of the same, 77 : 7th sternum, 78 : 8th sternum.

Male : Length 7.5-8.0 mm.

Integumental color : Mandibles reddened apically ; flagellum brownish beneath ; posterior margins of metasomal terga brownish subhyaline.

Pubescence : Hairs on head not long, not dense, mainly dull whitish to pale yellowish, mixed with sparse, fuscous hairs on paraocular area, frons and genal area near eye and vertex ; those on thorax long, dull whitish to pale yellowish, sometimes mixed with fuscous hairs ; those on legs pale yellowish ; those on metasomal terga 1-2 long, whitish to pale yellowish, on terga 3-5 short, whitish, on tergum 6 whitish; metasomal sterna 2-5 each with a sparse, curled fringe apically.

Structure : Mandible long, curved, falciform, with sharp apex ; basal projection of mandible nearly triangular, not long; malar space short, about 1/6 times as long as base of mandible ; process of labrum slightly emarginate, well convex, smooth and shiny ; clypeus transversely convex, narrowly tessellate on basal portion, smooth and shiny with moderate-sized, sparse punctures on the rest of clypeus ; clypeus without an impunctate, median space ; genal area not distinctly angled behind ; flagellum 1 about as long as broad, shorter than 2 or 3 ; mesoscutum densely tessellate, indistinctly punctate, dull ; propodeum roughened ; enclosure ill defined, rugose basally, tessellate



Figs. 79-80. Maps showing the distributions of *Andrena* (*Andrena*) *maukensis* Matsumura (79) and *A.* (*A.*) *lapponica shirozui* Hirashima (black) and *A.* (*A.*) *l. sumizome*, new subspecies (white) (80). A square indicates the type locality.

apically; mesepisternum densely tessellate with a little roughened punctures; metasoma tessellate, especially on basal terga, becoming smoother toward apical terga; posterior depressions of terga broad, not well indicated; genitalia and associated structures as illustrated (Figs. 74-78).

Distribution: Japan (Hokkaido, Honshu, and Kyushu) and Saghalien, the Kuril Is. (Etorofu and Shakotan Is.). This is the first record of this species from the Kuril Is. and Kyushu. (Fig. 79)

Floral records: Hirashima (1962) recorded *Vaccinium vitis-idea* L. from Honshu. We record 6 new plants as follows: Compositae: *Taraxacum officinale* Weber (5 ♀ 4 ♂); *Solidago virgaurea* subsp. *leiocarpa* Hulten (2 ♀). Salicaceae: *Salix taraiensis* Kimura (1 ♂). Leguminosae: *Lathyrus japonicus* Willd. (1 ♂). Rosaceae: *Rubus parvifolius* L. (1 ♂). Aquifoliaceae: *Ilex sugerokii* var. *brevipedunculata* S. Y. Hu (1 ♀).

Flight records: Female: mid July (Kyushu); early June to late August (Honshu); mid June to mid August (Hokkaido). Male: early June to early July (Honshu); late May to mid July (Hokkaido).

Specimens examined: We have examined 23 females and 11 males. Some of them are cited as follows: THE KURIL IS.: Etorofu Is.: 1 ♀ (S. Kuwayama & Y. Sugihara); 1 ♂ (Y. Sugihara); Shakotan Is.: 1 ♂ (S. Kuwayama & Y. Sugihara). HOK-

KAIDO : 2 ♀, Rausu-Seseki, Shiretoko, 16. vi. 1984 (O. Tadauchi) ; 2 ♂, Mt. Rausu, 1300 m, Shiretoko, 23. vi. 1963 (T. Saigusa) ; 1 ♂, Ochiishi, Nemuro, 28. vi. 1957 (R. Ishikawa) ; 1 ♂, Akkeshi, 24. vi. 1965 ; 1 ♂, Tokachimitsumata, Kamishihoro, 29. v. 1984 (O. Tadauchi) ; 2 ♀, same locality and collector as above, 12. vi. 1984 ; 1 ♀, Kogen Spa, Mt. Taisetsu, 15. viii. 1984 (O. Tadauchi) ; 1 ♀, Ginsendai, Mt. Taisetsu, 14. viii. 1984 (O. Tadauchi) ; Tennin-kyo, Mt. Taisetsu, 27. vi. 1967 (T. Saigusa) ; 1 ♂, Yudoo Lake, Tokachi, 3. vii. 1975 (H. Fukuda) ; 1 ♀, Yamabe, Furano, 23. v. 1984 (O. Tadauchi). HONSHU : 1 ♀, Mt. Iwaki, Aomori Pref., 17. vii. 1960 (S. Masaki) ; 1 ♂, Mt. Chokai, Akita Pref., 20. vi. 1978 (K. Baba & N. Kato) ; 3 ♂, Kanayama, Masutomi, Yamanashi Pref., 8. vi. 1975 (J. Emoto) ; 1 ♀, Karuizawa, Nagano Pref., 7. vi. 1952 (R. Ishikawa) ; Mt. Haku, Ishikawa Pref., 1 ♀, 22. vii. 1962 (I. Togashi) ; same locality and collector as above: 3 ♀, vii. 1973 ; 1 ♀, 2. vii. 1973 ; 1 ♀, 1. viii. 1973 ; 1 ♂, 19. vii. 1977 ; 1 ♀, 20. vii. 1977 ; 1 ♀, 21. vii. 1979 ; 1 ♂, 15. vii. 1981. KYUSHU: Mt. Tsurugi, Oita Pref., 12. vii. 1978 (Y. Hirashima).

Other localities (*new records) : *Wakkanai, Hamakoshimizu, *Akkeshi, *Mt. Memuro, Nukabira, *Mikuni-toge, Aizankei, Yukomanbetsu, *Sapporo (Hokkaido) ; Mt. Hakkoda (Aomori Pref.) ; Mt. Chokai, Oguni (Yamagata Pref.) ; Mt. Takizawa, Renge Spa (Niigata Pref.) ; Ohno (Fukui Pref.).

13a. *Andrena (Andrena) lapponica shirozui* Hirashima, new status
(Figs. 12, 27, 42, 80-85, 94, Table 4)

Andrena (Andrena) shirozui Hirashima, 1962, J. Fac. Agr., Kyushu Univ., 12: 143, female; Hirashima, 1966, *ibid.*, 14 : 103.

We treat here *Andrena (A.) shirozui* Hirashima as a subspecies of the European *Andrena lapponica* Zetterstedt. It is separable from the nominate subspecies by the clypeus less tessellate, broadly shiny and smooth with sparser and weaker punctures, the tibial scopa broadly whitish in front, pale brownish behind, the hairs on the thorax whitish, the caudal fimbria brownish, the male clypeus with much blackish hairs and the hairs on the thorax whitish.


This subspecies distributes in Hokkaido and northern Honshu (Tohoku Districts). The description of male, which is new to science is given below.

Male : Length about 7.5 mm.

Integumental color : Mandibles reddened apically ; flagellum brownish beneath ; posterior margins of metasomal terga yellowish brown subhyaline.

Pubescence : Hairs on head not long, not dense, dull whitish, mixed with fuscous hairs ; those on paraocular area, frons, and genal area and vertex fuscous ; hairs on thorax dull whitish to pale yellowish, not mixed with fuscous hairs above ; hairs on legs pale to yellowish ; those on metasomal terga 1-2 long, dull whitish, on 3-5 terga short, brownish, on tergum 6 whitish ; metasomal sterna 2-5 without long, curled fringes apically.

Structure : Mandible long, curved, falciform, with sharp apex ; basal projection of mandible not long ; malar space long, about 1/3 times as long as base of mandible ; process of labrum transverse, well convex, smooth and shiny ; clypeus longitudinally convex, narrowly tessellate on basal portion, smooth and shiny with moderate-sized,



Figs. 81-85. Genital capsule and associated structures of *Andrena* (*Andrena*) *lapponicashirozui* Hirashima, 81: dorsal view of genital capsule, 82: lateral view of the same, 83: ventral view of the same, 84: 7th sternum, 85: 8th sternum.

sparse punctures on the rest of clypeus; clypeus with a narrow, impunctate, median space; genal area not distinctly angled behind; flagellum 1 about 1.5 times as long as wide, slightly shorter than 2 or 3; mesoscutum densely tessellate, punctate, dull; propodeum roughened; enclosure ill defined, narrowly rugose basally, tessellate apically; mesepisternum nearly roughened with a little roughened punctures; metasomal terga tessellate, becoming smoother toward apical terga; posterior depressions of terga broad, not well indicated; genitalia and associated structures as illustrated (Figs. 81-85).

Distribution: Japan (Hokkaido and northern Honshu). (Fig. 80)

Floral records: Hirashima (1962) recorded 2 plants in northern Honshu. We newly record 2 plants as follows: Compositae: *Taraxacum officinale* Weber (1 ♂). Rosaceae: *Gerum pentapetalum* Makino (1 ♀).

Flight records: Female: early July (northern Honshu); mid June to late July (Hokkaido). Male: mid June to early July (Hokkaido).

Specimens examined: HOKKAIDO: 1 ♂, Mt. Rausu, Shiretoko, 23. vi. 1967 (T. Saigusa), 7 ♂, Mt. Taisetsu, 7. vii. 1964 (T. Matsumura); 1 ♂, Mikuni Pass, Mt. Taisetsu, 11. vi. 1974 (H. Fukuda); 1 ♂, Sugatami, Mt. Taisetsu, 10. vii. 1967 (T. Matsumura); 1 ♂, same locality and collector as above, 23. vi. 1967; 1 ♀, Mt. Yotei,

19. vi. 1958 (M. Shiokawa).

Other localities : Ashoro, Nukabira (*type locality*), Aizankei, Yukomanbetsu (Hokkaido) ; Mt. Hakkoda (Aomori Pref.) ; Mt. Zao (Miyagi Pref.).

13b. ***Andrena (Andrena) lapponica sumizome* Tadauchi, Hirashima et Matsumura,
new subspecies**
(Figs. 80, 94)

Andrena (Andrena) sp. (aff. *shirozui*) Yamauchi et al., 1982, A list of Insects in Gifu Pref., 422.

This new subspecies is found in the alpine regions in central Honshu, such as the North and South Japan Alps and Mt. Haku. The female of the new subspecies is characteristic in having the vertex and the antennal region with much fuscous hairs and not mixed with whitish ones, the mesepisternum mixed with much fuscous hairs, the metasomal terga 2-4 hairy and each with an obscure fringe of suberect to erect, sooty white hairs, and the wing darker.

Distribution : Japan (alpine region, central Honshu). (Fig. 80)

Floral record : Not available.

Flight records : Female : late June to mid August.

Type material: Holotype female (Type No. 2574, Kyushu Univ.), Gonoike, Mt. Ontake, 23. vi. 1976 (K. Yamauchi). Paratypes : same locality and collector as holotype: 1 ♀, 23. vi. 1976; 2 ♀, 18. vii. 1974; 1 ♀, 1. viii. 1976; 2 ♀, 16. vii. 1978; 2 ♀, Komagatake, Yamanashi Pref., 18. vii. 1956 (H. Kamiya); 1 ♀, Mt. Komagatake, Yamanashi Pref., 25. vii. 1936 (H. Masuda); 1 ♀, Mt. Senjo-Kitadake, Yamanashi Pref., 28-29. vii. 1959 (Y. Miyatake); 1 ♀, Mt. Haku, 2000 m, Ishikawa Pref., 11. viii. 1959 (I. Togashi); 2 ♀, Mt. Haku, Ishikawa Pref., 22. vii. 1962 (I. Togashi); 1 ♀, Midagahara, Mt. Haku, 29. vii. 1956 (S. Nakamura).

Type depository : The type is preserved in the collection of the Entomological Laboratory, Faculty of Agriculture, Kyushu University.

14. ***Andrena (Andrena) aburana* Hirashima**
(Figs. 13, 28, 43, 86, 94, Table 4)


Andrena (Andrena) aburana Hirashima, 1962, J. Fac. Agr., Kyushu Univ., 12 : 146, female & male ; Hirashima, 1966, *ibid.*, 14 : 103 and 116.

Distribution : Japan (Honshu, Sado Is., Shikoku, and Kyushu). (Fig. 86)

Floral records : Cruciferae : *Brassica Napus* L. (4 ♀). Rosaceae : *Prunus Mume* Sieb. et Zucc. (1 ♀). Salicaceae: *Salix* sp. (1 ♀).

Flight records : Female : late March to early May (Kyushu) ; late March to late May (Honshu). Male : early to mid March (Kyushu) ; early to late March (Shikoku) ; late March to late April (Honshu).

Specimens examined : We have examined 38 females and 22 males. Some of them are cited as follows : HONSHU : 1 ♂, Kurokawa, Niigata Pref., 10. iv. 1985 (K. Baba) ; 1 ♀ 2 ♂, Mt. Utatsu, Kanazawa, Ishikawa Pref., 10. iv. 1974 (Y. Yoneda) ; 1 ♀, Fukami, Monzen, Ishikawa Pref., 19. iv. 1972 (I. Togashi); 1 ♀, Imajo, Fukui Pref., 18. iv. 1981 (Y. Haneda). SADO IS. : 3 ♀, Suizu, 10. v. 1980 (K. Baba). KYUSHU: 1 ♀,



Figs. 86-87. Maps showing the distributions of *Andrena* (***Andrena***) *aburana* Hirashima (86) and ***A. (A.) benefica*** Hirashima (87). A square indicates the type locality.

Kuroubaru, Chikuhō, Fukuoka Pref., 7. iv. 1975 (O. Tadauchi) ; 1 ♀, Mt. Koshōzan, Fukuoka Pref., 21. iii. 1962 (S. Ide) ; 2 ♀, Mt. Korasan, Kurume, Fukuoka Pref., 14. iv. 1963 (S. Ide) ; 1 ♀, Kurume, Fukuoka Pref., 3. iv. 1963 (S. Ide) ; 1 ♀, Mt. Hikosan, Fukuoka Pref., 4. v. 1976 (K. Ōhara) ; 1 ♀, Marutani, Miyakonojo, Miyazaki Pref., 7. iv. 1976 (O. Tadauchi) ; 1 ♀, Tabuse, Murasho, Miyazaki Pref., 5. iv. 1976 (O. Tadauchi).

Other localities (*new records) : *Yorii (Saitama Pref.) ; *Kiyosumi (Chiba Pref.) ; Mt. Mitake (Tokyo) ; Iwagasaki, Sasagamine (Niigata Pref.) ; *Shirayama (Ishikawa Pref.) ; Fukui, Eiheiiji, Kamishihi, Katsuyama, Ohno, Ikeda, Mihama, Natasho (Fukui Pref.) ; "Kyoto (Kyoto Pref.) ; Mt. Minoo (Osaka Pref.) ; *Sasayama (Hyogo Pref.) ; *Matsue (Shimane Pref.) ; Hagi (Yamaguchi Pref.) ; Matsuyama (Ehime Pref.) ; Kashii, *Inunaki, Mt. Homan (***type locality***), *Dazaifu, Mt. Narutake (Fukuoka Pref.) ; Kumamoto (Kumamoto Pref.).

15. *Andrena* (***Andrena***) ***benefica*** Hirashima

(Figs. 14, 29, 44, 87, 94, Table 4)

Andrena (***Andrena***) ***benefica*** Hirashima, 1962, J. Fac. Agr., Kyushu Univ., 12 : 151, female & male ;

Hirashima, 1966, *ibid.*, 14: 104 and 115; Kim and Kim, 1983, *Kor. J. Ent.*, 13: 5.

Distribution : Japan (Hokkaido, Honshu, Sado Is., Shikoku, and Kyushu) and Korea. This is the first record of this species from Shikoku. (Fig. 87)

Floral records : Hirashima (1962) recorded 6 plants. We examined 439 females and 103 males collected on 17 plants as follows : Cruciferae : *Brassica Napus* L. (404 ♀ 33 ♂); *B. pekinensis* Pupr. (1 ♀); *Capsella Bursa-pastoris* Medic. (1 ♂); *Raphanus sativus* var. *hortensis* Backer (2 ♀). Aceraceae: *Acer ginnala* Maxim. (1 ♀); *A. sp.* (11 ♀ 1 ♂). Compositae : *Taraxacum officinale* Weber (3 ♀); *Lapsana apogonoides* Maxim. (1 ♀). Salicaceae : *Salix gracilistyla* Miq. (2 ♀); *S. sp.* (1 ♀ 67 ♂). Rosaceae : *Potentilla kleiniana* subsp. *anemonefolia* Murata (2 ♀); *Crataegus cuneata* Sieb. et Zucc. (1 ♀); *Pyrus pyrifolia* var. *culta* Nakai (1 ♀); *Malus pumila* Mill. (1 ♀); *Pourthiaea villosa* var. *laevis* Stapf (4 ♀). Ericaceae : *Pieris japonica* D. Don (1 ♀). Liliaceae : *Gagea lutea* Ker-Gawl. (1 ♀).

Flight records : Female : mid March to late May (Kyushu) ; mid April to early May (Shikoku); early April to early June (Honshu); late April to early July (Hokkaido). Male : mid March to late April (Kyushu) ; mid April (Shikoku) ; early April to late May (Honshu) ; late April to early May (Hokkaido).

Specimens examined : We have examined 520 females and 153 males. Some of them are cited as follows. HOKKAIDO : 1 ♀, Kyushu Univ. Forests, Ashoro, 11. vi. 1984 (O. Tadauchi) ; 2 ♀, Hokkaido Univ. Campus, Sapporo, 13. v. 1984 (O. Tadauchi). HONSHU : 15 ♀, Nishihiranaï, Aomori Pref., 30. v. 1976 (O. Tadauchi) ; 1 ♂, Shichiri, Nikko, Tochigi Pref., 11. v. 1976 (O. Tadauchi) ; 1 ♂, Arai, Niigata Pref., 24. v. 1976 (O. Tadauchi) ; 3 ♀ 17 ♂, Hatahoko, Mt. Norikura, Gifu Pref., 8. v. 1976 (O. Tadauchi) ; 2 ♀, Kabuto, Seki, Mie Pref., 13. v. 1975 (O. Tadauchi) ; 68 ♀, Masumizu, Mt. Daisen, Tottori Pref., 26. iv. 1975 (O. Tadauchi). SADO IS. : 1 ♂, Mt. Myoken, 12. v. 1985 (K. Baba). SHIKOKU : 4 ♀, Hashikura, Ikeda, Tokushima Pref., 1. v. 1975 (O. Tadauchi) ; 11 ♀ 21 ♂, Ochiai, Kuma, Ehime Pref., 15. iv. 1976 (O. Tadauchi) ; 1 ♀, Taisho, Kochi Pref., 16. iv. 1976 (O. Tadauchi). KYUSHU : 16 ♀, Kuroubaru, Chikuhō, Fukuoka Pref., 9. iv. 1975 (O. Tadauchi) ; 4 ♀, Seta, Ozu, Kumamoto Pref., 9. iv. 1976 (O. Tadauchi) ; 3 ♀, Yoshimatsu, Kagoshima Pref., 4. iv. 1976 (O. Tadauchi).

Other localities (*new records) : Hakodate (Hokkaido) ; *Moya, *Kogawara (Aomori Pref.) ; *Shibutami, *Kuriyagawa (Iwate Pref.) ; *Mt. Izumigatake (Miyagi Pref.) ; *Okitama, *Murayama (Yamagata Pref.) ; *Nakayama, *Nozawa (Fukushima Pref.) ; *Kuroiso, *Nasu (Tochigi Pref.) ; *Yorii, *Ochigawa (Saitama Pref.) ; *Nakano, Yoyogi, Egota (Tokyo) ; *Nakai, Kawasaki, Atsugi (Kanagawa Pref.) ; Koshoku, Ina (Nagano Pref.) ; Asahi, Kanamaru-Sekigawa, Kurokawa, Senami, Yamato, Kashiwazaki (Niigata Pref.) ; Gifu, *Sekigahara (Gifu Pref.) ; *Mt. Haku (Ishikawa Pref.) ; Mikuni, Fukui, Eiheiji, Katsuyama, Ohno, Izumi, Ikeda, Tsuruga (Fukui Pref.) ; *Ikaruga (Nara Pref.) ; *Fukuchiyama (Kyoto Pref.) ; *Sasayama (Hyogo Pref.) ; *Nagi (Tottori Pref.) ; *Mt. Dogo, *Hiba (Hiroshima Pref.) ; *Ato (Yamaguchi Pref.) ; Tachibanayama (*type locality*), Fukuoka, Futsukaichi, *Kurume, *Tanushimaru, Mt. Hikosan (Fukuoka Pref.) ; *Hokuzan-Dam (Saga Pref.) ; Kumamoto (Kumamoto Pref.) ; *Miyakonojo (Miyazaki Pref.) ; *Miyanojo (Kagoshima Pref.).

16. *Andrena* (*Andrena*) *babai* Tadauchi et Hirashima, new species

(Figs. 15, 30, 45, 88-94, Table 4)

Andrena (*Andrena*) sp. 1, Haneda, 1985, A list of Insects in Fukui Pref., **310**; Tadauchi & Hirashima, 1986, Trans. Essa Ent. Soc., (63) : 18.

This new species is similar to *Andrena* (**A.**) *benefica* Hirashima and the European *Andrena praecox* (Scopoli). But it is separable from *benefica* by the clypeus strongly convex with roughened, denser punctures, the hairs on the male clypeus silvery white, not intermixed with brownish or blackish ones, and the outer face of male propodeum with much fuscous hairs. It is also separable by *praecox* by the smaller size, the less tessellate clypeus with a shiny, smooth, impunctate, median space, the tibial scopa nearly whitish, the head and the thorax whitish, the hairs of the metasomal terga scanty, the male clypeus not flat, nearly smooth and shiny with larger, more distinct, sparser punctures, the male clypeus with whitish hairs not dense, the male metasomal sterna 2-5 each with a distinct, long, curled hair fringe.

So far as we know, the distribution of this species is restricted to Niigata, Ishikawa and Fukui Prefectures, central Honshu. It flies early in spring and visits the flowers of *Salix* sp.

Female : Length about 9.0 mm.

Integumental color : Flagellum brownish beneath ; posterior margins of metasomal terga reddish brown subhyaline.


Pubescence : Hairs on head and thorax short, not dense, those on head whitish, with brownish ones on paraocular area, frons, and genal area near eye and vertex ; facial fovea brownish above, yellowish below ; facial fovea with upper end occupying 3/4 of space between eye and ocellus ; hairs on thorax dull whitish mixed with pale brownish ; trochanteral and femoral floccus white ; tibial scopa whitish, narrowly brownish above ; hairs on metasomal terga scanty, white, on tergum 1 long, on terga 2-5 short ; caudal fimbria pale brownish.

Structure : Process of labrum transverse, slightly emarginate ; malar space about 1/4 times as long as base of mandible ; clypeus strongly convex, tessellate on basal half, smooth and shiny to very weakly tessellate on apical half with distinct, moderate-sized punctures ; clypeus with a median, impunctate space ; mesoscutum densely tessellate with indistinct, sparse punctures ; mesoscutellum smooth and shiny to weakly tessellate anteriorly, tessellate posteriorly ; propodeum densely tessellate, enclosure with basal half rugose, apical half tessellate ; mesepisternum shagreened with weak, indistinct punctures ; metasomal terga weakly tessellate, nearly impunctate, feebly shiny ; posterior depressions of terga not well indicated.

Male : Length 7.5-8.5 mm.

Integumental color : Mandibles reddened apically ; flagellum brownish beneath ; posterior margins of metasomal terga yellowish brown subhyaline.

Pubescence : Hairs on head not specially dense, long, silvery white, with fuscous ones on vertex, paraocular area, frons, and genal area near eye ; those on thorax long, whitish ; those on propodeum dull whitish, mixed with much fuscous ones ; those on legs whitish except for tarsi, which are covered with pale yellowish hairs ; those on metasomal terga 1-2 long, whitish, on terga 3-5 short, pale, on tergum 6 whitish to pale



Figs. 88-92. Genital capsule and associated structures of *Andrena (Andrena) babai*, new species, 88: dorsal view of genital capsule, 89: lateral view of the same, 90: ventral view of the same, 91 : 7th sternum, 92 : 8th sternum.

yellowish ; metasomal sterna 2-5 each with a fringe of long, curled, white hairs.

Structure : Mandible long, curved, falciform, with sharp apex ; basal projection of mandible slender ; malar space short, about 1/5 times as long as base of mandible ; clypeus slightly convex, narrowly tessellate basally, smooth and shiny apically with sparse, distinct punctures ; clypeus without an impunctate, median space ; genal area not distinctly angled behind ; flagellum 1 about one and half times as long as wide, slightly longer than 2, and nearly as long as 3 ; mesoscutum densely tessellate, sparsely and shallowly punctate ; mesoscutellum nearly smooth to very weakly tessellate anteriorly, tessellate and punctate posteriorly ; propodeum roughened, enclosure ill defined, basal half rugose, apical half tessellate ; mesepisternum roughened with indistinct punctures ; metasomal terga weakly tessellate, becoming smoother toward apical terga ; posterior depressions of terga broad, indistinct ; genitalia and associated structures as illustrated (Figs. 88-92).

Distribution : Japan (central Honshu (Niigata, Ishikawa, and Fukui Prefs.) and Sado Is.). (Fig. 93)




Fig. 93. A map showing the distribution of *Andrena* (*Andrena*) ***babai***, new species. A square indicates the type locality.

Floral record : Salicaceae : *Salix* sp. (8 ♀).

Flight records : Female : early April to early May. Male : late March to late April.

Type material : Holotype female (Type No. 2574, Kyushu Univ.), Senami, Niigata Pref., 9. iv. 1985 (K. Baba). Paratypes : same locality and collector as holotype : 3 ♂, 26. iv. 1984 ; 1 ♀, 9. iv. 1985 ; 1 ♀, 10. iv. 1985 ; 1 ♂, 13. iv. 1985 ; 1 ♀, Funado, Nakajo, Niigata Pref., 9. iv. 1985 (K. Baba) ; 5 ♀ 1 ♂, Hanami, Tsubame, Niigata Pref., 10. iv. 1985 (K. Baba) ; 7 ♀, Kusoozu, Nagaoka, Niigata Pref., 21. iv. 1985 (K. Baba) ; 1 ♀, Fukuoka, Kohchi-mura, Ishikawa Pref., 1. iv. 1982 (Y. Haneda) ; 1 ♂, Tatsunokuchi, Ishikawa, Pref., 2. iv. 1982 (Y. Haneda) ; Tomitsuka, Ohno, Fukui Pref. : 1 ♂, 13. iv. 1977 (Y. Haneda) ; 1 ♀, 16. iv. 1975 (Y. Haneda) ; Shingawara, Ohno, Fukui. Pref. : 1 ♀, 13. iv. 1972 (Y. Haneda) ; 1 ♀, 3. v. 1972 (Y. Haneda) ; Rokuroshi, Ohno, Fukui Pref. : 1 ♀, 3. v. 1976 (Y. Haneda) ; 1 ♂, 10. iv. 1977 (Y. Haneda) ; 1 ♀ 1 ♂, Fukui, Fukui Pref., 11. iv. 1981 (Y. Haneda) ; 1 ♂, Hokeiji, Ohno, Fukui Pref., 20. iv. 1975 (Y. Haneda) ; 2 ♂, Hiyoshi, Ohno, Fukui Pref., 30. iii. 1964 (T. Haneda). SADO IS. : Suizu, 10. v. 1980 (K. Baba).

Type depository : The type is preserved in the collection of the Entomological




Fig. 94. Flight records for the 16 Japanese species of *Andrena*(*Andrena*). A straight line indicates female and a dashed one male, K : Kyushu, S : Shikoku, Hn : Honshu, Hk : Hokkaido.

Table 4. Summary of floral records for the Japanese species of *Andrena* (*Andrena*). The numbers show as follows : the former indicates the number of plant species in the family the latter above in parentheses indicates the female number of *Andrena* collected ; and the below the male number collected.

Plant	Records of <i>Andrena</i> (Andrew)														
Family	<i>brevihirtiscopa</i>	<i>mali</i>	<i>mikado</i>	<i>ishiharai</i>	<i>nauai</i>	<i>esakii</i>	<i>sakagami</i>	<i>apicata aino</i>	<i>hondoica</i>	<i>longitibialis</i>	<i>maukensis</i>	<i>lapponica shirozui</i>	<i>aburama</i>	<i>benefica</i>	<i>babai</i>
Compositae	1 ⁽⁰⁾ ₍₂₎		1 ⁽¹⁾ ₍₀₎	4 ⁽⁴⁾ ₍₁₃₎	2 ⁽¹¹⁾ ₍₃₅₎		2 ⁽⁶⁶⁾ ₍₀₎		1 ⁽¹⁾ ₍₃₎	1 ⁽²⁾ ₍₀₎	2 ⁽⁷⁾ ₍₄₎	1 ⁽⁰⁾ ₍₁₎		2 ⁽⁴⁾ ₍₀₎	
Salicaceae	1 ⁽¹⁾ ₍₀₎		1 ⁽⁰⁾ ₍₁₎	3 ⁽³¹⁾ ₍₁₎	3 ⁽¹⁷⁾ ₍₅₎		1 ⁽⁵⁾ ₍₀₎	1 ⁽⁹⁾ ₍₂₎	5 ⁽¹⁷⁾ ₍₃₎	1 ⁽¹⁾ ₍₀₎	1 ⁽⁰⁾ ₍₁₎		1 ⁽¹⁾ ₍₀₎	2 ⁽³⁾ ₍₆₇₎	1 ⁽⁸⁾ ₍₀₎
Rosaceae	3 ⁽⁴⁾ ₍₀₎	1 ⁽²⁾ ₍₀₎			4 ⁽¹⁷⁾ ₍₀₎		1 ⁽²⁾ ₍₀₎			1 ⁽⁰⁾ ₍₁₎	1 ⁽⁰⁾ ₍₁₎	1 ⁽¹⁾ ₍₀₎	1 ⁽⁰⁾ ₍₁₎	5 ⁽⁹⁾ ₍₀₎	
Cruciferae	1 ⁽⁰⁾ ₍₅₎			1 ⁽⁰⁾ ₍₂₎	1 ⁽¹⁾ ₍₁₎					2 ⁽²⁾ ₍₂₎			1 ⁽⁴⁾ ₍₀₎	4 ⁽⁴⁰⁷⁾ ₍₃₄₎	
Aceraceae	1 ⁽⁴²⁾ ₍₆₎					1 ⁽²⁾ ₍₄₎								2 ⁽¹²⁾ ₍₁₎	
Ericaceae			5 ⁽⁵¹⁾ ₍₇₎				1 ⁽²⁾ ₍₀₎			6 ⁽¹⁷⁾ ₍₃₂₎				1 ⁽¹⁾ ₍₀₎	
Liliaceae							1 ⁽⁹⁾ ₍₀₎							1 ⁽¹⁾ ₍₀₎	
Caprifoliaceae				1 ⁽¹⁾ ₍₀₎			1 ⁽¹⁾ ₍₀₎								
Araliaceae				2 ⁽¹⁾ ₍₃₎			1 ⁽¹⁾ ₍₀₎								
Ranunculaceae				1 ⁽²⁾ ₍₁₎											
Umbelliferae				1 ⁽⁷⁾ ₍₁₂₎											
Polygonaceae				1 ⁽⁰⁾ ₍₁₎											
Saxifragaceae										1 ⁽¹⁾ ₍₀₎					
Scrophulariaceae									1 ⁽¹⁾ ₍₀₎						
Labiatae										2 ⁽²⁾ ₍₁₎					
Leguminosae											1 ⁽⁰⁾ ₍₁₎				
Aquifoliaceae											1 ⁽¹⁾ ₍₀₎				
Staphyleaceae							1 ⁽¹⁾ ₍₀₎								
Violaceae							1 ⁽¹⁾ ₍₀₎								
Totals	7 ⁽⁷⁴⁾ ₍₁₃₎	1 ⁽²⁾ ₍₀₎	7 ⁽⁵²⁾ ₍₈₎	14 ⁽⁴⁶⁾ ₍₃₃₎	10 ⁽⁴⁶⁾ ₍₄₁₎	1 ⁽²⁾ ₍₄₎	10 ⁽⁸⁸⁾ ₍₀₎	1 ⁽⁹⁾ ₍₂₎	7 ⁽¹⁹⁾ ₍₆₎	14 ⁽²⁵⁾ ₍₃₆₎	6 ⁽⁸⁾ ₍₇₎	2 ⁽¹⁾ ₍₁₎	3 ⁽⁵⁾ ₍₁₎	17 ⁽⁴³⁷⁾ ₍₁₀₂₎	1 ⁽⁸⁾ ₍₀₎

Laboratory, Faculty of Agriculture, Kyushu University.

This species is named for the honor of Dr. Kintaro Baba of Niigata Pref.

ACKNOWLEDGEMENTS

We are thankful to Dr. K. Baba of Niigata Pref., Mr. Y. Haneda of Ohno, Fukui Pref., Mr. M. Usui of Alberta, Canada, Dr. K. Yamauchi of Gifu Univ., Mr. E. Katayama of Tochigi Pref., Dr. Y. Maeta of Shimane Univ. for their gift or loan of valuable specimens. Many thanks are also due to the following entomologists for their gift or loan of specimens : Prof. S. F. Sakagami and Dr. H. Fukuda of Hokkaido Univ., Prof. M. Munakata of Hakodate College, Hokkaido Univ. of Educ., Mr. T. Nanbu of Saitama Pref., Prof. R. Ishikawa of Tokyo Metropolitan Univ., Prof. I. Togashi of Ishikawa Pref. College of Agr., Mr. K. Ôhara of Tokushima Pref., Prof. T. Saigusa, Assoc. Prof. M. T. Chûjô, Mr. K. Takeno and Mr. Y. Abe of Kyushu Univ., Mr. Y. Ono and Mr. H. Shimizu of Shimane Univ., Dr. K. Kanmiya and Mr. Y. Yoneda of Kurume Univ., Mr. Y. Harada of Okayama Pref. and Mr. S. Ide of Fukuoka Pref. We are also indebted to the collections of the Entomological Laboratory of Kyushu University and of Hokkaido National Agricultural Experiment Station. We are thankful to Assoc. Prof. S. Inoue of Kyushu Univ. Forests for his identification of some plants in Hokkaido.

REFERENCES

- Alfken, J. D. 1906 Die Gruppe der *Andrena varians* K. *Abh.naturw.ver. Bremen*, **18** : 129-131
- Alfken, J. D. 1914 Die Bienenfauna von Bremen. *Ibid.*, **22** : 1-220
- Alfken, J. D. 1929 Entomologische Ergebnisse der schwedischen Kamtschatka-Expedition 1920-1922. 21. Apidae, excl. Genus *Bombus*. *Arkiv f.Zoologi*, 20A (16) : 1-8
- Dalla Torre, C. G. de 1896 *Catalogus Hymenopterorum*, Vol. 10, Apidae (*Anthophila*). 643 pp
- Fukuda, H., S. F. Sakagami, K. Yamauchi, and T. Matsumura 1973 Biofaunistic survey of wild bees in Hamakoshimizu, Eastern Hokkaido. *Jap. J. Ecol.*, **23** : 160-170 (In Japanese)
- Haneda, Y. 1985 A list of Insects in Fukui Prefecture, Apoidea, In "A List of Insects in Fukui Prefecture", pp. 303-328 (In Japanese)
- Hedicke, H. 1933 Beiträge zur Systematik der Gattung *Andrena* F. (Hym. Apoid.). *Mitt.zool.Mus. Berlin*, 19: 199-220
- Hirashima, Y. 1953 The insect fauna of Mt. Ishizuchi and Omogo Valley, Iyo, Japan. Four new species of Apoidea (Hymenoptera). *Shikoku Ent.Soc.*, **3** : 132-138
- Hirashima, Y. 1957 Descriptions and records of bees of the genus *Andrena* from Eastern Asia III (Hymenoptera, Andrenidae). *Mushi*, **30** : 49-57
- Hirashima, Y. 1962 Systematic and biological studies of the family Andrenidae of Japan (Hymenoptera, Apoidea) Part 2. Systematics, 1. *J. Fac. Agr., Kyushu Univ.*, **12**: 117-154
- Hirashima, Y. 1966 Ditto. Part 2. Systematics, 7. *Ibid.*, **14**: 89-131
- Hirashima, Y. and O. Tadauchi 1979 A list of the genus *Andrena* (Hymenoptera, Andrenidae) of Niigata Prefecture. In "Insects in Niigata Prefecture", pp. 1-32 (In Japanese)
- Ikudome, S. 1980 Flower preference of twenty species of Andrenid bees in Kochi Plain. *Bull. Kugoshima Worn. J. Coll.*, (15) : 311-326 (In Japanese)
- Ikudome, S. 1983 Floral records of wild bees in Tosayama-mura, Kochi Prefecture, Shikoku, Japan (Hymenoptera, Apoidea). *Zbid.*, (18) : 137-148
- Kim, M. and C. Kim 1983 On the nine unrecorded Andrenidae from Korea (Hymenoptera : Apoidea).

Kor. J. Ent., 13: 5-9

- LaBerge, W. E. 1964 Prodrum of American bees of the genus *Andrena* (Hymenoptera, Apoidea). *Bull. Univ. Neb. St. Mus.*, 4: 279-316
- Lanham, L. J. N. 1949 A subgeneric classification of the new world bees of the genus *Andrena*. *Univ. Calif. Publ. Ent.*, 8: 183-238
- Osichnynk, G. Z. 1977 Bees-Andrenidae. *Fauna Ukraini*, Kiev, 12 : 1-213 (In Ukrainian)
- Matsumura, T. and M. Munakata 1969 Relative abundance, phenology and flower preference of andrenid bees at Hakodateyama, Northern Japan (Hymenoptera, Apoidea). *J. Fac. Sci., Hokkaido Univ., Ser. VI, Zool.*, 17: 106-126
- Matsuura, M., S. F. Sakagami, and H. Fukuda 1974 A wild bee survey in Kibi (Wakayama Pref.), Southern Japan. *Ibid.*, 19: 422-437
- Miyamoto, S. 1960 Flower-visiting habits of fourteen species of Andrenid bees (Biological studies on Japanese bees XIV). *Kontyû*, 28: 65-86 (In Japanese)
- Munakata, M. 1971 Relative abundance, phenology and flower preference of Andrenid bees at Akagawa near Hakodate, Northern Japan (Hymenoptera, Apoidea). *J. Hokkaido Univ. Educ., Sect. II, B*, 22: 26-39
- Munakata, M. and K. Kobayashi 1983 A wild bee survey at Kamekawa, Kikonai, southern Hokkaido. *Seibutsu-kyozai, Hakodate*, (18) : 15-25 (In Japanese)
- Munakata, M. and M. Kudo 1981 A wild bee survey at the Rishiri Is. in 1968 and 1970. *Ibid.*, (16) : 122-130 (In Japanese)
- Sakagami, S. F. and H. Fukuda 1972 Autumn bee fauna in Hokkaido University Uryu and Nakagawa Experiment Forests. *Res. Bull. Coll. Exp. Forests, Coll. Agr., Hokkaido Univ.*, 29: 1-24 (In Japanese)
- Sakagami, S. F. and H. Fukuda 1973 Wild bee survey at the Campus of Hokkaido University. *J. Fac. Sci., Hokkaido Univ., Ser. VI., Zool.*, 19: 190-250
- Sakagami, S. F. et R. Ishikawa 1969 Note préliminaire sur la répartition géographique des bourdons japonais, avec descriptions et remarques sur quelques formes nouvelles ou peu connues. *Ibid.*, 11: 152-196.
- Sakagami, S. F. and T. Matsumura 1967 Relative abundance, phenology and flower preference of andrenid bees in Sapporo, North Japan (Hymenoptera, Apoidea). *Jap. J. Ecol.*, 17 : 237-250
- Schmiedeknecht, O. 1930 *Die Hymenopteren nord-und mitteleuropas. Zweite Auflage.* 1062 pp
- Stoeckert, F. K. 1933 Die Bienen Frankens (Hym. Apid.). *Deut. ent. Zeit.*, 1932, Beiheft : 1-294
- Stoeckert, F. K. 1954 Fauna Apoideorum Germaniae. *Abh. Bayerischen Akad. Wissensch.*, Neue Folge, (65) : 1-87
- Strand, E. and K. Yasumatsu 1938 A new *Andrena*-species from Japan (Hymenoptera : Apoidea). *Mushi*, 11: 67-69
- Svensson, G. and J. Tengö 1976 *Andrena* (Hym., Apoidea) on the Island of Öland, Sweden, with key to species I Subgenus *Andrena* (s. s.) Fabricius. *Ent. Tidskr.*, 97 : 78-89
- Tadauchi, O. 1982 A numerical taxonomic study of the genus *Andrena* (Hymenoptera, Andrenidae) of Japan. *J. Fuc. Agr., Kyushu Univ.*, 26: 169-191
- Tadauchi, O. and Y. Hirashima 1986 A list of the superfamily Apoidea (Hymenoptera) of Niigata Prefecture I. Andrenidae. *Trans. Essa Ent. Soc.*, (63) : 15-30. (In Japanese)
- Tadauchi, O. and Y. Hirashima 1986 A list of the family Andrenidae (Hymenoptera, Apoidea) from Akita, Yamagata, and Toyama Prefectures collected by Dr. K. Baba. *Ibid.*, (63): 31-32. (In Japanese)
- Warncke, K. 1967 Beitrag zur klärung palaarktischer *Andrena*-Arten. *Eos*, 43 : 171-318
- Warncke, K. 1968 Die Untergattungen der westpalaarktischen Bienengattung *Andrena* F. *Mem. Est. Mus. Zool. Univ. Coimbra*, (307): 5-111
- Usui, M., Y. Nishijima, H. Fukuda, and S. F. Sakagami 1976 A wild bee survey in Obihiro, Eastern Hokkaido. *Res. Bull. Obihiro Univ.*, 10: 225-251

- Yamauchi, K., Y. Morimoto, K. Watanabe, S. F. Sakagami, and T. Matsumura 1982 A list of insects in Gifu Prefecture, Apoidea, *In* "Insects in Gifu Prefecture", pp. 415-430 (In Japanese)
- Yamauchi, K., Y. Murakumo, M. Ogura, and S. F. Sakagami 1974 Biofaunistic survey of wild bees in Minami (Gifu Prefecture), Central Japan. *Sci. Rep. Fac. Educ., Gifu Univ. (Nat. Sci.)*, 5: 220-232 (In Japanese)
- Yasumatsu, K. 1940-41 A list of the Far Eastern species of the genus *Andrena* (Hym., Apoidea). *Peking Nat. Hist. Bull.*, 15: 273-284