A Revision of the Genus *Tetragnatha* Latreille (Araneae, Tetragnathidae) of Asia, Part I

Okuma, Chiyoko
Entomological Laboratory, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23876>

出版情報：九州大学大学院農学研究院紀要. 32 (3/4), pp.165-181, 1988-03. Kyushu University
バージョン：
権利関係：

A Revision of the Genus *Tetragnatha* Latreille (Araneae, Tetragnathidae) of Asia, Part I*

Chiyoko Okuma

Entomological Laboratory, Faculty of Agriculture,
Kyushu University, Fukuoka 812

(Received September 24, 1987)

The spiders of Asian *Tetragnatha* are revised based on the specimens from Nepal, Bangladesh, Sri Lanka, Thailand, Malaysia, Indonesia, Philippines, Hong Kong, Taiwan, Korea and Japan. Twenty-nine species are recognized, which are divided into two major species groups, A and B. This paper (Part I) reports the species of the major group A. New records of distribution are as follows : *T. javana* from Nepal and Bangladesh and *T. ceylonica* from Philippines.

This paper reports the result of my systematic study on the genus *Tetragnath* Latreille of Asia. Twenty nine species in all, including nine species already described by me, are recognized from Nepal, Bangladesh, Sri Lanka, Thailand, Malaysia, Indonesia, Philippines, Hong Kong, Taiwan, Korea and Japan. Illustrations of the important characters of all the species are given, and keys are provided for the species groups and species. *Tetragnath* may be divided into two major species groups, which are tentatively called A and B groups. These groups may eventually be recognized as subgenera. The present paper (Part I) included 9 species which belong to the major species group A.

For abbreviations of the chelicer al aremature see Okuma (1987).

Before going further, I wish to express my hearty thanks to Prof. Y. Hirashima of Kyushu University for helpful suggestions and encouragement throughout the course of my study. I am grateful for my colleagues listed below who offered me valuable specimens for the present study : Mr. Joseph Koh (Singapore), Dr. J. Martens (Univ. Mainz, Germany), Dr. JoAnn M. Tenorio (Bishop Mus., Honolulu), prof. Yau-I Chu (National Taiwan Univ., Taiwan), Mr. S. Asama (Chiba Pref.), Mr. Y. Chikuni (Nagano Pref.), Dr. T. Goto (Kyushu Univ.), Dr. T. Kamura (Kyoto Pref. Univ.), Mr. H. Maki hara (Forestry and Forest Products Res. Ins., Tsukuba), Dr. Y. Nishikawa (Ohtemon Gakuin Univ., Osaka), Mr. K. Ohara (Tokushima Pref.), Dr. H. Ono (Nat. Sci. Mus., Tokyo), Mr. T. Sasaki (Univ. of the Ryukyus), Mr. A. Tanigawa (Yokohama City), Dr. K. Umeya (National Agr. Res. Center, Tsukuba) and Dr. T. Yaginuma (Ohtemon Gakuin Univ.). And I am also grateful to Assoc. Prof. K. Morimoto, Dr. K. Ogata and Dr. O. Tadauchi of Kyushu University for their kindness in many ways.

* Contribution from the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka (Ser. 3, No. 256).

Grouping of the species

The species of *Tetragnatha* may be divided into two major groups, A and B, according to the conductor of male palpus (Levi, 1981). In the group A, it is plicate laterally, while in the group B, it is entire. The major species groups may be divided further into (minor) species-groups. The group A includes the *bituberculata*, *caudicula*, *extensa*, *praedonia* and *shinanoensis* species-groups. The group B includes the *cylindrica*, *lauta*, *mandibulata*, *maxillosa*, *rubriventris*, *serra*, *squamata* and *striata* species-groups.

KEY TO THE SPECIES-GROUPS OF ASIAN TETRAGNATHA
BASED ON THE MALE

- 1 Conductor with pleats laterally 2 (group A)
- Conductor without pleats laterally 6 (group B)
- 2 Lateral eyes more separated than median eyes ; abdomen with a tail overhanging spinnerets *caudicula*-group
- Lateral eyes as close as or closer than median eyes ; abdomen not as above 3
- 3 Posterior row of eyes wider than anterior row of eyes 4
- Two rows of eyes equal or subequal in width 5
- 4 Lateral eyes closer than median eyes and situated on a tubercle *bituberculata*-group
- Lateral eyes slightly closer than median eyes and situated on each tubercle *praedonia*-group
- 5 Conductor bifurcated at tip *shinanoensis*-group
- Conductor not bifurcated at tip *extensa*-group
- 6 Fang with an outer cusp near base ; (Gl) robust *squamata*-group
- Fang usually without outer cusp near base 7
- 7 Posterior row of eyes evidently narrower than anterior row of eyes 8
- Posterior row of eyes not narrower than anterior row of eyes 9
- 8 Chelicera as long as or longer than carapace *rubriventris*-group
- Chelicera shorter than carapace ; lateral eyes more separated than median eyes *striata*-group
- 9 Abdomen more than 6 times as long as broad *cylindrica*-group
- Abdomen about 3-5 times as long as broad 10
- 10 Tip of (a) simply pointed 11
- Tip of (a) bifurcated 13
- 11 Legs entirely without spines ; generally weak surface *lauta*-group
- Legs with many spines 12
- 12 Fang geniculated *serra*-group
- Fang not as above *mandibulata*-group
- 13 Chelicera with three strong apical teeth 14
- Chelicera not as above *maxillosa*-group
- 14 Chelicera without (T) *nitens*-group
- Chelicera with (T) *josephi*-group

The caudicula-group

Diagnosis. Posterior row of eyes evidently narrower than anterior row of eyes ; lateral eyes more separated than median eyes ; abdomen with a tail overhanging spinnerets.

KEY TO THE SPECIES

- 1 Spinnerets situated beyond the middle of abdomen *caudicula*
 — Spinnerets situated about the middle of abdomen *javana*

***Tetragnatha caudicula* (Karsch, 1879)**

(Fig. 1, Table 1)

Eugnatha caudicula Karsch, 1879, Verh. Nat. Ver. Rheinl. Westf., 36 : 66 ; Simon, 1881, Arachn. de France, 5 (1) : 7.

Tetragnatha caudicula Boesenberg und Strand, 1906, Abh. Senckenbg. Ges., 30 (1-2) : 179 ; Paik, 1957, Korean J. Biol., 2 : 43 ; Yaginuma, 1960, Spiders of Japan in Colour : 74.

Fig. 1. *Tetragnatha caudicula* (Karsch). A : Left chelicera of male, upper view. B : Ditto, inner view. C : Ditto, upper inner view. D : Paracymbium of male. E and F : Conductor and embolus of male. G : Left chelicera of female, upper view. H : Ditto, inner view. I : Ditto, lateral view. J : Genital fold of female. K : Eye group of female. L : Abdomen of female, lateral view.

Table 1. *Tetragnatha caudicula* (Karsch). Relative lengths of total legs and femora.

Locality	n	length of mean±s.d.	1st legs r a n k	1st legs ratio-	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
Hokkaido	♂ 1	21.90		100	62	28	63
Kanagawa	♂ 2	21.78	20.30-23.25		60	27	62
Tottori	♂ 1	26.50			61	28	62
Tokushima	♂ 4	21.31f4.08	16.40-25.30		60 (0.998)	27 (0.999)	61 (0.999)
Ryukyu Is.	♂ 5	22.04f2.77	19.15-26.65		59 (0.996)	27 (0.980)	62 (0.999)
KOREA	♂ 3	24.1af1.33	22.70-25.25		61 (0.992)	27 (0.969)	61 (0.988)
Hokkaido	♀ 2	18.78	15.25-22.30		61	29	64
Kanagawa	♀ 2	22.68	20.00-25.35		61	28	62
Tokushima	♀ 4	21.55f2.20	18.85-23.80		60 (0.986)	28 (0.883)	64 (0.980)
Ryukyu Is.	♀ 2	20.18	19.50-20.85		59	28	67
KOREA	♀ 9	21.67f2.02	18.50-24.45		61 (0.978)	28 (0.980)	63 (0.977)
(Femora)							
Hokkaido	♂ 1	6.30			67	34	77
Kanagawa	♂ 2	6.10	5.80- 6.40		67	34	75
Tottori	♂ 1	7.40			68	35	76
Tokushima	♂ 4	5.90f1.07	4.60- 7.00		66 (0.999)	34 (0.999)	75 (0.998)
Ryukyu Is.	♂ 5	6.23f0.79	5.35- 7.50		65 (0.993)	34 (0.957)	75 (0.999)
KOREA	♂ 3	6.72f0.32	6.35- 6.90		68 (0.993)	35 (1)	75 (0.945)
Hokkaido	♀ 2	5.38f1.52	4.30- 6.45		65	34	78
Kanagawa	♀ 2	6.50f1.13	5.70- 7.30		65	34	75
Tokushima	♀ 4	6.20±0.72	5.30- 6.95		65 (0.990)	34 (0.885)	77 (0.991)
Ryukyu Is.	♀ 2	5.85f0.21	5.70- 6.00		65	34	81
KOREA	♀ 9	6.21f0.62	5.20- 7.05		67 (0.978)	34 (0.974)	77 (0.975)

n= Number of specimens.
r = coefficient of correlation.

Male. Body length, exclusive of chelicerae 6.0-11.0 mm ; carapacial length 1.6-3.0 mm, width LO-1.3 mm ; abdominal length 4.5-8.0 mm, width 0.6-1.0 mm ; cheliceral length 1.1-2.0 mm.

Eyes. Anterior row of eyes occupying full width of head ; posterior row of eyes evidently narrower than anterior one ; viewed from above, anterior row nearly straight, posterior row somewhat strongly recurved, so that lateral eyes more separated than median eyes.

Chelicerae. Basal segment about 0.7 times as long as carapace ; (a) large, bifurcated at tip ; (t) and (sl) somewhat directed backward ; (T) large, (rsu) about 5 in number ; (AXI) and (GI) present, (rsl) about 6 in number ; fang unarmed.

Legs. Leg formula 1. 4. 2.3 ; all legs with spines, first femur with about 10 spines ; relative lengths of legs as shown in Table 1.

Palpi. Paracymbium bluntly rounded at tip ; conductor twisted at the middle and terminating in a characteristic form as shown in Fig. 1, E and F.

Abdomen. Long, inclusive of the tail, about 6-8 times as long as broad.

Color in alcohol. Legs, carapace and chelicerae generally light brown ; maxillae, lip and sternum somewhat dusky brown ; abdomen covered with dusky silvery spangles dorsally, somewhat dusky brown ventrally.

Female. Body length, exclusive of chelicerae 8.0 - 15.0 mm ; carapacial length 2.3-3.0 mm, width 1.1-1.8 mm ; abdominal length 5.8-12.0 mm, width 1.3-2.0 mm ; cheliceral length 1.0-1.8 mm.

Eyes. Nearly as in male.

Chelicerae. Basal segment 0.4-0.6 times as long as carapace, outside near apex and middle of dorsum somewhat swollen and with a tubercle each ; (AXu) and (Gu) present, (rsu) about 5 in number ; (GI) present, (rsl) about 5 in number ; fang with (EX) near base.

Legs. Nearly as in male ; relative lengths of legs as shown in Table. 1.

Abdomen. Long, about 4.5-6.0 times as long as broad.

Color in alcohol. Similar to male.

Specimens examined : KOREA : 3♂, 9♀, Paddy field, Suweon, 18. VII - 4. VIII. 1975, N. Hokyō et al. JAPAN : 48 specimens from Hokkaido (Ashoro), Ibaragi (Tsukuba), Tokyo (Hachioji), Tochigi (Nikko, Chuzenji-kohan), Kanagawa (Hiratsuka), Tottori (Mt. Daisen), Tokushima (Kamiakui-cho), Ohita (Odanoike, Yuhuin-cho), Amami-oshima Is. (Mt. Yuwan), Ishigaki Is. (Arakawa), Iriomote Is., Minamidaito-zima.

Distribution : Japan and Korea.

***Tetragnatha javana* (Thorell, 1890)**

(Fig. 2, Table 2)

Eucta javana Thorell, 1890, Ann. Mus. Civ. Genova, 28 : 236.

Tetragnatha (Eucta) javana : Okuma, 1968, Mushi, 42 : 100; Okuma, 1970, Mushi, 44 : 73.

Fig. 2. *Tetragnatha javana* (Thorell). A : Left chelicera of male, upper view. B : Ditto, lower view. C : Ditto, inner view. D : Ditto, lateral view. E and F : Conductor and embolus of male. G : Paracymbium of male. H : Eye group of female. I : Genital fold of female. J : Left chelicera of female, lateral view. K : Ditto, inner view. L : Abdomen of female, lateral view. (After Okuma, 1984).

Table 2. *Tetragnatha javana* (Thorell). Relative lengths of total legs and femora.

Locality	n	length of 1st legs mean±s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
JAPAN	♂ 3	24.63±2.29	19.65-27.80	100	57 (0.969)	25 (0.918)	58 (0.959)
TAIWAN	♂ 10	20.07±1.75	18.05-21.10		56 (0.993)	24 (0.992)	56 (0.998)
THAILAND		24.63±2.80	19.90-30.10		54 (0.997)	23 (0.991)	55 (0.988)
JAPAN	♀ 10	24.95±2.14	20.75-27.55		55 (0.943)	25 (0.863)	57 (0.941)
TAIWAN	♀ 1	23.20			54	24	54
THAILAND	♀ 10	26.07±1.77	22.20-28.70		53 (0.960)	23 (0.964)	54 (0.963)
(Femora)							
JAPAN	♂ 10	7.01±0.69	5.55- 7.90		64 (0.970)	32 (0.897)	71 (0.958)
TAIWAN	♂ 1	5.63±0.55	5.00- 5.90		62 (0.996)	30 (0.996)	69 (0.989)
THAILAND	♂ 10	6.90±0.68	5.65- 7.85		61 (0.990)	30 (0.968)	67 (0.924)
JAPAN	♀ 10	7.32±0.64	6.00- 8.10		61 (0.958)	31 (0.894)	70 (0.925)
TAIWAN	♀ 1	6.70			60	31	66
THAILAND	♀ 10	7.57±0.51	6.40-8.35		59 (0.970)	29 (0.952)	66 (0.943)

Tetragnatha javana: Okuma, 1984, Esakia, (22): 88.

Additional specimens: NEPAL: 1♂, Chauni 1300-1400, Kathmandu Valley, end of July, 1973, J. Martens. BANGLADESH: 126, 13♀, Central Bangladesh, VIII-IX. 1978, H. D. Catring; 2♂, 3♀, Mymensingh, 12. IV. 1987, Y. Hirashima. JAPAN: 40 specimens from Iriomote Is. (Sonai), Ishigaki Is. (Kabira and Nakura), Okinawa (Hachi), Amami-Oshima (Yakkachi, Naze and Taira).

Distribution: Africa, Nepal (new record), Bangladesh (new record), SE Asia and Japan (the Ryukyus up to Amami-Oshima).

Note: This species is common in paddy fields of SE Asia; body length of male 8.0-12.0 mm, female 10.0-18.0 mm.

The bituberculata-group

Diagnosis. Posterior row of eyes wider than anterior row of eyes; lateral eyes closer than median eyes, and situated on a tubercle; anterior median eyes closer than posterior median eyes; male chelicerae with (a), (t), (sl) and (T); female chelicera with (e) between (Gu) and (T).

Tetragnatha bituberculata, an Australasian species, does not occur in Asia.

KEY TO THE SPECIES

- 1 Male paracymbium with tip pointed, and subapical margin serrate *ceylonica*
- Male paracymbium not as above 2
- 2 Male paracymbium with tip rounded *makiharai*
- Male paracymbium with tip rather squared *boninensis*

Tetragnatha ceylonica Cambridge, 1869

(Fig. 3, Table 3)

Tetragnatha ceylonica Cambridge, 1869, Journ. Linn. Soc. Zool., 10: 394; Okuma, 1987, Esakia, (25): 48.

Additional specimens: PHILIPPINES: 1♀, Mindanao, 12. VIII. 1985, Y. Nishikawa (NSMT-Ar. 1127); 1♂, 1♀, Mindanao, 13. VIII. 1985, Y. Nishikawa (NSMT-Ar. 1128); 1♀, Palawan Is., 1. IX. 1985, M. Sakai (NSMT-Ar. 1130); 1 young, Palawan Is.,

Fig. 3. *Tetragnatha ceylonica* Cambridge. A : Left chelicera of male, upper view. B : Ditto, lateral view. C : Ditto, lower view. D : Left palpus of male. E : Paracymbium of male. F : Conductor and embolus of male. G : Left chelicera of female, upper view. H : Ditto, lower inner view. I : Eye group of female. J : Genital fold of female. K : Abdomen of the young of very early stage, lateral view. L : Abdomen of female, lateral view.

Table 3. *Tetragnatha ceylonica* Cambridge. Relative lengths of total legs and femora.

Locality	n	length of mean±s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
JAPAN	♂ 5	23.66f2.35	20.85-26.10	100	58 (0.9921)	25 (0.939)	55 (0.977)
TAIWAN	♂ 8	26.15±4.34	18.95-31.95		57 (0.992)	24 (0.988)	55 (0.993)
THAILAND	♂ 1	21.90			54	23	53
JAPAN	♀ 11	22.18f1.71	19.70-25.9s		58 (0.976)	26 (0.949)	57 (0.976)
TAIWAN	♀ 10	23.00f2.17	17.90-25.1s		57 (0.988)	25 (0.969)	57 (0.983)
THAILAND	♀ 1	23.20			59	26	58
(Femora)							
JAPAN	♂ 5	6.76f0.68	5.95- 7.45		65 (0.991)	30 (0.924)	65 (0.998)
TAIWAN	♂ 8	7.44±1.23	5.50- 9.30		64 (0.996)	30 (0.993)	64 (0.997)
THAILAND	♂ 1	6.05			62	28	65
JAPAN	♀ 11	6.45f0.49	5.65- 7.40		65 (0.970)	30 (0.888)	68 (0.966)
TAIWAN	♀ 10	6.66±0.61	5.35- 7.30		64 (0.936)	30 (0.962)	68 (0.991)
THAILAND	♀ 1	6.85			66	31	69

26. VIII. 1985, M. Tomokuni (NSMT-Ar. 1129). THAILAND: 1♂, Kaoyai National Park, 26. XII. 1977, T. Irie ; 5♀, Chang mai, 11. XII. 1985, T. Sasaki ; 2♀, Khoscido, Chanthaburi, 20. XII. 1985, T. Sasaki ; 1 young, Kaoyai National Park, 21. VI. 1986, T. Sasaki ; 1♀, Bangkok, 13. II. 1986, T. Sasaki. TAIWAN : 32 specimens from (Tachi, Ilan Hsien), (Lienhuachih, Nanto Hsien), (Wulai, Taipei Hsien), (Hungye-hwenchuan, Hualien Hsien), (Chihpenwanchuan, Taitung Hsien), (Shihtoushan, Hsin-chu Hsien), (Lanhsu, Taitung Hsien). JAPAN : Yonaguni, Ishigaki Is. (Mt. Omoto), Iriomote Is. (Funaura, Mt. Goza and Ohara).

Distribution : New Guinea, Thailand, Philippines (new record), Taiwan and Japan

(the Ryukyus up to Okinawa).
Body length : male 6.3-11.0 mm, female 7.5-12.0 mm.

Tetragnatha makiharai Okuma, 1977
(Fig. 4, Table 4)

Tetragnathu makiharai Okuma, 1977, Acta Arachnol., 27 : 27.
Additional specimens : JAPAN : 22 specimens from Okinawa (Nago, Oku, Hen-

Fig. 4. *Tetragnatha makiharai* Okuma. A : Left chelicera of male, upper view. B : Ditto, lateral view. C : Ditto, lower view. D : Conductor and embolus of male. E : Paracymbium of male, F : Left palpus of male. G : Left chelicera of female, upper view. H : Ditto, lower view, I : Abdomen of male, lateral view. J : Eye group of male. K : Genital fold of female.

Table 4. *Tetragnatha makiharai* Okuma . Relative lengths of total legs and femora.

Locality	n	length of mean±s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
Tsushima Is.	♂ 4	24.99±1.54	23.54-27.15	100	55 (0.993)	24 (0.981)	51 (0.981)
Amami-oshima	♂ 4	23.11±1.64	21.20-25.20		56 (0.923)	25 (0.985)	54 (0.919)
Okinawa Is.	♂ 14	25.77±2.27	21.70-28.75		56 (0.982)	24 (0.965)	53 (0.987)
Kuma Is.	♂ 2	27.58	27.20-27.95		57	25	54
Tsushima Is.	♀ 7	23.84±1.44	21.65-26.10		55 (0.956)	24 (0.857)	52 (0.957)
Amami-oshima	♀ 5	23.03±0.78	22.20-23.90		56 (0.887)	26 (0.630)	54 (0.783)
Okinawa Is.	♀ 22	23.75±1.74	20.40-26.70		55 (0.885)	25 (0.833)	54 (0.892)
Kume Is.	♀ 2	22.75	21.85-23.65		53	23	52
(Femora)							
Tsushima Is.	♂ 4	6.96±0.46	6.50- 7.60		61 (0.971)	28 (0.973)	60 (0.977)
Amami-oshima	♂ 4	6.48±0.41	6.00- 7.00		64 (0.865)	30 (0.997)	64 (0.965)
Okinawa Is.	♂ 14	7.25±0.65	6.15- 8.20		62 (0.977)	29 (0.915)	63 (0.956)
Kume Is.	♀ 2	6.84±0.839	6.20- 7.85		63 (0.952)	28 (0.726)	62 (0.978)
Amami-oshima	♀ 5	6.70±0.23	6.50- 7.00		63 (0.944)	30 (0.899)	65 (0.812)
Okinawa Is.	♀ 22	6.98±0.46	6.15- 7.80		61 (0.981)	28 (0.860)	64 (0.898)
Kume Is.	♀ 2	6.60	6.40- 6.80		59	27	63

nogi and Yonaha dake), Kumezima Is., Tsushima Is. (Mt. Oboshi yama, Minemura).
Distribution : Japan (the Ryukyus and Tsushima Is).
Body length : male 6.5-8.6 mm, female 8.2-11.0 mm.

***Tetragnatha boninensis* Okuma, 1981**
(Fig. 5, Table 5)

Tetragnatha boninensis Okuma, 1981, Esakia, (17) : 141.
Additional specimens : 76, 3♀, Hahazima, Ogasawara Is., 1-7. VII. 1986, H. Makihara.
Distribution : Japan (the Ogasawara Is.).
Body length : male 5.2-6.6 mm, female 6.2-6.5 mm.

Fig. 5. *Tetragnatha boninensis* Okuma. A : Left chelicera of male, upper view. B : Ditto, lateral view. C : Ditto, lower view. D : Distal portion of conductor and embolus of male. E : Paracymbium of male. F : Left chelicera of female, upper view. G : Abdomen of female, lateral view. H : Eye group of female. (After Okuma, 1981).

Table 5. *Tetragnatha bobinensis* Okuma. Relative lengths of total legs and femora.

Locality	n	length Of means.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
ogasawara Is. ♂	4	16.85t2.27	14.65-19.65	100	63 (0.995)	29 (0.972)	57 (0.994)
ogasawara Is. ♀	4	10.43f0.42	10.00-10.95		63 (0.907)	31 (0.984)	60 (0.849)
(Femora)							
ogasawara Is. ♂	4	4.76f0.69	4.10-5.60		70 (1)	35 (0.958)	69 (0.999)
Ogasawara Is. ♀	4	2.95f0.13	2.00-3.10		67 (0.894)	37 (0.775)	71 co.9231

The praedonia-group

Diagnosis. Posterior row of eyes slightly wider than anterior row of eyes, lateral eyes somewhat closer than median eyes ; fang with an outer cusp near base in both sexes.

***Tetragnatha praedonia* L. Koch, 1878**

(Fig. 6, Table 6)

Tetragnatha praedonia L. Koch, 1878, Verh. zool. bot. Ges. Wien, 27 : 744 ; Boesenberg und Strand, 1906, Abh. Senckenbg. Ges., 30 (1-2) : 177 ; Saito, 1933, Trans. Sapporo Nat. Hist. Soc., 13 (1) : 47 ; Saito, 1934, Journ. Agr. Hokkaido Univ., 33 (5) : 335 ; Saito, 1939, Saito Ho-on Kai Mus. Res. Bull., 18 : 58 ; Yaginuma, 1960, Spiders of Japan in Colour, : 73.

Male. Body length, exclusive of chelicerae 4.8-12.0 mm ; carapacial length 1.8-4.3 mm, width 1.1-2.5 mm ; abdominal length 3.0-7.7 mm, width 1.0-2.2 mm ; cheliceral length 2.0-4.7 mm.

Eyes. Anterior row of eyes occupying full width of head, posterior row of eyes slightly wider than anterior one ; viewed from above, anterior row somewhat strongly recurved, posterior row somewhat gently recurved, so that lateral eyes slightly closer than median eyes.

Chelicerae. Basal segment as long as or slightly longer than carapace, (a) bifid at tip ; (t) and (sl) present, (T) large, (rsu) about 8-10 in number ; (AXI) and (GI) present, an additional blunt tubercle present on outer side of posterior margin of fang furrow, (rsl) 5-9 in number ; fang armed with (EX) near base.

Legs. Leg formula 1. 2. 4. 3 ; all legs with many spines, first femur with 13-21

Fig. 6. *Tetragnatha praedonia* L. Koch. A: Left chelicera of male, upper view. B: Ditto, lower view. C : Ditto lateral view. D : Paracymbium of male. E and F : Distal portion of conductor and embolus of male. G : Left palpus of male. H : Left chelicera of female, upper view. I : Ditto, lower view. J : Eye group of male. K : Genital fold of female. L : Abdomen of female, lateral view.

Table 6. *Tetragnatha praedonia* L. Koch. Relative lengths of total legs and femora.

Locality	n	length of mean±s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
JAPAN	♂ 15	31.15f3.63	25.15-36.45	100	63 (0.989)	29 (0.977)	59 (0.992)
TAIWAN	♂ 8	27.00f5.54	15.20-34.30		62 (0.991)	28 (0.985)	59 (0.997)
CHINA	♂ 1	30.73f3.22	27.05-33.00		62 (1)	28 (0.996)	59 (0.9961)
JAPAN	♀ 3	31.85f2.71	26.23-35.30		62 (0.986)	28 (0.947)	57 (0.972)
TAIWAN	♀ 7	28.66f2.75	24.35-33.10		60 (0.978)	27 (0.864)	56 (0.983)
CHINA	♀ 8	27.87f3.19	22.03-31.98		60 (0.992)	27 (0.967)	54 (0.735)
(Femora)							
JAPAN	♂ 15	8.63f1.04	7.15-10.25		70 (0.991)	36 (0.989)	70 (0.993)
TAIWAN	♂ 8	7.53f1.54	4.20- 9.40		70 (0.998)	35 (0.993)	70 (0.998)
CHINA	♂ 3	8.47±0.84	7.50- 9.00		69 (0.909)	35 (0.992)	70 (0.992)
JAPAN	♀ 15	9.10f0.81	7.30-10.10		68 (0.985)	35 (0.952)	69 (0.974)
TAIWAN	♀ 7	8.09f0.74	6.95- 9.30		67 (0.958)	34 (0.892)	69 (0.985)
CHINA	♀ 8	7.99f1.00	6.15- 9.20		66 (0.993)	34 (0.972)	68 (0.997)

spines ; relative lengths of legs as shown in Table 6.

Palpi. Paracymbium bifid at tip ; conductor and embolus as shown in Fig. 6, E and G.

Abdomen. About 3-4 times as long as broad ; distal end of abdomen slightly overhanging spinnerets.

Color in alcohol. Generally light brown to blackish brown ; abdomen with indistinct several pairs of dark patches dorsally, and with a central dark gray strip ventrally.

Female. Body length, exclusive of chelicerae 8.2-14 mm ; carapacial length 2.6-3.8 mm, width 1.7-2.4 mm ; abdominal length 5.5-10.5 mm, width 1.7-3.8 mm ; cheliceral length 2.0-3.3 mm.

Eyes. Nearly as in male.

Chelicerae. Basal segment slightly shorter than carapace ;(AXu) present, (Gu) somewhat robust, (e) present, (T) not so large, (rsu) 6-8 in number ;(AXI) present near (Gl),(Gl) and (AXI) nearly equal in size, (L2) very swollen at base, (rsI) 7-10 in number ; fang robust, armed with (EX) near base.

Legs. Leg formula 1. 2-4. 3 ; all legs with many spines, first femur with 11-20 spines ; relative lengths of legs as shown in Table 6.

Abdomen. About 3-4 times as long as broad; distal end of abdomen slightly overhanging spinnerets ; genital fold as shown in Fig. 6, K.

Color in alcohol. Similar to male but abdominal markings more conspicuous than in male.

Specimens examined : Many specimens of both sexes from Japan and Taiwan. This species is one of the commonest spiders in Japan, Taiwan and China.

Distribution : Japan, Korea, Taiwan and China.

The *shinanoensis*-group

Diagnosis. Anterior and posterior rows of eyes equal or subequal in width and arranged nearly parallel ; male chelicera with (a), (sl) and (T) ; conductor bifurcated terminally.

Tetragnatha shinanoensis Okuma and Chikuni, 1978
(Fig. 7, Table 7)

Tetragnatha shinanoensis Okuma and Chikuni, 1978, Acta Arachnol., 28 (1): 1.
Additional specimens : 2 ♀, Amakawa mura, Yoshino Gun, Nara Pref., 20. V. 1979, N. Kikuya ; 2 ♂, 1 ♀, Kitayama, Kyoto Pref., 3. VI. 1979, I. Kinoshita ; 1 ♂, 1 ♀, Miyama machi, Kitakuwata Gun, Kyoto Pref., 24. V. 1980, T. Kamura ; 1 ♀, Miyama machi, Kitakuwata Gun, Kyoto Pref., 25. V. 1985, T. Kamura.

Distribution : Japan (Honshu).
Note : This species is strikingly different from other known species of *Tetragnatha* by its bifurcate structure of the male conductor. In this species the pleats on the male conductor is indistinct. Body length of male 5.1-6.0 mm, female 5.9-7.3 mm.

Fig. 7. *Tetragnatha shinanoensis* Okuma. A : Right chelicera of male, lower view. B : Ditto lateral view. C : Ditto, upper view. D : Distal portion of conductor and embolus of male. E : Paracymbium of male. F : Right chelicera of female, lower view. G : Ditto, upper view. H : Eye group of female. I : Abdomen of female, lateral view. J : Genital fold of female. (After Okuma, 1978).

Table 7. *Tetragnatha shinanoensis* Okuma. Relative lengths of total legs and femora.

Locality	n	length of mean±s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)	7						
Nagano ♂	15	20.96±1.36	18.50-22.35	100	61 (0.989)	27 (0.806)	53 (0.964)
Nagano ♀	15	20.54±1.77	17.35-23.80		61 (0.982)	28 (0.962)	55 (0.957)
(Femora)							
Nagano ♂	7	5.44±0.37	4.90-5.80		68 (0.978)	36 (0.642)	66 (0.903)
Nagano ♀	15	5.47±0.27	5.00-6.00		68 (0.896)	37 (0.822)	68 (0.581)

The extensa-group

Diagnosis. Anterior and posterior rows of eyes subequal in width ; male chelicera with (a), (sl) and (T) ; conductor not bifurcated at tip.

KEY TO THE SPECIES

- 1 Basal segment of male chelicera about 0.8 times as long as carapace ;
 (a) bifid at tip *extensa*
 — Male chelicera about 0.7 times or less as long as carapace *pinicola*

***Tetragnatha extensa* (Linnaeus, 1758)**

(Fig. 8, Table 8)

Aranea extensa Linnaeus, 1758, Syst. Nat. (10 Ed.), : 621 ; Linnaeus, 1761, Faun. Svec., (2 Ed.) : 489 ;
 Fabricius, 1775, Syst. Ent., : 431 ; De Geer, 1783, Mem. p. s. Hist. Ins., 7: 236; Olivier, 1789,
 Encycl. Method., 4 : 204.

Aranea solandri Scopoli, 1763, Ent. Cam., : 397.

Aranea mouffetii Scopoli, 1763, Ent. Carn., : 398.

Aranea prima Herrich-Schaffer, 1804, Icon. Ins. Ratisb. Indig., : 49.

Aranea secunda Herrich-Schaffer, 1804, Icon. Ins. Ratisb. Indig., : 49.

Aranea quarta Herrich-Schaffer, 1804, Icon. Ins. Ratisb. Indig., : 49.

Tetragnatha rubra Risso, 1826, Hist. Nat. Princ. Prod. Europ. Merid., : 168.

Tetragnatha extensa : Sundevall, 1832, Vet. Ak. Handl., : 256 ; Hahn, 1834, Die Arachniden, 2 : 43 ; L.
 Koch, 1862, Corresp. Bl. Zool. Min. Ver. Regensbg., 16 (5) : 79 ; Westring, 1862, Aran. Svec., : 84 ;
 Blackwall, 1864, Hist. Spid. Gr. Brit. Irel., 2 : 367 ; Keyserling, 1865, Verh. zool. bot. Ges. Wien,

Fig. 8. *Tetragnatha extensa* (Linne). A : Left chelicera of male, upper view. B : Ditto, lower view. C : Ditto, lateral view. D : Paracymbium of male. E : Conductor and embolus of male. F : Left palpus of male. G : Left chelicera of female, upper inner view. H : Ditto, lower inner view. I : Genital fold of female. J : Eye group of male. K : Abdomen of female.

Table 8. *Tetragnatha extensa* (Linne). Relative lengths of total legs and femora.

Locality	n	length of means ± s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
JAPAN	♂	11	21.96f4.66	14.40-26.65	100	67 (0.995)	32 (0.970)
NEPAL	♂	1	17.00		68	33	65
JAPAN	♀	16	20.65f2.96	15.15-25.40		66 (0.983)	33 (0.929)
(Femora)							
JAPAN	♂	11	6.08f1.27	3.90- 7.30		74 (0.993)	39 (0.960)
NEPAL	♂	1	4.70		73	39	77
JAPAN	V	16	5.97f0.91	4.40-7.55		71 (0.988)	39 (0.923)
							76 (0.987)

15 : 844 ; Ohlert, 1867, Aran. Prov. Preussen, : 31 ; Thorell, 1875, Proc. Boston Soc. Nat. Hist., 17 : 493 ; Cambridge, 1881, Proc. Dorset F. Cl., 2 : 244 ; Dahl, 1883, Schrift. Nat. Ver. Schlesw. Holst. 5 (2): 32; Emerton, 1884, Trans. Connect. Ac. Sci., 6: 333; Hansen, 1885, Schiodte, 2001. Danica, : 105 ; Chyzer and Kulczynski, 1891, Aran. Hungar., 1: 144 ; McCook, 1893, Amer. Spid., 3: 259; Becker, 1896, Ann. Mus. Nat. Belg., 12 (1): 62; Emerton, 1902, Common Spid.,: 201 ; Kulczynski, 1903, S. B. Ak. Wiss. Wien, 112 (1) : 647 ; Lessert, 1910, Catal. Invert. Suisse ; Araign., : 288 ; Gravely, 1921, Rec. Ind. Mus. Calcutta, 22 (4 ; 24) : 425 ; Seely, 1928, Bull. N. York Stat. Mus., 278 : 113 ; Simon, 1929, Arachn. de France 6 (3) : 648, 651, 749 ; Wiehle, 1939, Nov. Act. Leopoldina (N. F.), 6 (41) : 376 ; Yaginuma, 1960, Spiders of Japan in Colour : 73.

Tetragnatha gibba C. L. Koch, 1837, Uebers. Arachn. Syst., 1 : 5.

Tetragnatha chrysochlora Walckenaer, 1841, Hist. Nat. Ins. Apt., 2 : 215 ; Walckenaer, 1847, Hist. Nat. Ins. Apt., 4 : 478.

Tetragnatha versicolor Walckenaer, 1841, Hist. Nat. Ins. Apt., 2 : 215.

Tetragnatha fluviatilis Keyserling, 1865, Verh. zool. bot. Ges. Wien, 15 : 852.

Tetragnatha nowickii L. Koch, 1870, Jahrb. Gel. Ges. Krakau, 41 : 13.

Tetragnatha extensa solandri Boesenberg, 1901, Zoolgica, 13 (1) : 58, 59.

Tetragnatha solandri Lessert, 1910, Catal. Invert. Suisse ; Araign., : 292.

Male. Body length, exclusive of chelicerae 5.1-9.0 mm ; carapacial length 2.0-3.0 mm, width 1.3-2.0 mm ; abdominal length 3.1-6.0 mm, width 1.1-1.6 mm ; cheliceral length 1.5-2.5 mm.

Eyes. Anterior row of eyes occupying full width of head, posterior row of eyes slightly wider or narrower than anterior one ; viewed from above, two rows nearly parallel and moderately recurved.

Chelicerae. Basal segment about 0.8 times as long as carapace, (a) bifurcated at tip, (AXu) present, (Gu) absent, (sl) sloping, (T) large, (rsu) 5-6 in number ; (AXI) and (GI) present, (GI) somewhat long, (rsl) about 7-8 in number ; fang unarmed.

Legs. Leg formula 1. 2. 4. 3; all legs with spines, first femur with 6-11 spines; relative lengths of legs as shown in Table 8.

Palpi. Tip of paracymbium generally rounded and with small distal notch; conductor and emborus as shown in Fig. 8, E and F.

Abdomen. Moderately long, about 3-4 times as long as broad, not extended posteriorly to spinnerets.

Color in alcohol. Generally light yellowish brown; abdomen with covered by sparse silvery spangles dorsally, somewhat dark brown ventrally.

Female. Body length, exclusive of chelicerae 7.3- 12.0 mm ; carapacial length 2.6-3.1 mm, width 1.7-2.1 mm ; abdominal length 4.7-9.0 mm, width 1.9-3.4 mm ; cheliceral length 1.7-2.2 mm.

Eyes. Nearly as in male.

Chelicerae. Basal segment about 0.7 times as long as carapace ; (AXu) and (Gu)

present, (rsu) about 10 in number ; (AXI) absent, (Gl) present (rsl) about 10 in number ; a keel present between (Gu) and (L2) as shown in Fig. 8, G and H ; fang unarmed.

Legs. Nearly as in male ; relative lengths of legs as shown in Table 8.

Abdomen. About 2.5 times as long as broad ; distal end of abdomen not extended posteriorly to spinnerets.

Color in alcohol. Nearly as in male.

Specimens examined : NEPAL : 1♂, 2♀, Mts. Pir Panjal 2,400 m, Tangmarg, 21. V. 1976, J. Martens. JAPAN : 69 specimens from Hokkaido (Ashoro and Koshimizu Kaen), Aomori (Mt. Hakkoda-san), Iwate (Mt. Hayachine), Tochigi (Nikko), Nagano (Minamiazumi-Gun and Shioziri City).

Distribution : Holarctic region.

Note: Roberts (1985) remarks as follows: "The sternal patterns separate *T. extensa* and *T. pinicola* from the other species, but it may sometimes be difficult to distinguish between small or immature females of *T. extensa* and adult females of *T. pinicola*. This problem is reliably overcome by measuring the ratio of femur I width/femur III length. This is greater than 0.205 in females of *T. extensa* (adult and subadult) and less than 0.176 in *T. pinicola*".

***Tetragnatha pinicola* L. Koch, 1870**

(Fig. 9, Table 9)

Tetragnatha pinicola L. Koch, 1870, Jahrb. Gel. Ges. Krakau, 41 : 11 ; Dahl, 1883, Scharif. Nat. Ver.

Fig. 9. *Tetragnatha pinicola* L. Koch. A : Left chelicera of male, upper view. B : Ditto, lower view. C : Ditto, lateral view. D : Conductor and embolus of male. E : Paracymbium of male. F : Left palpus of male. G : Left chelicera of female, upper view. H : Ditto, lower view. I : Eye group of male. J : Abdomen of male, lateral view. K : Genital fold of female.

Table 9. *Tetragnatha pinicola* L. Koch Relative lengths of total legs and femora.

Locality	n	length of means ± s.d.	1st legs range	1st legs ratio	2nd legs ratio (r)	3rd legs ratio (r)	4th legs ratio (r)
(Legs)							
Ilokkaido	♂ 7	15.36±1.37	13.65-16.80	100	69 (0.996)	35 (0.980)	61 (0.981)
Tochigi	♂ 8	16.28±2.04	12.45-18.95		68 (0.996)	34 (0.987)	60 (0.997)
Ehime	♂ 1	18.35			66	35	59
Kumamoto	♀ 2	16.83	16.55-17.10		67	34	59
Hokkaido	♀ 6	12.96±1.39	11.40-14.55		70 (0.985)	36 (0.921)	63 (0.994)
Tochigi	♀ 10	16.66±2.29	13.05-19.25		68 (0.999)	36 (0.997)	62 (0.997)
Ehime	♀ 5	15.17±0.72	14.10-16.05		69 (0.969)	36 (0.846)	62 (0.919)
Kumamoto	♀ 1	17.80			67	35	60
(Femoral)							
Ilokkaido	♂ 7	4.21±0.37	3.80- 4.60		75 (0.993)	43 (0.980)	73 (0.990)
Tochigi	♂ 8	4.52±0.60	3.40- 5.25		74 (0.995)	42 (0.980)	72 (0.992)
Chime	♂ 1	5.10			72	43	72
Kumamoto	♀ 2	4.55	4.50- 4.60		73	41	70
Hokkaido	♀ 6	3.59±0.37	3.20- 4.00		76 (0.983)	44 (0.842)	77 (0.991)
Tochigi	♀ 10	4.65±0.67	3.60- 5.35		75 (0.999)	44 (0.990)	76 (0.997)
Ehime	♀ 5	4.23±0.24	3.85- 4.50		75 (0.988)	44 (0.954)	76 (0.931)
Kumamoto	♀ 1	4.90			74	42	74

Schlew. Holst., 5 (2): 33; Chyzer and Kulczynski, 1891, Aran. Hungar., 1: 144; Boesenberg, 1901, Zoologica, 13 (1): 61; Kulczynski, 1903, S.B. Ak. Wiss. Wien, 112 (1): 647; Lessert, 1910, Catal. Invert. Suisse; Araign.,: 291; Simon, 1929, Arachn. de France, 6 (3): 651, 749; Wiehle, 1939, Nov. Act. Leopoldina (N. F.), 6 (41): 376; Yaginuma, 1960, Spiders of Japan in Colour: 73.

Eugnatha picta Lendl, 1886, Math. Term. Kozlem., 22: 129.

Male. Body length, exclusive of chelicerae 4.1-8.2 mm; carapacial length 1.5-2.2 mm, width 1.0-1.4 mm; abdominal length 2.5-6.0 mm, width 0.9-2.1 mm; cheliceral length 1.1-1.4 mm.

Eyes. Anterior row of eyes occupying full width of head, posterior row of eyes slightly wider or narrower than anterior one; viewed from above, two rows nearly parallel and moderately recurved.

Chelicerae. Basal segment about 0.7 times as long as carapace, (a) not bifid at tip, (AXu) present, (Gu) absent, (sl) sloping, (T) large, (rsu) about 5 in number; (AXI) and (GI) present, (rsl) about 5 in number; fang unarmed.

Legs. Leg formula 1. 2. 4. 3; all legs with spines, first femur with 5-7 spines; relative lengths of legs as shown in Table 9.

Palpi. Paracymbium bluntly rounded at tip; distal end of conductor bird's head-like and somewhat twisted as shown in Fig. 9, E and F.

Abdomen. About 3 times as long as broad; distal end of abdomen not extended posteriorly to spinnerets.

Color in alcohol. Generally light yellowish brown; dorsum of abdomen with covered by silvery spangles and with several pairs of small black dots along central line, venter somewhat dark brown.

Female. Body length, exclusive of chelicerae 6.4-9.5 mm; carapacial length 1.9-2.6 mm, width 1.2-1.6 mm; abdominal length 4.5-6.9 mm, width 1.8-2.5 mm; cheliceral length 1.0-1.7 mm.

Eyes. Nearly as in male.

Chelicerae. Basal segment 0.5-0.7 times as long as carapace; without particular features.

Legs. Nearly as in male; relative lengths of legs as shown in Table 9.

Abdomen. About 2.5 times as long as broad; distal end of abdomen not extended posteriorly to spinnerets.

Color in alcohol. Nearly as in male, but abdominal small black dots often absent.

Specimens examined : JAPAN : 113 specimens from Hokkaido (Nukabira and Ashoro), Tochigi (Nikko), Gifu (Mt. Shin-hodaka-San), Ehime (Mt. Ishizuchi-san), Kumamoto (Mt. Shiratori-yama).

Distribution : Palearctic Region.