

## Carbohydrate Enzymes in House Fly *Musca domestica* Larvae : Biochemical Properties and Toxicological Significance

Hirashima, Akinori

Laboratory of Pesticide Chemistry, Faculty of Agriculture, Kyushu University

Oyama, Kazuhiko

Laboratory of Pesticide Chemistry, Faculty of Agriculture, Kyushu University

Ueno, Ryohei

Laboratory of Pesticide Chemistry, Faculty of Agriculture, Kyushu University

Ishaaya, Isaac

Laboratory of Pesticide Chemistry, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/23886>

---

出版情報 : 九州大学大学院農学研究院紀要. 34 (1/2), pp.37-42, 1989-11. Kyushu University  
バージョン :  
権利関係 :



## Carbohydrate Enzymes in House Fly *Musca domestica* Larvae : Biochemical Properties and Toxicological Significance

Akinori Hirashima, Kazuhiko Oyama, Ryohei Ueno,  
Isaac Ishaaya\* and Morifusa Eto

Laboratory of Pesticide Chemistry, Faculty of Agriculture,  
Kyushu University 46-02, Fukuoka 812, Japan

(Received March 23, 1989)

The optimum temperatures and pHs for trehalase, amylase and invertase enzymes of *Musca domestica* were 52, 37 and 42°C, and 5.5, 6.5 and 6.0, respectively. The activity was linear up to 90 min and 60 min was chosen as the reaction time.  $K_m$  values for invertase, trehalase and amylase were 0.0185 M, 0.00676 M and 0.294%, respectively. *Musca domestica* larvae fed for two days on dietary concentrations of 5 and 10 ppm 2-methoxy-5-phenyl-1, 3, 2-oxazaphospholidine 2-sulfide (5-PMOS) resulted in weight gain inhibition of 39 and 70%, relative to control, correlating well with the suppression in the soluble gut trehalase activity. The digestive amylase and invertase enzymes were not affected by this compound. Hence, the trehalase system, which plays an important role in energy supply in insects, could be or closely relates to the primary site for the observed weight gain inhibition after 5-PMOS treatment. Similar results were observed with 4-isobutyl-2-methoxy-1, 3, 2-oxazaphospholidine 2-sulfide (iBMOS). This system is suggested as a biochemical parameter to assess retardation of growth caused by various insecticides.

### INTRODUCTION

Enzyme systems affecting insect growth such as trehalase which in insects degrades trehalose for internal energy supply (Ishaaya, 1986 : Wyatt, 1967) and digestive enzymes such as protease, amylase and invertase (Ishaaya, 1986) have been used to assess retardation of larval growth caused by secondary compounds present in plants or by treatment of insecticides exhibiting antifeeding activity (Ascher and Ishaaya, 1973 : Ishaaya and Casida, 1975). This paper describes the optimization of the trehalase, amylase and invertase enzymes in *M. domestica* larvae along with their toxicological significance using potential insecticides, such as 4-isobutyl-2-methoxy-1, 3, 2-oxazaphospholidine 2-sulfide (iBMOS) and 2-methoxy-5-phenyl-1, 3, 2-oxazaphospholidine 2-sulfide (5-PMOS).

### MATERIALS AND METHODS

#### Chemicals

4-Isobutyl-2-methoxy-1,3,2-oxazaphospholidine 2-sulfide (iBMOS) was prepared in the presence of triethylamine from methoxy phosphorodichloridothionate and

---

\*Permanent address : Department of Entomology, Agricultural Research Organization, The Volcani Center, Bet Dagan 50-250, Israel.

L-leucinol (Eto *et al.*, 1981). 2-Methoxy-5-phenyl-1,3,2-oxazaphospholidine Z-sulfide (5-PMOS) was similarly synthesized from 2-amino-1-phenylethanol (Wu *et al.*, 1988).

### **Rearing and bioassay**

A house fly diet (10 g, Oriental Kobo, K. K., Tokyo, Japan) was mixed with 35 ml acetone solution containing the test compound, or with 35 ml acetone alone as a control. After the solution had been evaporated under a hood for 2 days at room temperature, 12 g of water was added to the diet and thoroughly mixed. The diet was distributed in 1.5 g portions of ten replicates in test tubes (13 x 150 mm). Three larvae of susceptible (SRS) house fly, 1-2 days old weighing  $2 \pm 1$  mg each, were introduced into each test tube along with a thin piece of cotton placed over the diet to enable pupation. The tubes were kept at 28°C until pupation and emergence were complete. For weight gain measurement and enzyme activity determination, 2 larvae weighing  $12 \pm 1$  mg each were introduced into 5 replicates of 6 test tubes each along with 1.3 g portions of diet and the tubes were held at 28°C for 2 days. The weight gain along with the larval trehalase, amylase and invertase activities were determined. Assays run with solvent treatments containing no chemicals were used as control. The data for both biological and enzyme tests are reported as means with SE values.

### **Preparation of larval enzymes and enzyme assays**

The guts of 15-25 *M. domestica* larvae (weighing  $19 \pm 1$  mg each) were dissected free from other tissues by cutting off the larval head and removing the alimentary canal onto a wet filter paper (Hirashima *et al.*, 1989a ; Ishaaya and Casida, 1975) and then homogenized in cold distilled H<sub>2</sub>O (10-fold relative to larval weight), using a chilled glass-Teflon Potter-Elvehjem tissue grinder tube. The homogenate was centrifuged for 15 min at 10,000 g at 2°C, the supernatant fraction being used for the enzyme assays. The protein level (Lowry *et al.*, 1951) in the enzyme solution was  $755 \pm 12$  µg for the amylase assay, and  $378 \pm 6$  µg for the trehalase and the invertase assays. Trehalase, amylase and invertase activities were assayed colorimetrically on the basis of liberated glucose as previously described (Hirashima *et al.*, 1989a ; Ishaaya and Casida, 1975) under conditions found to be the optima for *M. domestica* larvae.

The reaction mixtures for amylase consisted of 0.2 ml of 4% (w/v) starch, 0.1 ml of 0.2 M phosphate buffer (pH 6.5) and 0.1 ml of enzyme solution. The optimum reaction mixture for trehalase and invertase consisted of 0.2 ml of 4% (w/v) trehalose (for trehalase) or 0.2 ml of 4% (w/v) sucrose (for invertase), 0.1 ml of 0.2 M phosphate buffer (pH 5.5 and 6.0 for trehalase and invertase, respectively) and 0.2 ml enzyme solution. After 60 min incubation at 37, 42 and 52°C for amylase, invertase and trehalase, respectively, 0.8 ml of 3, 5-dinitrosalicylic acid (DNSA) reagent (Noelting and Bernfeld, 1948) was added. The mixture was heated for 5 min at 100°C, and then immediately cooled in an ice-bath. The absorbancy at 550 nm was determined in E units with a Shimadzu UV-2100 spectrophotometer.

## **RESULTS AND DISCUSSIONS**

### **Biochemical properties of *M. domestica* larval gut enzymes**

The optimum conditions for enzyme assays were evaluated in a series of prelimi-

nary experiments, in which individual factors were varied, all others being kept at the optimum.

The optimum temperatures and pHs for trehalase, amylase and invertase were 52, 37 and 42°C (Fig. 1), and 5.5, 6.5 and 6.0 (Fig. 2), respectively. The activity was linear up to 90 min (Fig. 3) and 60 min was chosen as the reaction time for all the enzyme assays.  $K_m$  values for invertase, trehalase and amylase were 0.0185 M, 0.00676 M and 0.294%, respectively (Fig. 4). One E unit of the absorbancy at 550 nm was 0.60 mg of glucose for trehalase and invertase, and 0.45 mg of glucose for amylase, respectively (Fig. 5).

### Toxicological significance of carbohydrate enzymes of *M. domestica*

At sublethal concentrations, the five-membered cyclic phosphorothionates (iBMOS and 5-PMOS) resulted in a strong retardation of larval growth of the house fly larvae.




Fig. 1. Temperature optimum curves for trehalase, amylase and invertase of house fly *M. domestica* larvae.






Fig. 2. pH Optimum curves for trehalase, amylase and invertase of house fly larvae.



**Fig. 3.** Rate curves for trehalase, amylase and invertase of house fly larvae. Amylase activity was measured after dilution with water (1 : 1 volume) at the end of the reaction with DNSA reagent.



**Fig. 4.** Effect of substrate concentrations on trehalase, amylase and invertase of *M. domestica* larvae.

Hence, the digestive amylase and invertase activities which are important in food digestion and the trehalase enzyme in degradation of trehalose to glucose for energy supply (Ishaaya et al., 1986 : Wyatt, 1967) were assayed in larvae fed on diet containing these compounds.

House fly larvae fed for two days on dietary concentrations of 5 and 10 ppm 5-PMOS resulted in larval weight gain inhibition of 39 and 70%, correlating well with the suppression of the activity of the soluble larval gut trehalase (Table 1). 5-PMOS




Fig. 5. Standard curve for glucose. Amylase activity was determined after dilution with water (1 : 1 volume) at the end of the reaction with DNSA reagent.

Table 1. Effect of dietary iBMOS and 5-PMOS on the larval weight gain, the gut trehalase, and on the digestive amylase and invertase activity of *M. domestica* 2 days after treatment.

| Compound and dietary conc., ppm | Relative to control, % $\pm$ SE <sup>a</sup> |                         |                         |                         |
|---------------------------------|--|-------------------------|-------------------------|-------------------------|
|                                 | Weight                                       | Trehalase               | Amylase                 | Invertase               |
| iBMOS                           |  |                         |                         |                         |
| 50                              | 75 $\pm$ 14                                  | 69 $\pm$ 4 <sup>b</sup> | 101 $\pm$ 1             | 84 $\pm$ 1 <sup>b</sup> |
| 60                              | 40 $\pm$ 5 <sup>b</sup>                      | 60 $\pm$ 1 <sup>b</sup> | 81 $\pm$ 1 <sup>b</sup> | 82 $\pm$ 0 <sup>b</sup> |
| 5-PMOS                          |  |                         |                         |                         |
| 5                               | 61 $\pm$ 8 <sup>b</sup>                      | 59 $\pm$ 0 <sup>b</sup> | 121 $\pm$ 2             | 105 $\pm$ 3             |
| 10                              | 30 $\pm$ 5 <sup>b</sup>                      | 56 $\pm$ 1 <sup>b</sup> | 119 $\pm$ 1             | 107 $\pm$ 0             |

<sup>a</sup> Larval weight gains are averages with their SE values of 5 replicates of 6 test tubes (13  $\times$  150 mm) including 2 larvae each. Larvae 1-2 days old (12  $\pm$  1 mg) were introduced into the medium and maintained for 2 days. The enzyme assay data are average of 3 replicates. In the control, larval weight gain was 8.7  $\pm$  0.3 mg, and the trehalase, amylase and invertase activities were 0.73  $\pm$  0.01, 1.50  $\pm$  0.02 and 0.88  $\pm$  0 mg glucose/reaction, respectively.

<sup>b</sup> Differ significantly from the untreated control at  $P=0.05$ .

showed no inhibitory effect on the digestive amylase and invertase. Similar phenomena were observed when 50 and 60 ppm iBMOS used. These results indicate that iBMOS and 5-PMOS have at least two actions, the first of which affects AChE activity (Hirashima et al., 1989b), similar to other conventional acyclic organophosphorus compounds at lethal concentrations, and the second of which affects biochemical processes leading to a reduced activity of trehalase at sublethal concentrations. Hence, the trehalase is an important system to assess retardation of growth caused by iBMOS, 5-PMOS and other related compounds.

## ACKNOWLEDGEMENT

This work was supported in part by a Grant-In-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan, and the JSPS (Japan Society for the Promotion of Science) Fellowship to I. Ishaaya for research in Japan.

## REFERENCES

- Ascher, K. R. S. and I. Ishaaya 1973 Antifeeding and protease and amylase-inhibiting activity of fentin acetate in *Spodoptera littoralis* larvae. *Pestic. Biochem. Physiol.*, 3 : 326-336
- Eto, M., A. Hirashima, S. Tawata and K. Oshima 1981 Structure-insecticidal activity relationship of five-membered cyclic phosphoramidates derived from amino acids. *J. Chem. Soc. Jpn.*, 5 : 705-711
- Hirashima, A., I. Ishaaya, R. Ueno, K. Oyama and M. Eto 1989a Effect of salithion enantiomers on the trehalase system and on the digestive ptotease, amylase, and invertase of *Tribolium castaneum*. *Pestic. Biochem. Physiol.*, in press.
- Hirashima, A., R. Ueno, K. Oyama, I. Ishaaya and M. Eto 1989b Effect of five-membered cyclic phosphorothionates on larval growth, trehalase, digestive enzymes, acetylcholinesterase, and cyclic adenosine 3',5'-monophosphate level of *Tribolium castaneum* and *Musca domestica*. *Pestic. Biochem. Physiol.*, in press.
- Ishaaya, I. 1986 Nutritional and allelochemic insect-plant interactions relating to digestion and food intake : Some examples. In "Insect-Plant Interactions" ed. by J. R. Miller and T. A. Miller, Springer-Verlag, New York, pp. 191-223
- Ishaaya, I. and J. E. Casida 1975 Phenyltin compounds inhibit digestive enzymes of *Tribolium confusum* larvae. *Pestic. Biochem. Physiol.*, 5 : 350-358
- Lowry, O. H., N. J. Rosenbrough, A. L. Farr and R. L. Randall 1951 Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, 193 : 265-275
- Noelting, G. and P. Bernfeld 1948 Sur les enzymes amylolytiques. III. La  $\beta$ -amylase : Dosage d'activité et contrôle de l'absence d' $\alpha$ -amylase. *Helv. Chim. Acta*, 31: 286-290
- Wu, S.-Y., A. Hirashima, R. Takeya and M. Eto 1988 Synthesis and insecticidal activity of oxazaphospholidines, oxathiaphospholanes and thiazaphospholidines. *Agric. Biol. Chem.*, 52 : 2911-2917
- Wyatt, G. R. 1967 The biochemistry of sugars and polysaccharides in insect. *Adv. Insect Physiol.*, 4: 287-360