Anthocyanin Pigments in the Grape Skins of Several Red *Vitis vinifera* Cultivars bred in Japan

Shiraishi, Shin-ichi
University Farm, Faculty of Agriculture, Kyushu University

Watanabe, Yuka
University Farm, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/23969>

出版情報：九州大学大学院農学研究院紀要. 36 (1/2), pp.63-67, 1991-10. Kyushu University
バージョン：
権利関係：

Anthocyanin Pigments in the Grape Skins of Several Red *Vitis vinifera* Cultivars bred in Japan

Shin-ichi Shiraishi and Yuka Watanabe

University Farm, Faculty of Agriculture,
Kyushu University, Kasuyamachi, Fukuoka 811-23, Japan.

(Received July 1, 1991)

The anthocyanin pigments of red grapes *Vitis vinifera* cultivars, 'Kaiji', 'Sekirei', 'Morgenshön', 'Mario' and 'Unicorn' were examined by thin-layer chromatography and reflection densitometry. Seven spots were separated, and they were identified to pigments by *Rf* values, molybdate shift and color reactions, except for one unidentified spot. In 'Kaiji', the pigments in decreasing order of concentration, were cyanidin-3-monoglucoside (92%), delphinidin-3-monoglucoside (8%), and traces of peonidin-3-monoglucoside and an unidentified pigment. 'Ruby Okuyama', 'Morgenshön' and 'Sekirei' had constitutions similar to 'Kaiji'. Pigments of 'Rizamat', 'Mario' and 'Unicorn', in decreasing order of concentration, were malvidin-, cyanidin-, petunidin-, delphinidin- and peonidin-3-monoglucosides, 'Mario' lacking peonidin-3-monoglucoside. In 'Kaiji', the aglycones were cyanidin (92%) and delphinidin (8%). In 'Ruby Okuyama' and 'Morgenshön', the aglycones were in about the same order, and plus petunidin in 'Sekirei'. 'Rizamat' and 'Unicorn' contained five aglycones, in decreasing order of concentration, malvidin, cyanidin, petunidin, delphinidin and peonidin, and peonidin was absent in 'Mario'.

INTRODUCTION

Investigation by Ribéreau-Gayon of *Vitis vinifera* cultivars have indicated that this species is characterized by 3-monoglucosides for five common anthocyanins (Ribéreau-Gayon et al., 1955). Rankine et al. and Albach et al. reported that the pigment composition of the majority of red wine cultivars of *V. vinifera* appeared to conform to a general eight-band pattern when analyzed by paper chromatography. The 3-monoglucosides of peonidin and of cyanidin have been recently identified as major components of certain *V. vinifera* table grape cultivars (Albach et al., 1963). Anthocyanins isolated from two *V. vinifera* cultivars 'Koshu' and 'Gros Colman' were identified as 3-monoglucosides only (Akuta et al., 1977).

A fresh red skin color is very attractive for table grapes. In Japan, viticulture of *vinifera* cultivars in open fields is very difficult because of the rainy climate during the growing period. But viticulture under structures enables the production of *vinifera* cultivars such as 'Flame Tokay' or 'Rizamat' which are fresh red and of high quality. The new *V. vinifera* cultivars which were recently bred in Japan are well suitable for viticulture under structures.

The present paper is concerned with the identification and qualitative assessment of the anthocyanins of seven red table grapes grown in Japan.

MATERIALS AND METHODS

Ripe grapes of 'Kaiji', 'Sekirei', 'Morgenshön' and 'Unicorn' were supplied by N. Uehara, Director of the Uehara Grape Institute in Yamanashi. Ripe grapes of 'Ruby Okuyama', 'Rizamat' and 'Mario' were supplied by T. Simi of the Akitsu Branch Station of the Fruit Tree Research Station.

The methods of sample treatment, isolation of anthocyanins and measurement of the optical densities of the separated spots were the same as in previous report (Shiraishi *et al.*, 1986).

RESULTS AND DISCUSSION

The parentages and skin colors of grape cultivars examined were shown in Table 1. The skin colors of 'Kaiji', 'Sekirei', 'Morgenshön' and 'Ruby Okuyama' were almost fresh red, but those of 'Rizamat', 'Mario' and 'Unicorn' were dull red or dark red.

Figures 1 and 2 show a tracing from a two-dimensional thinlayer chromatograms of the pigments from 'Kaiji' and 'Rizamat'. The identities of the pigments, as

Table 1. Parentages and skin color of grape cultivars examined.

Cultivar	Parent of cultivar	Skin color
Kaiji	'Flame Tokay' × 'Neo Muscat'	Red
Sekirei	A bud mutation of 'Kaiji'	Fresh purple-red
Morgenshön	'Katta Kurgan' × 'Kaiji'	Fresh red
Ruby Okuyama	A bud mutation of 'Itaria'	Fresh red
Rizamat	'Katta Krugan' × 'Parkentskij'	Purple-red
Mario	'Rizamat' × 'Neo Muscat'	Reddish purple-black
Unicorn	Seedling of <i>V. vinifera</i> cultivar	Dull red

Fig. 1. A two-dimensional thin-layer chromatogram of the anthocyanins in 'Kaiji'

Fig. 2. A two-dimensional thin-layer chromatogram of the anthocyanins in 'Rizamat'.

indicated in Tables 2 and 3, are based on the position of the pigments on the two-dimensional chromatograms molybdate shift and color reactions, and co-chromatography with known pigment samples isolated in prior studies with *V. vinifera* pigments mixtures using column chromatography method (Ohta et al., 1978). Table 4 lists the percent distribution of the pigments of seven cultivars as determined by chromatogram scanner. Table 5 shows the aglycone constitution of seven cultivars.

In 'Kaiji', cyanidin-3-monoglucoside is by far the major anthocyanin pigment. The cyanidin-derived anthocyanin accounts for 92% of the total pigments; the delphinidin derivative is a distant second, 8%; and the peonidin-derived anthocyanin accounts for only a trace. Only an unknown spot at R_f 0.35 on *i*BAW and 0.18 in *i*AHW was also recognized and its color changed with molybdate. This spot was also recognized in a trace from 'Ruby Okuyama'. The percent distribution of pigments in 'Ruby Okuyama' and 'Morgenshön' was similar to that of 'Kaiji', the former lacking a trace amount of peonidin-3-monoglucoside, the latter lacking an unknown spot. In 'Sekirei' the major anthocyanin pigment, cyanidin-3-monoglucoside, accounted for 76% of the total pigments; the delphinidin derivative for 22%; and the petunidin-derived anthocyanin accounted for only a trace. The ratio of the absorption density of crude extract in 0.1% HCl-MeOH from 'Kaiji' and 'Sekirei' was 10.4 : 28.8 at 535 nm. This result shows the improvement of red color development by a bud mutation in 'Kaiji'.

Major aglycone in 'Rizamat' is malvidin, followed by cyanidin, petunidin, peonidin and delphinidin. The major aglycone of 'Mario' is also malvidin, and in decreasing order of importance are petunidin, cyanidin and delphinidin, this cultivar lacking peonidin. Therefore, seven cultivars examined here are able to be grouped as the 'Kaiji' group, 'Kaiji', 'Sekirei', 'Ruby Okuyama' and 'Morgenshön' and the 'Rizamat' group, 'Rizamat', 'Mario' and 'Unicorn'. The 'Sekirei' group shows fresh red skin

Table 2. Comparison of *Rf* values and colors of individual anthocyanins in examined grapes

Pigment No.	<i>Rf</i> value ($\times 100$) at 25°C		Color in		Molybdate shift
	<i>i</i> BAW	AHW	visible light	UV light	
1	11	8	Purple	Dull purple	+
2	17	11	Reddish purple	Dull purple	+
3	21	14	Red	Dull purple	+
4	27	17	Red-purple	Dull purple	
5	32	19	Pink	Dull red-purple	
9	49	15	Red-purple	Dull purple	
21	35	18	Red	Dull purple	+

Table 3. Identification of the anthocyanins in examined grapes.

Pigment No.	Identification	Abbreviation
1	Delphinidin-3-monoglucoside	(Dl-3G)
2	Petunidin-3-monoglucoside	(Pt-3G)
3	Cyanidin-3-monoglucoside	(CY-3G)
4	Malvidin-3-monoglucoside	(Mv-3G)
5	Peonidin-3-monoglucoside	(Pn-3G)
9	Malvidin-3-monoglucoside acylated with p-coumarate	(Mv-3G-Cm)
21	Unidentified	-

Table 4. Percentages of constituent anthocyanins in skin of the examined grapes.

Pigment No.	Abbreviation	Estimated percent of total anthocyanin content							
		Kaiji	Sekirei	Ruby	Okuyama	Morgenshön	Rizamat	Mario	Unicorn
1	Dl-3G	8	22		3	2	5	11	17
2	Pt-3G		2				12	14	14
3	Cy-3G	92	76		97	98	13	11	28
4	Mv-3G						36	32	26
5	Pn-3G	trace	trace				8		14
9	Mv-3G-Cm	—					26	32	
21	Unknown	trace	—		trace		—		

Table 5. Aglycone constitution of the examined grapes.

Aglycone	Estimated percent of total aglycone content							
	Morgenshön	Ruby	Okuyama	Kaiji	Sekirei	Rizamat	Mario	Unicorn
Cyanidin	98		97	92	76	13	11	28
Peonidin				trace	trace	8		14
Delphinidin	2		3	8	22	5	11	17
Petunidin					2	12	14	14
Malvidin						62	64	26
Unknown			trace	trace				

color, and the 'Rizamat' group shows dull or dark red, namely, in the former cyanidin-derived anthocyanin was the major pigment, and they did not contained methylated anthocyanins, malvidin, petunidin and peonidin, or only a little. In breeding of fresh red cultivars, parents should be made use of 'Kaiji' group which contain high amounts of cyanidin and delphinidin lack methylated anthocyanins, malvidin, petunidin and peonidin.

REFERENCES

- Akuta, A., H. Ohta, Y. Osajima, N. Matsudomi and K. Kobayashi 1977 Studies on anthocyanin pigments in fruits harvested in Japan. V. Anthocyanin pigments in *Vitis vinifera* varieties, 'Koshu' and 'Gros colman' grapes *J. Food Sci. Technol.*, 24 : 521-523
- Albach, R., R. E. Kepner and A. D. Webb 1959 Comparison of anthocyanin pigments of red *vinifera* grapes. II. *Amer. J. Enol. Viticult.*, 10 : 164
- Albach, R. F., R. E. Kepner and A. D. Webb 1963 Peonidin-3-monoglucoside in *vinifera* grapes. *J. Food Sci.*, 28 : 55-58
- Benitley, K. W. 1960 The national pigment. In "The chemistry products. Vol. IV Interscience Publishers, New York, pp. 28
- Ohta, H., S. Akuta, S. Shiraishi and Y. Osajima 1978 Anthocyanin Pigments of Three Y-Grape Strains Backcrossed by *Vitis vinifera* varieties. *J. Food Sci. Technol.*, 25: 421-425
- Rankine, S., R. E. Kepner and A. D. Webb 1958 Comparison anthocyanin pigments of *vinifera* grapes. *Amer. J. Enol. Viticult.*, 9: 105-110
- Ribérea-Gayon, J., P. Sudraud and P. M. Durquety 1955 Relation entre genetique et nature chimique des pigments anthocyaniques de la baie dans le gere *Vitis*. *Revue General de Botanique*, 62 : 995-1005
- Shiraishi, S., Y. Watanabe, H. Okubo and S. Uemoto 1986 Anthocyanin Pigments of Black-Purple Grapes Related to Variety 'Kyoho' (*Vitis vinifera* L. $\times$ *V. Labrusca* L.). *J. Japan. Soc. Hort. Sci.*, 55: 123-129