

Stimulation of Spore Formation, Mycelial Growth and Thiostrepton Production by several Thiol Compounds and β -NAD in *Streptomyces azureus* ATCC 14921 and Its Derivatives

Ogata, Seiya

Microbial Genetics Division, Institute of Genetic Resources, Faculty of Agriculture, Kyushu University

Tawara, Yuka

Microbial Genetics Division, Institute of Genetic Resources, Faculty of Agriculture, Kyushu University

Tokunaga, Satoko

Microbial Genetics Division, Institute of Genetic Resources, Faculty of Agriculture, Kyushu University

Mise, Osamu

Microbial Genetics Division, Institute of Genetic Resources, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/24026>

出版情報：九州大学大学院農学研究院紀要. 37 (3/4), pp.331-340, 1993-03. Kyushu University
バージョン：
権利関係：



**Stimulation of Spore Formation,
Mycelial Growth and Thiostrepton Production
by several Thiol Compounds and β -NAD
in *Streptomyces azureus* ATCC 14921 and Its Derivatives**

**Seiya Ogata, Yuka Tawara, Satoko Tokunaga, Osamu Mise,
Hitoshi Matsubara, Mangkorn Rodprapakorn, Toshio Hara,
Chizuru Kinoshita* and Woo-Yond Choi****

Microbial Genetics Division, Institute of Genetic Resources, Faculty of Agriculture,
Kyushu University 46, Fukuoka 812, Japan

(Received December 11, 1992)

Sporulation-stimulating compounds were screened by using mutant PKIOOa of *Streptomyces azureus* ATCC 14921 (wild-type strain PKO), in which spore formation was inhibited by a pock-forming plasmid pSA1.1. On agar media, L- and D-cysteines, bacitracin, glutathione, and β -NAD induced the formation of colored spores or spore mass of the strain PKIOOa. These compounds also stimulated the spore formation of strain PKO and its good spore-forming (plasmid-free) strain PK100C, and the growth of aerial and submerged mycelia of these three strains. L- and D-cysteines allowed 1.3 to 1.4-fold increase in the production of antibiotic thiostrepton in strains PKO and PK100C. The thiostrepton production of strain PKIOOa, which hardly produced it, was also increased by L- and D-cysteines by as much as 20% of that of strain PKO.

INTRODUCTION

Streptomyces species are prokaryotes that differentiate and produce spores with species- (or strain-) specific color. In the typical differentiation on agar media, the substrate (or vegetative) mycelium develops and later gives rise to an aerial mycelium, which is finally converted in part into chains of spores. The morphological differentiation, in particular, during the aerial mycelial formation is often associated with the production of antibiotics and other secondary metabolites (Chater, 1989; Hopwood, 1988). Great interests have been focused on the screening of substances which affect morphological differentiation, spore formation and aerial mycelial formation of streptomycetes, and also on their mode of action (Chater, 1989; Coleman and Ensign, 1982; Hopwood, 1988; Kondo et al., 1988; Khokholov, 1988; Maruo, 1987; Natsume et al., 1989; Ochi, 1990; Ochi et al., 1984; Ogata, 1991; Ogata et al. 1985b; 1991; Vargha and Szabo, 1984). An abundant production of spores is necessary for the cultivation of industrial streptomycetes, because they are generally used as the first inoculum (starter).

Much interests are also focused on the inhibition of spore formation and aerial mycelial growth by pock-forming plasmids (conjugative sex plasmids) in *Streptomyces* spp. (Bibb et al., 1977; Hopwood et al., 1973; 1984; Kieser et al., 1982; Ogata, 1991).

* Nakamura Gakuen College, Fukuoka 812, Japan.

** College of Agriculture, Chungnam University, Korea.

The spore formation and thiostrepton production in *Streptomyces azureus* were markedly decreased in the strains which carried a pSA1.1 (8.8 kb), a multicopy derivative of pock-forming plasmid pSA1 of the wild-type strain (Miyoshi *et al.*, 1986; Ogata *et al.*, 1988). The parent pSA1 also injured its host (Ogata *et al.*, 1981; 1989; Ogata, 1991). In this paper, we screened cysteine, bacitracin, glutathione and β -NAD (nicotinamide adenine dinucleotide) as restorers of spore formation inhibited by pSA1.1. We also investigated the effects of these compounds on spore formation, aerial mycelial growth, submerged mycelial growth and thiostrepton production in *S. azureus* ATCC 14921 (wild-type strain) and its derivatives, pSA1.1-carrying and plasmid-free (good spore-forming) strains, and discuss the function of these reagents.

MATERIALS AND METHODS

Bacterial strains

Streptomyces azureus ATCC 14921 (wild-type strain PKO) and its derivatives were used in this work (Miyoshi *et al.*, 1986). The wild-type strain PKO carried a pock-forming plasmid pSA1 (1 or less copy/host genome). A plasmid-free strain PKIOOC was obtained from strain PKIOO spontaneously, which carried a multicopy plasmid pSA1.1 (20 to 30 copies/host genome), a derivative of pSA1. Strain PKIOOC formed large numbers of spores. Strain PKIOOa was constructed by mating between strains PKIOO and PKIOOC, and carried plasmid pSA1.1 (20 to 30 copies/host genome), as described previously. The construction of strain PKIOOa was performed, because strain PKIOO derived the plasmid-free cells at a high frequency. Spore formation of strain PKIOOa was less than that of its parent PKIOO. Some characteristics of strains PKO, PKIOO and PKIOOC had been reported in the previous papers (Miyoshi *et al.*, 1986; Ogata *et al.*, 1981; 1988).

Media and cultural conditions

Rye flakes agar and Bennett broth and agar supplemented with or without 1% corn starch were used for the growth of organisms (Ogata *et al.*, 1981). The spores of strains PKO and PKIOOC were started as the first inoculation, as described previously. For the starter of strain PKIOOa, the submerged mycelia stocked at -30°C were used, because its spore formation was very poor. Equal concentrations of starter were used. The cultural temperature was 28°C. The liquid culture was carried out on the rotary shaker at 250 rpm in the flask with glass beads.

Measurement of number of spores

The spores were scrapped from 7-day-old slant cultures of rye flakes agar and suspended in 0.1M Tris-HCl buffer (pH 7.2). Chains of spores were broken down with a mixer or homogenizer as previously described (Ogata *et al.*, 1981). The mycelia were removed by repeated filtration through a filter paper (Toyo roshi no. 2). The number of spores was estimated by microscopic counts of ten random haemocytometer fields.

Measurement of length of aerial mycelium

The length of aerial mycelium was measured by the method of Schauer *et al.*

(1988). Lawn of mycelia grown for 5 days on Petri plates of rye flakes agar was excised and sliced from top to bottom, and then placed sideways on a cover slip. The cover slip was placed on a microscope, and the length of the mycelium was microscopically measured.

Weighing of mycelial mass in liquid culture

Precultivation was carried out for 24h in Bennett broth without 1% corn starch, and then the culture broth was transferred (5% v/v) to a fresh broth with 1% corn starch. The mycelia grown for 5 days were centrifuged at 3,000xg for 10 min, and then the mycelial mass was weighed (wet weight).

Thiostrepton productivity

Thiostrepton productivity in the solid and liquid cultures was determined as previously described (Ogata et al., 1985a). The mycelia grown for 5 days in the liquid culture containing 1% corn starch were gathered by centrifugation and then applied to DMSO-extraction. The concentration of thiostrepton in the extract was determined by paper disk method with *Bacillus subtilis* ATCC 6633 as test organisms: paper disks of 8 mm in diameter were used and a standard curve was obtained with known concentration of thiostrepton (supplied by Asahi Kasei-Kogyo Ltd).

In the case of solid culture, plugs (8 mm in diameter) of agar were excised with a cork borer from the Bennett agar of Petri plates containing 1% corn starch and placed on the assay plates with the same test organisms. Thiostrepton production was indicated by a clear zone of inhibition around the agar plugs, and the concentrations of thiostrepton produced were determined by the comparison of the diameter of inhibition zones with those of paper disk (8 mm in diameter) with known concentrations of thiostrepton.

RESULTS

Detection of stimulators of spore formation

Strain PK100a which carried 20 to 30 copies of a multicopy plasmid pSA1.1, of which spore formation was strongly inhibited, was inoculated onto the surface of rye flakes agar of Petri plates as an indicator strain. Paper disks containing compounds such as amino acids, nucleic acid-related compounds, vitamins and antibiotics at 1 to 1000 $\mu\text{g/ml}$ were placed on the plates. Cysteine (Sigma Chemical Co.), bacitracin (Sigma Chemical Co. and Asahi Kasei-Kogyo Ltd.), glutathione (Sigma Chemical Co.) and β -NAD (Kohjin Ltd.), which induced a faint bluish green color (indicating spore formation) around the paper disk were detected after 5 days or more cultivation.

The effects of these compounds on spore formation of strain PK100a, the wild-type strain PKO and the good spore-forming strain PK100C were tested. The maximum sporulation efficiency was obtained during 7-days cultivation of each strain. The results at the several concentration of each compound are shown in Table 1. The number of spores on the strain PK100a was increased 100-fold or more by these compounds compared with the control with no additions. They were also effective on the spore formation of the wild-type strain PKO and plasmid-cured strain PK100C, with 1.5 to 5-fold increment in the number of spores at the optimum concentration.

The effect on the plasmid pSA1-carried strain PKO was more distinct than on the plasmid-cured strain PK100C. Therefore, the sporulation-stimulating effect of these compounds seems to be in proportion to be the inhibitory effect of plasmids. Bacitracin and P-NAD showed marked effects on the promotion of the spore formation in all the strains used. They also advanced the onset of coloring (maturation age) of spores or spore mass, as shown in Fig. 1. The effect of L-cysteine on spore formation was similar to that of D-cysteine.

Effect of cysteine, bacitracin, glutathione and P-NAD on aerial mycelium

The stimulatory effects of cysteine, bacitracin, glutathione and P-NAD on the formation of aerial mycelium were also observed. The length of the aerial mycelium estimated was shown in Table 2. The effect of bacitracin on the wild-type strain PKO was the most marked with about 3.5-fold stimulation. The growth of aerial mycelium was so obvious as to be recognized with a glance at the colonies or the lawn of plate culture. P-NAD also showed a strong effect on PKO with about a 1.8-fold increase. The effect of others rather resembled and slight. These results suggested that the stimulation of the aerial mycelium formation was closely correlated with the stimulation of the spore formation.

Table 1. Effect of compounds on the formation of spores in *Streptomyces azureus* ATCC 14921 (wild-type PKO) and its derivatives.

Compounds ($\mu\text{g/ml}$)	Strain		
	PKO	PK100a (spores/slant)	PK100C
No addition	1.8×10^8	4.6×10^7	8.5×10^8
L-cys (10)	1.9×10^8	5.4×10^7	9.0×10^8
L-cys (50)	3.0×10^8	1.5×10^7	1.2×10^9
L-cys (100)	2.7×10^8	1.0×10^7	9.5×10^8
D-Cys (50)	3.0×10^8	9.8×10^7	1.1×10^9
Bacitracin (10)	2.8×10^8	1.0×10^7	1.0×10^9
Bacitracin (50)	8.6×10^8	2.5×10^7	2.0×10^9
Bacitracin (100)	2.1×10^8	1.4×10^7	1.3×10^9
Glutathione (10)	1.9×10^8	4.8×10^6	8.8×10^8
Glutathione (50)	2.4×10^8	7.2×10^6	9.7×10^8
Glutathione (100)	1.0×10^8	8.7×10^5	8.0×10^8
β -NAD (100)	3.1×10^8	2.3×10^6	9.2×10^8
P-NAD (500)	8.8×10^8	1.8×10^7	1.7×10^9
β -NAD (1000)	1.7×10^8	8.5×10^6	1.0×10^9

PKO, wild-type and plasmid pSA1-carrying strain; PK100a, plasmid pSA1.1-carrying strain; PK100C, plasmid-free strain; Cys, cysteine; NAD, nicotinamide adenine dinucleotide.

Each value is the average of five cultures.

Table 2. Effect of compounds on the length of aerial mycelium in the plate culture of *Streptomyces azureus* ATCC 14921 (PK0) and its derivatives.

Compounds ($\mu\text{g/ml}$)		Strain		
		PK0	PK100a (μm)	PK100C
No addition		69	67	80
L-cys	(10)	72	70	81
L-cys	(50)	100	86	106
L-cys	(100)	98	83	98
D-Cys	(50)	98	74	98
Bacitracin	(10)	124	102	92
Bacitracin	(20)	153	108	107
Bacitracin	(50)	243	96	123
Bacitracin	(100)	170	80	114
Glutathione	(10)	70	68	80
Glutathione	(50)	72	88	88
Glutathione	(100)	67	66	76
β -NAD	(500)	125	80	111
β -NAD	(1000)	111	73	94

Explanations of PK0, PK100a, PK100C, Cys and NAD are the same as described in Table 1.

Each value is the average of about 15 specimens were prepared from five plates.




Fig. 1. Stimulation of maturation age of spores.

The coloring of spores or spores mass around the disk contained β -NAD was advanced. Wild-type strain PK0 was used in this experiment. Paper disk: 8mm in diameter.

Effect of cysteine, bacitracin, glutathione and P-NAD on the growth of submerged mycelium

Bacitracin and P-NAD showed a growth stimulating effect accompanied by an increase in the submerged mycelium mass, as shown in Table 3. The effect was remarkable and an increase in the amount of mycelium of the strain PKO was 1.3-fold or more than that with no addition at the optimum concentration. The slight or negligible effect was observed for cysteine and glutathione.

Effect of cysteine, bacitracin, glutathione and P-NAD on production of thio-strepton

The effects of cysteine, bacitracin, glutathione and P-NAD on the production of thio-strepton in plate culture were tested. As shown in Table 4, only L- and D-cysteine caused an increase in thio-strepton production for all the strains tested. The effect of L-cysteine was similar to that of D-cysteine. Thio-strepton production of the strain PK100a was very low, but it was increased 1.3-fold with L- or D-cysteine at the optimum concentration compared to that with no addition.

Although thio-strepton production of the strain PK100a in liquid culture was hardly detectable, it was increased by 10 to 20% of that of strain PKO when the optimal concentration of L- or D-cysteine was 100 $\mu\text{g/ml}$ (Table 5). L- and D-cysteine

Table 3. Effect of compounds on the growth of mycelium in the liquid culture of *Streptomyces azureus* ATCC 14921 (PKO) and its derivatives.

Compounds ($\mu\text{g/ml}$)		Strain		
		PKO	PK100a (mg/ml)	PK100C
No addition		52.3	44.6	52.1
L-cys	(10)	53.9	45.7	52.1
L-cys	(50)	59.2	46.0	52.6
L-cys	(100)	53.2	45.2	50.9
D-Cys	(50)	58.5	45.8	52.1
Bacitracin	(10)	58.5	49.2	56.9
Bacitracin	(20)	70.0	50.3	61.7
Bacitracin	(50)	72.0	50.8	58.1
Bacitracin	(100)	51.5	43.1	45.7
Glutathione	(10)	52.4	44.0	52.2
Glutathione	(50)	52.8	44.7	52.5
Glutathione	(100)	52.5	44.3	52.0
P-NAD	(100)	52.3	45.7	61.0
P-NAD	(500)	67.9	51.7	66.7
β -NAD	(1000)	61.6	51.5	67.1

Explanations of PKO, PK100a, PK100C, Cys and NAD are the the same as described in Table 1.
Each value is the average of five cultures.

Table 4. Effect of compounds on the production of thiostrepton in the plate culture of *Streptomyces azureus* ATCC 14921 (PKO) and its derivatives.

Compounds (μ g/ml)		Strain		
		PKO	PK100a (μ g/agar plug)	PK100C
No addition		6.4	3.0	4.5
L-cys	(10)	6.5	3.5	4.7
L-Cys	(50)	8.1	4.2	5.3
L-cys	(100)	7.8	3.5	5.3
L-cys	(200)	6.0	3.2	4.3
D-Cys	(50)	7.7	4.2	5.3
D-Cys	(100)	7.5	4.0	5.0
Bacitracin	(10)	6.4	3.0	4.5
Bacitracin	(50)	6.6	3.0	4.7
Bacitracin	(100)	4.0	1.9	2.5
Glutathione	(10)	6.1	3.0	4.5
Glutathione	(50)	6.4	3.0	4.6
Glutathione	(100)	5.1	3.0	3.4
P-NAD	(100)	6.3	3.0	4.5
P-NAD	(500)	6.4	3.0	4.5
P-NAD	(1000)	6.1	3.0	4.1

Explanations of PKO, PK100a, PK100C, Cys and NAD are the the same as described in Table 1. Plug: 8 mm in diameter and 5 mm in thickness. Each value is the average of five plates.

Table 5. Effect of compounds on the production of thiostrepton in the liquid culture of *Streptomyces azureus* ATCC 14921 (PKO) and its derivatives.

Compounds (μ g/ml)		Strain		
		PKO	PK100a	PK100C
		(w.w.mg of mycelium)		
No addition		3.0	ND	2.8
L-cys	(10)	3.2	ND	3.0
L-cys	(50)	4.3	0.5	3.5
L-cys	(100)	4.2	0.9	3.6
L-Cys	(200)	2.4	0.3	2.2
D-Cys	(50)	4.0	0.2	3.0
D-Cys	(100)	4.1	0.4	3.3
Bacitracin	(50)	3.0	ND	2.8
Glutathione	(50)	3.0	ND	2.8
β -NAD	(500)	3.0	ND	2.8

Explanations of PKO, PK100a, PK100C, Cys and NAD are the the same as described in Table 1. ND, not detectable; w.w.mg, wet weight of cells in milligram.

Each value is the average of five cultures.

allowed about 1.4- and 1.3-fold increase in thiostrepton production in strains PKO and PKIOOC, respectively. Bacitracin and β -NAD had no stimulatory effect on thiostrepton production per mycelium. But the production was enhanced in the cultural mass, because the total amount of mycelium was increased in the media (Table 3).

DISCUSSION

Cysteine, bacitracin, glutathione and β -NAD were detected as spore formation-stimulating compounds, which also stimulated the aerial mycelium formation, and the growth of submerged mycelium not only in the pSA1.1-carrying strain PK100a but in the wild-type strain PKO and plasmid-free strain PKIOOC. The first target of each of these four compounds on the cells might differ from one another. However, judging from the known properties or functions of these thiol compounds (Aronson *et al.*, 1976; Cheng *et al.*, 1973; Garcia-Patrone, 1985) and NAD (Penyige *et al.*, 1990), we supposed that they enhanced or changed the cell membrane function, and then stimulated mycelial growth or spore formation in the three strains used. It was also considered that the stimulation of function of the cell membrane overcame the inhibitory effect of pSA1.1, which allowed the strain PK100a to form spores. Similar overcoming effect would be done on the strain PKO, which carried the parent plasmid pSA1, because the stimulatory effect on the strain PKO was greater than that on the plasmid-cured strain PKIOOC. We are eagerly continuing our efforts to clarify the mode of action of bacitracin on the wild-type strain PKO, because this compound showed a distinctly stronger effect on its cell differentiation.

Only cysteine was effective on the stimulation of the thiostrepton production. We understood that it was due to the fact that the structural component of thiostrepton contained one cysteine residue and five thiazole rings derived from cysteine (Anderson *et al.*, 1970; Foss, 1988). But bacitracin and glutathione could not be the direct sources of cysteine, although they were cysteine-containing compounds. Racemization was supposed to have occurred from the observation that D-cysteine had a similar effect as that of L-cysteine. Furthermore, the stimulatory effects of mycelial growth and morphological differentiation such as aerial mycelium and spore formation were considered to affect the production of thiostrepton, a secondary metabolite.

We had screened chemicals which overcame the inhibition of spore formation. The chemicals detected were also stimulated both spore formation and mycelial growth (or thiostrepton production) of all strains used. So, the screening method used in this work appears to be an effective method for the screening of substances which stimulate spore formation and mycelial growth or production of antibiotics or secondary metabolites of streptomycetes or which overcome the inhibitory effect of pock-forming plasmids.

ACKNOWLEDGEMENTS

This work was partially supported by a grant-in-aid for scientific research from the Ministry of Education of Japan.

REFERENCES

- Anderson, B., D. C. Hogkin and M. A. Viswamitra 1970 The structure of thiostrepton. *Nature*, 225: 233-235
- Aronson, A. I., and P. Fitz-James 1976 Structure and morphogenesis of the bacterial spore coat. *Bacteriol. Rev.*, 40: 360-402
- Bibb, M. J., R. F. Freeman and D. A. Hopwood DA 1977 Physical and genetical characterization of a second sex factor, SCP2, for *Streptomyces coelicolor* A3(2). *Mol. Gen. Genet.*, 154: 155-156
- Chater, K. F. 1989 Sporulation in *Streptomyces*. p. 277-300. In: I. Smith, R. A. Slepecky and P. Setlow (ed.) Regulation of prokaryotic development. American Society for Microbiology, Washington, D. C.
- Cheng, H. M., A. L. Aronson and S. C. Holt 1973 Role of glutathione in the morphogenesis of the bacterial spore coat. *J. Bacteriol.*, 113: 1134-1143
- Coleman, R. H., and J. C. Ensign 1982 Regulation of formation of aerial mycelia and spores of *Streptomyces viridochromogenes*. *J. Bacteriol.*, 149: 1102-1111
- Floss, H. G. 1988 Biosynthesis of the antibiotics nosiheptide and thiostrepton. p. 401-405. In: K. Okami, T. Beppu and H. Ogawara (ed.) Biology of actinomycetes '88. Japan Scientific Societies Press, Tokyo.
- Garcia-Patrone, M. 1985 Bacitracin increases size of parasporal crystals and spores in *Bacillus thuringiensis*. *Mol. Cell. Biochem.*, 68: 131-137
- Hancock, R. and P. C. Fitz-James 1964 Some differences in the action of penicillin, bacitracin and vancomycin on *Bacillus megaterium*. *J. Bacteriol.*, 87: 1044-1050
- Hopwood, D. A 1988 Towards an understanding of gene switching in *Streptomyces*, the basis of sporulation and antibiotic production. *Proc. R. Soc. Lond.*, B235: 121-138
- Hopwood, D. A., K. F. Chater, J. E. Donding and A. Vivian 1973 Recent advances in *S. coelicolor* genetics. *Bacteriol. Rev.*, 37:371-405
- Hopwood, D. A., D. J. Lydiate, F. Malpartida, H. M. Wright 1984 Conjugative sex plasmids of *Streptomyces*. p. 615-634. In: D. R. Helinski, S. N. Cohen, D. B. Clewell, D. A. Jackson and A. Hollaender (ed.) Plasmids in bacteria. Plenum Publishing Corp., New York.
- Kieser, T., D. A. Hopwood, H. M. Wright and C. J. Thompson 1982 pIJ101, a multicopy broad host -range *Streptomyces* plasmid: function analysis and development of DNA cloning vectors. *Mol. Gen. Genet.*, 185: 223-238
- Kondo, S., K. Yasui, M. Natsume, M. Katayama and S. Marumo 1988 Isolation, physico-chemical properties and biological activity of pamamycin-607, an aerial mycelium-inducing substance from *Streptomyces alboniger*. *J. Antibiot.*, 41: 1198-1204
- Khokhlov, A. S. 1988 Results and perspectives of actinomycete autoregulators studies. p. 338-345. In: K. Okami, T. Beppu, H. Ogawara (ed.) Biology of actinomycetes '88, Japan Scientific Society Press, Tokyo.
- Maruo, S. 1987 The life cycle-regulating substances of microorganisms. *Nippon Nogeikagaku Kaishi*, 62: 1616-1621
- Miyoshi, Y-K., S. Ogata and S. Hayashida 1986 Multicopy derivative of pock-forming plasmid pSA1 in *Streptomyces azureus*. *J. Bacteriol.*, 168: 452-454
- Natsume, M., K. Yasui and S. Marumo S 1989 Calcium ion regulates aerial mycelium formation in actinomycetes. *J. Antibiot.*, 42: 440-447
- Ochi, K. 1990 *Streptomyces griseus* as an excellent object for studying microbial differentiation. *Actinomycetol.*, 4: 23-30
- Ochi, K., Y. Saito, K. Umehara, I. Ueda and M. Kohsaka 1984 Restoration of aerial mycelium and antibiotic production in a *Streptomyces griseoflavus* arginine auxotroph. *J. Gen. Microbiol.*, 130: 2007-2013
- Ogata, S. 1991 Morphological differentiation and pock formation in *Streptomyces*. *J. Antibact.*

- Antifung. Agents, 19: 405-412
- Ogata, S., Y-K. Miyoshi and S. Hayashida 1988 Characterization of a mutant thiostrepton-producing *Streptomyces azureus* ATCC 14921 and its plasmids. J. Ferment. Technol., 66: 213-217
- Ogata, S., H. S. Park, H. Suenaga and S. Yoshino 1989 Factors in the formation of spontaneously developing plaquelike pocks of *Streptomyces azureus* ATCC 14921. p. 115-119. In: Y. Koyama (ed.) Trends in Actinomycetology in Japan, Society for Actinomycetes, Japan.
- Ogata, S., S. Yamada, K. Doi, S. Tokunaga, H. Matsubara and T. Hara 1991 Inhibition of spore formation and thiostrepton production by accumulated intermediates of purine biosynthesis and the suppression of AICA (5-amino-4-imidazole carboxamide) in purine auxotrophic mutants of *Streptomyces azureus* ATCC 14921. J. Fac. Agr. Kyushu Univ., 35: 161-167
- Ogata, S., S. Yamada and S. Hayashida S 1985a Genetic recombination in *Streptomyces azureus* by protoplast fusion and high production of thiostrepton by the recombinants. J. Gen. Appl. Microbiol., 31: 187-191
- Ogata, S., S. Yamada and S. Hayashida 1985b Stimulation and inhibition of thiostrepton production in *Streptomyces azureus* by 5-amino-4-imidazole carboxamide (AICA). J. Gen. Microbiol., 31: 193-196
- Ogata, S., S. Yoshino, S. Suenaga, K. Aoyama, N. Kitajima and S. Hayashida 1981 Specific lysogenicity in *Streptomyces azureus*. Appl. Environ. Microbiol., 42: 135-141
- Penyige, A., G. Barabas, I. Szabo and J. C. Ensign 1990 ADP-ribosylation of membrane proteins of *Streptomyces griseus* strain 52-1. FEMS Microbiol. Lett., 69: 293-298
- Schauer, A., M. Ranes, R. Santamaria, J. Guijarro, E. Lawlor, C. Mendez, K. Chater and R. Losick 1988 Visualizing gene expression in time and space in the filamentous bacterium *Streptomyces coelicolor*. Science, 240: 768-772
- Vargha, G. Y. and G. Szabo 1984 Nutritional requirements for differentiation in a *Streptomyces fradiae* mutants. Zeit. Allgem. Mikrobiol., 24: 189-199