

## Effect of Intergeneric Grafting on Growth and Photosynthesis in *Momordica dioica* Roxb.

Mian, Mohammed Abdul Khaleque

Laboratory of Practical Botany, Faculty of Agriculture, Kyushu University

Morokuma, Masahiro

Laboratory of Practical Botany, Faculty of Agriculture, Kyushu University

Ali, Mohammad

Laboratory of Practical Botany, Faculty of Agriculture, Kyushu University

Agata, Waichi

Laboratory of Practical Botany, Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/24038>

---

出版情報：九州大学大学院農学研究院紀要. 38 (1/2), pp.65-71, 1993-12. Kyushu University  
バージョン：  
権利関係：

## Effect of Intergeneric Grafting on Growth and Photosynthesis in *Momordica dioica* Roxb.

Mohammed Abdul Khaleque Mian\*, Masahiro Morokuma', Mohammad Ali\*, Waichi Agata<sup>1</sup> and Kunimitsu Fujieda<sup>2</sup>

1 Laboratory of Practical Botany, Faculty of Agriculture, Kyushu University, 46-02, Fukuoka 812, Japan

2 Laboratory of Horticulture, Faculty of Agriculture, Kyushu University, 46-02, Fukuoka 812, Japan

(Received July 5, 1993)

Plants of *Momordica dioica* grafted onto rootstocks of five different Cucurbita species were evaluated with respect to growth and photosynthetic activities. Growth rate of *M. dioica* plants grafted on rootstocks of *C. ficifolia*, *C. moschata* 'No. 8', 'Tetsukabuto' (F<sub>1</sub> hybrid of *C. maxima* X *C. moschata*) and 'Shintosa 1 gou' (F<sub>1</sub> hybrid of *C. maxima* X *C. moschata*) was greater than that of non-grafted ones. Differential influence of rootstocks upon growth characteristics was also observed. The plants using the rootstocks except the Bangladesh local produced 2-3 times higher leaf area and dry matter than those of non-grafted plants at both early vegetative and flowering stages. The rootstocks played an important role in the suppressing tuberization of roots and in the enhancement of the scions. The results suggest that there is ample scope for modifying dry matter partitioning to increase of economic yield of this crop through grafting on other Cucurbita species.

### INTRODUCTION

*Momordica dioica* Roxb., a Cucurbitaceous crop, has edible fruits and is grown in Bangladesh and the north eastern part of India as a summer vegetable (Singh, 1990). It is generally propagated by root tuber although seed propagation is also possible (Ali et al., 1991). The plants propagated either by sexual or asexual means produce inedible roots for future propagation. Normally, the plants complete their life cycle within one year but the tuberous roots remain viable until the next summer when high temperatures and the high moisture content of the soil stimulate sprouting of new plants (Ali et al., 1991). Since the tuberous roots are a storage organ, the plants partition a large amount of photosynthates to the roots which limits the availability of the metabolic products for economic yield of fruits. It is assumed that, if the formation of root tubers is suppressed there would be an increase in the total above-ground biomass and an improvement in economic yield. It is not possible to achieve this goal through conventional breeding as there is a lack of non-tuber forming genotypes within *Momordica dioica* and there is incompatibility between this species and other closely related *Momordica* genotypes (Trivedi and Roy, 1972). One possible solution is to suppress tuberization by grafting *M. dioica* plants onto other non-root tuber forming plants. There are reports of successful intergenetic grafts in many cucurbits

---

\*Present address: Department of Genetics and Breeding, Institute of Postgraduate Studies in Agriculture Gazipur 1703, Bangladesh

(Imatsu, 1949; Lee, 1989; Uffelen, 1983) but there are no reports of grafting of *Momordica dioica* with any other crop species. In the present investigation, the effect on growth and photosynthetic activities of grafting *Momordica dioica* on other Cucurbita species was studied.

## MATERIALS AND METHODS

### Development of grafted plants

A cultivar of *M. dioica* from Bangladesh was grafted on five different rootstocks, namely 'Tetsukabuto' (an F<sub>1</sub> hybrid of *C. maxima* Duch X *C. moschata* Duch), 'Shintosa 1 gou' (another F<sub>1</sub> hybrid of *C. maxima* X *C. moschata*), *C. moschata* ('Bangladesh local'), *C. moschata* 'No. 8' and *C. ficifolia* Boucha. The stock plants were grown in vermiculite in mid-August, 1990 under glasshouse conditions at the laboratory's field of Horticulture, Kyushu University, Cleft grafting was made when the stock plants were at the cotyledonary leaf stage. Young shoot tips having three unopened leaves were collected from opposite sides of the lower end with a sharp blade. A straight cut, about 1 cm deep, was made between the two leaves of the rootstocks plant and the cut end of the scion was inserted. A plastic clip was attached at the joint to achieve a proper graft union. Fifty grafts were made with each of the rootstocks. The grafted plants were transplanted to small plastic pots (9 cm diameter) containing vermiculite, covered with polyethylene sheet and kept under shade for one week and sprayed with a mist water 3-4 times per day. The clips were removed after two weeks. When the graft union was established, shades were also removed so as to harden the plants. After one month the plants were transferred to larger pots (15 cm diameter) filled with a mixture of sand and volcanic gravels. For comparison, 20 non-grafted *M. dioica* plants were propagated as stem cuttings. All plants were grown in a plastic house at a temperature of 25-30°C. Growth characteristics and photosynthetic activities were measured at two stages, (41 days after grafting) and at flower bud initiation (81 days after grafting).

### Measurement of growth characteristics

Plant growth parameters including the number of fully expanded leaves (EL), length of vines (VL), number of internodes per vine (NI), and number of branches per vine (NB), were recorded on 10 plants selected randomly. Shoot dry weight (SDW) and root dry weight (RDW) were recorded on three plants from each treatment, with the exception of plants grafted on *C. moschata* ('Bangladesh local') as insufficient plants were available at the flowering stage. The shoots and roots of the sample plants were separated and oven dried at 80 °C for 48 hours. Total leaf area per plant (TLA), and mean leaf area (MLA) on a single plant per treatment were measured using an automatic leaf area meter (model AAM-8).

### Measurement of photosynthetic activities

Net photosynthetic rate (Pn), transpiration rate (E), stomatal conductance (Gs) and intercellular CO<sub>2</sub> concentration (Ci) were determined by a portable infrared gas analyzer and data logger (ADC, model-SPS-H, UK). Measurements were made on at least three fully expanded mature leaves between the 5th and the 9th nodes from the

top of five different plants, under a solar photon flux of about  $1500 \mu\text{E m}^{-2}\text{s}^{-1}$ , and the parameters were calculated using the method of Von Caemmerer and Farquhar (1981). Root respiration was also measured by infrared gas analyzer at a temperature of 30 °C.

## RESULTS

All the growth parameters of *M. dioica* at both the early vegetative and flower initiation stages were strongly influenced by grafting (Table 1 and 2). At the early stage, the shoot dry weight of the grafted plants was significantly greater than on the non-grafted plants. The difference between non-grafted and grafted plants was even greater at the flower bud initiation stage except for those using *C. moschata* ('Bangladesh local') as a rootstock. The plants grafted on Bangladesh rootstock grew very poorly and at flowering, all the plants became yellow and ultimately died through rotting of the roots. The grafted plants produced significantly more biomass than the non-grafted ones. At the early growth stage, the difference in above-ground dry matter production between grafted and non-grafted plants was marked, although the root dry matter weights were similar at that stage.

At flower initiation, the grafted plants had the highest shoot dry weight but the non-grafted plants had the highest root dry weight (Table 2). Some root tubers were observed in non-grafted plants, while tuber formation was completely suppressed in plants grafted onto rootstocks having a normal root forming habit.

Both total leaf area per plant and mean leaf area per one leaf were greatly influenced by the rootstocks at the early flowering stages but the effect was more pronounced at the later stage (Table 2).

Grafting did not have a consistent effect on net photosynthetic rate (Table 3). The rootstock, 'Tetsukabuto' produced the highest rate of net photosynthesis at both the early vegetative and flowering stages. In general, irrespective of treatment, the rate of photosynthesis was higher at the early stage. At the flowering stage, the plants using rootstock *C. moschata* ('Bangladesh local') gave aberrant results because of the abnormal growth of these plants. The effect of rootstocks on transpiration rate and stomatal conductance were similar to those for photosynthesis. The relative effect of rootstock on net photosynthetic rate, respiration rate and stomatal conductance were similar at both the stages. The intercellular  $\text{CO}_2$  concentration was not influenced by rootstock at the developmental stage. However, at the flowering stage, the  $\text{CO}_2$  concentration in grafted plants was comparatively higher than that in non-grafted ones in spite of the concentration was lower. The rate of root respiration, particularly in non-grafted plants, was the lowest during tuber forming stage though it was the highest at the early vegetative stage (Table 3).

## DISCUSSION

The results obtained in this experiment revealed that *M. dioica* was graft compatible. Those rootstocks that are used extensively for grafting Cucumis in Japan and other countries (Lee, 1989) also performed well with *M. dioica*. The poor

**Table 1.** Effect of grafting on growth characteristics in *Momordica dioica* at early vegetative stage (41 days after grafting).

Rootstocks	Scion	EL	NI	NB	VL	TLA	MLA	SDW	RDW	% of RDW
		(No./plant)			(cm)	(cm <sup>2</sup> )	(cm <sup>2</sup> )	(g)	(g)	to TDW
Non-grafted	<i>M. dioica</i>	9.5* ± 1.9	9.7 ± 1.6	0	38.7 ± 8.3	332	25.5 ± 9.6	1.23 ± 0.22	0.15	10.37 f0.06
'Tetsukabuto' <sup>1</sup>	''	20.3 ± 6.0	16.7 ± 4.0	0.8 ± 1.0	70.1 ± 20.3	916	38.1 ± 15.9	2.42 ± 0.56	0.22	8.33 ± 0.02
'Shintosa 1 gou' <sup>2</sup>	''	18.6 ± 4.8	16.9 ± 3.1	0.6 ± 0.7	75.5 ± 20.8	700	30.7 ± 14.3	1.98 ± 0.55	0.17	7.91 ± 0.04
<i>C. moschata</i> <sup>3</sup>	''	19.0 ± 4.6	18.1 ± 2.0	0.8 ± 1.0	77.2 f18.5	445	21.1 ± 10.8	1.68 ± 0.60	0.13	7.18 ± 0.01
<i>C. moschata</i> 'No. 8'	''	20.3 ± 4.4	16.4 ± 2.6	1.4 ± 0.9	60.6 f17.2	799	39.3 ± 15.5	1.98 ± 0.37	0.22	10.00 f0.04
<i>C. ficifolia</i>	''	20.3 ± 4.0	17.5 ± 2.4	0.7 ± 0.8	73.9 ± 21.1	585	27.9 ± 11.8	2.39 ± 0.47	0.25	7.86 ± 0.04

1 and 2: F<sub>1</sub> hybrids of *C. maxima* × *C. moschata* 3: 'Bangladesh local' EL: Number of expanded leaves, NI: Number of internode, NB: Number of branches, VL: Vine length, TLA: Total leaf area, MLA: Mean leaf area, SDW: Shoot dry weight, RDW: Root dry weight, TDW: Total dry weight, \*: mean ± S.E.

**Table 2.** Effect of grafting on growth characteristics in *Momordica dioica* at flower initiation stage (81 days after grafting).

Rootstocks	Scion	EL	NI	NB	VL	TLA	MLA	SDW	RDW	% of RDW
		(No./plant)			(cm)	(cm <sup>2</sup> )	(cm <sup>2</sup> )	(g)	(g)	to TDW
Non-grafted	<i>M. dioica</i>	28.3* ± 5.6	25.7 ± 1.3	0.3 ± 0.5	148.0 ± 4.3	909	43.3 f15.1	5.25 ± 0.47	2.10	28.67 f0.54
'Tetsukabuto''	''	63.0 ± 13.6	35.0 ± 2.9	3.0 ± 0.8	192.3 ± 11.5	2542	72.7 f20.1	16.73 ± 2.60	0.65	3.74 10.12
'Shintosa 1 gou' <sup>2</sup>	''	80.7 f19.4	34.3 ± 5.5	3.3 ± 2.1	221.0 t21.2	1896	75.9 f19.5	14.57 ± 2.50	0.74	4.83 f0.24
<i>C. moschata</i> <sup>3</sup>	''	20.0	18.0	1.0	74.0	318	21.2 ± 7.0	2.50	0.13	4.94
<i>C. moschata</i> 'No. 8'	''	67.7 F14.4	30.7 ± 6.2	2.7 ± 1.0	151.3 ± 16.7	1812	47.7 ± 16.0	11.53 ± 3.56	0.91	7.32 ± 0.08
<i>C. ficifolia</i>	''	54.7 ± 2.6	31.3 ± 8.9	3.0 ± 2.5	153.3 ± 24.2	2012	49.1 t17.2	11.90 ± 0.65	0.89	6.96 ± 0.34

1 and 2: F<sub>1</sub> hybrids of *C. maxima* × *C. moschata* 3: 'Bangladesh local' EL: Number of expanded leaves, NI: Number of internode, NB: Number of branches, VL: Vine length, TLA: Total leaf area, MLA: Mean leaf area, SDW: Shoot dry weight, RDW: Root dry weight, TDW: Total dry weight, \*: mean ± S.E.

Table 3. Effect of grafting on net photosynthetic rate (Pn), transpiration rate (E), stomatal conductance (Gs), intercellular CO<sub>2</sub> concentration (Ci) and root respiration (RR) at early vegetative and flower initiation stages of *Momordica dioica*.

		Early vegetative stage (41 days after grafting)				
Rootstocks	Scion	Pn ( $\mu\text{molm}^{-2}\text{s}^{-1}$ )	E ( $\text{mmolm}^{-2}\text{s}^{-1}$ )	Gs ( $\text{molm}^{-2}\text{s}^{-1}$ )	Ci (ppm)	RR ( $\text{CO}_2\text{mgDWg}^{-1}\text{h}^{-1}$ )
Non-grafted	<i>M. dioica</i>	11.20* ± 0.55	6.99 ± 0.75	0.59 ± 0.16	296.7 ± 5.9	19.42
'Tetsukabuto'	"	12.09 ± 1.38	6.55 ± 0.80	0.72 ± 0.27	296.5 ± 8.6	10.78
'Shintosa 1 gou' <sup>2</sup>	"	8.66 ± 0.80	4.08 ± 0.52	0.31 ± 0.07	292.2 ± 11.2	17.25
<i>C. moschata</i> <sup>3</sup>	"	8.82 ± 2.27	4.40 ± 1.39	0.36 ± 0.19	288.1 ± 21.6	4.98
<i>C. moschata</i> 'No. 8'	"	9.42 ± 2.73	4.36 ± 1.01	0.36 ± 0.15	291.0 ± 9.8	7.47
<i>C. ficifolia</i>	"	10.20 ± 0.88	5.20 ± 0.43	0.51 ± 0.41	299.3 ± 4.3	8.82
		Flower initiation stage (81 days after grafting)				
Non-grafted	<i>M. dioica</i>	8.05* ± 2.70	2.00 ± 0.60	0.10 ± 0.04	194.5 ± 6.7	4.41
'Tetsukabuto'	"	9.11 ± 1.50	2.98 ± 0.76	0.18 ± 0.06	239.7 ± 12.3	7.50
'Shintosa 1 gou' <sup>2</sup>	"	6.84 ± 1.67	1.99 ± 0.37	0.09 ± 0.02	213.3 ± 1.8	7.50
<i>C. moschata</i> <sup>3</sup>	"	-0.09	0.24	0.01	362.2	2.56
<i>C. moschata</i> 'No. 8'	"	6.75 ± 2.50	2.20 ± 0.90	0.11 ± 0.07	223.2 ± 14.3	6.66
<i>C. ficifolia</i>	"	4.44 ± 2.99	1.66 ± 1.10	0.07 ± 0.06	223.9 ± 10.5	11.42

1 and 2: F<sub>1</sub> hybrids of *C. maxima* × *C. moschata*, 3: 'Bangladesh local' \*: mean ± S.E.

performance of *C. moschata*, a 'Bangladesh local' rootstock might be due to post-graft incompatibility. Such differential graft compatibility was also reported by Marukawa and Onuma (1967) and Marukawa and Yamamuro (1967) who grafted melon and water melon on different Cucurbita species. The plants with Japanese rootstocks produced 2 - 3 times more total leaf area and dry matter production per plant than non-grafted ones at both the early and flowering stages.

The most interesting results was that root dry matter accumulation increased dramatically at the flowering stage in non-grafted plants (Table 2). At that stage, the non-grafted plants diverted about 29% of the total photosynthates to the roots for tuber formation, while the grafted plants diverted only 3-7%. Root respiration in non-grafted plants also declined markedly in spite of increasing of root dry weight. Such partitioning of photosynthates and the low root respiration at the flowering stage of the non-grafted plants helped to accumulate greater dry weight in the roots. Tuber formation was completely suppressed in grafted plants, while there was the opportunity to transport more than 90% of the photosynthates to the shoot. It is also possible that the rootstocks are inherently more efficient in absorbing and supplying nutrients to the scions.

The suppression of the tuber forming habit, might be the cause of the higher growth performance as well as more dry matter production in grafted plants.

This is the first report of grafting *M. dioica* grafted on other Cucurbita species. It is clearly important to know whether grafting improves yield of *M. dioica*. So the effect of grafting on economic yield will be dealt with in the next paper. Lee (1989) has observed that Cucumis and Cucurbita species performed better when grafted on Cucurbita rootstocks and Uffelen (1983) reported superior performances when Cucurbita species were used as rootstocks. There is other evidence that Thorne and Evans (1964) obtained higher net assimilation rate in spinach beet plants grafted on sugar beet rootstocks.

In conclusion, the findings of the present investigation suggest that there is the potential for improving the productivity of *M. dioica* through grafting with other Cucurbita species.

#### ACKNOWLEDGEMENT

We are grateful to Dr. Fumitake Kubota, Associate Professor, Laboratory of Practical Botany, Kyushu University for his cooperation and assistance during this study.

#### REFERENCES

- Ali, M., H. Okubo, T. Fujii and K. Fujieda 1991 Techniques for propagation and breeding of kakrol (*Momordica dioica* Roxb.). *Scientia Hort.*, 47: 335-343
- Imatsu, T. 1949 On the symbiotic affinity caused by the grafting among Cucurbitaceous species. *J. Japan Soc. Hort. Sci.*, 19: 36-42 (In Japanese with English summary)
- Lee, J. M. 1989 On cultivation of grafted plants of Cucurbitaceous vegetables. *J. Kor. Soc. Hort.*

- Sci.*, 30(3):169-179
- Marukawa, S. and K. Onuma 1967 Studies on the selection of Cucurbita spp. as melon stock. I. On the problems of a) approach-grafting, b) the handling of the foliage on the stock and the roots of the melon top and c) whether the melon top is influenced by the growing ability of Cucurbita spp. as stock in low temperatures. *Bull. Ibaragi Hort. Expt. Stn.*, No. 2: 35-42 (In Japanese with English summary)
- Marukawa, S. and K. Yamamuro 1967 Studies on the selection of Cucurbita spp. as water melon stock. II. Compatibility of Cucurbita spp. and varieties. *Bull. Ibaragi Hort. Expt. Stn.*, No. 2: 29-34. (In Japanese with English summary)
- Singh, A. K. 1990 Cytogenetics and evolution in the Cucurbitaceae. In "Biology and utilization of the Cucurbitaceae". by D. M. Bates, R. W. Robinson and C. Jeffrey, Cornell University Press, Ithaca and London, pp. 18-19
- Thorne, G. N. and A. F. Evans 1964 Influence of tops and roots on net assimilation rate of sugar-beet and spinach beet grafts between them. *Ann. Bot.*, 28: 499-508
- Trivedi, R. N. and R. P. Roy 1972 Cytological studies in some species of Momordica. *Genetica*, 43: 282-291
- Uffelen, J. A. M. Van 1983 Rootstocks for grafting cucumbers. *Hort. Abst.*, 53(9): 631
- Von Caemmerer, S. and G. D. Farquhar 1981 Some relationships between the biochemistry of photosynthesis and gas exchange of leaves. *Planta*, 153: 376-387