

DISCOVERY OF THE NEARCTIC GENUS CHASMATONOTUS  
LOEW (DIPTERA, CHIRONOMIDAE) FROM JAPAN,  
WITH DESCRIPTIONS OF THREE NEW SPECIES

Yamamoto, Masaru

<https://doi.org/10.5109/2404>

---

出版情報 : ESAKIA. 15, pp.79-96, 1980-10-31. Entomological Laboratory, Faculty of Agriculture,  
Kyushu University  
バージョン :  
権利関係 :

**DISCOVERY OF THE NEARCTIC GENUS CHASMATONOTUS LOEW  
(DIPTERA, CHIRONOMIDAE) FROM JAPAN, WITH  
DESCRIPTIONS OF THREE NEW SPECIES\***

MASARU YAMAMOTO

Entomological Laboratory, Faculty of Agriculture  
Kyushu University, Fukuoka 812, Japan

**Abstract**

The chironomid genus *Chasmatonotus*, which has been known from the Nearctic Region, is reported from Japan for the first time. Three new Japanese species, *Chasmatonotus unilobus*, *Chasmatonotus parabolicolor*, and *Chasmatonotus saigusai*, are described and figured.

The genus *Chasmatonotus* was proposed by Loew in 1864 for a single species *C. unimaculatus* from New Hampshire, U. S. A. After that, Osten Sacken described *C. bimaculatus* in 1877 from Quebec. Coquillett described *C. univittatus* in 1900 from Alaska, and *fascipennis* from British Columbia and *hyalinus* from California in 1905. In 1937 Rempel revised the genus and described 3 additional species, i.e., *atripes* from New York, *bicolor* from Tennessee and *maculipennis* from Washington. He pointed out that the species identified by Johanssen (1905) and Malloch (1915) as *bimaculatus* O. S. was *atripes*. He also stated that *maculipennis* is probably a Palaearctic species, because the specimens on which his description was based were those intercepted in quarantine at Seattle in Washington on Japanese maple from Japan. Thus the genus is hitherto represented by eight species which are distributed exclusively in the Nearctic Region.

Some twenty years ago Prof. T. Saigusa of Kyushu University collected several specimens of curious chironomid midges at the subalpine to alpine zones of Central Mountain Region of Honshu, and he granted them to me. In 1977 I obtained many specimens of a similar midge at the subalpine zone of Amô Tôge, Gifu Pref., Honshu. When he visited Kyushu University in 1978, Prof. J. E. Sublette of Eastern New Mexico University suggested me that these midges belong to the genus *Chasmatonotus*. After a close examination of

---

\* Contribution from the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka (Ser. 3, No. 79).

the male genitalia of these midges, I found out that they represent three species. All of these Japanese species are closely allied to *bicolor* not only in the wing marking but also in the general structure of the male genitalia, but are very distinct in detailed structure of the latter. Therefore I will describe them as new species in the following lines. I do not still find *maculipennis* in Japan which was suggested to be introduced from Japan to Washington, as stated before. However, thanks to Prof. J. E. Sublette, it became clear that the genus *Chasmatonotus* is also distributed in the Eastern Palaearctic Region.

The genus *Chasmatonotus* is readily distinguishable from any other chironomid genera by having a median longitudinal fissure on the mesoscutum and for Japanese species by having the five antennal flagellomeres in both sexes. The latter character is also found in male of African genus *Harrisonina* Freeman.

Japanese species of the genus *Chasmatonotus* are found in the mountainous regions of the Japan Alps and neighbouring mountains, especially at the moist herbaceous places in woods. Prof. T. Saigusa and I observed that the midges were actively walking about on the ground and on the herb, but did not observe them on wings, although they have well developed wings. I also observed the species collected at Am8 Tôge was mating on the ground or on the herb near the high moor in woods. So far as I know, the immature stages of this genus have not been clarified but I think they are probably terrestrial considering the environment in which they live, so far as the Japanese species are concerned.

Before going further I express my grateful thanks to Prof. Y. Hirashima and Assoc. Prof. K. Morimoto, Entomological Laboratory, Faculty of Agriculture, Kyushu University for their constant guidances. I wish to express my cordial thanks to Prof. T. Saigusa, Biological Laboratory, College of General Education, Kyushu University, for his kind guidance in preparing this paper and giving me an opportunity to examine the interesting materials. My thanks are also due to Prof. J. E. Sublette, Eastern New Mexico University, U. S. A., for his kind suggestion on the systematic position of the midges. I am much indebted to Mr. K. Ôhara, Kyushu University, for valuable specimens.

#### Genus *Chasmatonotus* Loew

*Chasmatonotus* Loew, 1864: *Bed. ent. Z.* 8: 50. Type-species : *Chasmatonotus unimaculatus* Loew, 1864.

Three new Japanese species of *Chasmatonotus* described below unexceptionally bear the generic characters mentioned by Rempel (1937) excepting the presence of the acrostichal setae and 5 flagellomeres, instead of 6, of antennae. The acrostichal setae which are found in Japanese species are so minute that I suppose Rempel probably overlooked them on the Nearctic species.

The characteristics of Japanese species are summarized as follows:

Head (Figs. Z-A, -B) small, placed under antepronotum; compound eye small, semicircular, widely separated from each other; prefrons and area surrounding antennal scape membranous; antennal flagellum composed of 5 flagellomeres in both sexes; gena with 0-6 setae; maxillary palpus with 5 segments but 1st segment membranous and indistinct.

Thorax (Figs. 3-A, -B) rather long; antepronotum well developed, deeply divided in middle; scutum with a deep median longitudinal fissure which sparsely bears very minute acrostichal setae; dorsocentral setae irregularly uni- to triserial, stouter and longer on prescutellar area; supra-alar setae present or absent; scutellar setae in a double row; preepisternum with 3-18 setae on upper area; anapleural suture disappearing in middle.

Legs long and stout; fore-, middle and hind tibiae with 1, 2 and 2 apical spurs respectively which bear scales at basal 1/2; apical spurs of middle tibia subequal in length; outer spur of hind tibia longer than inner one; basal two tarsomeres of middle and hind legs provided with a pair of spines apically which are bare and subequal in length; tibial comb of hind leg composed of 10-16 spines; pulvilli well developed and pad-like; empodium long and filiform.

Wing black with a white basal marking which extend into a narrow stripe; anal lobe not produced; squama unfringed but with 0-5 sparse setae;  $R_{2+3}$  obsolete or obvious, variable within a species; M ending slightly beyond tip of  $R_{4+5}$ ; fCu below r-m; costa, R,  $R_1$  and  $R_{4+5}$  with prominent suberect setae.

Male genitalia with gonopod variable in shape; epandrium flattened, with anal point short and stout but prominent; gonocoxite with or without basal lobes, inner surface of ventrodiscal portion of gonocoxite with some short setae or long curved setae; gonostylus finger-shaped with a basal projection or subtriangular, and with a few stout spines near apex. Ninth abdominal tergite of female deeply emarginate in middle; laterosternite with a prominent apical projection. Spermatheca 2 in number, large.

#### KEYS TO JAPANESE SPECIES OF THE GENUS *Chasmatonotus*

##### MALES

- Gonostylus slightly tapering, almost parallel-sided in lateral aspect, with a ventrally produced basal projection; gonocoxite with prominent basal lobes; anal point short, with dorsal margin arched or almost straight ..... 2
- Gonostylus subtriangular in shape; basal lobe of gonocoxite only slightly swollen; anal point moderately long, blade-shaped, with dorsal margin weakly emarginate ..... *Saigusai* sp. nov.
2. Gonocoxite in ventral aspect wide and nearly parallel-sided, with inner margin much expanding inwardly, almost straight; basal projection of gonostylus bilobed at tip; anal point with dorsal margin arched ..... *unilobus* sp. nov.
- Gonocoxite in ventral aspect tapering towards tip, with inner margin not expanding inward but strongly angulate in middle; basal projection of gonostylus simple; anal point with dorsal margin almost straight ..... *parabicolor* sp. nov.

## FEMALES

1. Basal part of laterosternite simple, without a basal lobe; apical projection of laterosternite long, almost extending to posterior margin of tergite IX ..... 2  
 - Basal part of laterosternite strongly projected into a lobe, apical projection of laterosternite short, ending much before posterior margin of tergite IX .....  
 ..... **unilobus** sp. nov.
2. Apical projection of laterosternite stout, 8.7~ as long as thick in lateral aspect ..... **saiguai** sp. nov.  
 - Apical projection of laterosternite slender, 17.8× as long as thick in lateral aspect ..... **parabicolor** sp. nov.





Fig. 1. *Chasmatonotus unilobus* sp. nov., ♂.

***Chasmatonotus unilobus* sp. nov.**

♂. Colouration : Head subshining brownish black; antennal pedicel brownish black, thinly grey pollinose, flagellum brown; clypeus, maxillary and labial palpi brownish black. Thorax with antep pronotum brown with light yellow tinge to brownish black, postpronotum brown, sometimes with yellowish tinge; mesonotum subshining brownish black; scutum yellowish brown at anterolateral corner or entirely brownish black; scutellum and postnotum entirely brownish black. Thoracic pleura subshining brownish black, thinly grey pollinose.


Legs including coxae predominantly subshining brownish black, thinly grey pollinose ; all trochanters brown, slightly tinged with yellow; extreme bases of all femora slightly tinged with yellow; spurs of all tibiae black, spines of 1st to 2nd tarsomeres of middle and hind legs black. Wing strongly infuscated, with blackish veins ; apical 2/5 of cell  $R_s$ , apical 1/4 of cell M, apical 1/3 of cell Cu, apical 1/2 of cell A behind 2A much darkened, blackish; basal 1/4 of wing whitish from M to hind margin; this white area produced into a narrow stripe which extends to apical 3/4 of wing. Halter yellow. Abdomen predominantly brownish black, thinly grey pollinose; basal 2 terga and basal 3 sterna yellow. Genitalia brownish black, thinly grey pollinose.

Head: Antennal flagellomere lengths ( $\mu$ ) : 96, 43, 38, 40, 66. First to 5th flagellomeres with 12-17, 5-6, 5-6, 6-7, 5-7 setae respectively, Antennal ratio, range 0.26-o. 36, mean 0.30. Second to 5th palpal segment lengths ( $\mu$ ): 64, 126, 123, 171, and the palpal segments with 7-13, 37-49, 28-42, 20-31 setae respectively, Vertex, clypeus and gena with 24-42, 13-23, 1-3 setae respectively.



**Fig. 2.** *Chasmatonotus unilobus* sp. nov. A: head, frontal aspect, ♂. B: same, lateral aspect. C: antenna, ♂. D: same, ♀. E: tip of hind tibia, ♂.

Thorax : Lateral anteprenotals 7-11, median anteprenotals 7-12 ; dorsocentrals 11-23, irregularly biserial ; acrostichals 11-17, prealars 5-11, supra-alars 0-2 ; scutellars 17-20, biserial. Preepisternum with 6-10 setae.



**Fig. 3.** Thorax of *Chasmatonotus unilobus* sp. nov., ♂. A: lateral aspect. B: dorsal aspect.

Legs : Fore-, middle and hind coxae with 28-36, 12-23, 23-32 apical setae respectively. Middle coxa with 9-10 minute sensory hairs on middle portion of outer surface. Fore-, middle and hind trochanters with 26-34, 23-32, 23-31 apical setae respectively. Outer spur of hind tibia 1.6-l. 9× as long as inner one. Tibial comb of hind leg composed of 13-14 spines.

Lengths (means, 0.01 mm in unit) and proportions (ranges and means) of legs :

	F	T	Tal	Ta2	Ta3	Ta4	Ta5	BV	SV	LR
P <sub>1</sub>	116	116	74	43	30	20	15	2.74-2.96, 2.87	2.94-3.37, 3.12	0.61-o.66, 0.64
P <sub>2</sub>	109	113	48	28	21	14	13	3.33-3.79, 3.52	4.26-4.89, 4.57	0.40-O.49, 0.43
P <sub>3</sub>	127	135	70	37	28	15	14	3.44-3.61, 3.52	3.62-3.96, 3.72	0.50-O.54, 0.52

Wing (Fig. 4-A) : Length 1.9-2.1 mm. Width 0.6-O. 7 mm. L/WR, range 3.04-3.38, mean 3.23. VR, range 0.94-1.03, mean 1.00. R, R<sub>1</sub> and R<sub>4+5</sub> with 31-53, 11-22, 26-41 setae respectively. Radialis with 7-9 annular organs on its basal part, 1-2 annular organs on middle anteriorly, 5-9 annular organs on apical part, and with 3-6 setae near middle. Squama with 0-3 setae.


Genitalia (Figs. 6-A, -B, -C, -D) : Anal point (Fig. 5-A) short, arched dorsally at basal 1/3. Gonocoxite comparatively wide and nearly parallel-sided in ventral aspect, apical to inner margin of its ventral surface much expanded

and covering its ventrodiscal surface and base of gonostylus; inner surface of ventrodiscal portion with about 10 short setae; dorsal part of basal lobe only slightly swollen, ventral part of basal lobe projected into a short, slender process. Gonostylus moderately long, weakly bent inwardly in middle; basal part strongly projected inwardly, bilobed at tip; extreme tip with a strong spine.

Length of body: 1.8-2.8 mm.

♀. Colouration almost as in male but thorax including antepnotum entirely brownish black. All trochanters and basal parts of all femora darker than in male. Cercus yellow.

Head: Antennal flagellomere lengths ( $\mu$ ): 116, 45, 40, 41, 76. First to 5th flagellomeres with 10-21, 5-6, 6-7, 6-7, 5-10 setae respectively. Antennal ratio, range 0.28-0.33, mean 0.31. Second to 5th palpal segment lengths ( $\mu$ ): 77, 150, 150, 186, and the palpal segments with 8-15, 42-56, 38-53, 25-41 setae



**Fig. 4.** Male wings of *Chasmatonotus* spp. A: *Chasmatonotus unilobus* sp. nov. B: *Chasmatonotus parabolicolor* sp. nov. C: *Chasmatonotus saigusai* sp. nov.






Fig. 5. Anal point (A-C) and gonostylus (D) of *Chasmatonotus* spp. A: *Chasmatonotus unitobus* sp. nov., lateral aspect. B: *Chasmatonotus parabolicolor* sp. nov., lateral aspect. C: *Chasmatonotus saigusai* sp. nov., lateral aspect. D: *Chasmatonotus saigusai* sp. nov., dorsal aspect.

respectively. Vertex, clypeus and gena with 28-62, 19-30, 2-6 setae respectively.


Thorax: Lateral anteprenotals 11-19, median anteprenotals 7-14, dorsocentrals 26-31, irregularly bi- to triserial; acrostichals 9-21, prealars 5-13, and no supra-alar; scutellars 22-32, biserial. Preepisternum with 4-18 setae.

Legs: Fore-, middle and hind coxae with 33-50, 17-28, 32-47 apical setae respectively. Middle coxa with 9-11 minute sensory hairs on its middle portion of outer surface. Fore-, middle and hind trochanters with 30-45, 26-42, 30-37 apical setae respectively. Outer spur of hind tibia 1.6-1.8x as long as inner one. Tibia1 comb of hind leg composed of 14-16 spines.


Lengths (means, 0.01 mm in unit) and proportions (ranges and means) of legs:

	F	T	Ta1	Ta2	Ta3	Ta4	Ta5	BV	s v	LR
P <sub>1</sub>	123	127	75	41	30	21	16	2.95-3.24, 3.04	3.24-3.39, 3.32	0.57-0.61, 0.60
P <sub>2</sub>	119	123	48	27	23	15	14	3.36-3.80, 3.65	5.00-5.28, 5.10	0.37-0.40, 0.39
P <sub>3</sub>	139	150	75	40	31	17	12	3.04-3.69, 3.53	3.00-4.16, 3.85	0.45-0.60, 0.50

Wing: Length 2.2-2.7 mm. Width 0.7-0.9 mm. L/WR, range 2.76-3.10, mean 2.96. VR, range 0.90-1.06, mean 1.00. R, R<sub>1</sub> and R<sub>4+5</sub> with 42-64, 13-20, 31-42 setae respectively. Radialis with 7-9 annular organs on its basal part,



**Fig. 6.** Male genitalia of *Chasmatonotus unilobus* sp. nov. A: epandrium, gonocoxite and gonostylus in dorsal aspect. B: gonocoxite and gonostylus in dorsal aspect. C: same, inner lateral aspect. D: hypandrium, gonocoxite and gonostylus in ventral aspect.



**Fig. 7.** Female genitalia of *Chasmatonotus unilobus* sp. nov. **A:** lateral aspect. **B:** ventral aspect.

1-3 annular organs on middle anteriorly, 4-7 annular organs on apical part, and with 4-7 setae near middle. Squama with 1-5 setae.

Genitalia (Figs. 7-A, -B) : Laterosternite strongly lobate basally, its apical part with comparatively short projection which bears about 10 setae on apical 1/2. Coxite VIII with 10-17 long setae.

Length of body: 1.9-2.8 mm.

DISTRIBUTION : Japan (Honshu) .

TYPE MATERIAL: Holotype ♂ (Type No. 2176, Kyushu Univ.), Amô (1400 m), Hida-Kawai, Gifu Pref., Honshu, 12. vi. 1977, M. Yamamoto leg. Paratypes : 59 ♂♂, 32 ♀♀, same data as holotype ; 9 ♂♂, 11 ♀♀, same locality and date, K. Ôhara leg.

The holotype is deposited in the collection of the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka.

REMARKS: This species can be distinguished at a glance from two other Japanese species in having the widened and parallel-sided gonocoxite, very simple basal lobe and apically bilobed basal projection of the gonostylus in male. The female genitalia of the new species is very characteristic in having the strong basal lobe on the laterosternite.

The type-specimens were found at the moist herbaceous place near the high moor in woods.

*Chasmatonotus parabolicolor* sp. nov.

♂. Colouration: Head, thorax and legs almost as in *Chasmatonotus unilobus* sp. nov., but anteprenotum paler. Wing marking resembling *unilobus*, but the ground colour of dark area more uniformly brown, apical 1/5 of cell M and middle of cell A slightly darker. Abdomen brownish black, thinly grey pollinose; basal 4 segments yellow to yellowish brown, darker posteriorly. Genitalia brownish black, thinly grey pollinose.

Head : Antennal flagellomere lengths ( $\mu$ ) : 80, 38, 33, 35, 78. First to 5th flagellomeres with 11-12, 4, 4, 5, 7-9 setae respectively. Antennal ratio 0.42. Second to 5th palpal segment lengths ( $\mu$ ) : 73, 139, 143, 184, and the palpal segments with 9-16, 37-43, 26-36, 16-25 setae respectively. Vertex, clypeus and gena with 16-21, 16-23, 0-2 setae.

Thorax : Lateral anteprenotals 11-12, median anteprenotals 5-7 ; dorsocentrals 21-27, irregularly biserial ; acrostichals S-15, prealars 6-9, no supra-alar ; scutellars 20-21, biserial. Preepisternum with 3-6 setae.


Legs : Fore-, middle and hind coxae with 20-30, 11-17, 21-26 apical setae respectively. Middle coxa with 9 minute sensory hairs on its middle portion of outer surface. Fore-, middle and hind trochanters with 36-48, 30-39, 31-35 apical setae respectively. Outer spur of hind tibia 1.5-l. 6x as long as inner one. Tibial comb of hind leg composed of 12-14 spines.

Lengths (means, 0.01 mm in unit) and proportions (ranges and means) of legs :

	F	T	Tal	Ta2	Ta3	Ta4	Ta5	BV	sv	LR
P <sub>1</sub>	117	116	76	42	29	19	15	2.84-3.02, 2.93	3.07-3.20, 3.15	0.62-0.65, 0.64
P <sub>2</sub>	108	112	47	27	21	14	13	3.34-3.79, 3.59	4.53-4.79, 4.70	0.41-0.43, 0.42
P <sub>3</sub>	129	136	70	39	29	15	14	3.30-3.63, 3.45	3.71-3.90, 3.78	0.50-0.53, 0.52

Wing (Fig. 4-B) : Length 1.8-2.0 mm. Width 0.5-0.6 mm. L/WR, range 3.20-3.45, mean 3.35. VR, range 0.94-1.00, mean 0.98. R, R<sub>1</sub> and R<sub>4+5</sub> with 26-37, 9-14, 21-28 setae respectively. Radialis with 8-10 annular organs on its basal part, 2 annular organs on middle anteriorly, 3-7 annular organs on apical part, and with 3-7 setae near middle. Squama without setae.

Genitalia (Figs. 8-A, -B, -C, -D) : Anal point (Fig. 5-B) short, with dorsal margin almost straight. Gonocoxite strongly tapering towards apex, inner margin of its ventral surface simple on apical 1/2, not expanding inwardly, therefore inner surface of ventrodistal portion visible from beneath, the surface bearing some long curved setae ; inner margin of ventral surface strongly angulate in middle and with a transverse elevation running dorsally from the angle; dorsal part of basal lobe slightly developed and with some setae, ventral part of basal lobe projected into a short, rather broad process which



**Fig. 8.** Male genitalia of *Chasmatonotus parabolicolor* sp. nov. A: epandrium, gonocoxite and gonostylus in dorsal aspect. B: gonocoxite and gonostylus in dorsal aspect. C: same, inner lateral aspect. D: hypandrium, gonocoxite and gonostylus in ventral aspect,

bears some setae near apex. Gonostylus resembling that of *unilobus* in general appearance, but basal projection not bilobed, a longitudinal elevation running from tip of the basal projection; extreme tip with a strong spine.

Length of body: 2.0-2.3 mm.

♀. Colouration almost as in male. Abdomen pale brown; basal 2 segments yellow, following segments pale brown, darkened towards tip; cercus yellow.

Head: Antenna1 flagellomere lengths ( $\mu$ ): 103, 45, 34, 43, 85. First to 5th flagellomeres with 11, 5, 6, 5, 7 setae respectively. Antenna1 ratio 0.38. Second to 5th palpal segment lengths ( $\mu$ ): 40, 75, 98, 120, and the palpal segments with 16, 50, 43, 18 setae respectively. Vertex, clypeus and gena with 20, 22, 2 setae respectively.

Thorax : Lateral anteprenotals 16, median anteprenotals 10 ; dorsocentrals 32, irregularly biserial ; acrostichals 11, prealars 12, no supra-alar; scutellars 23, biserial. Preepisternum with 14 setae.

Legs: Fore-, middle and hind coxae with 32, 17, 25 apical setae respectively. Middle coxa with 8 minute sensory hairs on its middle portion of outer surface. Fore-, middle and hind trochanters with 48, 39, 33 apical setae respectively. Outer spur of hind tibia  $1.7\times$  as long as inner one. Tibial comb of hind leg composed of 15 spines.

Lengths (means, 0.01 mm in unit) and proportions (means) of legs:

	F	T	Ta1	Ta2	Ta3	Ta4	Ta5	BV	sv	LR
P <sub>1</sub>	123	125	70	38	30	20	15	3.10	3.54	0.56
P <sub>2</sub>	118	123	43	28	23	15	15	3.53	5.65	0.35
P <sub>3</sub>	143	150	75	43	30	18	15	3.50	3.90	0.50

Wing: Length 2.4 mm. Width 0.8 mm. L/WR 3.00. VR 1.00. R, R<sub>1</sub> and R<sub>4+5</sub> with 46, 25, 35 setae respectively. Radialis with 8 annular organs on its basal part, 1 annular organ on middle anteriorly, 8 annular organs on apical part, and with 6 setae near middle. Squama with 1 seta.

Genitalia (Figs. 9-A, -B) : Laterosternite completely separated from 9th tergite, and with an apical projection which is long, slender and bears more than 10 setae on its apical 1/2. Coxite VIII with 19 long setae.


Length of body: 2.3 mm.

DISTRIBUTION : Japan (Honshu).

TYPE MATERIAL : Holotype ♂ (Type No. 2177, Kyushu Univ.), Mt. Shiroumadake (1600-2600 m), Kita-Azumi-Gun, Nagano Pref., Honshu, 16. viii. 1962, T. Saigusa leg. Paratypes : 10♂♂, 2♀♀, same data as holotype.

The holotype is deposited in the collection of the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka.

REMARKS : In general appearance this new species much resembles Nearctic



**Fig. 9.** Female genitalia of *Chasmatonotus parabolicolor* sp. nov. A: lateral aspect. B: ventral aspect.


*C. bicolor* Rempel, 1937, in which the anal point is stouter, the basal lobe of the gonocoxite well developed and much strongly expanded posteriorly, and the gonocoxite strongly bent subbasally.

The female of the new species may be distinguished from *C. saigusai* sp. nov. by the longer apical projection of the laterosternite.

#### ***Chasmatonotus saigusai* sp. nov.**

♂. Colouration : Head paler than in *C. parabolicolor* sp. nov. Anteprenotum yellow to yellowish brown, proepisternum brown, proepimeron yellow to yellowish brown, postpronotum yellowish brown. Mesonotum subshining deep reddish brown ; scutum yellow on anterior 1/6 and around postpronotum, scutellum and postnotum subshining dark brown. Thoracic pleura subshining yellowish brown to dark brown, thinly grey pollinose. Legs almost as in two preceding species but slightly paler. Wing marking resembling that of *C. unilobus* sp. nov., cell Cu slightly tinged with yellowish brown. Halter yellow. Abdomen predominantly yellowish brown to dark brown; basal 2-3 segments yellow to pale yellowish brown, darkened posteriorly. Genitalia dark brown, thinly grey pollinose.

Head : Antennal flagellomere lengths ( $\mu$ ): 115, 38, 40, 43, 85. First to 5th flagellomeres with 14, 6, 6, 6, 8 setae respectively. Antennal ratio 0.36. Sec-



**Fig. 10.** Male genitalia of *Chasmatonotus saigusai* sp. nov. A: epandrium, gonocoxite and gonostylus in dorsal aspect. B: gonocoxite and gonostylus in dorsal aspect. C: same, inner lateral aspect. D: hypandrium, gonocoxite and gonostylus in ventral aspect.



ond to 5th palpal segment lengths ( $\mu$ ): 70, 133, 118, 163, and the palpal segments with 8-9, 30-34, 20-25, 12-15 setae respectively. Vertex, clypeus and gena with 23-25, 16-21, O-2 setae respectively.

Thorax : Lateral anteprenotals 14-15, median anteprenotals 9-10; dorso-centrals 17-26, uniserial; acrostichals 13-14, prealars 7-8, no supra-alar; scutellars 21, biserial. Preepisternum with 5-7 setae. Posterior anepisternum with 1-3 setae.

Legs : Fore-, middle and hind coxae with 26-37, 11-21, 23-29 apical setae respectively. Middle coxa with 7-9 minute sensory hairs on middle portion of outer surface. Fore-, middle and hind trochanters with 26-31, 29-31, 19-26 apical setae respectively. Outer spur of hind tibia 1.6~ as long as inner one. Tibial comb of hind leg composed of 13 spines.

Lengths (means, 0.01 mm in unit) and proportions (ranges and means) of legs :

	F	T	Tal	Ta2	Ta3	Ta4	Ta5	BV	sv	LR
P <sub>1</sub>	118	119	75	46	33	23	15	2.67-2.68, 2.68	3.06-3.26, 3.15	0.62-0.65, 0.63
P <sub>2</sub>	109	111	48	30	23	16	13	3.19-3.41, 3.29	4.48-4.82, 4.63	0.42-0.44, 0.43
P <sub>3</sub>	128	140	75	41	31	18	14	3.24-3.37, 3.30	3.56-3.57, 3.57	0.53-o.54, 0.54

Wing (Fig. 4-C) : Length 1.9-2. Omm. Width 0.6 mm. L/WR, range 3.24-**3.35**, mean 3.30. VR, range 1.00-1.04, mean 1.02. R, R<sub>1</sub> and R<sub>4+5</sub> with 29-44, 7-16, 24-30 setae respectively. Radialis with 10 annular organs on its basal part, 2 annular organs on middle anteriorly, 7-8 annular organs on apical part, and with 4-5 setae near middle. Squama with O-2 setae.


Genitalia (Figs. 10-A, -B, -C, -D) : Anal point (Fig. 5-C) moderately long, blade-shaped, with weakly emergent dorsal margin. Gonocoxite resembling that of *parabicolor* but transverse elevation of middle area weaker, both basal lobes only slightly swollen; inner surface of ventrodial portion with some long curved setae. Gonostylus (Figs. 5-D) very characteristic, moderately large, subtriangular in shape, inner area strongly concaved and with a weak longitudinal eminence near dorsal edge, with 1-2 stout spines near tip.

Length of body : 2.0-2.3 mm.

♀. Colouration almost as in male, but in darker specimens thorax as in *parabicolor*. Colouration of abdomen as in *parabicolor*.

Head: Antenna1 flagellomere lengths ( $\mu$ ): 118, 45, 26, 26, 76. First to 5th flagellomeres with 12, 6, 6, 6, 6 setae respectively. Antennal ratio, range 0.29-o.35, mean 0.31. Second to 5th palpal segment lengths ( $\mu$ ): **80**, 147, 147, 217, and the palpal segments with 10-11, 33-39, 24-28, 19-22 setae respectively. Vertex, clypeus and gena with 19-31, 16-18, O-1 setae respectively.

Thorax : Lateral anteprenotals 13-14, median anteprenotals 5-9 : dorso-centrals 17-34, irregularly biserial ; acrostichals 9-12, prealars 11-13, no supra-alar;



**Fig. 11.** Female genitalia of *Chasmatonotus saigusai* sp. nov. A: lateral aspect. B: ventral aspect.

scutellars 22-31, biserial. Preepistenum with 3-7 setae.

Legs : Fore-, middle and hind coxae with 30-36, 13-23, 29-31 apical setae respectively. Middle coxa with 9 minute sensory hairs on middle portion of outer surface. Fore-, middle and hind trochanters with 30-31, 24-26, 20-21 apical setae respectively. Outer spur of hind tibia 1.5~ as long as inner one. Tibial comb of hind leg composed of 12 spines.

Lengths (means, 0.01 mm in unit) and proportions (ranges and means) of legs :

	F	T	Tal	Ta2	Ta3	Ta4	Ta5	BV	s v	LR
P <sub>1</sub>	129	128	79	48	33	24	16	2.69-2.93, 2.79	3.14-3.39, 3.25	0.60-o. 64, 0.62
P <sub>2</sub>	128	136	64	36	28	16	15	3.40-3.50, 3.45	3.69-4.73, 4.14	0.42-o. 52, 0.47
P <sub>3</sub>	130	130	66	38	29	18	14	3.34-3.35, 3.35	3.49-4.78, 3.94	0.43-o. 56, 0.51

Wing: Length 2.6 mm. Width 0.9 mm. L/WR 2.92. VR 0.98. R, R<sub>1</sub> and R<sub>4+5</sub> with 48, 14, 33 setae respectively. Radialis with 11 annular organs on its basal part, 3 annular organs on middle anteriorly, 8 annular organs on apical part, and with 6 setae near middle. Squama with 1 seta.

Genitalia (Figs. 11-A, -B) : Structure much resembling that of *parabicolor*, but apical projection of laterosternite stouter than in *parabicolor*, and with about 20 setae on its apical 1/2. Coxite VIII with 14 long setae.

Length of body: 2.3 mm.

DISTRIBUTION: Japan (Honshu).

TYPE MATERIAL: Holotype ♂ (Type No. 2178, Kyushu Univ.), Mt. Kinpusan

(1800-2596 m), Yamanashi Pref., 1. vii. 1963, T. Saigusa leg. Paratypes : 2♂♂, same data as holotype; 1♂, 3♀♀, Mt. Senjô-Dake, Nagano Pref., 4. vii. 1963, T. Saigusa leg. ; 1♂, Tokugô-Tôge (2135 m), 9. vii. 1963, T. Saigusa leg. ; 1♂, 1♀, Karasawa (2600 m), Nagano Pref., 9. vii. 1963, T. Saigusa leg., 4♀♀, same locality and same collector, 10. vii. 1963.

The holotype is deposited in the collection of the Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka.

REMARKS: This species is distinguished from all the known species in having the gonocoxite with slightly swollen basal lobes and the gonostylus subtriangular,

### References

- Coquillett, D. W., 1905. New nematocerous Diptera from North America. *J.N. Y. ent. Soc.* 13: 66.
- Loew, H., 1864. Diptera Americae septentrionalis indigena. *Bed. ent. Z.* 8: 50.
- Malloch, J. R., 1915. The Chironomidae, or midges, of Illinois, with particular references to the species occurring in the Illinois River. *Bull. Ill. State Lab. nnt. Hist.* 10: 499.
- Rempel, J. G., 1937. Note on the genus *Chasmatonotus* with descriptions of three new species (Diptera, Chironomidae). *Can. Ent.* 69: 250-255.