

Localization of Aluminum in Root Tip Tissues of Wheat Varieties Differing in Aluminum Tolerance

Andrade, Leide R. M. de

Plant Nutrition Laboratory, Faculty of Agriculture, Kyushu University

Ikeda, Motoki

Plant Nutrition Laboratory, Faculty of Agriculture, Kyushu University

Ishizuka, Junji

Plant Nutrition Laboratory, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/24140>

出版情報：九州大学大学院農学研究院紀要. 41 (3/4), pp.151-156, 1997-03. Kyushu University
バージョン：
権利関係：

Localization of Aluminum in Root Tip Tissues of Wheat Varieties Differing in Aluminum Tolerance

Leide R. M. de Andrade, Motoki Ikeda and Junji Ishizuka

Plant Nutrition Laboratory, Faculty of Agriculture,
Kyushu University, Fukuoka 812-81, Japan

(Received October 21, 1996; Accepted December 17, 1996)

The purpose of this study was to investigate the mechanisms of Al tolerance in wheat varieties by the direct observation of Al distribution in root cells using the hematoxylin stain method. Five-day-old seedlings of Al-tolerant varieties, Atlas 66 and Shirosanjyaku, and an Al-sensitive variety, Chikushikomugi, were treated with 150 μM Al in 0.4 mM CaCl₂, pH 4.5 for 24 hours under aseptic conditions; afterwards the roots were stained with hematoxylin. Comparison of hematoxylin stain of root tips among the varieties showed that the root cap of Al-sensitive seedlings was stained intensively, reflecting high accumulation of Al in this region. The roots developed symptoms of Al toxicity such as increased vacuolation, and swelling and rupture of the cells of the epidermis. On the other hand, in Al-tolerant varieties the central cap cells and the cortical region of the meristem and elongation zone were not stained. The site of the earliest Al accumulation in Al-tolerant varieties seemed to be the edge cells of the root cap. In the elongation zone, Al accumulated mainly in the cell wall and nuclei. Those differences in the staining pattern were probably due to immobilization of Al in the cytosol that could not react with hematoxylin and/or due to the ability of Al-tolerant varieties to better exclude Al from cytoplasm.

INTRODUCTION

To elucidate the mechanisms of Al toxicity in plants, the characterization of the primary site for Al uptake in the roots is of primary importance. In order to localize Al in root tissues, various methods have been used, from the use of Al dyes (hematoxylin, aluminon, morin) to such sophisticated techniques as X-ray microanalysis (Ownby, 1993; Naidoo et al., 1978) or secondary-ion mass spectrometry (Lazof *et al.*, 1994). The hematoxylin stain method (Polle *et al.*, 1978) is of very popular use in evaluation of Al tolerance in plants (Takagi *et al.*, 1981; Wallace and Anderson, 1984). The method is based on the colorimetric property of hematoxylin to give a blue-purple stain when it is complexed with Al (Baker, 1962; Rincon and Gonzalez, 1992). With this technique it has been possible to indicate the differences in Al tolerance among wheat varieties (Takagi *et al.*, 1981; Wheeler *et al.*, 1992a) and histochemical localization of Al in root surface and the cell wall and nuclei of root cells.

The aim of the present study was to investigate the mechanisms of Al tolerance in wheat varieties differing in Al tolerance by the direct observation of distribution of Al in root tissues and cells by using the hematoxylin method.

MATERIALS AND METHODS

Seedlings of Al-tolerant varieties, Atlas 66 (AT) and Shirosanjyaku (SH), and an Al-

sensitive variety, Chikushikomugi (CK), of wheat (*Triticum aestivum* L.) were grown aseptically. The seeds were disinfected by soaking in NaClO solution (1.5% active chlorine) for 90 minutes, rinsed several times with sterile water, and then germinated on moistened paper in sterile Petri dishes in the dark for 48 hours. Twenty seedlings were transferred onto a stainless-steel screen in a test tube (20 cm in length, 4 cm in diameter) over 50 ml of sterile nutrient solution (pH 4.5) containing macronutrients (mM): 0.50 NH₄⁺, 1.0 NO₃⁻, 0.50 Ca²⁺, 0.50 K⁺, 0.30 Mg²⁺, 0.40 Cl⁻, 0.40 SO₄²⁻, 0.30 Na⁺, 0.0025 PO₄³⁻, and micronutrients (μ M): 0.91 Mn²⁺, 0.03 Cu²⁺, 4.63 BO₃³⁻, 0.08 Zn²⁺, 3.2 Fe³⁺ (Na-EDTA). The tubes were placed on a rotary shaker (120 rpm) at 20 °C for 4 days in a growth chamber with light intensity of 150 μ mol m⁻²s⁻¹ and a photoperiod of 12 hours. On the fifth day, after the solution was aspirated from the tubes the seedlings were treated with 150 μ M Al (AlCl₃ or Al₂(SO₄)₃+0.4 mM CaCl₂) and without Al (0.4 mM CaCl₂) at pH 4.5 for 24 hours. Each tube contained 40 ml of the treatment solution. Al was added after Al stock solution was passed through 0.45 μ m membrane filter.


After exposure to Al, the seedlings were stained in 50 ml of hematoxylin solution (2 g hematoxylin (Nacalai Tesque, Kyoto) and 0.2 g NaIO₃ in 1 liter of deionized water) for 20 minutes according to the method of Polle *et al.* (1978). Thereafter, the seedlings were rinsed in running water for 1 minute and soaked in 50 ml of deionized water, which was changed 3 times at each interval of 20 minutes. The terminal 2 – 3 cm segments of roots were excised and root apices were mounted on glass slides and examined with a Nikon Optiphot light microscope. Photographs were taken with a Nikon FX-35 WA camera by using a Nikon HFX-IIA system automatic exposure unit.

RESULTS

A root tip of AT that was not exposed to Al but stained with hematoxylin is shown in Fig. 1A. The root showed lack of staining on the root cap and epidermis. The pattern of the hematoxylin staining of the roots treated with Al showed that Al accumulated mainly in the apical region (root cap and meristem) in both Al-tolerant AT and Al-sensitive CK. Nevertheless, clear differences in the staining intensity were observed between tolerant and sensitive root tips (Fig. 1B), which is in agreement with results of other varieties in previous reports (Polle *et al.*, 1978; Wheeler *et al.*, 1992a; Rincon and Gonzalez, 1992; Ownby, 1993).

The primary site for accumulation in roots of Al-tolerant SH and AT appeared to be the edge cells (sloughed-off cells) of the root cap that were intensely dyed (Fig. 1C and 1D). No staining was observed in the central cells of the root cap and the cells of meristem zone were only slightly stained. In the cortical and epidermal cells of semi-apical region of root tips (4 – 6 mm), Al accumulated mainly in the cell walls, intercellular spaces, and nuclei (Fig. 1E).

In Al-sensitive CK, root tips was intensively stained in all the experiments, showing a deeper staining pattern in the cells of the root cap (Fig. 1F). The meristematic cells were swollen and extensively vacuolated (Fig. 1G and 1H).



Photographs of hematoxylin staining of Al-treated and Al-untreated roots of wheat varieties differ

Fig. 1.

ing in aluminum tolerance **Plate A** An apex of Atlas 66 root unexposed to Al. Bar:0.1mm Symbols: CC:center cap cells, CT:cortex, E:epidermis, M:meristem, RC:root cap. **Plate B** Apices of Shirosanjyaku (left) and Chikushikomugi (right) roots treated with $75 \mu\text{M Al}_2(\text{SO}_4)_3$ for 24 hours. Magnification is the same as in Plate A. **Plate C** An apex of Shirosanjyaku root treated with $75 \mu\text{M Al}_2(\text{SO}_4)_3$ for 24 hours. Magnification is the same as in Plate A. **Plate D** An apex of Atlas 66 root treated with $75 \mu\text{M Al}_2(\text{SO}_4)_3$ for 24 hours. Magnification is the same as in Plate A. **Plate E** Semi-apical region of the tip of Shirosanjyaku root treated with $150 \mu\text{M AlCl}_3$ for 24 hours. Bar: $50 \mu\text{m}$. **Plate F** An apex of Chikushikomugi root treated with $150 \mu\text{M AlCl}_3$ for 24 hours. Magnification is the same as in Plate A. **Plate G** Cells of the root cap of Chikushikomugi root treated with $150 \mu\text{M AlCl}_3$ for 24 hours. Magnification is the same as in Plate E. **Plate H** Cells of the meristematic region of Chikushikomugi root treated with $150 \mu\text{M AlCl}_3$ for 24 hours Bar: $100 \mu\text{m}$.

DISCUSSION

The distribution of Al, as visualized with hematoxylin stain, showed that Al first accumulated in the outer cells of the root cap and epidermis. The symptoms of Al toxicity such as tissue lesions and swollen epidermal cells were similar to those described previously (Wagatsuma *et al.*, 1987; Kinraide, 1988; Rincon and Gonzalez, 1992; Ikeda and Tadano, 1993; Ryan *et al.*, 1993).

In Al-tolerant varieties, hematoxylin stain was absent from the central cap cells and the cortical regions of the meristem and elongation zones (Fig. 1B, 1C and 1D). The site of the earliest Al accumulation in those root tips seems to be the edge cells of the root cap, which were intensely dyed. Rincon and Gonzalez (1992) measured the amounts of Al accumulated in root segments of Al-tolerant (Atlas 66) and Al-sensitive (TAM 105) wheat varieties and observed that TAM 105 accumulated more Al in the 0 – 2 mm root segments than Atlas 66. The results suggested that Al-tolerant roots had a mechanism to prevent accumulation of Al in this region. Delhaize *et al.* (1993) actually reported that excretion of organic acids having potential to chelate Al took place in a 1-3 mm root portion. The same staining pattern as the present study was observed in root tips of maize by Ryan *et al.* (1993), but it was independent of Al tolerance of those cultivars.

In the semi-apical region (3 – 6 mm) of Al-tolerant varieties, Al accumulated mainly in the cell wall and nuclei (Fig. 1E). The almost complete absence of stain in the cytoplasm of those cells indicates that small amounts of Al reached this compartment but they were immobilized or precipitated in such a way as they were not able to make a complex with hematoxylin. Before reaching nuclei, Al necessarily passes through the neutral cytoplasm, permeates the plasmalemma, and passes through the neutral cytoplasm and then the nuclear envelope. In the cytoplasm with a pH of approximately 7.5, most of the soluble Al is supposed to be negatively charged $\text{Al}(\text{OH})_4^-$ that appears to be nontoxic (Kochian 1995). The unmordanted hematoxylin which is a very weak anionic dye is inclined to make a complex with positively charged Al (Baker, 1962), thus being unlikely to stain the $\text{Al}(\text{OH})_4^-$ in the cytoplasm. The level of Al^{3+} that may exist in the cytoplasm is estimated to be in the picomolar to nanomolar ranges (Kochian 1995). Based on the work of Baker (1960; 1962) it is presumed that once Al enters into the cell, the positively charged Al might be driven towards sites of negatively charged phosphoric groups of the nucleic acid molecule, associating the Al with chromatin. Then, the bound Al might attract hematoxylin and the color is developed. Accumulation of Al in the nuclei is often reported (Matsumoto *et al.*, 1976; Morimura *et al.*, 1978; Rincon and Gonzalez, 1992).

Al-sensitive root tips presented increased vacuolation and swelling and rupture of the cells of the epidermis, and were completely stained at higher concentrations of Al (Fig. 1F, 1G and 1H). These symptoms are associated with Al toxicity (Wheeler *et al.*, 1992b; de Lima and Copeland, 1994; Marienfeld *et al.*, 1995). Although vacuolation is commonly observed in cells intoxicated by Al, the reason for the increased vacuolation was probably not a direct effect of Al but an effect of growth retardation.

The complete staining of the root apices of Al-sensitive CK reflects high accumulation of Al in this portion. This pattern of hematoxylin staining could result from solid phase precipitation of Al at high pH's or due to Al fixed in root tissues as AlPO_4 as suggested by Ownby (1993). In fact the Al-hematoxylin complex precipitates at pH's higher than 4.8.

Ownby (1993) demonstrated by using X-ray microanalysis that Al and P were co-localized in the cell wall regions of the outer cortex of Al-stressed roots which is in accordance with early observations of Clarkson (1967) and Naidoo *et al.* (1978). This solid phase Al-P might be able to react with hematoxylin. Based on those results, Ownby (1993) postulated that cultivars whose root cells were damaged by Al allowed cellular phosphate to leak into the cell wall region and precipitate as AlPO_4 and consequently were intensely stained with hematoxylin. This hypothesis seems to be plausible because it is unlikely that Al-hematoxylin complex would be formed under the neutral cytoplasmic pH conditions as described above. Moreover, Al-sensitive CK generally showed greater damage in the tissue and it would facilitate the leakage of P from the cells.

ACKNOWLEDGMENT

We thank Dr. Tomohiko Yoshida, Laboratory of Crop Science Kyushu University, for his help in collecting wheat seeds.

REFERENCES

- Baker, J. R. 1960 Experiments on the action of mordants. 1. "Single-bath" mordant dyeing. *Quart. J. Micr.Sci.*, **101**: 255-272
- Baker, J. R. 1962 Experiments on the action of mordants. 2. Aluminum-haematin. *Quart. J. Micr.Sci.*, **103**: 493-517
- Clarkson, D. T. 1967 Interactions between aluminum and phosphorus on root surfaces and cell wall material. *Plant Soil*, **27**: 347-356
- Delhaize, E., P. R. Ryan, and P. J. Randall 1993 Aluminum tolerance in wheat. II Aluminum-stimulated excretion of malic acid from root apices. *Plant Physiol.*, **103**: 695-702
- de Lima, M. L. and L. Copeland 1994 Changes in the ultrastructure of the root tip of wheat following exposure to aluminum. *Aust. J. Plant Physiol.*, **21**: 85-94
- Ikeda, H. and T. Tadano 1993 Ultrastructural changes of the root tip cells in barley induced by a low concentration of aluminum. *Soil Sci. Plant Nutr.*, **39**: 109-117
- Kinraide, T. B. 1988 Proton exclusion by wheat roots exhibiting severe aluminum toxicity symptoms. *Plant Physiol.*, **88**: 418-423
- Kochian, L. V. 1995 Cellular mechanisms of aluminum toxicity and resistance in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, **46**: 237-260
- Lazof, D. B., J. G. Goldsmith, T. W. Ruffy, and R. W. Linton 1994 Rapid uptake of aluminum into cells of intact soybean root tips: A microanalytical study using secondary ion mass spectrometry. *Plant Physiol.*, **106**: 1107-1114
- Marienfeld, S., H. Lehmann, and R. Stelzer 1995 Ultrastructural investigations and EDX-analysis of Al-treated oat (*Avena sativa*) roots. *Plant Soil*, **171**: 167-173
- Matsumoto, H., E. Hirasawa, H. Torikai, and E. Takahashi 1976 Localization of absorbed aluminium in pea root and its binding to nucleic acids. *Plant Cell Physiol.*, **17**: 127-137
- Morimura, S., E. Takahashi, and H. Matsumoto 1978 Association of aluminum with nuclei and inhibition of cell division in onion (*Allium cepa*) roots., *Z. Pflanzenphysiol.*, **88**: 395-401
- Naidoo, G., J. McD. Stewart, and R.J. Lewis 1978 Accumulation sites of Al in snapbean and cotton roots. *Agr. J.*, **70**: 489-492
- Ownby, J. D. 1993 Mechanisms of reaction of hematoxylin with aluminium-treated wheat roots. *Physiol. Plant.*, **87**: 371-380
- Polle, E., C. F. Konzak, and J. A. Kittrick 1978 Visual detection of aluminum tolerance levels in wheat by hematoxylin staining of seedling roots. *CropSci.*, **18**: 823-827

- Rincon, M. and R. A. Gonzalez 1992 Aluminum partitioning in intact roots of aluminum-tolerant and aluminum-sensitive wheat (*Triticum aestivum* L.) cultivars. *Plant Physiol.*, 99: 1021-1028
- Ryan, P. R., J. M. Ditomaso, and L. V. Kochian 1993 Aluminium toxicity in roots: An investigation of spatial sensitivity and the role of root cap. *J. Exp. Bot.*, 44: 437-446
- Takagi, H., H. Namai, and K.-I Murakami 1981 Evaluation of the hematoxylin staining method for detecting wheat tolerance to aluminum. *Jpn. J. Breed.*, 31: 152-160 (in Japanese)
- Wagatsuma, T., M. Kaneko, and Y. Hayasaka 1987 Destruction process of plant root cells by aluminum. *Soil Sci. Plant Nutr.*, 33: 161-175
- Wallace, S.U. and I. C. Anderson 1984 Aluminum toxicity and DNA synthesis in wheat roots. *Agr. J.*, 76: 5-8
- Wheeler, D. M., D. C. Edmeades, R. A. Christie, and R. Gardner 1992a Comparison of techniques for determining the effect of aluminium tolerance in wheat. *Plant Soil*, 146: 1-8
- Wheeler, D. M., D. J.C. Wild, and D. C. Edmeades 1992b Preliminary results from a microscopic examination on the effects of aluminium on the root tips of wheat. *Plant Soil*, 146: 83-87