Detection of Alien *Oryza punctata* Kotschy Chromosomes in Rice, *Oryza sativa* L., by Genomic in situ Hybridization

Yasui, Hideshi

Plant Breeding Laboratory, Faculty of Agriculture, Kyushu University

Nonomura, Ken-ichi

Plant Breeding Laboratory, Faculty of Agriculture, Kyushu University

Iwata, Nobuo

Plant Breeding Laboratory, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/24192>

出版情報：九州大学大学院農学研究院紀要. 42 (1/2), pp.63-68, 1997-12. Kyushu University
バージョン：
権利関係：

Detection of Alien *Oryza punctata* Kotschy Chromosomes in Rice, *Oryza sativa* L., by Genomic *in situ* Hybridization

Hideshi Yasui, Ken-Ichi Nonomura¹ and Nobuo Iwata

Plant Breeding Laboratory, Faculty of Agriculture,
Kyushu University, Fukuoka 812-81

(Received July 30, 1997 and accepted August 25, 1997)

Genomic *in situ* hybridization (GISH) using total *Oryza punctata* Kotschy genomic DNA as a probe was applied to detect alien chromosomes transferred from *O. punctata* (W1514: $2n=2x=24$: BB) to *O. sativa* Japonica cultivar, Nipponbare ($2n=2x=24$: AA). Only 12 chromosomes in the interspecific hybrids ($2n=3x=36$: AAB) between autotetraploid of *O. sativa* cultivar Nipponbare and a diploid strain of *O. punctata* (W1514) showed intense staining by FITC in mitotic metaphase spreads. Only one homologous pair of chromosomes out of the 12 pairs of *O. sativa* chromosomes, which is probably chromosome 8, contained similar repetitive sequences as present in *O. punctata* and were frequently slightly stained by GISH. Monosomic alien addition line(MAAL)s 7 ($2n=2x+1=25$: AA+7B: Type II) and 10 ($2n=2x+1=25$: AA+10B) were characterized by an additional single chromosome which had been intensely stained by GISH. Thus, GISH using *O. punctata* total genomic DNA allowed the detection of a specific chromosome transferred into cultivated rice. This method can be useful to search introgressed chromosomal segments or translocated chromosomes in the progenies of the MAALs each carrying an alien chromosome of *O. punctata* in rice.

INTRODUCTION

Repetitive sequences are well known to rapidly amplify in genomes after isolation from a common ancestral species and evolve toward species-specificity in animals (Schmid and Shen, 1986; Batzer and Deininger, 1991) and plants (Walbot and Cullis, 1985; Vedel and Delesny, 1987). These repetitive sequences, therefore, tend to be conservative within a genome, but differentiated among genomes. "Chromosome painting", a recent technique in molecular cytology, has made it possible to detect genome-specific chromosomes using repetitive sequences as probe. Genomic *in situ* hybridization (GISH) using total genomic DNA as a probe is a similar approach because total genomic DNA is rich in repetitive sequences, and can also detect genome-specific chromosomes in hybrids derived from a cross between closely or distantly related species (Ananthawat-Jonsson *et al.*, 1990; Mukai and Gill, 1991; Mukai *et al.*, 1991). This method can be useful in identifying specific alien chromosomes transferred into monosomic alien addition lines (MAALs) carrying a single chromosome of *O. punctata* in addition to normal chromosome complement of *O. sativa*. This study was designed to establish if GISH using total *O. punctata* DNA as a probe would allow for the intense and specific fluorescent staining of *O. punctata* chromosomes in prometaphase spreads and interphase nuclei of the progeny from an interspecific cross between *O. sativa* and *O.*

¹ Present address: Experimental-Farm, National Institute of Genetics Yata 1, 111, Mishima, Shizuoka-Ken, 411

punctata. In addition, the chromosome rearrangement of *O. punctata* chromosomes transferred into rice will be discussed.

MATERIALS AND METHODS

Plant materials

The Japonica rice cultivar Nipponbare ($2n=2x=24$: AA), a diploid strain of *O. punctata* (W1514: $2n=2x=24$: BB), their interspecific hybrid ($2n=3x=36$: AAB), and two MAALs 7 ($2n=2x+1=25$: AA+7B) (Type II) and 10 ($2n=2x+1=25$: AA+10B) were used in the present study. The diploid strain of *O. punctata* (W1514) was collected in Kenya in 1964 (Tateoka, 1965) and the seeds have been supplied from the National Institute of Genetics, Japan. This strain has been studied cytologically to determine genomic relationship between *O. sativa* and *O. punctata* (Katayama, 1967; Katayama and Ogawa, 1974; Ogawa and Katayama, 1974). The original interspecific cross was made between Nipponbare autotetraploids ($2n=4x=48$: AAAA) and W1514. The interspecific hybrids were obtained through embryo rescue. A series of MAALs were developed and isolated from the backcrossed progeny (Yasui and Iwata, 1991). MAAL 7 (Type II) and MAAL 10 had been identified by morphological resemblance to related trisomics (Yasui and Iwata, 1991) and RFLP analysis (Yasui *et al.*, 1992). Total genomic DNA of W1514 was isolated from leaves of a single plant. DNA was extracted from the ground tissue using the slightly modified potassium acetate method (Dellaporta *et al.*, 1983).

Genomic *in situ* hybridization (GISH)

Chromosome preparation: Prometaphase spreads from the plants were prepared from freshly emerged root tip tissues by using a modification of the procedure described by Kurata and Omura (1978). These root tip tissue was fixed in 3:1 ethanol/acetic acid, followed by 0.002 M 8-hydroxyquinoline treatment at 20 °C for 90 minutes, and macerated with the solution containing 4% (w/v) Cellulase Onozuka RS and 1% (w/v) Pectolyase Y-23 in 7.5 mM KCl at 37 °C for 30–45 minutes. Prometaphase spreads were prepared by flame drying following fixation in 3:1 methanol/acetic acid. Well spread chromosomes were stained with 10% Giemsa solution diluted by Sørensen phosphate buffer and observed for karyotype.

Hybridization: Genomic *in situ* hybridization was conducted as described by Hori *et al.* (1990) with a slight modification. Total *O. punctata* genomic DNA was used as a probe to detect the chromosome transferred from *O. punctata*. The DNA in the target cells was denatured by immersion in 70% formamide/0.3 M NaCl/0.03 M sodium citrate ($2\times$ SSC) for 3 minutes at 80 °C. The slides were rapidly dehydrated in an ethanol series (70%, 100%) and air-dried. The hybridization mixture [270 ml total volume consisting of 10% formamide, $2\times$ SSC, 20% dextran sulfate, 1 mg sonicated herring DNA per ml and 700 ng digoxigenin (DIG) labeled total *O. punctata* DNA] was then heated as above and applied. Hybridization was at 37 °C overnight (15 hours). Slides were washed in 50% formamide/ $2\times$ SSC for 15 minutes at 37 °C, and rinsed in $2\times$ SSC, $1\times$ SSC, $4\times$ SSC at room temperature each for 15 minutes.

Detection: For detection of hybridization sites, the slides were counterstained using the FITC-PI method. The slides were treated with a solution of a complex of anti-DIG-

FITC in 1% BSA/4×SSC. After rinsing, the slides were counterstained with 1.25 mg/ml propidium iodide for 30 minutes. Photographs of GISH prometaphase spreads were taken with Zeiss Axioplan photomicroscope using Fuji Fujichrome films.

RESULTS

There were no intense signal in metaphase spreads of *O. sativa* cultivar Nipponbare as a result of GISH using total *O. punctata* genomic DNA. The centromeric regions of chromosomes were frequently slightly stained (Data are not shown).

Among the prometaphase spreads of interspecific hybrids, only 12 chromosomes in many nuclei were intensely stained by FITC (Fig. 1a). In some prometaphase nuclei, however, more than 12 (13–14) chromosomes were intensely stained (Fig. 1b). These frequently stained additional chromosomes seemed to be a couple of homologous chromosome 8 of *O. sativa* because of their karyotypic feature.

One chromosome had been intensely stained in each prometaphase and interphase spreads in MAAL 7 (Type II) and MAAL 10 (Fig. 2). Both plates a (top and bottom) show GISH nuclei, while both plates b (top and bottom) show Giemsa staining nuclei. The intensely stained chromosome in MAAL 7 (Type II) was medium and submetacentric and it was different from chromosome 7 of *O. sativa* which was small and metacentric. The intensely stained chromosome in MAAL 10 was small and submetacentric and was similar

Fig. 1. GISH of the interspecific hybrids ($2n=3x=AAB$) with *O. punctata* total DNA as a probe, showing the twelve (a) and thirteen (b) chromosomes intensely stained by FITC. Note the intensely stained sites which appear to be heterochromatic regions.

to chromosome 10 of *O. sativa* which was small and with heterochromatic region in the entire short arm (Fig. 3).

Fig. 2. GISH with *O. punctata* total DNA as a probe (a) and Giemsa staining (b) of MAAL 7 (Type II) and MAAL 10, showing the hybridization sites on chromosomes (Arrowheads). Upper and bottom plates show MAAL 7 (Type II) and MAAL 10, respectively.

Fig. 3 Karyotype of MAAL 10 revealed by using GISH. Chromosome 10B characterized by its small size, sub-metacentric type and heterochromatic region in entire short arm was similar to *O. sativa* chromosome 10. The capital letters A and B show the chromosome(s) of *O. sativa* and *O. punctata*, respectively.

DISCUSSION

There were no intense signal in metaphase spreads of *O. sativa* cultivar Nipponbare as a result of GISH using total *O. punctata* genomic DNA. This suggests that the genome of *O. punctata* is differentiated from that of *O. sativa*. Only 12 chromosomes were intensely stained by FITC in the interspecific hybrids carrying 24 chromosomes of *O. sativa* and 12 chromosomes of *O. punctata*. Among 12 pairs of *O. sativa* chromosomes, only one homologous pair of chromosomes, presumably chromosome 8, was likely to have the specific repetitive sequences so that this pair of chromosomes were slightly stained by GISH frequently.

MAAL 7 (Type II) and MAAL 10 were characterized by their additional chromosomes which were intensely stained by FITC. The intensely stained chromosome in MAAL 7 (Type II) was medium and submetacentric and the karyotypic feature was different from chromosome 7 of *O. sativa* (characterized by small size and metacentric type). MAAL 7 (Type II) with brown pericarp had been derived from the progenies of MAAL 7 (Type I) with red pericarp inherited from *O. punctata* (Yasui and Iwata, 1991). Moreover, RFLP characterization of Type II had shown deficiency of several RFLP loci detected in Type I (Yasui *et al.*, 1992). These indicate that the alien chromosome in MAAL 7 (Type II) had been modified by chromosome rearrangement such as deletion or translocation. The intensely stained chromosome in MAAL 10 was similar to chromosome 10 of *O. sativa*. The results of karyotyping showed that karyological structure of chromosome 10 in both genomes were conservative.

GISH using *O. punctata* total DNA as a probe allowed the detection of a specific *O. punctata* chromosome transferred into cultivated rice. GISH, therefore, can be used in searching introgressed chromosomal segments or translocated chromosomes in the progenies of the MAALs each carrying an alien chromosome of *O. punctata* Kotschy.

ACKNOWLEDGEMENTS

The authors thank Dr. A. Yoshimura for helpful discussion and Dr. J. P. Gustafson for critical reading of the manuscript. We are grateful to Dr. K. Koga for reviewing the manuscript. Work in the laboratory was supported by grant from the Ministry of Education 08876003.

REFERENCES

- Ananthawat-Jonsson, K., T. Schwarzacher, A. R. Leitch, M. D. Bennett and J. S. Heslop-Harrison 1990 Discrimination between closely related *Triticeae* species using genomic DNA as a probe. *Theor. Appl. Genet.*, **79**: 721-728
- Batzer, M. A. and P. L. Deininger 1991 A human-specific subfamily of *Alu* sequences. *Genomics*, **9**: 481-487
- Dellaporta, S. L., J. Wood and J. B. Hicks 1983 A plant DNA miniprep: Version II. *Plant Molecular Biology Reporter*, **1**(4): 19-21
- Hori, T., E. Takahashi, D. Ayusawa, K. Takeishi, S. Kaneda and T. Seno 1990 Regional assignment to the human thymidylate synthase (TS) gene to chromosome band 18p11.32 by nonisotopic *in situ* hybridization. *Hum. Genet.*, **85**: 576-580
- Katayama, T. 1967 Cytogenetical studies on *Oryza*. F₁ hybrids of the crosses BBCC×CC, BBCC×a diploid strain of *O. punctata* and CC×a diploid strain of *O. punctata*. *Proc. Japan. Acad.*, **43**: 327-331

- Katayama, T. and T. Ogawa 1974 Cytological studies on the genus *Oryza*. VII. Cytogenetical studies on F₁ hybrids between diploid *O. punctata* and diploid species having C genome. *Japan. J. Breed.*, **24**: 165-168
- Kurata, N. and T. Omura 1978 Karyotype analysis in rice. I. A new method for identifying all chromosome pairs. *Japan. J. Genet.*, **53**: 251-255
- Mukai, Y. and B. S. Gill 1991 Detection of barley chromatin added to wheat by genomic *in situ* hybridization. *Genome*, **34**: 448-452
- Mukai, Y., B. Friebe, J. H. Hatchett and B. S. Gill 1991 Detection of rye chromatin in wheat specifying resistance to Hessian fly by *in situ* hybridization using total rye genomic DNA probes. In "Proc. 2nd Int. Symp. Chromosome Engineering in Plants" ed. by G. Kimber, Columbia, MO, pp. 184-188
- Ogawa, T. and T. Katayama 1974 Cytogenetics on the genus *Oryza*. Chromosome pairing in the interspecific hybrid between diploid *O. punctata* and *O. officinalis*. *Japan. J. Genet.*, **49**: 257-260
- Schmid, C. W. and C. K. J. Shen 1986 The evolution of interspersed repetitive DNA sequences in mammals and other vertebrates. In "Molecular Evolutionary Genetics" ed. by R. J. MacIntyre, Plenum, New York, pp. 323-358
- Tateoka, T. 1965 Taxonomy and chromosome numbers of African representatives of the *Oryza officinalis* complex. *Bot. Mag. Tokyo*, **78**: 198-201
- Vedel, F. and M. Delesny 1987 Repetitiveness and variability of higher plant genomes. *Plant Physiol. Biochem.*, **25**: 191-210
- Walbot, V. and C. A. Cullis 1985 Rapid genomic change in higher plants. *Annu. Rev. Plant Physiol.*, **36**: 367-399
- Yasui, H. and N. Iwata 1991 Production of monosomic alien addition lines of *Oryza sativa* having a single *O. punctata* chromosome. In "Rice Genetics II", International Rice Research Institute, Philippines, pp. 147-155
- Yasui, H., A. Yoshimura and N. Iwata 1992 Characterization of monosomic alien chromosomes of *O. punctata* transferred to *O. sativa* using RFLP markers. *Rice Genet. Newsl.*, **9**: 138-142