

Inhibitory Effect of Abscisic Acid on Shoot Growth and Flowering Induced by Gibberellic Acid in Nonprecooled Derooted Bulbs of Tulip (*Tulipa gesneriana* L.)

Saniewski, Marian
Research Institute of Pomology and Floriculture

Kawa-Miszczak, Ludwika
Research Institute of Pomology and Floriculture

Wegrzynowicz-Lesiak, Elzbieta
Research Institute of Pomology and Floriculture

Okubo, Hiroshi
Laboratory of Horticultural Science, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/24302>

出版情報：九州大学大学院農学研究院紀要. 44 (1/2), pp.25-32, 1999-11. Kyushu University
バージョン：
権利関係：

Inhibitory Effect of Abscisic Acid on Shoot Growth and Flowering Induced by Gibberellic Acid in Nonprecooled Derooted Bulbs of Tulip (*Tulipa gesneriana* L.)

Marian Saniewski¹, Ludwika Kawa-Miszczak¹, Elzbieta Wegrzynowicz-Lesiak¹ and Hiroshi Okubo²

¹ Research Institute of Pomology and Floriculture, Pomologiczna 18, 96–100 Skierniewice, Poland

² Laboratory of Horticultural Science, Faculty of Agriculture, Kyushu University 46–01, Fukuoka 812–8581, Japan

(Received July 2, 1999 and accepted August 24, 1999)

Effect of gibberellic acid (GA₃) at a concentration of 200 mg l⁻¹, abscisic acid (ABA) at a concentration of 20 mg l⁻¹ and their mixture on shoot growth and flowering in nonprecooled derooted tulip bulbs cv. Apeldoorn was investigated. As it was found previously GA₃ induced shoot growth and flowering in nonprecooled derooted bulbs of tulip. The present studies showed that the simultaneous application of GA₃ with ABA totally inhibited shoot growth and flowering induced by the gibberellin in nonprecooled derooted tulip bulbs. These results suggest that hormonal balance, gibberellins–abscisic acid, is the important factor in control of dormancy and dormancy release of tulip bulbs, causing shoot growth and flowering.

INTRODUCTION

It is well known that during the cooling of the tulip bulbs the amount of free gibberellins increases (Aung and De Hertogh, 1967; 1968; De Hertogh *et al.*, 1971; Van Bragt, 1971; Aung and Rees, 1974; Hanks and Rees, 1980). Exogenously applied gibberellins cannot substitute for cold treatment of nonprecooled tulip bulbs but do for only partially cooled bulbs and stimulate shoot growth and flowering (Van Bragt and Zijlstra, 1971; Rudnicki *et al.*, 1976; Van Bragt and Van Ast, 1976; Cocozza Talia and Stellacci, 1979; Hanks, 1984; 1985; Jones and Hanks, 1984). Saniewski *et al.* (1999) showed that in nonprecooled derooted bulbs flower bud blasting occurred and no growth of stem was observed but GA₃ greatly induced shoot growth and flowering. The stimulatory effect of gibberellin on shoot growth and flowering in nonprecooled derooted bulbs was much stronger than in case of nonprecooled rooted bulbs (Rudnicki *et al.*, 1976). Kawa and Saniewski (1986) showed that gibberellic acid had a strong stimulatory effect in increasing the length and fresh weight of pistil isolated from nonprecooled tulip bulbs but to a lesser degree in the case of cooled bulbs, cultured *in vitro*.

The presence of abscisic acid in tulips is well documented (Syrtanova *et al.*, 1975; Aung and De Hertogh, 1979; Singh *et al.*, 1979; Terry *et al.*, 1982) and low temperature treatment decreases the amount of ABA (Syrtanova *et al.*, 1973; Rakhimbayev *et al.*,

^{1,2}Correspondence to either author. ¹Tel: +48-46-833-2021, Fax: +48-46-833-3228, E-mail: msaniew@insad.isk.skierniewice.pl ²Tel: +81-92-642-2828, Fax: +81-92-642-2828, E-mail: hokubo@agr.kyushu.u.ac.jp

1978; Aung and De Hertogh, 1979).

In the present study the inhibitory effect of abscisic acid on tulip shoot growth and flowering induced by gibberellic acid in nonprecooled derooted tulip bulbs is documented.

MATERIALS AND METHODS

Tulip bulbs cv. Apeldoorn, with a circumference of 10–11 cm, after lifting were stored at 18–20°C until the end of October and then were stored at 17°C until used for experiments on January 22 (nonprecooled bulbs). After removal of dry scales all roots were excised and bulbs were kept on petri dishes in following solutions of growth regulators: 1) control, distilled water only, 2) GA₃ at a concentration of 200 mg l⁻¹, 3) ABA at a concentration of 20 mg l⁻¹, and 4) GA₃+ABA in above concentrations.

Fifteen bulbs per treatment were used and newly appeared roots were removed daily. The solutions were changed every 2 or 3 days. On February 12 all treatments were replaced by distilled water. During the first two weeks of experiment the length of sprouts (to the top of first leaf) were measured. The length of all internodes and leaves were recorded on February 8 and February 24, when experiment ended. The length of the first (basal) internode was measured from the basal plate to the first leaf node and the fourth internode from the third leaf to the base of the flower, after the scales from 5 bulbs per treatment were removed.

On January 22, 20 intact nonprecooled tulip bulbs (storage conditions are presented above) and 20 bulbs dry cooled at 5°C from October 15 were individually planted in the pots (after removal of dry scales), for comparison their growth and development.

The data were subjected to an analysis of variance and the Duncan's t-test was used to estimate the difference between means at P=0.05.

RESULTS AND DISCUSSION

The shoot growth of nonprecooled derooted tulip bulbs was very small and only sprouting of leaves was observed and flower bud blasting took place (Figs. 1, 2, 3). As it was found previously, GA₃ at a concentration of 200 mg l⁻¹ stimulated sprouting and finally induced shoot growth and caused flowering (Figs. 1, 2, 3). Both the growth of stem and all leaves was evidently stimulated by GA₃ (Fig. 3). Abscisic acid applied alone inhibited sprouting of nonprecooled derooted tulip bulbs (Figs. 1, 2) and it was caused through inhibition of growth of all intact leaves (Fig. 3). Since in the control – nontreated nonprecooled derooted tulip bulbs, the growth of stem was very small, the effect of ABA applied alone on its growth was not possible for observation (Figs. 2, 3). However, simultaneous application of GA₃ and ABA showed that abscisic acid greatly inhibited shoot growth (stem and leaves) and flowering induced by gibberellic acid in nonprecooled derooted tulip bulbs (Figs. 2, 3).

It should be mentioned that nonprecooled tulip bulbs used in our experiments were in deep dormancy stage since after planting of the intact bulbs (with roots) only small sprouting took place and no shoot growth was observed but in cooled bulbs normal growth and development of shoot occurred (Fig. 4).

Leaf explants isolated from nonprecooled tulip bulbs and cultured on





Fig. 1. Effect of GA₃ 200 mg l⁻¹, ABA 20 mg l⁻¹ and their mixture on sprouting of shoots (leaves) of nonprecooled derooted bulbs cv. Apeldoorn. Treatments were made on January 22. The photograph (Upper. Left to right: control, GA₃, ABA and GA₃+ABA) was taken on February 4 and the length of sprouts was measured from beginning of experiment until February 5 (Lower). Different letters indicate significant differences according to Duncan's t-test (P=0.05), values are calculated separately for each day of measurements.




Fig. 2. Effect of GA₃ 200 mg l⁻¹, ABA 20 mg l⁻¹ and their mixture on shoot growth and flowering of nonprecooled derooted bulbs cv. Apeldoorn. Treatments were made on January 22 and the plants were photographed on February 23, either with leaves (Upper) or after removal of all leaves before photography (Lower). From left to right: control (distilled water) – no growth of stem and flower bud blasting can be observed, GA₃ – stimulatory effect on shoot growth and flowering, ABA – no growth of stem and flower bud blasting, GA₃+ABA – inhibitory effect of ABA on stem growth and flowering induced by GA₃.




Fig. 3. Effect of GA, 200mg l⁻¹, ABA 20mg l⁻¹ and their mixture on stem (Upper) and leaves (Lower) growth of nonprecooled derooted bulbs cv. Apeldoorn (see Fig. 2). Treatments were made on January 22 and the measurements were made on February 8 and 24. Different letters indicate significant differences according to Duncan's t-test (P=0.05), values are calculated separately for each day of measurements; for total stem length (Upper) and separately for each leaf (Lower).



Fig. 4. Comparison of growth and development of tulips cv. Apeldoorn after planting of nonprecooled (Left) and cooled (Right) bulbs. Bulbs were planted on January 22, photographed on February 10.

Murashige–Skoog (MS) medium (without plant growth regulators) showed slight growth, however, the addition of gibberellic acid to the medium substantially stimulated an increase of length and fresh and dry weight of the explants, cultured in both normal and inverted positions (Kawa and Saniewski, 1990a). The growth of leaf explants isolated

from uncooled bulbs and stimulated by GA₃ was greatly inhibited by simultaneous application of ABA to the medium (Kawa and Saniewski, 1990a). The growth of stem explants (in inverted position), isolated from uncooled and cooled tulip bulbs and induced by auxin (IAA), was also inhibited by ABA applied to the medium together with the auxin (Saniewski and Gabryszewska, 1983; Kawa and Saniewski, 1990b). Thus, it seems that the presence of ABA in uncooled bulbs would inhibit the growth of stem and leaves and keep in dormancy the tulip bulbs. Aung and De Hertogh (1979) suggested that probable sites of abscisic acid biosynthesis in tulip bulbs are the developing bulblets, basal plate and roots, since the ABA level is high in these organs and low in scales and shoot.

The results of present studies showing the inhibitory effect of ABA on shoot growth and flowering induced by GA₃ in nonprecooled tulip bulbs suggest that the hormonal balance, abscisic acid – gibberellins, is the most important factor in control of dormancy release of tulip bulbs, causing shoot growth. The shoot growth is controlled by interaction of auxin with gibberellins as was suggested previously (Okubo and Uemoto, 1985; 1986; Okubo *et al.*, 1986; Saniewski, 1989).

REFERENCES

- Aung, L. H. and A. A. De Hertogh 1967 The occurrence of gibberellin-like substances in tulip bulbs (*Tulipa* sp.). *Plant Cell Physiol.*, **8**: 201–205
- Aung, L. H. and A. A. De Hertogh 1968 Gibberellin-like substances in non-cold and cold treated tulip bulbs (*Tulipa* sp.). In "Biochemistry and Physiology of Plant Growth Substances" ed. by F. Wightman and G. Setterfield, Runge Press, Ottawa, pp. 943–956
- Aung, L. H. and A. A. De Hertogh 1979 Temperature regulation of growth and endogenous abscisic acid-like content of *Tulipa gesneriana* L. *Plant Physiol.*, **63**: 1111–1116
- Aung, L. H. and A. R. Rees 1974 Changes in endogenous gibberellin levels in *Tulipa* bulblets during ontogeny. *J. Exp. Bot.*, **25**: 745–751
- Cocoza Talia, M. and P. Stellacci 1979 A research on the effect of gibberellin upon tulip flowering. *Acta Hort.*, **91**: 167–172
- De Hertogh, A. A., L. H. Aung, A. E. Einert and G. L. Staby 1971 Aspects of extractable gibberellin-like substances from *Tulipa gesneriana* L. *Acta Hort.*, **23**: 90–95
- Hanks, G. R. 1984 Factors affecting the response of tulips to gibberellins. *Scientia Hort.*, **23**: 379–390
- Hanks, G. R. 1985 The response of 9°C-tulips to gibberellins. *Scientia Hort.*, **27**: 153–161
- Hanks, G. R. and A. R. Rees 1980 Growth substances of tulip: the activity of gibberellin-like substances in field-grown tulips from planting until flowering. *Z. Pflanzenphysiol.*, **98**: 213–223
- Jones, S. K. and G. R. Hanks 1984 Treatment of tulips with gibberellic acid by vacuum infiltration. *J. Hort. Sci.*, **59**: 241–252
- Kawa, L. and M. Saniewski 1986 The effect of gibberellic acid and abscisic acid on tulip pistil growth in vitro. *Acta Hort.*, **177**: 129–133
- Kawa, L. and M. Saniewski 1990a The effect of plant growth regulators on tulip leaf growth in vitro (in Polish). *Prace Inst. Sad. i Kwiac., Ser. B – Rostliny Ozdobne*, **15**: 73–82
- Kawa, L. and M. Saniewski 1990b The effect of plant growth regulators on tulip stalk elongation in vitro (in Polish). *Prace Inst. Sad. i Kwiac., Ser. B – Rostliny Ozdobne*, **15**: 83–94
- Okubo, H. and S. Uemoto 1985 Changes in endogenous gibberellin and auxin activities during first internode elongation in tulip flower stalk. *Plant Cell Physiol.*, **26**: 709–719
- Okubo, H. and S. Uemoto 1986 Changes in endogenous gibberellin during internode elongation of tulip flower stalk affected by light conditions. *J. Japan. Soc. Hort. Sci.*, **55**: 215–220
- Okubo, H., S. Shiraishi and S. Uemoto 1986 Factors controlling elongation of the last internode in tulip flower stalk. *J. Japan. Soc. Hort. Sci.*, **55**: 320–325

- Rakhimbayev, I. R., G. A. Syrtanova and V. F. Solomina 1978 The effect of cold treatment on the level of biological activity of endogenous growth regulators of tulip bulbs (in Russ.). *Plant Physiol. (Russ.)*, **25**: 249–253
- Rudnicki, R. M., J. Nowak and M. Saniewski 1976 The effect of gibberellic acid on sprouting and flowering of some tulip cultivars. *Scientia Hortic.*, **4**: 387–397
- Saniewski, M. 1989 The use of paclobutrazol, an inhibitor of gibberellin biosynthesis, for study of hormonal control of tulip stem elongation. *Bull. Pol. Acad. Sci. Biol. Sci.*, **37**: 55–64
- Saniewski, M. and E. Gabryszewska 1983 Hormonal control of tulip stalk elongation in vitro. *Acta Hortic.*, **131**: 275–282
- Saniewski, M., L. Kawa-Miszczak, E. Węgrzynowicz-Lesiak and H. Okubo 1999 Gibberellin induces shoot growth and flowering in nonprecooled derooted bulbs of tulip (*Tulipa gesneriana* L.). *J. Fac. Agr., Kyushu Univ.*, **43**: 411–418
- Singh, B. N., E. Galston, W. Dashek and D. C. Walton 1979 Absciscic acid levels and metabolism in the leaf epidermal tissue of *Tulipa gesneriana* L. and *Commelina communis* L. *Planta*, **146**: 1435–1438
- Syrtanova, G. A., I. R. Rakhimbayev and W. I. Kefeli 1975 Native growth inhibitor of tulip (in Russ.). *Plant Physiol. (Russ.)*, **22**: 165–169
- Syrtanova, G. A., R. Turetzkaja and I. R. Rakhimbayev 1973 Natural auxin and inhibitors in dormant and growing tulip bulbs (in Russ.). *Plant Physiol. (Russ.)*, **20**: 1133–1136
- Terry, P. H., L. H. Aung and A. A. De Hertogh 1982 Identification of absciscic acid in *Tulipa gesneriana* L. by gas-liquid chromatography and mass spectroscopy. *Plant Physiol.*, **70**: 1574–1576
- Van Bragt, J. 1971 Endogenous gibberellin levels and floral stalk elongation in tulip cv. Apeldoorn. *Landbouwwetenschappen Gent*, **36**: 1301–1307
- Van Bragt, J. and K. J. Van Ast 1976 Substitution of the cold requirement of tulip cv. Apeldoorn by GA₃. *Scientia Hortic.*, **4**: 117–122
- Van Bragt, J. and F. A. Zijlstra 1971 Effects of gibberellins on flowering of tulip cv. Apeldoorn. *Z. Pflanzenphysiol.*, **64**: 139–144