Catastrophic flood in eastern Slovakia

Majtan, Stefan

Laboratory of Soil and Water Conservation, Faculty of Agriculture, Kyushu University

Omura, Hiroshi

Laboratory of Soil and Water Conservation, Faculty of Agriculture, Kyushu University

Jelinek, Robert

Geological Institute of Slovak Republic

<https://doi.org/10.5109/24322>

出版情報 : 九州大学大学院農学研究院紀要. 44 (1/2), pp.213-217, 1999-11. Kyushu University
バージョン :
権利関係 :

Catastrophic flood in eastern Slovakia

Štefan Majtán*, Hiroshi Omura*, Róbert Jelínek**

*Laboratory of Soil and Water Conservation, Faculty of Agriculture, Kyushu University

**Geological Institute of Slovak Republic, Banská Bystrica, Slovakia

(Received July 21, 1999 and accepted August 24, 1999)

During last year's July, a significant flood struck areas in northwest Eastern Slovakia, 560 kilometers northeast of the capital of Slovakia Bratislava. The affected region is located 30 kilometers northwest of Prešov, the center town of Prešov District.

On 20 July 1998 a heavy rain storm struck the villages of Jarovnice, Užovské Pekľany and Renčíšov which are located in the upper part of a 12 km valley. The heavy rainfall which fell over 90 minutes caused an unexpected catastrophic flood wave.

INTRODUCTION

In the last years, extensive and heavy rains caused significant flooding in several areas of eastern Slovakia in the Slovak Republic. The most tragic floods have been recorded in July 1997 and also in July 1998.

The most affected areas in Slovak Republic were especially the villages Jarovnice, Užovské Pekľany and Renčíšov in 1998 and Tichý Potok in 1997. In parts of valley where the width is only from 40 to 60 meters, the flood wave reached a height from 4 to 5 meters. There were 55 confirmed deaths. The estimation of general damages has reached 1.2 billion Sk.

Key words: *disaster, flooding, flood wave.*

NATURAL SETTING OF DISASTER AREA

Slovak Republic is located in the Central Europe. Climate in Slovakia is continental with hot summers and cold winters. Average annual precipitation in Slovakia Republic is about 800 mm, in disaster area is between 550-650 mm. The annual temperature of disaster area is 7°C at south part of Torysa's valley and 4°C at peaks of mountain. The coldest month is January with average minimum temperature -4°C. The warmest one is July with average maximum temperature 18°C (http SHMU). The highest precipitation month is July and the lowest one is February.

A main river Torysa with its main feeds Sekčov and Delna drains disaster area. The second main river is Svinka on which a catastrophic flow occurred.

Location of disaster

Renčíšov village is a small hamlet located in the uppermost part of valley. One week after the flood the village was accessible only by helicopters and heavy army mashinery. The village has 180 inhabitants. Approximately 90% of the village roads were destroyed as well as the homes and property of the inhabitants.

Užovské Pekľany village has suffered the greater damage to the property. Twenty percent of the 640 inhabitants are Roma families. The village is located in the narrowest part of the afflicted valley.

In *Jarovnice* village, with a population of approximately 4000, the majority of residents are Roma families. The majority of deaths occurred in this village and mostly among Roma people.

Meteorological situation

The center of rainfall was above upstream part of Malá Svinka watershed (Fig. 1). According to meteorologist, there was a thunderstorm with two, three cores respectively. The course of flooding was very dramatic. The thunderstorm, with which caused the most catastrophic flood wave, starts from about 3.30 p.m. to 5 p.m., and the first flood occurred within an hour since rain started.

Fig. 1. Schematic map of core rainfall with marked profile selected for calculation of overflow and volume of flood (http Region of Prešov).

RESULT AND DISCUSSION

Simulation of flood wave

The most affected village was Užovské Pekľany. The flood wave started growing up one or one and half hour after rainfall started. The peak of flood wave was reached approximately after two or two and half hour after starting rain. The eligible velocity of flood wave was about $2\text{--}2.5\text{ m}\cdot\text{s}^{-1}$, that means the critical time of flooding in Užovské Pekľany should have value 80–90 minutes.

From these conjectures, hydrologists consider, that the overflow in this village was due to whole rainfall and runoff from the whole watershed (Fig. 2). Also, they are considering the culmination overflowing should be reached up to $190\text{ m}^3\cdot\text{s}^{-1}$ (Abaffy, Kadubec 1998). In village Dubovica, the same situation should be also, but with lower culmination of overflow about $160\text{ m}^3\cdot\text{s}^{-1}$ (Abaffy, Kadubec 1998).

Fig. 2. Calculated course of flooding wave in Užovské Pekľany village.

Causes of flooding

Such catastrophic result of flooding signed especially these factors: rainfall, saturation of watershed by previous rainfall and also lower retentive capacity of area.

Rainfall – as mentioned above, the trigger of flood was heavy rainfall. In the center of thunder cloud it should be reached up to 130 mm. In the short intervals, the intensity of rainfall reached to $3\text{ mm}\cdot\text{min}^{-1}$ and more, which is representative maximum intensity of rainfall in Slovakia.

Saturation of watershed – in 20 July in same area the intensive rainfall occurred.

Lower retentive capacity of area – from this point of view the most important factor is hydrogeological structure of area. From an engineering geological point of view, the disaster area is located in the region of the Flysch. Simply, the Flysch complexes are

composed of beds of sandstones and shales. The complexes with a steep slopes which consist of Flysch formation are incoherent for sliding, break loose following by sliding respectively, after the saturation by water.

Secondary, but also significant catastrophic factors were breach waves. The original places of these waves, were barriers formed by wafting materials (soil, uproot trees and timber, building materials etc.).

Brief account of flooding waves

The hydrologists from Slovak hydrometric institute reconstructed development of flood waves. Based on this simulation, they are deduced, that in villages Renčiov, Užovské Pekľany, Jarovce and Dubovica were in the local stream more than 1000 years recurrence overflows as shown Tab. 1 and Fig. 3 respectively. Their destructive energy was powered by above mentioned breach waves.

Fig. 3. Shows overflow in selected profiles in order 100 years recurrence, 1000 years recurrence and calculated overflow.

Table 1. Calculated overflow and volume of flood in selected profiles (taken from Majerčáková, Škoda 1998). The location of profile shows Fig. 1.

Profile	Length [km]	Area [km²]	Forest [%]	Slope watershed [%]	Slope stream [‰]	Calculated overflow [m³.s⁻¹]	Calculated volume of flood [m³]
1	4.2	6.452	< 50	23	8	90	400000
2	4.3	7.065	> 50	28	9	95	425000
3	6.5	13.52	> 50	25	8	140	825000
4	7.8	24.26	60	26	6	190	1330000
5	13.3	35.39	50	20	8	230	1900000
6	4.45	10.9	50	24	9	120	650000
7	7.35	15.244	30	23	6	160	850000

Facts and numbers about catastrophic flood

The flooding wave on stream Malá Svinka, Svinka, Žehrica, Torysa and their feeds caused big damages along their course. This water induced disaster killed 55 lifes and 5804 animals. The flood directly affected 10850 people in 82 settlements. Shelter 756 people were damaged. During the rescue works 3618 people were evacuated. In the befallen area 2059 houses were submerged, 272 houses were destroyed. Also, 13881 ha agriculture land and 4398 ha meadow were flooded. Table 2. shows brief account of damages caused by flooding in July 1998 at Prešov and Košice district.

Table 2. Main catastrophic affect of flood occurred on 20. July in eastern part of Slovakia (taken from Abaffy, Kadubec 1998).

<i>Direct affect of flood</i>	<i>District</i>		<i>Together</i>
	Prešov	Košice	
City and villages	71	11	82
Overflow houses	1484	575	2059
Overflow sources of drinking water	991	476	1467
Damaged roads [km]	87.74	22.8	110.54
Damaged bridges and foot – bridges	165	32	197
Affected population	8379	2471	10850
Thence without shelter	747	9	756
Evacuee	3608	10	3618
Death persons	55	0	55
Injured persons	60	1	61

CONCLUSION

From intensity, extremity, extension of damages points of view, the flood in July 1998 was absolutely extraordinary hydrological phenomena in this century in Slovakia.

Sad to say, that so extraordinary phenomena have not been registered by an existing net of meteorological station. Concentrated rainfall in certain limited area, like in this cause, indicates that point measurement of rainfall can not record such extreme storm. Therefore, alternative area measurement methods should be used to get the best result.

REFERENCES

- Abaffy, Kadubec 1998 – Fakty a čísla o priebehu a následkoch povodní v Prešovskom a Košickom kraji v júli 1998. Vodohospodárstvo 8/98, pp. 13–17. in Slovak.
 http Region of Prešov – Information of Prešov district, Slovakia – <http://www.presov.sk>.
 http SHMU – Web page of Slovak Hydrometeorological Institute in Bratislava, Slovakia – <http://www.shmu.sk>.
 Majerčáková, Skoda 1998 – Prívalové povodne na severovýchodnom Slovensku. Vodohospodárstvo 8/98, pp 18–19. in Slovak.