Effects of Calcium Supplement to Purified Diets on Growth and Bone Mineralization in Juvenile Red Sea Bream and Black Sea Bream

Hossain M., Amzad

Laboratory of Fish Production Technology, Division of Fish Production Technology, Department of Animal and Marine Bioresource Science, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Furuichi, Masayuki

Laboratory of Fish Production Technology, Division of Fish Production Technology, Department of Animal and Marine Bioresource Science, Faculty of Agriculture, Kyushu University

<https://doi.org/10.5109/24398>

出版情報：九州大学大学院農学研究院紀要. 45 (2), pp.481-486, 2001-02-28. Kyushu University
バージョン：
権利関係：

Effects of Calcium Supplement to Purified Diets on Growth and Bone Mineralization in Juvenile Red Sea Bream and Black Sea Bream

M. Amzad Hossain* and Masayuki Furuichi

Laboratory of Fish Production Technology, Division of Fish Production Technology,
Department of Animal and Marine Bioresource Science, Faculty of Agriculture,
Kyushu University, Fukuoka 812–8581, Japan

(Received October 30, 2000 and accepted November 10, 2000)

Juvenile red sea bream (av. body wt. 1.4 g) and black sea bream (av. body wt. 0.5 g) were fed two purified casein based diets containing 0.34% (control) and 0.03% (unsupplemented) dietary calcium (Ca) for 10 weeks at $27.0 \pm 1.4^\circ\text{C}$. Final body weight, condition factor and hepatosomatic index were not affected by dietary Ca in both species. Feed efficiency in black sea bream was almost same between two dietary groups, however, feed efficiency in red sea bream fed the Ca unsupplemented diet were slightly lower than that in fish fed the control diet. Dietary Ca levels did not affect blood characteristics of red sea bream and black sea bream. In both species Ca and other mineral contents of the bone were independent of a dietary Ca supplement except that K and Mn contents of red sea bream fed the Ca unsupplemented diet were significantly lower than those in the control. It appeared that Ca supplements to the purified diets are not necessary in juvenile red sea bream and black sea bream.

INTRODUCTION

In the previous studies we observed that red sea bream with initial body weight of 11.5 g and black sea bream with initial body weight of 8.0 g did not require a dietary calcium (Ca) supplement (Hossain and Furuichi, 1999a, b). However, a dietary Ca supplement was necessary in tiger puffer, redlip mullet and Japanese flounder with initial body weight of 1.6 g, 0.6 g and 0.7 g, respectively (Hossain and Furuichi, 1998, 2000a, b). From those results it was considered that the unnecessary of Ca supplement to the diet detected in red sea bream and black sea bream might have been due to the larger initial body weight. Satoh *et al.* (1987) observed that whole body mineral contents of rainbow trout did not change after body weight reached 2.0 g, and recommended to conduct the feeding experiment for mineral study with fish less than 2.0 g body weight. The present study was conducted to investigate whether dietary Ca supplements were necessary to juvenile red sea bream and black sea bream.

* Laboratory of Fish Production Technology, Division of Fish Production Technology, Department of Animal and Marine Bioresource Science, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University. Corresponding author. Present address: Marine Fisheries and Technology Station, Bangladesh Fisheries Research Institute, Cox's Bazar–4700, Bangladesh.
E-mail: m_amzad_hossain@hotmail.com.

MATERIALS AND METHODS

Experimental Diets and Feeding Regime

Two purified diets with or without Ca supplement were formulated (Table 1). Vitamin-free milk casein, dextrin and α -starch (gelatinized potato starch), and pollack liver oil were used as protein, digestible carbohydrate and lipid sources, respectively. A mixture of vitamins (Halver, 1957) was added equally to the diets. The diet containing

Table 1. Composition of the experimental diets for red sea bream and black sea bream

Diet	Control	No Ca supplement
<i>Ingredient (%)</i>		
Casein	50	50
Amino acid mix.* ¹	4	4
α -Potato starch	7	7
Dextrin	10	10
Pollack liver oil	10	10
Vitamin mix.* ²	3	3
Mineral mix.* ³	6	6
Carboxymethylcellulose	4	4
Ca-lactate	2.308	—
α -Cellulose	3.692	6

*¹ Amino acid mixture (g/kg diet): arginine·HCl, 10; alanine, 10; glycine, 10; aspartate·Na, 10.

*² Vitamin mixture (Halver, 1957).

*³ Mineral mixture (mg/kg diet): KCl, 3840; MgSO₄·5H₂O, 4080; NaH₂PO₄·2H₂O, 34260; Fe-citrate, 1200; AlCl₃·6H₂O, 45; ZnSO₄·7H₂O, 132; MnSO₄·5H₂O, 877; CuCl, 7.9; KI, 1.9; CoCl₂·6H₂O, 0.7.

Table 2. Proximate and mineral compositions of the experimental diets for red sea bream and black sea bream

Diet	Control	No Ca supplement
<i>Proximate composition (% dm)*</i>		
Moisture	20.9	21.5
Crude protein	51.3	52.0
Crude lipid	9.1	9.3
Crude ash	5.1	5.0
<i>Mineral composition (dm)</i>		
Ca (%)	0.34	0.03
P (%)	1.00	1.05
K (%)	0.19	0.18
Mg(μ g/g)	400	370
Fe (μ g/g)	280	270
Zn(μ g/g)	48	52
Mn(μ g/g)	16	16
Cu(μ g/g)	11	12

* dm, dry matter.

0.3% Ca from Ca-lactate was control. The method of diet preparation was the same as that reported previously (Hossain and Furuichi, 1999b). The proximate and mineral compositions of the diets are shown in Table 2.

Fish and Rearing Methods

Juvenile red sea bream and black sea bream were accustomed to a casein diet in 100-ℓ circular tanks for 2 weeks before the experiments. At the start of the experiment, the fish of each species were weighed individually, selected, and distributed to four tanks (duplicate per treatment) in such a manner that average body weight and size variation of fish in each tank were similar. The number of fish in a tank was 30 (average body weight 1.4 g) for red sea bream and 40 (average body weight 0.5 g) for black sea bream. The fish were reared for 10 weeks, during which the experimental diets were fed twice a day at 09:00 and 15:30 h. Diets were offered to the fish until satiation. Each tank was provided with adequate oxygen by aeration and continuous sea water (2–3 ℓ/min) supplies. Water temperature was $27.0 \pm 1.4^\circ\text{C}$ during the rearing period. A Ca level in the rearing water was around 400 mg/l. Other rearing conditions and methods were the same as those reported previously (Hossain and Furuichi, 1999b).

Sampling and Analysis

At the end of the feeding trial, the body weight and body length of the fish were measured individually to calculate the condition factor. Blood samples were collected from the cuvierian duct with a 1 ml heparinized syringe. Each liver was weighed to calculate the hepatosomatic index. Pooled samples of all the livers and whole carcasses were stored at -20°C for chemical analysis and bone collection, respectively.

Analyses of blood, bone and diet samples were the same as those described previously (Hossain and Furuichi, 1999b).

Student's T-test was applied to determine the significance in difference between the treatment means ($P < 0.05$).

RESULTS

Red Sea Bream

Growth performances and feed utilization of red sea bream are shown in Table 3. The survival rates at the end of the experiment were 96.7% and 100% in fish fed the control and the Ca unsupplemented diets, respectively. No significant difference was observed in the final body weight between two treatment groups. Feed efficiency in fish fed the Ca unsupplemented diet was slightly lower than that in fish fed the control diet. Condition factor and hepatosomatic index were not affected by dietary Ca levels. Hematocrit values, hemoglobin and plasma protein contents were similar in both dietary groups (Table 4).

The results of bone analyses are presented in Table 5. Bone lipid and ash contents were similar in both groups. Ca and P contents of the bone appeared to be independent of a dietary Ca supplement. A deletion of supplemental Ca from the diet did not affect the bone Mg, Fe, Zn and Cu contents. However, lower bone K and Mn contents were detected in fish fed the Ca unsupplemented diet compared to the control diet.

Table 3. Growth and feed utilization of red sea bream and black sea bream fed the experimental diets with or without Ca supplement

Diet	Control	No Ca supplement
<i>Red sea bream</i>		
Av. body wt. (g)		
Initial	1.42	1.42
Final* ¹	14.4	13.9
Weight gain (%)	910	880
Feed efficiency (%)	83.4	75.7
Condition factor* ^{1,2}	3.28	3.37
HSI* ^{1,3}	2.24	2.34
Survival rate (%)	96.7	100
<i>Black sea bream</i>		
Av. body wt. (g)		
Initial	0.54	0.54
Final* ¹	12.6	12.3
Weight gain (%)	2230	2180
Feed efficiency (%)	104.5	105.6
Condition factor* ^{1,2}	2.87	2.82
HSI* ^{1,3}	1.84	1.85
Survival rate (%)	92.5	96.3

*¹ No significant difference ($P>0.05$).*² Body weight (g) $\times$ 100/(body length in cm)³.*³ Hepatosomatic index: Liver weight (g) $\times$ 100/body weight (g).**Table 4.** Hematocrit, hemoglobin and plasma protein of red sea bream and black sea bream fed the experimental diets with or without Ca supplement

Diet	Control	No Ca supplement
<i>Red sea bream</i>		
Hematocrit (%)*	38.5	38.0
Hemoglobin (g/100 ml)*	3.3	3.4
Plasma protein (g/100 ml)*	4.4	3.9
<i>Black sea bream</i>		
Hematocrit (%)*	38.0	38.0
Hemoglobin (g/100 ml)*	8.6	8.9
Plasma protein (g/100 ml)*	5.4	5.5

* No significant difference ($P>0.05$; n=10).

Black Sea Bream

Survivals of black sea bream were 92.5% and 96.3% in fish fed the control diet and Ca unsupplemented diet, respectively (Table 3). Final body weight was not statistically different between two dietary groups. Feed efficiency, condition factor and hepatosomatic index were not affected by a Ca unsupplementation to the diet. Effects of a deletion of dietary Ca were not observed in hematocrit, hemoglobin and plasma protein

Table 5. Lipid, ash, and mineral composition of the bone of red sea bream and black sea bream fed the experimental diets with or without Ca supplement*¹

Diet	Control	No Ca supplement
<i>Red sea bream</i>		
Crude lipid (%)	20.5	20.3
Crude ash (%)	55.8	56.0
Ca (%)	27.3	27.4
P (%)	10.5	10.6
Mg(%)	0.44	0.42
K (μg/g)	58	39* ²
Fe (μg/g)	208	200
Zn(μg/g)	106	104
Mn(μg/g)	48	38* ²
Cu(μg/g)	7.1	6.7
<i>Black sea bream</i>		
Crude lipid (%)	19.2	19.0
Crude ash (%)	62.0	63.0
Ca (%)	25.6	25.5
P (%)	10.2	10.3
Mg(%)	0.67	0.63
K (μg/g)	82.8	90.5
Fe (μg/g)	200	198
Zn(μg/g)	155	159
Mn(μg/g)	39.0	40.3
Cu(μg/g)	7.9	7.3

*¹ Dry matter basis.*² Significant difference ($P < 0.05$).

values of the fish (Table 4). Ash, lipid and mineral contents of the bone were identical between the control and Ca unsupplemented diet groups (Table 5).

DISCUSSION

Since the growth of fish was almost identical between the control and Ca unsupplemented diets in both red sea bream and black sea bream, these two fish species may absorb adequate Ca from sea water for their metabolic requirements. Previous studies showed that dietary Ca supplements were not necessary for relatively larger sized red sea bream and black sea bream (Hossain and Furuichi, 1999a, b). The present study is further confirmation of the previous studies, and revealed that these two species did not need a Ca supplement to a purified diet irrespective of the fish size. The similarity in Ca requirements in these two species may be for the reason that both species belong to the same family and have a similar food habit. The present results also are in accordance with a well accepted concept that fish have a capability to absorb Ca from surrounding water and a dietary supplement may not be necessary (Love, 1980). Sakamoto and Yone (1976) also reported that a Ca supplement was dispensable in a purified diet for red sea bream. It appeared from the growth, hematological, and liver and bone mineralization

data in the present study that Ca absorption from sea water might be adequate for red sea bream and black sea bream. In the study with radio isotope, we observed that red sea bream (average body weight 0.7 g) and black sea bream (average body weight 0.8 g) had increased the absorption of ^{45}Ca from sea water when Ca was deleted from a purified diet and overall absorption of Ca was very high in both species (Hossain and Furuichi: Fish. Res. Lab., Kyushu Univ., unpubl. data). Sakamoto (1981) also reported that the absorption of radioactive Ca in red sea bream increased with the decrease of dietary Ca levels. Therefore, the present investigation suggests that a dietary Ca supplement may not be necessary for red sea bream and black sea bream. On the other hand, environmental Ca was found not to be sufficiently available in tiger puffer, redlip mullet and Japanese flounder (Furuichi *et al.*, 1997, 1999; Hossain and Furuichi, 1998, 2000a, b). Therefore, further investigations are important to find out why Ca absorption from sea water is different among marine fish species.

ACKNOWLEDGEMENTS

We would like to express our sincere thanks to Dr. Seiichi Matsui and Dr. Takao Yoshimatsu for their kind help in conducting this research. This is the contribution no. 250 from Fishery Research Laboratory, Kyushu University.

REFERENCES

- Furuichi, M., Y. Furusho, M. A. Hossain, S. Matsui and R. Azuma 1997 Essentiality of Ca supplement to white fish meal diet for tiger puffer. *J. Fac. Agr., Kyushu Univ.*, **42**: 77–85
- Furuichi, M., M. A. Hossain and S. Matsui 1999 Effect of dietary supplemental calcium levels on growth and mineral composition of bone in tiger puffer. *Suisanzoshoku*, **47**: 409–413
- Halver, J. E. 1957 Nutrition of salmonid fishes. III. Water-soluble vitamin requirements of chinook salmon. *J. Nutr.*, **62**: 225–243
- Hossain, M. A. and M. Furuichi 1998 Availability of environmental and dietary calcium in tiger puffer. *Aquacult. Int.*, **6**: 121–132
- Hossain, M. A. and M. Furuichi 1999a Effect of deletion of calcium supplement from purified diet on growth and bone mineralization in red sea bream. *J. Fac. Agr., Kyushu Univ.*, **44**: 91–97
- Hossain, M. A. and M. Furuichi 1999b Necessity of dietary calcium supplement in black sea bream. *Fisheries Sci.*, **65**: 893–897
- Hossain, M. A. and M. Furuichi 2000a Essentiality of dietary calcium supplement in redlip mullet *Liza haematocheila*. *Aquacult. Nutr.*, **6**: 33–38
- Hossain, M. A. and M. Furuichi 2000b Necessity of calcium supplement to the diet of Japanese flounder. *Fisheries Sci.*, **66**: 660–664
- Love, R. M. 1980 *The Chemical Biology of Fishes*, Vol. 2, Academic Press, New York
- Sakamoto, S. 1981 Requirements and deficiency symptoms of dietary minerals in red sea bream. *Rep. Fish. Res. Lab., Kyushu Univ.*, **5**: 1–99
- Sakamoto, S. and Y. Yone 1976 Requirement of red sea bream for dietary Ca. *Rep. Fish. Res. Lab., Kyushu Univ.*, **3**: 59–64
- Satoh, S., T. Takeuchi and T. Watanabe 1987 Changes of mineral compositions in whole body of rainbow trout during growing stages. *Nippon Suisan Gakkaishi*, **53**: 273–279