Change in phenoloxidase and its precursor during silkworm (a80 strain) development

Yamamoto, Kohji
The University of Texas Health Center at Tyler

Fujii, Hiroshi
Laboratory of Silkworm Genetics, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Kusakabe, Takahiro
Laboratory of Sericultural Science, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Koga, Katsumi
Laboratory of Sericultural Science, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

他

<https://doi.org/10.5109/24399>

出版情報：九州大学大学院農学研究院紀要. 45 (2), pp.487-493, 2001-02-28. Kyushu University
バージョン：
権利関係：

Change in phenoloxidase and its precursor during silkworm (a80 strain) development

Kohji Yamamoto^{1,4)}, Hiroshi Fujii²⁾, Takahiro Kusakabe³⁾, Katsumi Koga³⁾,
Yoichi Aso⁴⁾, and Masatsune Ishiguro^{*4)}

¹⁾ The University of Texas Health Center at Tyler, 11937 HWY 271 Tyler TX 75708, U. S. A.

²⁾ Laboratory of Silkworm Genetics, ³⁾ Laboratory of Sericultural Science, and

⁴⁾ Laboratory of Protein Chemistry and Engineering, Graduate School of
Bioresource and Bioenvironmental Sciences, Kyushu University,
6–10–1 Hakozaki, Higashi-ku, Fukuoka 812–8581, Japan.

(Received October 31, 2000 and accepted November 10, 2000)

The changes in phenoloxidase and its precursor, so-called prophenoloxidase, of the silkworm, *Bombyx mori* (a80 strain) were examined using detection assay with or without detergent to distinguish PO and pro-PO in hemolymph (Yamamoto, *et al.*, 1999). Little or no phenoloxidase activity was detected in hemolymph at the silkworm development during the final larval instar and pupal stage. On the other hand, prophenoloxidase was detected for same time frame. In males, after prophenoloxidase increased slightly at day 4, it decreased to a minimum level at spinning and increased to a maximum level on the day of pupation and decreased gradually to day 5 after pupation. In females, prophenoloxidase decreased gradually to a minimum level on day of spinning and a slight increase was observed at day 8 and it increased to maximum level between day 2 and 3 after pupation and decreased thereafter. Prophenoloxidase mRNA expression was preceded a day or two before, when comparing the change of phenoloxidase mRNA with its product.

INTRODUCTION

Phenoloxidase (PO: monophenol, dihydroxyphenylalanine; oxygen oxidoreductase; EC 1. 14. 18. 1) catalyzes two sequential reactions; the hydroxylation of monophenol to *o*-diphenol and the oxidation of *o*-diphenol to *o*-quinone (Ashida and Yamazaki, 1990). In insects, PO is generally believed to be crucial for the cuticular melanization and sclerotization (Söderhäll, 1982; Hiruma and Riddiford, 1988; Sugunaran *et al.*, 1992).

The existence of a precursor protein, called prophenoloxidase (pro-PO), was reported from many insects: e.g., the fruit fly *Drosophila melanogaster* (Fujimoto *et al.*, 1993), the silkworm *Bombyx mori* (Yasuhara, *et al.*, 1995; Yamamoto, *et al.*, 1999), the tobacco hornworm *Manduca sexta* (Hall, *et al.*, 1995), the cockroach *Blaberus discoidalis* (Durrant, *et al.*, 1993), the wax moth *Galleria mellonella* (Kopacek, *et al.*, 1995), and a coleopteran insect *Holotrichia diomphalia* (Kwon, *et al.*, 1997). The activation of pro-PO have been shown by the prophenoloxidase-activating factor *in vitro* (Lee, *et al.*, 1998; Jiang, *et al.*, 1998; Satoh, *et al.*, 1999) and by organic compounds such as sodium dodecyl sulfate (Funatsu and Inaba, 1962; Inaba and Funatsu, 1964), cetylpyridinium chloride (Hall, *et al.*, 1995), 2-propanol (Asada, *et al.*, 1993) and dimethylbenzylmyristylammonium chloride (DBMA) (Yamamoto, *et al.*, 1999). Furthermore, the

* Corresponding author. FAX number: 092–642–3051, E-mail: ishiguro@agr.kyushu-u.ac.jp.

cDNA encoding pro-PO have been cloned and sequenced recently from the mosquito *Armigeres subalbatus* (Cho, *et al.*, 1998), the malaria vector *Anopheles gambiae* (Jiang, *et al.*, 1997), the fall webworm *Hypantria cunea* (Park, *et al.*, 1997), the coleopteran insect *Tenebrio molitor* (Lee, *et al.*, 1999), the fruit fly *D. melanogaster* (Fujimoto, *et al.*, 1995), the tobacco hornworm *M. sexta* (Hall, *et al.*, 1995), and the silkworm *B. mori* (Kawabata, *et al.*, 1995; Yamamoto, *et al.*, 2000).

Although many knowleges about pro-PO have been accumulated as described above, the biochemical and genetical aspects of the regulation of PO activity during development is still poorly understood. In housefly, *Musca domestica*, the change of PO activity was examined from larvae to pupae (Funatsu and Inaba, 1962). The authors found that the noticeable change in PO activity occurred at the stage of pupation. Almost the highest PO activity in the homogenate of the final instar larvae disappeared suddenly in the homogenate of the prepupae and then it is appeared again in the homogenate of aged pupae. However, the disappearance of PO activity in the prepupae did not imply that the enzyme vanished because the homogenate of the prepupae exhibited PO activity upon the addition of an anionic detergent such as sodium oleic acid or an extract from the aged pupae. It means that phenoloxidase is present as inactive form in prepupae and there is natural activator, which converts inactive form to active form, in aged pupae.

The fluctuation of PO activity during metamorphosis in the silkworm has never investigated. In order to get a better understanding of the regulatory events involved in the metamorphosis of the silkworm, we analyzed the change in PO activity and pro-PO during developmental stages. The comparison of these fluctuations with that for pro-PO mRNA was also included.

MATERIALS and METHODS

Insects and preparation of hemolymph and hemocytes

The a80 strain of *B. mori* was reared as described previously (Yamamoto, *et al.*, 1999). Females and males were distinguished by the imaginal bud on the abdominal surface. The hemolymph specimens collected into liquid nitrogen, were lyophilized and the powder obtained was used for the determination of total PO activity.

Measurement of PO activity in the crude extract

PO activity was measured spectrophotometrically with or without a cationic detergent, DBMA, as an activator (Yamamoto, *et al.*, 1999). The lyophilized hemolymph (5 mg) from various days was dissolved in 1 ml of 10 mM potassium phosphate buffer (pH 7.0). Fifty μ l of this mixture was added to the reaction mixture (1 ml) containing 0.05 M potassium phosphate buffer (pH 6.5), 5 mM L-3-(3,4-dihydroxyphenyl) alanine (L-dopa) and with or without 0.4 mM DBMA. While incubating at 30°C for 0.5 min, the absorbance at 475 nm was monitored with a Hitachi U-3210 spectrophotometer (Tokyo, Japan). One unit of PO was defined as the amount of active PO capable of producing 1 μ mole of dopachrome per min.

Measurement of pro-PO mRNA titer

Northern hybridization was done according to previous paper (Yamamoto, *et al.*,

2000). Level of pro-PO mRNA was assessed by scanning of the electrophoregram by densitometry using NIH image software and replotted.

RESULTS and DISCUSSION

Change in PO and pro-PO at developmental stages

The fluctuation of PO activity during metamorphosis in the silkworm has never investigated. Therefore, we measured the fluctuation of PO or pro-PO and we made better use of detection assay with or without DBMA to distinguish PO and pro-PO in hemolymph.

By using this method, little or no PO was detected, even if the metamorphosis from larva to pupar was occurred (Fig. 1). In housefly, however, the endogeneous PO activity was detected in the homogenate of the final instar larvae, and then it was disappeared suddenly in the homogenate of the prepupae (Funatsu and Inaba, 1962). The presence of active PO in hemolymph is thought to be harmful to the insects health and it is necessary to activate it during a need. Therefore, insects need to control active PO. Recently, endogeneous PO inhibitors were identified in the housefly, *M. domestica* (Daquinag, *et al.*, 1995) and in the tobacco hornworm, *M. sexta* (Sugumaran and Nellaippan, 2000) and both were shown to inhibits directly the activity of PO. The PO inhibitor of housefly could inhibit PO activity of silkworm (Yamamoto, *et al.*, 2000, unpublished results). This fact suggests that there might be the strong inhibitor in the silkworm that involves in the regulation system of PO activity, because active PO cannot be detected.

When the changes in pro-PO was determined in hemolymph at developmental stages from fifth larval instar to pupa, pro-PO was detected for thirteen days from larva to pupa (Fig. 1). In males, after pro-PO increased slightly at day 4, it decreased to a minimum level (1.39×10^{-3} unit/mg lyophilized hemolymph) at spinning (day 6) and increased to a maximum level (7.54×10^{-3} unit/mg lyophilized hemolymph) on the day of pupation and decreased gradually to day 5 after pupation. In females, pro-PO decreased gradually to a minimum level on day of spinning (2.41×10^{-3} unit/mg lyophilized hemolymph) and a slight increase was observed at day 8 and it increased to maximum level ($9.80\text{--}9.74 \times 10^{-3}$ unit/mg lyophilized hemolymph) between day 2 and 3 after pupation and decreased thereafter. Pro-PO is always present in this period, whereas no active PO was detected. As shown in Fig. 1, the large amount of pro-PO is stored during a few days after pupation. It is postulated that pro-PO in hemolymph was thought to be stored in case of emergency. Xiao-Qiang *et al.* demonstrated that after injection of bacteria, novel lectin mRNA appeared and lead to stimulate the activation of pro-PO (Yu, *et al.*, 1999).

There is a possibility that some chymotrypsin inhibitors had the ability to inhibit pro-PO activating factor of the same species (Aso, *et al.*, 1994). It was reported some inhibitors found from the locust *Locusta migratoria* (Brehelin, *et al.*, 1991) and *M. sexta* (Saul, *et al.*, 1986) could control pro-PO activation. Pro-PO is always present from larva to pupa, whereas no PO activity is detected (Fig. 1). This result indicates that chymotrypsin inhibitor is likely to control pro-PO activation.

We have recently investigated the developmental changes of mRNA titer in the silkworm, *B. mori* (Yamamoto, *et al.*, 2000). The strong signal of mRNA was detected on day 3 of the fifth instar and one day before pupation in males, whereas females exhibited those on day 4 of the fifth instar and on the day of pupation. Similar results were obtained

Fig. 1. Developmental changes in pro-PO content evaluated with PO activity and Pro-PO mRNA

PO activity of crude extract was measured after activation with DBMA as described in Materials and Methods. Closed circles represent the assay conducted in the presence of DBMA. Open circles represent the assay conducted in the absence of DBMA. Closed squares represent the level of mRNA. S and P represent spinning and pupation, respectively. Bars indicate the standard deviation ($n=3$).

when the pro-PO concentrations were measured by enzyme-linked immunosorbent assay.

It is interesting that the fluctuation of pro-PO mRNA of the silkworm was different from that of the fall webworm, *H. cunea* (Park, *et al.*, 1997). They demonstrated that the highest level of pro-PO mRNA was detected in the mid-instar larvae and there was no signal in pupae, adult and egg. There is also the difference between the fluctuation of the silkworm PO and that of housefly PO (Daquinag, *et al.*, 1995). In the housefly, *M. domestica*, by a Western blot detection method using polyclonal antibodies raised against housefly PO, no significant change in PO protein content was observed in the homogenates of the pharate adult compared with that of the homogenates of other stages of aged pupae. They suggested that phenoloxidase inhibitor functions as a regulator of the necessary level of PO.

As shown in Fig. 1, pro-PO mRNA expression was preceded a day or two before, when comparing the change of phenoloxidase mRNA with its product. This finding suggests that the expression of pro-PO could be controlled at transcriptional step. Insect development is known to be regulated by ecdysteroids and juvenile hormones (JH). Possible roles of these hormones in the control of pro-PO expression during development will be studied in detail. The effects of JH on the synthesis of PO were reported. The granular PO that is responsible for cuticular melanization in *M. sexta* larva was controlled by JH (Hiruma, K. and Riddiford, L. M., 1984, 1985 and 1988). Recently, the transcription of PO in *A. gambiae* was regulated by 20-hydroxyecdysone (Martin, *et al.*, 1999). It is also possible for PO synthesis in silkworm to be controlled by 20-hydroxyecdysone at transcription level. Taken together, there are three possibilities of the regulation mechanism of PO activity, that is, transcriptional control, activation control and the inhibition of active PO. In future research, it is important for us to look for PO inhibitor and we will endeavor to understand how this hormone regulates the expression of PO.

REFERENCES

- Asada, N., Fujimoto, K., Tanaka, M., and Ohnishi, E. 1993 Genetic polymorphism of phenoloxidase A1 in *Drosophila melanogaster*. *Jpn. J. Genet.*, **68**: 217-219
- Ashida, M., Ochiai, M., and Niki, T. 1988 Immunolocalization of prophenoloxidase among hemocytes of the silkworm, *Bombyx mori*. *Tissue Cell*, **20**: 599-610
- Ashida, M., and Yamazaki, Y. I. 1990 In "Molting and Metamorphosis", ed. by Ohnishi, E. and Ishizaki, H., Japan Scientific Societies Press, Tokyo, pp. 239-265
- Aso, Y., Yamashita, T., Meno, K., and Murakami, M. 1994 Inhibition of the prophenoloxidase-activating enzyme from *Bombyx mori* by endogenous chymotrypsin inhibitors. *Biochem. Mol. Biol. Int.*, **33**: 751-758
- Brehelin, M., Boigegrain, R. A., Drif, L., and Coletti-Previero, M. A. 1991 Purification of a protease inhibitor which controls prophenoloxidase activation in hemolymph of *Locusta migratoria* (insecta). *Biochem. Biophys. Res. Commun.*, **179**, 841-846
- Cho, W. L., Liu, H. S., Lee, C. H., Kuo, T. Y., Chang, T. Y., Liu, C. T., and Chen, C. C. 1998 Molecular cloning, characterization and tissue expression of prophenoloxidase cDNA from the mosquito *Armigeres subalbatus* inoculated with *Dirofilaria immitis microfilariae*. *Insect Mol. Biol.*, **7**: 31-40
- Daquinag, A. C., Nakamura, S., Takao, T., Shimonishi, Y. and Tsukamoto, T. 1995 Primary structure of a potent endogenous dopa-containing inhibitor of phenol oxidase from *Musca domestica*. *Proc. Natl. Acad. Sci. USA*, **92**: 2964-2968
- Durrant, H. J., Ratcliffe, N. A., Hipkin, C. R., and Söderhäll, K. 1993 Purification of the pro-phenol

- oxidase enzyme from hemocytes of the cockroach *Blaberus discoidalis*. *Biochem. J.*, **289**: 87–91
- Fujimoto, K., Masuda, K., Asada, N., and Ohnishi, E. 1993 Purification and characterization of prophenoloxidases from pupae of *Drosophila melanogaster*. *J. Biochem.*, **113**: 285–291
- Fujimoto, K., Okino, N., Kawabata, S., Iwanaga, S., and Ohnishi, E. 1995 Nucleotide sequence of the cDNA encoding the proenzyme of phenol oxidase A1 of *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA*, **92**: 7769–7773
- Funatsu, M. and Inaba, T. 1962 Studies on tyrosinase in housefly. Part I. Protyrosinase in the pupae of housefly and its activation, *Agric. Biol. Chem.*, **26**: 535–540
- Hall, M., Scott, T., Sugumaran, M., Söderhäll, K., and Law, J. 1995 Proenzyme of *Manduca sexta* phenol oxidase: purification, activation, substrate specificity of the active enzyme, the molecular cloning. *Proc. Natl. Acad. Sci. USA*, **92**: 7764–7768
- Hiruma, K., and Riddiford, L. M. 1988 Granular phenoloxidase involved in cuticular melanization in the tobacco hornworm: regulation of its synthesis in the epidermis by juvenile hormone. *Dev. Biol.*, **130**: 87–97
- Inaba, T. and Funatsu, M. 1964 Studies on tyrosinase in the housefly. Part III. Activation of protyrosinase by natural activator. *Agric. Biol. Chem.*, **28**: 206–215
- Jiang, H., Wang, Y., Korochkina, S. E., Benes, H., and Kanost, M. R. 1997 Molecular cloning of cDNAs for two pro-phenol oxidase subunits from the malaria vector, *Anopheles gambiae*. *Insect Biochem. Mol. Biol.*, **27**: 693–699
- Jiang, H., Wang, Y., and Kanost, M. R. 1998 Pro-phenol oxidase activating proteinase from an insect, *Manduca sexta*: A bacteria-inducible protein similar to *Drosophila easter*. *Proc. Natl. Acad. Sci. USA*, **95**: 12220–12225
- Kawabata, T., Yasuhara, Y., Ochiai, M., Matsuura, S., and Ashida, M. 1995 Molecular cloning of insect pro-phenol oxidase: A copper containing protein homologous to arthropod hemocyanin. *Proc. Natl. Acad. Sci. USA*, **92**: 7774–7778
- Kwon, T. H., Lee, J. H., Lee, J. S., Kawabata, S., Iwanaga, S., and Lee, B. L. 1997 Purification and characterization of prophenoloxidase from the hemolymph of coleopteran insect, *Holotrichia dimophalia* larvae. *Mol. Cells*, **28**: 90–97
- Kopacek, P., Weise, C., and Gotz, P. 1995 The prophenoloxidase from the wax moth *Galleria mellonella*: purification and characterization of the proenzyme. *Insect Biochem. Mol. Biol.*, **25**: 1081–1091
- Lee, S. Y., Kwon, T. H., Hyun, J. H., Choi, J. S., Kawabata, S., Iwanaga, S., and Lee, B. L. 1998 *In vitro* activation of pro-phenol-oxidase by two kinds of pro-phenol-oxidase-activating factors isolated from hemolymph of coleopteran, *Holotrichia dimophalia* larvae. *Eur. J. Biochem.*, **15**: 50–57
- Lee, H. S., Cho, M. Y., Lee, K. M., Kwon, T. H., Homma, K., Natori, S., and Lee, B. L. 1999 The pro-phenoloxidase of coleopteran insect, *Tenebrio molitor*, larvae was activated during cell clump/cell adhesion of insect cellular defense reactions. *FEBS Lett.*, **444**: 255–259
- Martin, A. D., Manetti, A. G. O., Han, S. -J., Lee, W. -J., Mathiopoulos, K. D., Muller, H. -M., Kafatos, F. C., Raikhel, A., and Brey, P. Y. 1999 Genomic structure and ecdysone regulation of the prophenoloxidase 1 gene in the malaria vector *Anopheles gambiae*. **96** (26), 14796–14800
- Park, D., Shin, S. W., Kim, M. G., Park, S. S., Lee, W., Brey, P. T., and Park, H. 1997 Isolation and characterization of the cDNA encoding the prophenoloxidase of fall webworm, *Hyphantria cunea*. *Insect Biochem. Mol. Biol.*, **27**: 983–992
- Satoh, D., Horii, A., Ochiai, M., and Ashida, M. 1999 Prophenoloxidase-activating enzyme of the silkworm, *Bombyx mori*. Purification, characterization, and cDNA cloning. *J. Biol. Chem.*, **274**: 7441–7453
- Saul, S. J. and Sugumaran, M. 1986 Protease inhibitor controls prophenoloxidase activation in *Manduca sexta*. *FEBS Lett.*, **208**: 113–116
- Söderhäll, K. 1982 Prophenoloxidase activating system and melanization– a recognition mechanism of arthropods? A review. *Dev. Comp. Immunol.*, **6**: 601–611
- Sugumaran, M., Giglio, L., Kundzicz, H., Saul, S., and Semensi, V. 1992 Studies on the enzymes involved in puparial cuticle sclerotization in *Drosophila melanogaster*. *Arch. Insect Biochem. Physiol.*, **19**: 271–283
- Sugumaran, M. and Nellaiappan, K. 2000 Characterization of a new phenoloxidase inhibitor from the cuticle of *Manduca sexta*, *Biochem. Biophys. Res. Commun.*, **268**: 379–383

- Yamamoto, K., Sugioka, M., Fujii, H., Aso, Y., and Ishiguro, M. 1999 Isolation and characterization of prophenoloxidase isoforms from the silkworm, *Bombyx mori* (a80 strain). *J. Seric. Sci. Jpn.*, **68**: 65–72
- Yamamoto, K., Yakiyama, M., Fujii, H., Kusakabe, T., Koga, K., Aso, Y., and Ishiguro, M. 2000 Expression of prophenoloxidase mRNA during silkworm hemocyte development. *Biosci. Biotechnol. Biochem.*, **64**: (6), 1197–2000
- Yasuhara, Y., Koizumi, Y., Katagiri, C., and Ashida, M. 1995 Reexamination of properties of prophenoloxidase isolated from larval hemolymph of the silkworm *Bombyx mori*. *Arch. Biochem. Biophys.*, **320**: 14–23
- Yu, X. -Q., Gan, H., and Kanost, M. R. 1999 Immunectin, an inducible C-type lectin from an insect, *Manduca sexta*, stimulates activation of plasma prophenol oxidase. *Insect Biochem. Mol. Biol.*, **29**: 585–597