Respiration and Quality of Fresh-cut Cabbages in Modified Atmosphere Packaging

Hu, Wenzhong

Laboratory of Postharvest Science, Division of Bioproduction System Science, Department of
Bioproduction Environmental Science, Faculty of Agriculture, Kyushu University

Uchino, Toshitaka

Laboratory of Postharvest Science, Division of Bioproduction System Science, Department of
Bioproduction Environmental Science, Faculty of Agriculture, Kyushu University

Nakamura, Takeshi

Laboratory of Postharvest Science, Division of Bioproduction System Science, Department of
Bioproduction Environmental Science, Graduate School of Bioresource and Bioenvironmental
Sciences, Kyushu University

Furuse, Naomi

Laboratory of Postharvest Science, Division of Bioproduction System Science, Department of
Bioproduction Environmental Science, Graduate School of Bioresource and Bioenvironmental
Sciences, Kyushu University

他

<https://doi.org/10.5109/24468>

出版情報：九州大学大学院農学研究院紀要. 47 (1), pp.149-158, 2002-10-30. Kyushu University
バージョン：
権利関係：

Respiration and Quality of Fresh-cut Cabbages in Modified Atmosphere Packaging

Wenzhong HU^{*}, Toshitaka UCHINO, Takeshi NAKAMURA^{},
Naomi FURUSE^{**}, Daisuke HAMANAKA^{**},
Hussain SOROUR^{***}, Yoshiaki HORI
and Shun-ichiro TANAKA**

Laboratory of Postharvest Science, Division of Bioproduction System Science,
Department of Bioproduction Environmental Science, Faculty of Agriculture,
Kyushu University, Fukuoka 812-8581, Japan

(Received June 28, 2002 and accepted July 12, 2002)

The respiration rates of fresh-cut cabbage were measured by pouch method with 25 and 30 μm OPP films in thickness at 5 and 20 °C. The appearance, flavor and ascorbic acid content were also determined during storage. The results indicated that respiration rates of fresh-cut cabbages were suppressed under the condition of decreasing O_2 and increasing CO_2 concentrations at 5 and 20 °C. It was not found the effects of vertical cutting (VC), parallel cutting (PC) and thickness of film on the respiration at 5 °C. But the respiration rate were much higher in VC treatments than that in PC treatments, especially for the CO_2 production rate at 20 °C regardless film thickness. The results also showed that a shift from aerobic to anaerobic respiration was occurred under 1.2–1.5% O_2 level. Total ascorbic acid (TAA), ascorbic acid (AA) and dehydroascorbic acid (DHAA) were increased by AA biosynthesis initially and then decreased by higher rate of AA oxidation. Moreover, it was shown the results of sensory evaluation that there were no differences in appearance and flavor for cutting mode and thickness of the film at 5 °C. The better appearance and flavor appeared in VC treatments than PC treatments for cutting mode and thickness of film at 20 °C. It suggests that better quality could be obtained from the combination of VC treatments and low temperature of 5 °C regardless of thickness of film.

INTRODUCTION

Fresh-cut or minimally processed fruits and vegetables are a rapidly growing segment of retail and food service horticultural fresh industry (Gorny, *et al.*, 1998). Although fresh-cut products have met the consumers' desire for convenience and freshness, the shelf life is still important challenges for fresh-cut products. There are two basic problems confront the extension of shelf life of fresh-cut fruits and vegetables products. First, processing procedures of cutting cause intermixing of polyphenol oxidase with phenolics which undergo enzymatic browning to product an undesirable brown color (Senesi, *et al.*, 2000). Second, wounding tissue induces a high respiration rate, which triggers faster tex-

^{**} Laboratory of Postharvest Science, Division of Bioproduction System Science, Department of Bioproduction Environmental Science, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

^{***} Postdoctoral Fellow of Japan Society for the Promotion of Science, Laboratory of Postharvest Science, Division of Bioproduction System Science. Department of Bioproduction Environmental Science, Kyushu University

^{*} Corresponding author (E-mail:huxu@agr.kyushu-u.ac.jp)

ture deterioration and decrease in nutrients as like ascorbic acid compared to intact tissues (Rosen and Kader, 1989). These lead to a reduced quality and shorter shelf life than that can normally be expected from the fruits and vegetables. To overcome these problems and to have suitable shelf life for fresh-cut product, optimum lower temperature and relative humidity needs to be maintained during whole chain from harvest processing to consumption. The modified atmosphere packaging (MAP) is considered to be the most effective method for extending the shelf life of fresh and minimally processed product by depressing the respiration rate of product and preventing the moisture loss and microbial growth. However, O₂ concentration under lower oxygen limit or CO₂ concentration over the upper limit of aerobic condition that leads to fermentative metabolism can cause off-flavor and flavor loss (Watada, *et al.*, 1996; Kato-Noguchi, *et al.*, 1997).

A major metabolic process occurring in fresh-cut product is respiration. The respiration rate of product is an excellent indicator of metabolic activity of tissue; that is, it is a useful guide to the potential storage life of the fresh-cuts. But little information is available on respiration rate of fresh-cut cabbage in MAP during the storage and market. The objectives of this study were to determine the respiration rate of fresh-cut cabbage in MAP with two kinds of films in thickness. The evaluation of quality such as ascorbic acid content was also investigated.

MATERIALS AND METHODS

Plant material

Cabbages (*Brassica oleracea* L.) were obtained from the local supermarket in Fukuoka city, Japan. Samples were transported to the laboratory without refrigeration and immediately processed. Cabbages were selected for uniform size and appearance. Outer and damaged leaves were removed.

Cabbage slice preparation

The cabbages were rinsed with 0.02% sodium hypochlorite solution to reduce the surface microbial load before cutting. Knife and cutting board were washed and treated with 75% ethanol prior to use. The cabbage was chopped by hand with a sharp stainless steel knife. The cutting modes were divided into two forms: vertical cutting (VC) and parallel cutting (PC) to vein of cabbage with 2 mm wide by 5–8 cm in length. The slices were washed for 6 min with tap water and then washed with 5 °C distilled water for 4 min. The slices were centrifuged for 3 min at 1,720 (rpm) to remove the water on the cabbage surface by washing machine. And then slices were selected at random and samples of 100 ± 1 g were placed in the pouches (10 × 20 cm) and heat sealed. A silicone cap was applied on each package to permit analysis of package headspace with a syringe. The void volume in the package was determined by subtracting the volume of fresh-cut cabbage from that of the entire package. The volume of fresh-cut cabbage and the entire package were measured based on water displacement. All processing operations were conducted under 5–8 °C. The film used for cabbage slices packaging was oriented polypropylene (OPP) film (Sumitomo Bakelite Co., Ltd, Tokyo Japan), with two kinds of 25 and 30 µm film in thickness. The film permeability was determined by the Akimoto's method (Akimoto, *et al.*, 1997). The pouches were stored at 5 and 20 °C, respectively.

Gas analysis

Gas sample was withdrawn by gas-tight 1.0 ml syringe to measure gas composition from the inside the package. The gas concentrations were measured by injecting 1.0 ml gas sample into gas chromatography (GL Sciences GC-390, Tokyo Japan) equipped with thermal conductivity detector (TCD) and D2000 integrator (Hitachi, Ltd. Tokyo Japan). Helium was used as carrier gas and the flow rate was 30 ml·min⁻¹. The injector and column temperatures were 80 and 50 °C, respectively. The column was WG100 with molecular sieve 5 A and Porapak Q 80/100 mesh.

Mathematical model for respiration

To determine the respiration rate of fresh-cut cabbage under MAP, the basic material balance given in equations can be used (Akimoto *et al.*, 1997). The changes in the free volume of gas within package at the short time were shown in following equations.

$$\frac{dV_s}{dt} = \frac{dV_c}{dt} + \frac{dV_o}{dt} + \frac{dV_N}{dt} \quad (1)$$

$$\frac{dV_c}{dt} = \frac{A}{L} K_c (P_{Ca} - P_c) + R_c W \quad (2)$$

$$\frac{dV_o}{dt} = \frac{A}{L} K_o (P_{Oa} - P_o) + R_o W \quad (3)$$

$$\frac{dV_N}{dt} = \frac{A}{L} K_N (P_{Na} - P_N) \quad (4)$$

where V is the volume of gas in pouch (ml), A is surface area of pouch (m²), L is thickness of film (m), K_c , K_o and K_N are the permeability of film for CO₂, O₂ and N₂ (ml·m·m⁻²·atm⁻¹·h⁻¹), R is respiration rate (ml·kg⁻¹·h⁻¹), W is weight of product (kg), P_{Ca} , P_{Oa} and P_{Na} are partial pressure of CO₂, O₂ and N₂ outside pouch (atm), P_c , P_o and P_N are partial pressure of CO₂, O₂, N₂ inside pouch (atm). These equations show that changes rate of gas volume of package was determined by permeability of film and respiration of product. According to gas partial pressure between internal and external package, this method can be used to determine CO₂ production and O₂ consumption rates of product based on CO₂, O₂ and N₂ levels within package. It has been demonstrated that the method is very effective to calculate the respiration rate under steady and unsteady atmospheric conditions.

Sensory evaluations

Visual appearance and flavor were evaluated by five trained people at cutting and during storage period for every day. Evaluation was carried out in two items of appearance and flavor. They were both scored on 4-point scale: excellent (−); very good (+); fair (++); poor (+++).

Ascorbic acid

Ascorbic acid content was measured by high performance liquid chromatography (HPLC) equipped with Shim-pack SCR-101N column (7.9 mmφ × 30 cm) (Shimadzu

Corp., Tokyo Japan). The column is packed with a cation exchanger resin which is sulfonated polystyrene-divinylbenzene copolymer. The mobile phase was mixed solution of 10 mM oxalic acid dihydrate, 15 mM NaOH and 1 mM EDTA at a flow rate of 1.0 ml min^{-1} . Temperature is 40°C and detector is RI. Cabbage sample was extracted from excised slices of 5 g cabbage with 5 times the volume of 5% metaphosphoric acid in a mortar with pestle. The homogenate was filtered through filter paper and centrifuged at 3,000 (rpm) for 15 min. The supernatant was taken as sample to measure the ascorbic acid in cabbage.

RESULTS AND DISCUSSION

Respiration rate

Changes in respiration rate, respiratory quotient (RQ) and gas concentration of fresh-cut cabbage in VC treatments were shown in Fig. 1. The changes in respiration rate was approximately same at the conditions of increasing CO_2 and decreasing O_2 concentrations for 5 days storage at 5°C (Fig. 1A and 1B). The respiration rates were slightly decreased in MAP during storage of 5 days. There was no significant difference between the two kinds of 25 and $30 \mu\text{m}$ thickness films in depressing respiration during storage period. The ratio of CO_2 produced to O_2 consumed, known as RQ, is normally assumed to be equal to 1.0 if the metabolic substrates are carbohydrates. In the present study, RQ was fluctuated around 1.0, indicating that anaerobic respiration did not occur for 3-day storage. The O_2 concentration continued to decrease and a certain point of O_2 concentration was reached, O_2 consumption rates were remarkably decreased and CO_2 production rates were increased conversely. It was indicated a shift to anaerobic respiration confirmed by higher RQ values at 1.2–1.5% O_2 levels. This effect of O_2 concentration was more marked at higher temperature (Fig. 1C and 1D). At 20°C , respiration rates were suppressed markedly with quickly increasing CO_2 and decreasing O_2 levels at first 1 day, and then shift from aerobic to anaerobic respiration occurred with higher RQ at 1.2–1.5% O_2 levels. It was also demonstrated by off-flavor of fermentation when the package was opened for sensory evaluation. For changes in gas concentrations, O_2 level was decreased and CO_2 level was increased quickly at initial period and then CO_2 level was increased slowly and O_2 level was reached about 1.2–1.5%, indicating that O_2 and CO_2 levels were regulated by product respiration and permeability of film during storage (Fig. 1A–1D). The results suggest that MAP could be used as a cheap and easy method to extend the storage life of fresh-cut cabbage.

Changes in respiration rate, RQ and gas concentration of fresh-cut cabbage in PC treatments were shown in Fig. 2. The trends of changes in respiration rate were similar to the results of VC treatments (Fig. 1). It was also indicated that the thickness of film has less effect on the changes in gas concentration and respiration of fresh-cut cabbages. The storage life of horticultural products has well been related to the respiration rate of products, the suppression of respiratory metabolism will extend the shelf life of stored fresh product (Lopez-Briones, *et al.*, 1992; Kader, 1986). But O_2 and CO_2 levels beyond its tolerance limits can induce anaerobic respiration and CO_2 damage to result in browning and development of off-flavors. It was suggested that better keeping quality for fresh-cut cabbages could be obtained from low O_2 storage atmosphere above 1.2–1.5% lower limit

Fig. 1. Changes in gas concentration, respiration rate and RQ of fresh-cut cabbage in VC treatments at 5 and 20°C. (●) represents CO₂ production rate, (○) O₂ consumption rate, (▲) respiratory quotient (RQ). Solid and dashed lines denote the CO₂ and O₂ concentrations, respectively.

oxygen. For the effect of cutting mode, there were no significant differences in changes in gas concentration and respiration rate between PC and VC treatments at 5°C during storage (Fig. 1A–1B and Fig. 2A–2B). At 20°C, the respiration rate were much higher in VC treatments than that in PC treatments, especially for the CO₂ production rate (Fig. 1C–1D and Fig. 2C–2D).

MAP can be beneficial in maintaining quality of the fresh-cut product (Gorny, 1997). The suitable gas mixture for MAP has been based on that recommended for the whole commodity (Saltveit, 1997). Fresh-cut products probably can tolerate more extreme levels of O₂ and CO₂, because they do not have as much cuticle or skin to restrict gas diffusion, and the distance of gas diffusion from center to outside of fresh-cut product is much less than that for the whole commodity (King, *et al.*, 1989). It was also reported that an additional benefit of MAP may be attained by actively flushing the package with the desired gas rather than allowing the MAP to develop naturally since the marketing period of fresh-cut product is relatively short (Bai, *et al.*, 2001). In the present study, fresh-cut cabbage has much higher respiratory activity to consume O₂ and produce CO₂ quickly.

Fig. 2. Changes in gas concentration, respiration rate and RQ of fresh-cut cabbage in PC treatments at 5 and 20°C. ($\bullet$) represents CO_2 production rate, ($\circ$) O_2 consumption rate, ($\blacktriangle$) respiratory quotient (RQ). Solid and dashed lines denote the CO_2 and O_2 concentrations, respectively.

Anaerobic respirations were occurred after 3 days of storage at 5°C and after 1 day of storage at 20°C. It induced anaerobic respiration and higher CO_2 damage to result in browning and development of off-flavors. Respiration rates of fresh-cuts generally were higher than the intact product. Although benefits may occur with modified atmosphere, the extinction point of the fresh-cut products must be recognized to avoid anaerobic respiration (Ko, *et al.*, 1996).

Fresh-cuts generally are much more perishable than intact products because they have been subjected to severe physical stress, such as peeling, cutting, slicing, shredding, trimming and removal of protective cells. Consequently, fresh-cuts probably should be held at a lower temperature than that recommended for intact commodities. Temperature of 0°C is in most case preferable. However, this is in most cases economically not achievable. Temperature between 5 and 10°C is more commonly found in practice. The temperature effects on the tissue metabolism and biochemical reactions significantly, it is also of major important to maintain low temperature to prevent microbial growth. It suggests that it is important for handling and storing fresh-cut cabbage at lower temperature or near 0°C, if the product is not sensitive to chilling injury.

Ascorbic acid

Fig. 3 shows changes in ascorbic acid contents in fresh-cut cabbage at 5°C during storage. TAA, AA and DHAA were increased in PC treatments for 3 days, and then decreased. In the VC treatments, the results were similar to PC treatments, they increased for 4 days and then decreased. But the increase of TAA from $32.4 \text{ mg} \cdot 100\text{g}^{-1}$ fresh weight (FW) to $43.3\text{--}45.8 \text{ mg} \cdot 100\text{g}^{-1}$ FW was higher in PC treatments than that from $30.6 \text{ mg} \cdot 100\text{g}^{-1}$ FW to $34.7\text{--}36.3 \text{ mg} \cdot 100\text{g}^{-1}$ FW in VC treatments.

Fig. 4 shows changes in ascorbic acid contents in fresh-cut cabbage at 20°C during storage. TAA, AA and DHAA were increased and then decreased. The increase of TAA was less than that at 5°C (Fig. 3). But increase of DHAA was more significant than the increase at 5°C. AA level is the result of the difference between AA production (biosynthesis or regeneration from DHAA) and AA oxidation. If the rate of AA oxidation is higher than the rate of AA production, then AA content decreases (Zerbini, *et al.*, 2002). In the present study, increase of DHAA was significant at 20°C, indicating that the oxidation rate was higher than that at 5°C. It demonstrated that activity of ascorbate oxidase was higher at 20°C than that at 5°C. The significant decrease of AA after storage 3 days, it was due to oxidation of AA by ascorbate oxidase following tissue damage or by the reaction of AA with phenolic substances during storage. It could be also assumed that the role of AA as a reducing power in respect to *de nova* synthesis of polyphenols is pos-

Fig. 3. Changes in ascorbic acid contents in VC and PC treatments at 5°C

Fig. 4. Changes in ascorbic acid contents in VC and PC treatments at 20°C

sible. Ascorbic acid plays an important part in the defence mechanism against free radicals which induce peroxidation (Barth, *et al.*, 1993). AA can be easily oxidised to dehydroascorbic acid, the latter in turn can either be reduced.

Sensory evaluation

Sensory evaluation of fresh-cut cabbage at 5°C was shown in Table 1. Visual appearances were not changed much during storage of 4 days in all treatments. Off-flavor was slightly occurred after 3 days by anaerobic respiration. At 20°C (Table 2), appearances were not changed in VC treatments, but browning was occurred after 3 days. In PC treatments, the browning was occurred much significantly. For the flavor, off-flavor was occurred after 1 day and appeared more serious in PC treatments after storage of 3 days. It seems that the changes in appearance and flavor depend on physiological changes associated with cutting direction. It was considered that the callus in wounded place of cabbage was formed quicker in VC treatments than PC treatments. Because the respiration rate were much higher in VC treatments than that in PC treatments, especially for the CO₂ production rate at 20°C. Moreover, the cutting wound area was large and tissue sap seeped much from vascular bundle of cabbage in PC treatments. The seeped tissue sap in PC treatments can induce obvious browning and off-flavor. It seems that thickness

Table 1. Sensory evaluation of fresh-cut cabbage at 5 °C

Item	Cutting treatment	Storage period (day)				
		0	1	2	3	4
Appearance	VC-25	—	—	—	—	—
	VC-30	—	—	—	—	—
	PC-25	—	—	—	—	—
	PC-30	—	—	—	—	—
Flavor	VC-25	—	—	—	+	+
	VC-30	—	—	—	+	+
	PC-25	—	—	—	+	+
	PC-30	—	—	+	+	+

Table 2. Sensory evaluation of fresh-cut cabbage at 20 °C

Item	Cutting treatment	Storage period (day)				
		0	1	2	3	4
Appearance	VC-25	—	—	—	—	—
	VC-30	—	—	—	—	—
	PC-25	—	—	—	++	+++
	PC-30	—	—	—	++	+++
Flavor	VC-25	—	+	+	+	+
	VC-30	—	+	++	++	++
	PC-25	—	+	+	++	+++
	PC-30	—	+	+	++	+++

of film has less effect on the respiration and quality of fresh-cut cabbages. It was suggested that the better quality can be obtained from VC treatments at 5 °C.

CONCLUSIONS

The respiration rates, quality attributes of fresh-cut cabbage were determined in MAP with 25 and 30 μm OPP films in thickness at 5 and 20 °C during storage. The respiration rates of fresh-cut cabbages were suppressed under the condition of decreasing O_2 and increasing CO_2 concentrations. It was not found the effects of cutting mode (VC and PC treatments) and thickness of film on the respiration at 5 °C. But the respiration rate were much higher in VC treatments than that in PC treatments, especially for the CO_2 production rate at 20 °C. The results also showed that a shift from aerobic to anaerobic respiration was occurred under 1.2–1.5% O_2 levels for fresh-cut cabbage. TAA, AA and DHAA were increased by AA biosynthesis initially and then decreased by higher rate of AA oxidation. For sensory evaluation, there were no differences in appearance and flavor for cutting mode and thickness of film at 5 °C. The better appearance and flavor appeared in VC than PC treatments for cutting mode and thickness of film at 20 °C. It suggests that better quality could be obtained from the combination of VC treatments and low

temperature of 5 °C regardless of thickness of OPP film during storing and marketing

ACKNOWLEDGEMENTS

The authors would like to thank Sumitomo Bakelite Co., Ltd for providing the OPP film.

REFERENCES

- Akimoto, K. and S. Maezawa 1997 A new method for estimating respiration rate of fruits and vegetables in modified atmosphere packaging. *J. of the J. Soc. Agri. Machinery*, **59**(1): 109–116
- Bai, J.H., R.A. Saftner, A.E. Watada and Y.S. Lee 2001 Modified atmosphere maintains quality of fresh-cut cantaloupe (*Cucumis melo* L.). *J. Food Sci.*, **66**: 1207–1211
- Barth, M. M., E. L. Kerbel, S. Broussard and S. J. Schmidz 1991 Modified atmosphere packaging protects market quality in broccoli spears under ambient temperature storage. *J. Food Sci.*, **58**: 1070–1072
- Gorny, J. R., M. I. Gil and A. A. Kader 1998 Postharvest physiology and quality maintenance of fresh-cut pears. *Acta Hort.*, **464**: 231–236
- Gorny, J. R. 1997 Summary of CA and MA requirements and recommendations for fresh-cut (minimally processed) fruits and vegetables. *Proceedings of Seventh International Controlled Atmosphere Conference*, **Vol. 5. Postharvest Outreach Program**, University of California, Davis, CA: 30–66
- Guerzoni, M. E., A. Gianotti and M. R. Corbo and M. Sinigalia 1996 Shelf-life modeling for fresh-cut vegetables. *Postharvest Biol. Technol.*, **9**: 195–207
- Kader, A. A. 1986 Biochemical and physiological basis for effects of controlled and modified atmospheres on fruits and vegetables. *Food Technol.*, **5**: 99–104
- Kato-Guchi, H. and A. E. Watada 1997 Effects of low-oxygen atmosphere on ethanolic fermentation in fresh-cut carrots. *J. Am. Soc. Hort. Sci.*, **122**: 107–111
- King, A. D. and H. R. Bolin 1989 Physiological and microbiological storage stability of minimally processed fruits and vegetables. *Food Technol.*, **2**: 132–135
- Ko, N. P., A. E. Watada, D. V. Schlimme and J. C. Bouwkamp 1996 Storage of spinach under low oxygen atmosphere above the extinction point. *J. Food Sci.*, **61**: 398–400+406
- Lopez-Briones, G., P. Varoquaux, Y. Chambroy, J. Bouquant, G. Bureau and B. Pascat 1992 Storage of common mushroom under controlled atmospheres. *Inter J Food Sci Technol.*, **27**: 493–505
- Rosen, J. C. and A. A. Kader 1989 Postharvest physiology and quality maintenance of slice pear and strawberry fruits. *J. Food Sci.*, **54**: 656–659
- Senesi, C., M. Prinziavalli, K. Sala and M. Gennari 2000 Physicochemical and microbiological changes in fresh-cut green bell peppers as affected by packaging and storage. *Ital. J. Food Sci.*, **12**: 55–64
- Slitveit, T. J. 1997 Summary of CA and MA requirements and recommendations for harvested vegetables. *Proceedings of Seventh International Controlled Atmosphere Conference*, **Vol. 4. Postharvest Outreach Program**, University of California, Davis, CA: 98–117
- Watada, A. E., N. P. Ko and D. A. Minott 1996 Factors affecting of fresh-cut horticultural products. *Postharvest Bio. Technol.*, **9**: 115–125
- Zerbini, P. E., A. Rizzolo, A. Brambilla and P. Cambiaghi 2002 Loss ascorbic acid during storage of conference pears in relation to the appearance of brown heart. *J. Sci. Food Agric.*, **82**: 1007–1013