Anti-melanogenesis properties of quercetin- and its derivative-rich extract from *Allium cepa*

Arung, Enos Tangke

Department of Forest and Forest Products Sciences, Faculty of Agriculture, Kyushu University |
Department of Forest Product Technology, Faculty of Forestry, Mulawarman University

Furuta, Shoko

Department of Nutrition and Health Science, Faculty of Human Environmental Science, Fukuoka
Women's University

Ishikawa, Hiroya

Department of Nutrition and Health Science, Faculty of Human Environmental Science, Fukuoka
Women's University

Kusuma, Irawan Wijaya

Department of Forest Product Technology, Faculty of Forestry, Mulawarman University

他

<http://hdl.handle.net/2324/25500>

出版情報 : Food Chemistry. 124 (3), pp.1024-1028, 2011-02-01. Elsevier
バージョン :
権利関係 : (C) 2010 Elsevier Ltd.

Anti-melanogenesis properties of quercetin- and its derivative-rich extract from
Allium cepa

Enos Tangke Arung ^{1,2}, Shoko Furuta ³, Hiroya Ishikawa ³, Irawan Wijaya Kusuma ²,
Kuniyoshi Shimizu ^{1*}, Ryuichiro Kondo ¹

1.Department of Forest and Forest Products Sciences, Faculty of Agriculture, Kyushu
University, Fukuoka, 812-8581, Japan

2.Department of Forest Product Technology, Faculty of Forestry, Mulawarman
University, Samarinda, 75123, Indonesia

3.Department of Nutrition and Health Science, Faculty of Human Environmental
Science, Fukuoka Women's University, Fukuoka, 813-8529, Japan

* Corresponding author:

Kuniyoshi Shimizu

Address: 6-10-1 Hakozaki, Higashi-ku, Fukuoka, Japan, 812-8581

E-mail: shimizu@agr.kyushu-u.ac.jp

Phone/Fax: +81-92-642-3002

Abstract

In an effort to find a new whitening agent, we have found that the methanol extract of the dried skin of *Allium cepa* showed inhibition of melanin formation. Bioassay-guided fractionation led to the isolation of quercetin (**1**) and quercetin 4'-*O*- β -glucoside (**3**) from *A. cepa* as the inhibitors of melanin formation in B16 melanoma cells with an IC₅₀ of 26.5 and 130.6 μ M, respectively. In addition, we evaluated the effect of some quercetin derivatives, such as isoquercitrin (**2**); quercetin 3, 4'-*O*-diglucoside (**4**); rutin (**5**) and hyperin (**6**) on B16 melanoma cells. These quercetin derivatives did not show any inhibition of melanin formation. Furthermore, the ORAC values of compounds **1-6** were 7.64, 8.65, 4.82, 4.32, 8.17 and 9.34 μ mol trolox equivalents/ μ mol, respectively. Dried skin of red onion showed inhibitory activity of melanin formation in B16 melanoma cells as well as antioxidant properties.

Keywords : *Allium cepa*, red onion, quercetin derivatives, anti melanogenesis, antioxidant

1. Introduction

Melanin pigments are formed in specialized pigment-producing cells known as

melanocytes, which originate in the neural crest during embryogenesis and are distributed throughout the embryo during its development (Sánchez-Ferrer, Rodríguez-López, & García-Carmona, 1995).

Melanin biosynthesis occurs in a cascade of enzymatic and spontaneous reactions that convert tyrosine to melanin pigments. The initial and rate-limiting step in melanin synthesis is the hydroxylation of tyrosine to dihydroxyphenylalanine or DOPA (Yoon et al., 2007). DOPA-oxidation produces a highly reactive intermediate that is further oxidized to form melanin by a free radical-coupling pathway. If free radicals are inappropriately processed in melanin synthesis, hydrogen peroxide (H_2O_2) is generated, leading to the production of hydroxyl radicals ($HO\bullet$) and other reactive oxygen species (ROS) (Perluigi et al., 2003).

The onion is a versatile vegetable that is consumed fresh as well as in the form of processed products. The regular consumption of onions in food is associated with a reduced risk of neurodegenerative disorders, cancer, cataract, ulcer, osteoporosis, vascular disease and heart disease (Kaneko & Baba, 1999; Kawaii, Tomono, Katase, Ogawa, & Yano, 1999; Sanderson, Mclauchlin, & Williamson, 1999; Shutenko et al., 1999). Onion is one of the major sources of various biologically active phytochemicals e.g., phenolic acids, flavonoids, cepaenes, thiosulfinates and anthocyanins (Singh et al.,

2009). The major flavonoids found in the dry peel of the onion, which has usually been considered to be waste, contain large amounts of quercetin, quercetin glycoside and their oxidative product which are effective antioxidants against the lethal effect of oxidative stress (Gulsen, Makris, & Kefalas, 2007; Prakash, Upadhyay, Singh, & Singh, 2007).

Onion (*Allium* species) plays an important role in traditional medicine in Indonesia; it is used as a diuretic, febrifuge, and poultice to cure wounds and to remove scars from the skin, and it suppresses the blood sugar level and platelet aggregation, (de Padua, Bunyaphrathatsara, & Lemmens, 1999).

Based on our preliminary screening data (not shown), the methanol extract of the dried skin of the red onion (*Allium cepa*) from Indonesia showed a potent melanin biosynthesis inhibitory activity on B16 melanoma cells. These findings led us to focus on the active compounds in the dried skin of the red onion.

2. Materials and Methods

2.1. Chemicals

NaOH, DMSO, L-tyrosine, L-DOPA, and rutin (**5**) purchased from Wako (Osaka, Japan). Mushroom tyrosinase, Fluorescein sodium salt (FL), 2,2'-azobis (2-methylpropionamidine) dihydrochloride (AAPH), and

6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox) and 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyl tetrazolium bromide or MTT were obtained from Sigma (St. Louis, MO, USA). Quercetin-3, 4'-*O*-diglucoside (**4**) and hyperin (**6**) were from Tokiwa Phytochemical (Tokiwa, Japan). EMEM was from Nissui Chemical Co (Osaka, Japan). The ethylenediaminetetraacetic acid or EDTA was from Dojindo (Kumamoto, Japan). The 1, 1-diphenyl-2-picrylhydrazyl (DPPH) was from TCI (Tokyo, Japan) and isoquercitrin (**2**) was from Fluka (Steinheim, Germany). Other chemicals are of the highest grade commercially available.

2.2. Plant materials

Red onion (*A. cepa*) was purchased from traditional market in Jakarta, Indonesia, on September in 2008. Voucher specimen (ETA-CW-6) was deposited in Wood Chemistry Laboratory, Department of Forest Product Technology, Faculty of Forestry, Mulawarman University, Indonesia.

2.3. Preparation of plant extracts

Plant materials (dried skin or flesh part of *A. cepa*) were dried at room temperature and powdered. The dried materials (17.38 g) were extracted with methanol (650 mL) at

room temperature with shaker at 150 rpm during 48 h. The extract solutions were filtered and concentrated *in vacuum*, to obtain the crude methanol extracts. The crude extracts were 1.75 g.

2.4. Isolation of Quercetin (**1**) and Quercetin 4'-O- β -glucoside (**3**)

The dried skin of methanol extract of *A. cepa* (1.4 g) was applied to silica gel column chromatography (71 g of Wakogel C-200, 3.5 x 50 cm). The chromatography was eluted with *n*-hexane/EtOAc [10:0 (100 mL), 9:1 (50 mL), 7:3 (50 mL), 5:5 (200 mL), 3:7 (200 mL), 1:9 (100 mL)] and EtOAc/MeOH [9:1 (100 mL), 8:2 (100 mL), 7:3 (250 mL), 6:4 (50 mL), 5:5 (100 mL), 4:6 (50 mL), 3:7 (50 mL), 2:8 (50 mL), 1:9 (100 mL), 0:10 (100)] to give thirty three fractions (Fr 1 to Fr 33). The fractions Fr. 3 (66.9 mg) and Fr. 15 (84.7 mg) was analyzed by TLC (Thin Layer Chromatography) and HPLC (High Performance Liquid Chromatography). Based on NMR (Nuclear Magnetic Resonance) analysis, Fr. 3 was identified as quercetin (**1**) and Fr. 15 was quercetin-4'-O-glucoside (**3**) by comparison of previous NMR data (Alfonso & Kapetanidis, 1994; Tanabe, Ogawa, Tesaki & Watanabe, 1997), respectively (Figure 1). The NMR spectra of compounds were recorded at 400 MHz on JNM-AL400 FT NMR spectrometer (Jeol). All compounds were dissolved in DMSO-*d*₆, or methanol-*d*₄, and

chemical shifts were referred to deuterated solvents. The compounds were assigned for

^1H , ^{13}C , HMQC, and HMBC.

2.5. HPLC analysis

All the crude extracts of dried skin and flesh of onion were dissolved in HPLC grade methanol, filtered through sterile 0.22 μm millipore filter and subjected to qualitative analysis by using Waters 600 HPLC instrument. The instrument is equipped with a photodiode array (detector Waters 996), controller (Waters 600s), pump (Water 626), and auto sampler injector Model 231 (Gilson). A ODS Inertsil C18 (4.6 mm i.d. x 250 mm) was used as a column. Data were integrated by Empower Build 1154-J series software (Waters). Separation was achieved by flow rate of 1 mL/min with methanol (60%)/water (40%) containing 1% trifluoroacetic acid in isocratic program by monitoring the absorbance at 250 nm.

2.6. Oxygen radical absorbance capacity (ORAC) assay

Samples were directly dissolved in acetone/water/acetic acid (70:29.5:0.5, v/v/v), and diluted with 75 mM potassium phosphate buffer (pH 7.4) for analysis. Trolox, FL and AAPH solutions were prepared with 75 mM phosphate buffer (pH 7.4). The ORAC

assay was performed as described by Ou, Hampsch-Woodill, & Prior (2001) with some modification as follows: 300 μ L of the standard (Trolox) or sample solution is mixed with 1.8 mL of 48 nM FL solution, and then they were incubated independently at 37°C for 15 min. AAPH solution (900 μ L; 12.9 mM, final concentration) was added to the mixture and vortexed for 10s. Then, it immediately placed in a fluorescence spectrophotometer (Model FP-6500, JASCO Co., Ltd., Tokyo, Japan), and measured every 5s for 60 min at 37°C (Ex: 485 nm, Em: 520 nm). A blank (FL + AAPH) using phosphate buffer, standard solutions (6.25-50 μ M Trolox), and sample solutions were measured at same conditions. Three independent assays were performed for each sample. The area under the fluorescence decay curve (AUC) was calculated as

$$AUC = 1 + \sum \frac{f_i}{f_0}$$

where f_0 is the initial fluorescence reading at 0 min and f_i is the fluorescence reading at time i.

The ORAC values were calculated as Prior et al. (2003) by using a equation (Y) a + b(X) between Trolox concentration (Y)(μ M) and the net area under the FL decay curve (X). Linear regression was used in the range of 6.25-50 μ M Trolox. Data are expressed as micromoles of Trolox equivalents (TE) per micromole of sample (μ mol TE/ μ mol).

2.7. Cell culture

A mouse melanoma cell line, B16, was obtained from RIKEN Cell Bank. The cells were maintained in EMEM supplemented with 10% (v/v) fetal bovine serum (FBS) and 0.09 mg/mL theophylline. The cells were incubated at 37°C in a humidified atmosphere of 5% CO₂.

2.8. Inhibitory effect of melanin biosynthesis using cultured B16 melanoma cells

This assay was determined as described by Arung, Shimizu, & Kondo (2007). Briefly, confluent cultures of B16 melanoma cells were rinsed in phosphate-buffered saline (PBS) and removed from the plastic using 0.25 % trypsin/EDTA. The cells were placed in two plates of 24-well plastic culture plates (1 plate is for determining of melanin and other is for cell viability) at a density of 1×10^5 cells/well and incubated for 24 h in media prior to being treated with the samples. After 24 h, the media were replaced with 998 µL of fresh media and 2 µL of DMSO was added with or without (control) the test sample at various concentrations (n=3). Arbutin was used as a positive control. The cells were incubated for an additional 48 h, and then the medium was replaced with fresh medium containing each sample. After 24 h, the remaining adherent cells were assayed (see below).

2.9. Determination of melanin content in B16 melanoma cells

The melanin content of the cells after treatment was determined as follows. After removing the medium and washing the cells with PBS, the cell pellet was dissolved in 1.0 mL of 1N NaOH. The crude cell extracts were assayed using a micro plate reader (Bio-Tek, USA) at 405 nm to determine the melanin content. The results from the cells treated with the test samples were analyzed as a percentage of the results from the control culture.

2.10. Cell viability

Cell viability was determined by use of the micro culture tetrazolium technique (MTT). The MTT assay provides a quantitative measure of the number of viable cells by determining the amount of formazan crystals produced by metabolic activity in treated versus control cells. Culture was initiated in 24-well plates at 1×10^5 cells per well. After incubation, 50 μ L of MTT reagent [3-(4, 5-dimethyl-2-thiazolyl)-2, 5-diphenyl-2H-tetrazolium bromide in PBS (5 mg/mL)] was added to each well. The plates were incubated in a humidified atmosphere of 5% of CO₂ at 37°C for 4 h. After the medium was removed, 1.0 mL isopropyl alcohol (containing 0.04 N HCl) was added into the plate, and the absorbance was measured at 570 nm relative to 630 nm.

3. Results and discussion

3.1. The effect of the methanol extract of the dried skin and flesh of *A. cepa* on B16 melanoma.

The diameter of Indonesian red onion (*A. cepa*) is approximately 1.5 – 2.0 cm. The thickness of the dried skin is approximately 0.5 - 1.0 mm. Figures 2 and 3 show the effect of the methanol extract of the dried skin and flesh of *A. cepa* on B16 melanoma cells, respectively. The dried skin extract dose dependently inhibited melanin formation in B16 melanoma cells. The inhibition of 40-50% of melanin formation was evident at concentrations of 50 and 100 µg/mL without any cytotoxicity. Arbutin, which is used in skin-whitening cosmetics, was used as a positive control (Virador, Kobayashi, Matsunaga & Hearing, 1999). In contrast, the extract of the flesh of the onion did not lead to melanin inhibition even at concentrations up to 250 and 500 µg/mL. According to these data, the dried skin extract is more potent than the flesh extract. This result may be related to the amount of active compound in the dried skin, as shown in Figures 4 and 5. Figure 4 depicts HPLC chromatograms obtained by the injection of 20 µL of solution which contained methanol extract of the dried skin of *A. cepa* at the concentration of 1 mg/mL. Based on the methanol extract of the dried skin, the content of quercetin, isoquercitrin; quercetin 4'-*O*-glucoside; and quercetin 3,4'-*O*-diglucoside

were 13.8, 10.3, 6.4 and 8.3%, respectively. We have tried to analyze the flesh extract at a concentration of 1 mg/mL but each peak of the compounds was too small to be detected; therefore, we increased the concentration for HPLC analysis. Figure 5 shows the HPLC chromatogram obtained by the injection of 20 μ L of solution which contained methanol extract of the flesh of *A. cepa* at the concentration of 11 mg/mL. Based on the methanol extract of the flesh, the content of quercetin; isoquercitrin; quercetin 4'-*O*-glucoside, and quercetin 3,4'-*O*-diglucoside were 0.01, 0.59, 0.35 and 0.50%, respectively. As seen on the Y axis of both figures, the content of quercetin derivatives in the methanol extract of the dried skin of *A. cepa* was apparently much higher than that in the extract of the flesh. Also, it was shown that the extract of dried skin contained quercetin (**1**) and isoquercitrin (**2**) dominantly, but in the flesh extract, isoquercitrin (**2**) and quercetin 3,4'-*O*-diglucoside (**4**) were contained abundantly. It was reported that the content of quercetin, quercetin 4'-*O*-glucoside and quercetin 3,4'-*O*-diglucoside in different parts of the onion bulb (*Allium cepa* L.) found difference as well as the various cultivars (Beesk, Perner, Schwarz, George, Kroh, & Rohn, 2010). They found the outer layer of *A. cepa* content approximately 2.2, 2.7, 1.7% of quercetin, quercetin 4'-*O*-glucoside, and quercetin 3,4'-*O*-diglucoside in the dry weight of outer layer, respectively. In the inner parts, they reported that the content of quercetin, quercetin

4'-*O*-glucoside, and quercetin 3,4'-*O*-diglucoside was 0, 0.53 and 0.72% in the dry weight of inner parts, respectively. In our opinion, the difference of quercetin and its derivatives content should affect the difference in their melanin inhibitory activity as depicted in Figures 2 and 3.

3.2. Isolated compounds

The biologically-guided fractionation of methanol extracts of *A. cepa* on B16 melanoma cells led us to focus on the active fraction of the dried skin extract. As a result of fractionation, Fr. 3 was identified as one of the active compounds as well as Fr. 15. Thus, NMR assignment was performed to elucidate the structure of these fractions by comparison with previous data (Alfonso & Kapetanidis, 1994; Tanabe, Ogawa, Tesaki & Watanabe, 1997). It was revealed that Fr. 3 and Fr. 15 were **1** (69.9 mg, 4.9 % yield in the extract) and **3** (84.7 mg, 6.1 % yield in the extract), respectively.

3.3. Anti melanogenesis properties

The anti melanogenesis effects of isolated compounds **1** and **3** (Figure 1) from the dried skin of *A. cepa* were determined by using B16 melanoma cells. Table 1 depicted the IC₅₀ values for the melanin biosynthesis inhibition of quercetin and its derivatives. The activities of the tested compounds, in terms of IC₅₀ values, decreased in the following order: **1** (<50 µM), **3** (100–150 µM), **4**, **5**, **2** and **6** (>150 µM). Notably,

isolated compounds **1** and **3** from *A. cepa*, were found to be more-potent inhibitors of melanin formation in B16 melanoma cells than the positive control, arbutin (198 μ M). It should be noted that the opposite results were reported previously, namely that **1** enhanced melanogenesis in human melanoma cells and normal epidermal melanocyte (Nagata, Takekoshi, Takeyama, Homma, & Osamura, 2004). Kubo, Nitoda, & Nihei (2007) also reported that **1** enhanced the total melanin content in B16 melanoma cells. Recently, it was reported that **1** suppressed melanin formation in B16 melanoma cells with decreased intracellular tyrosinase activity and its protein expression (Fujii & Saito, 2009). The reason for the difference of the effect of **1** on melanin production in cells remains unclear.

Based on our analysis, the IC₅₀ of **1** to inhibit melanin formation was 26.5 μ M with 88% cell viability, while that of **3** was 130.6 μ M with 82% cell viability. The attachment of a glucoside moiety in quercetin tended to reduce the ability of quercetin to inhibit melanin formation in B16 melanoma cells. These results led us to evaluate other quercetin derivatives such as **2**, **4**, **5**, and **6** (Figure 1) in B16 melanoma cells. Compounds **2**, **5** and **6**, which represent different glycosyl moieties (glucoside, rhamnoside, and galactoside) at C-3, displayed no inhibition of melanin formation. Concerning this result, a previous report showed that quercetin 3-*O*-L-arabinofuranoside

did not suppress melanin formation (Fujii & Saito, 2009). These results revealed that the attachment of a glycosyl moiety at the C-3 position may decrease the ability of quercetin to reduce the melanin formation. We concluded that the hydroxyl group at the C-3 position is essential to the inhibition of melanin formation in B16 melanoma cells. To our knowledge, this is the first report showing that the dried skin of *A. cepa* and its isolated compound, **3**, showed the ability to act as a potent skin-whitening agent by inhibiting the melanin formation in B16 melanoma cells similarly to arbutin, which was used as a positive control. Therefore, further experiments are needed to determine the exact mechanism of this compound.

3.4. Antioxidant properties

Skin is a major candidate for and target of oxidative stress caused by reactive species (RS), including reactive oxygen species and reactive nitrogen species. RS are major and significant contributors to skin hyperpigmentation and skin aging (Kim, Kang, & Yokozawa, 2008). It has been generally believed that agents having antioxidant activity show anti-aging, whitening, and anti-inflammatory activities (Choi, Song, Hur & Sim, 2008). Since compounds **1** and **3** have shown promising results as whitening agents using B16 melanoma cells, we conducted an antioxidant assay (ORAC) in order to determine their ability to counteract oxidative stress from UV

radiation. Table 2 shows the ORAC value results of compounds **1** – **6**, which were 7.64, 8.65, 4.82, 4.32, 8.17 and 9.34 $\mu\text{mol TE}/\mu\text{mol}$, respectively. Compounds **1**, **2**, **5** and **6** showed more potent antioxidant activity than **3** and **4**. In the ORAC assay, quercetin showed an ORAC value similar to that reported by Kohri et al., (2009). However, we did not find any previous report of an ORAC assay for **3**. Based on the results in Table 1, it was revealed that 4'-glycosylation of quercetin decreased its ability as an antioxidant as represented by ORAC values.

The potency of **1** and **3** as antioxidants might be related to the presence of hydroxyl groups in the form of a catechol moiety as reported by Cao, Soffic, & Price (1997). Moreover, the relationship between flavonoid structure and antioxidant activity has been studied, and it has been found that 3',4'-dihydroxy substitution in the B ring, as in quercetin, increases the antioxidant activity substantially compared with a mono-hydroxy substituent, as in kaempferol (Rice-Evans, Miller, Bolwell, Bramley, & Pridham, 1995; Cotelle, Bernier, Catteau, Pommery, Wallet, & Gaydou, 1996).

4. Conclusion

Our study demonstrated that the melanin inhibition ability of quercetin derivatives with glycosyl moieties at C-3 is reduced. Also, our results showed that quercetin

derivatives with a glycosyl moiety at C-4' reduced its antioxidant activity. In addition, compounds **1** and **3** from the dried skin of *A. cepa*, are promising compounds that could be useful for treating hyperpigmentation as skin-whitening agents and as antioxidants. However, it should be noted that safety is a primary consideration for its practical use in humans.

Acknowledgement

This research supported by Japan Society for the Promotion of Science (JSPS).

REFERENCES

- Alfonso, D., & Kapetanidis, I. (1994). Flavonoids from *Iohcroma gesneriodes*. *Pharmaceutica Acta Helvetiae*, 68, 211-214.
- Arung, E.T., Shimizu, K., & Kondo, R. (2007). Structure-activity relationship of prenyl-substituted polyphenols from *Artocarpus heterophyllus* as inhibitors of melanin biosynthesis in cultured melanoma cells. *Chemistry and Biodiversity*, 4, 2166-2171.
- Beesk, N., Perner, H., Schwarz, D., George, E., Kroh, L.W., Rohn, S. (2010). Distribution of quercetin-3,4'-*O*-diglucoside, quercetin-4'-*O*-monoglucoside, and

308 quercetin in different parts of the onion bulb (*Allium cepa* L.) influenced by genotype.
 309 *Food Chemistry*, 122, 566-571.

310 Cao, G., Sofic, E., & Prior, R.L. (1997). Antioxidant and prooxidant behavior of
 311 flavonoids: structure-activity relationships. *Free Radical Biology and Medicine*, 22,
 312 749–760.

313 Choi, M.Y., Song, H.S., Hur, H.S., & Sim, S.S. (2008). Whitening activity of luteolin
 314 related to the inhibition of cAMP pathway in alpha-MSH-stimulated B16 melanoma
 315 cells. *Archives of Pharmacol Research*, 31, 1166–1171.

316 Cotellet, N., Berbier, J.L., Catteau, J.P., Pommery, J., Wallet, J.C., & Gaydou, E.M.
 317 (1996). Antioxidant properties of hydroxyl-flavones. *Free Radical Biology and*
 318 *Medicine*, 20, 35-43.

319 de Padua, L.S., Bunyaphatsara, N., & Lemmens, R.H.M.J. (1999). Plant resources of
 320 South-East Asia: Medicinal and poisonous plants, vol. 12 (1), pp. 93-97. Bogor,
 321 Prosea Press.

322 Fujii, T., & Saito, M. (2009). Inhibitory effect of quercetin isolated from rose hip (*Rosa*
 323 *canina* L) against melanogenesis by mouse melanoma cells. *Bioscience,*
 324 *Biotechnology, and Biochemistry*, 73, 1989-1993

325 Gulsen, A., Makris, D.P., & Kefalas, P. (2007). Biomimetic oxidation of quercetin:

326 isolation of a naturally occurring quercetin heterodimer and evaluation of its in vitro
 327 antioxidant properties. *Food Research International*, 40, 7–14.

328 Kaneko, T., & Baba, N. (1999). Protective effect of flavonoids on endothelial cells
 329 against linoleic acid hydroperoxide-induced toxicity. *Bioscience, Biotechnology, and*
 330 *Biochemistry*, 63, 323–328.

331 Kawaii, S., Tomono, Y., Katase, E., Ogawa, K., & Yano, M. (1999). Antiproliferative
 332 activity of flavonoids on several cancer cell lines. *Bioscience, Biotechnology, and*
 333 *Biochemistry*, 63, 896–899.

334 Kim, Y. J., Kang, K. S., & Yokozawa, T. (2008). The anti-melanogenic effect of
 335 pycnogenol by its anti-oxidative actions. *Food and Chemical Toxicology*, 46,
 336 2466–2471.

337 Kohri, S., Fujii, H., Oowada, S., Endoh, N., Sueishi, Y., Kusakabe, M., et al. (2009). An
 338 oxygen radical absorbance capacity-like assay that directly quantifies the
 339 antioxidant's scavenging capacity against AAPH-derived free radicals. *Analytical*
 340 *Biochemistry*, 386, 167–171

341 Kubo, I., Nitoda, T., & Nihei, K. (2007). Effects of Quercetin on Mushroom Tyrosinase
 342 and B16-F10 Melanoma Cells. *Molecules*, 12, 1045-1056

343 Nagata, H., Susumu, T., Takeyama, R., Homma, T., & Osamura, R.Y. (2004).

344 Quercetin Enhances Melanogenesis By Increasing the Activity and Synthesis of
 345 Tyrosinase in Human Melanoma Cells and in Normal Human Melanocytes. *Pigment*
 346 *Cell Research*, 17, 66–73.

347 Ou, B., Hampsch-Woodill, M., & Prior, R.L. (2001). Development and validation of an
 348 improved oxygen radical absorbance capacity assay using fluorescein as the
 349 fluorescent probe. *Journal of Agricultural and Food Chemistry*, 49, 4619–4626.

350 Perluigi, M., De Marco, F., Foppoli, C., Coccia, R., Blarzino, C., Marcante, M.L., et al.
 351 (2003) Tyrosinase protects human melanocytes from ROS-generating compounds.
 352 *Biochemical and Biophysical Research Communications*, 305, 250–256.

353 Prakash, D., Upadhyay, G., Singh, B.N., & Singh, H.B. (2007). Antioxidant and free
 354 radical-scavenging activities of seeds and agri-wastes of some varieties of soybean
 355 (Glycine max). *Food Chemistry*, 104, 783–790.

356 Prior, R. L., Hoang, H., Gu, L., Wu, X., Bacchioca, M., Howard, L., et al. (2003).
 357 Assays for hydrophilic and lipophilic antioxidant capacity (oxygen radical
 358 absorbance capacity (ORACFL) of plasma and other biological and food samples.
 359 *Journal of Agricultural and Food Chemistry*, 51, 3273–3279.

360 Rice-Evans, C.A., Miller, N.J., Bolwell, P.G., Branley, P.M., & Pridham, J.B. (1995).
 361 The relative antioxidant activities of plant-derivatived polyphenolic flavonoids. *Free*

362   *Radical Research*, 22, 375-383.

363   Sánchez-Ferrer, A., Rodríguez-López, J.N., & García-Carmona, F. (1995). Tyrosinase: a
364   comprehensive review of its mechanism. *Biochimica Biophysica Acta*, 1247, 1-11.

365   Sanderson, J., Mclauchlin, W., & Williamson, G. (1999). Quercetin inhibits hydrogen
366   peroxide-induced oxidization of the rat lens. *Free Radical Biology and Medicine*, 26,
367   639–645.

368   Shutenko, Z., Henry, Y., Pinard, E., Seylaz, J., Potier, P., Berthet, F., et al. (1999).
369   Influence of the antioxidant quercetin in vivo on the level of nitric oxide determined
370   by electron paramagnetic resonance in rat brain during global ischemia and
371   reperfusion. *Biochemical Pharmacology*, 57,199–208.

372   Singh, B.N., Singh, B.R., Singh, R.L., Prakash, D., Singh, D.P., Sarma, B.K., et al
373   (2009). Polyphenolics from various extracts/fractions of red onion (*Allium cepa*) peel
374   with potent antioxidant and antimutagenic activities. *Food and Chemical Toxicology*,
375   47, 450–452.

376   Tanabe, S., Ogawa, N., Tesaki, S., & Watanabe, M. (1997). Isolation and identification
377   of an onion blacking-active component by chelate formation with ferric ions. *Nippon*
378   *Kasei Gakkaishi*, 48, 339–342.

379   Virador, V.M., Kobayashi, N., Matsunaga, J., & Hearing, V.J. (1999). A standardized

380 protocol for assessing regulators of pigmentation. *Analytical Biochemistry*, 270,

381 207-219

382 Yoon, J.H., Shim, J.S., Cho, Y., Baek, N.I., Lee, C.W., Kim, H.S., et al. (2004).

383 Depigmentation of melanocytes by isopanduratin A and 4-hydroxypanduratin A

384 isolated from *Kaempferia pandurata* Roxb. *Biological and Pharmaceutical Bulletin*,

385 30, 2141-2145.

Legend of Figures

Figure 1. The structure of quercetin and its derivatives.

Figure 2. Effect of the methanol extracts of the dried skin of *A. cepa* on B16 melanoma cells [Arbutin : 100 µg/mL]. Each column represents the mean ± SD of three independents test (Student's *t*-test). Significantly different from the control value : P<0.05 (*), P<0.01 (**).

Figure 3. Effect of the methanol extracts of the flesh of *A. cepa* on B16 melanoma cells. Each column represents the mean ± SD of three independents test (Student's *t*-test). Significantly different from the control value : P<0.05 (*), P<0.01 (**).

Figure 4. HPLC analysis of the methanol extracts of dried skin of *A. cepa* [1. Quercetin, 2. Isoquercitrin, 3. Quercetin 4'-*O*-glucoside, 4. Quercetin 3,4'-*O*-diglucoside, Flow rate : 1 mL/min of MeOH (60): Water (40); Sample concentration : 1 mg/mL, Wave length 250 nm, Injection volume: 20 µL]

Figure 5. HPLC analysis of the methanol extract of flesh of *A. cepa* [1. Quercetin, 2. Isoquercitrin, 3. Quercetin 4'-*O*-glucoside, 4. Quercetin 3,4'-*O*-diglucoside Flow rate : 1 mL/min of MeOH (60): Water (40); Sample concentration : 11 mg/mL, Wave length 250 nm, Injection volume: 20µL]

409   **Legend of Tables**

410   **Table 1.** Effect of quercetin and its derivatives on B16 melanoma cells (n = 3)

411   **Table 2.** Effect of quercetin and its derivatives on ORAC assay (n = 3)

412

413

414

- 1: $R_1 = R_2 = R_3 = R_4 = R_5 = H$
2: $R_1 = R_2 = R_3 = R_4 = H$, $R_5 = \text{glc}$
3: $R_1 = \text{glc}$, $R_2 = R_3 = R_4 = R_5 = H$
4: $R_1 = \text{glc}$, $R_2 = R_3 = R_4 = H$, $R_5 = \text{glc}$
5: $R_1 = R_2 = R_3 = R_4 = H$, $R_5 = \text{rha}$
6: $R_1 = R_2 = R_3 = R_4 = H$, $R_5 = \text{gal}$

415

416

Figure 1.

417

418

419

420 **Figure 2.**

421

422

423

424 **Figure 3.**

Figure 4.

Figure 5.