

Regulation of Angiotensin II receptor signaling by cysteine modification of NF- κ B

Nishida, Motohiro

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University

Kitajima, Naoyuki

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University

Saiki, Shota

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University

Nakaya, Michio

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University

他

<https://hdl.handle.net/2324/25612>

出版情報 : Nitric Oxide. 25 (2), pp.112-117, 2011-08-01. Elsevier

バージョン :

権利関係 : (C) 2010 Published by Elsevier Inc.

Review Article

Regulation of Angiotensin II receptor signaling by cysteine modification of NF- κ B

Motohiro Nishida, Naoyuki Kitajima, Shota Saiki, Michio Nakaya and Hitoshi Kurose

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

Corresponding authors:

Motohiro Nishida

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan.

TEL&FAX: +81-92-642-6878

nishida@phar.kyushu-u.ac.jp

Hitoshi Kurose

Department of Pharmacology and Toxicology, Graduate School of Pharmaceutical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan.

TEL&FAX: +81-92-642-6884

kurose@phar.kyushu-u.ac.jp

Abstract

Angiotensin II (Ang II) is a major vasoactive peptide of the renin-angiotensin system. Ang II is originally found as one of potent vasoconstrictors, but is now attracted attention as an essential mediator of many cardiovascular problems, including endothelial dysfunction, arrhythmia and structural remodeling of cardiovascular systems. Most of the known pathophysiological effects of Ang II are mediated through Ang type1 receptors (AT₁Rs), and the up-regulation of AT₁Rs is one of important causes by which Ang II can contribute to cardiovascular diseases. A growing body of evidence has suggested that reactive oxygen species (ROS) and reactive nitrogen species (RNS) play important roles in the regulation of AT₁R signaling. In cardiac fibroblasts, stimulation with cytokines or bacterial toxins induces AT₁R up-regulation through NADPH oxidase-dependent ROS production. In contrast, nitric oxide (NO) decreases AT₁R density through cysteine modification (S-nitrosylation) of a transcriptional factor, nuclear factor κ B (NF- κ B). The difference between the effects of ROS and NO on AT₁R expression may be caused by the difference between intracellular location of ROS signaling and that of NO signaling, as the agonist-induced S-nitrosylation of NF- κ B requires a local interaction between NO synthase (NOS) and NF- κ B in the perinuclear region. Thus, the spatial and temporal regulation of cysteine modification by ROS or RNS may underlie the resultant changes of AT₁R signaling induced by agonist stimulation.

Keywords: Angiotensin, AT₁ receptor, heart, reactive oxygen species, nuclear factor- κ B, S-nitrosylation

1. Ang II signaling in the heart

The renin-angiotensin system (RAS) plays an important role in maintaining blood pressure homeostasis and salt balance in mammals [1, 2]. Ang II, a key regulator of RAS and exerts biological functions [3, 4], is generated to the following pathway. The protease renin first cleaves angiotensinogen to generate Ang I, which is composed of 10 amino acid polypeptide (Ang (1-10)). Next, the Ang converting enzyme (ACE) cleaves Ang I to generate Ang II (Ang (1-8)). Although there exists alternative Ang II-generating enzymes, such as cathepsins and chymase [5, 6], ACE is believed to be the sole enzyme in the regulation of Ang II production in the RAS [1, 2]. Ang II is then degraded to the heptapeptide Ang (1-7) by ACE2 [7]. Several reports have shown a counter-regulatory role for Ang (1-7) by opposing many AT₁R-mediated actions [8, 9]. The Ang (1-7) is further degraded by aminopeptidase to generate Ang III (Ang (2-8)) and Ang IV (Ang (3-8)). These Ang-derived metabolites exert biological activities through binding to G protein-coupled receptors (GPCRs) including AT₁R, AT₂R, and Mas [10, 11]. Among the Ang-derived metabolites, Ang II is a key regulator of cardiovascular functions, and most of biological effects of Ang II are mediated via AT₁R [3, 4, 10-12].

AT₁R-mediated signaling is divided into two groups: G protein-dependent and G protein-independent signaling pathways [13]. AT₁Rs are known to couple with three G protein subfamilies; G_q, G_i, G₁₂ [13, 14]. Among them, G_q and G₁₂ family proteins-mediated signaling pathways participate in structural remodeling of the heart (Figure 1). The G_q family protein induces activation of phospholipase C (PLC), which in turn induces increase in intracellular Ca²⁺ concentration ([Ca²⁺]_i) through production inositol-1,4,5-trisphosphate (IP₃) and diacylglycerol (DAG) by hydrolyzing

phosphatidylinositol 4,5-bisphosphate (PIP₂). Sustained increase in [Ca²⁺]_i in cardiomyocytes induces activation of calmodulin (CaM)-dependent protein phosphatase 2B, calcineurin, which leads to increase in expression of hypertrophic genes, such as atrial natriuretic peptide (ANP), β-myosin heavy chain (β-MHC), and α-skeletal muscle actin (α-SKA), through activation of nuclear factor of activated T cells (NFAT) [15]. We have recently reported that DAG-activated transient receptor potential canonical (TRPC) channels (TRPC3 and TRPC6) mediate Ang II-induced Ca²⁺ influx through voltage-dependent Ca²⁺ channel (VDCC) and cardiac hypertrophy *in vivo* and *in vitro* [16, 17]. In addition, activation of extracellular signal-regulated kinase (ERK) signaling pathway also participates in G_q-mediated cardiac hypertrophy. Lorenz *et al.* [18] has reported that Gβγ released from Gα_q induces autophosphorylation of ERK1/2 at Thr¹⁸⁸ residue and subsequent nuclear translocation of ERK1/2, resulting in inducing expression of hypertrophic genes. Ang II also induces ROS production through NADPH oxidase activation [19, 20]. Interestingly, Ang II-induced ROS production through Nox2 activation attenuates basal endothelial NOS (eNOS) activity through proline-rich tyrosine kinase (Pyk2)-dependent phosphorylation of eNOS at Tyr657 in endothelial cells [21], suggesting the mechanism of endothelial dysfunction observed in cardiovascular diseases associated with increased activity of the RAS. Using rat neonatal cardiomyocytes, we found that G₁₂ family proteins mediate Ang II-induced ROS production and activation of c-Jun NH₂-terminal kinase (JNK) and p38 MAPK [22, 23]. The activation of apoptosis signal-regulating kinase (ASK) and protein tyrosine kinase (PTK) by cysteine oxidation of thioredoxin and protein tyrosine phosphatase 1B may participate in ROS-mediated activation of JNK and p38MAPK [3, 4]. These results suggest that ROS function as a second messenger of G_{12/13} signaling in

cardiomyocytes. Furthermore, the inhibition of $G\alpha_{12/13}$ signaling in cardiomyocytes attenuates pressure overload-induced collagen deposition of the heart (fibrosis), using transgenic mice with cardiomyocytes-specific overexpression of regulator of G protein signaling (RGS) domain of p115 Rho guanine nucleotide exchange factor (p115RhoGEF) [24]. These results suggest that $G\alpha_{12/13}$ -mediated ROS production contributes to cardiac fibrosis induced by pressure overload. As cardiac fibrosis is one of major causes of left ventricular diastolic dysfunction, and the inhibition of $G\alpha_{12/13}$ -mediated ROS signaling attenuated diastolic dysfunction, $G\alpha_{12/13}$ -mediated ROS signaling may be a novel therapeutic target for the treatment of heart failure.

2. Redox regulation of AT_1R proteins

Expression cloning from bovine adrenal and rat smooth muscle cells has revealed that both AT_1R s are typical seven transmembrane domain proteins, composed of 359 amino acids and a molecular mass of about 41 kDa [25]. As the AT_1R possesses three N-glycosylation sites and four cysteine residues in the extracellular regions, the native glycosylated AT_1R has a molecular mass of about 65 kDa [26]. Although the rodent AT_1R s exist as two distinct subtypes, $AT_{1A}R$ and $AT_{1B}R$, that are 95% identical in their amino acid sequences, these structural features are present in several other mammalian and nonmammalian AT_1R s [10, 27]. The AT_1R s are not only activated by Ang-derived metabolites, but also activated by mechanical stretch [28, 29]. The disulfide bridges between the N-terminal region and third extracellular loop, and between the first and second extracellular loops are formed to maintain the conformation of the AT_1R protein [30]. As the disulfide bridge between the N-terminal region and third extracellular loop is not present in AT_2R , the AT_2R is resistant to inactivation by reducing agents.

Furthermore, Zhang *et al.* [31] has recently reported that nitro-oleic acid specifically binds to the AT₁R, reduces G_q protein coupling, and inhibits inositol-1,4,5-trisphosphate production and Ca²⁺ mobilization, without inhibiting Ang II binding to the receptor. Thus, the AT₁R-mediated signaling is negatively regulated by the oxidation or reduction of the AT₁Rs.

3. Regulation of AT₁R expression level

The gene expression of AT₁R is regulated by various transcriptional factors, such as Sp1, Sp3, myocyte enhancing factor (MEF)-2, peroxisome proliferator-activated receptor (PPAR)- γ , and NF- κ B [32-36]. It has been reported that Sp1 and Sp3 are predominantly responsible for regulating the basal expression of the human AT₁R gene [33, 37]. In contrast, MEF-2 heterodimers and Sp1 are required for basal expression of the rat AT_{1A}R gene [34]. PPAR- γ , an anti-inflammatory transcriptional factor, and Sp1 directly interact with each other on a CG-box-related sequence within the -58/-34 bp region of the rat AT_{1A}R promoter, which leads to suppression of AT_{1A}R transcription by inhibiting Sp1 binding activity [32]. On the other hand, NF- κ B, an inflammatory transcriptional factor, has been reported to participate in up-regulation of rat AT_{1A}R density induced by proinflammatory cytokines [36, 38, 39]. Cowling *et al.* [36] has reported that NF- κ B binds 2 *cis*-response elements located at -365/-355 bp and -2540/-2530 bp and induce transactivation of a minimal promoter. They have indicated that proinflammatory cytokines, such as tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β), increase the expression level of AT_{1A}R (Figure 2). Transcription of the rat AT_{1A}R is also enhanced by Toll-like receptor 4 (TLR4) ligands, such as lipopolysaccharide (LPS) and oxidized low-density lipoprotein (LDL), both of

which strongly increase NF- κ B activity. In contrast, NO, growth factors, statins and 15-deoxy- δ -PGJ₂ decrease AT₁R density [32, 40-43]. This mechanism may be explained by the decrease in DNA binding affinity of transcriptional factors through oxidative modification [43]. In addition, Ang II, cAMP stimulating agents and estrogens increase AT₁R mRNA decay rates, leading to decrease in AT₁R density (Figure 2). The AT₁R mRNA stability is regulated by calreticulin. The phosphorylated calreticulin binds to AUUUUA sequence localized in 3'-UTR region of AT₁R mRNA, leading to increase in AT₁R mRNA stability [44]. Thus, both transcriptional and posttranscriptional mechanisms are involved in the regulation of AT₁R expression.

4. ROS-mediated up-regulation of AT₁Rs

Ichiki *et al.* [45] have previously reported that exogenous exposure of hydrogen peroxide (H₂O₂) decreases AT₁R density through AT₁R mRNA destabilization. As the Ang II-induced decrease in AT₁R mRNA stability was diminished by the treatment with diphenyleneiodonium (DPI), an NADPH oxidase inhibitor, they suggest that endogenous ROS derived from NADPH oxidase negatively regulate AT₁R mRNA stability. In contrast, treatment with IL-1 β and TNF- α induces NADPH oxidase-dependent ROS production and increases AT₁R density [38, 39, 46]. Therefore, the role of endogenous ROS in the regulation of AT₁R expression may depend on the location and timing of ROS production induced by agonist stimulation.

We have recently found that endogenous ROS increase AT₁R density in rat cardiac fibroblasts [47]. *Pertussis* toxin (PTX), a major virulence factor of Gram-negative bacillus *Bordetella pertussis* which causes whooping cough, is well established as a

pharmacological tool for a specific inhibitor of receptor-G_{i/o} protein coupling. PTX is composed of A-protomer and B-oligomer, and A-protomer exerts ADP-ribosyltransferase activity on the α -subunit of heterotrimeric G_i proteins (G α_i), leading to inhibition of receptor-G protein coupling, whereas B-oligomer of PTX recognizes and binds carbohydrate-containing receptors that deliver A-protomer into the cytosol. As the treatment of rat cardiac fibroblasts with PTX induces up-regulation of AT₁Rs, indicating that PTX increases AT₁R density independently of ADP ribosylation. We also found that PTX induces TLR4 stimulation, which leads to activation of a small GTP-binding protein, Rac, through Syk tyrosine kinase (Figure 3). Activation of Rac subsequently induces NF- κ B activation through I κ B α phosphorylation and degradation. Activation of NF- κ B then increases expression of IL-1 β , which in turn activates Rac through IL-1 β receptor stimulation. Thus, IL-1 β -dependent amplification of ROS-mediated NF- κ B signaling is required for the PTX-induced AT₁R up-regulation. It is still unclear how endogenous ROS induce phosphorylation and degradation of I κ B α , but recent studies have suggested that stress-induced Ca²⁺ mobilization induces nuclear accumulation of I κ B α , leading to I κ B α degradation without N-terminal phosphorylation of I κ B α [48, 49]. As the stimulation of TLR4 induces ROS production independently of MyD88 signaling pathways has also been reported in macrophages [50], ROS-mediated signaling induced by TLR4 stimulation may play a critical role in TLR4-induced inflammatory responses (i.e., AT₁R up-regulation) in mammalian cells.

5. Suppression of AT₁R signaling by S-nitrosylation

It has been reported that NO and nitro-unsaturated fatty acids decrease Ang II

signaling in the heart [31, 43]. We have recently found that long-term treatment of rat neonatal cardiac fibroblasts with adenosine 5'-triphosphate (ATP) decreases AT₁R density. This phenomenon is so-called 'heterologous down-regulation', which indicates that stimulation of one GPCR reduces expression levels of different GPCR [51]. Cross-talk between different GPCR signaling pathways may serve to fine-tune cell signaling [52], but the molecular mechanism(s) underlying heterologous down-regulation is largely unknown. The ATP-induced AT₁R down-regulation was canceled by the treatment with cyclosporine A or siRNAs for P2Y₂R, indicating that P2Y₂R-stimulated sustained increase in intracellular Ca²⁺ concentration and subsequent activation of calcineurin-NFAT signaling participate in ATP-induced AT₁R down-regulation. Interestingly, the ATP-induced suppression of AT₁R signaling was completely diminished by 1400W, a selective inhibitor of inducible NO synthase (iNOS). As iNOS promoter region contains some NFAT-binding sequences and activation of calcineurin-NFAT signaling pathway has been reported to induce expression of iNOS proteins in cardiomyocytes and *in vivo* mouse hearts [53]. In fact, ATP increased expression of iNOS proteins in a concentration-dependent manner, and ATP-induced iNOS expression was completely suppressed by cyclosporine A. The expression of constitutively active mutant of NFAT also decreased AT₁R density, which was canceled by 1400W. Thus, NFAT-dependent iNOS expression is required for ATP-induced suppression of AT₁R signaling. We focused on the conserved cysteine (Cys) residue in the Rel homology domain of NF-κB [54, 55]. The Cys wedged between acidic and basic amino acids is nucleophilic, acidic and redox active, and numerous reactions may occur on this Cys thiol side chain [56]. As the reactive Cys locates in the Rel homology domain that is involved in DNA binding or dimer

formation of p65, the NO-mediated Cys modification (S-nitrosylation) of p65 may decrease transcriptional activity of NF- κ B [54, 55]. Using biotin-switch assay method as shown in Fig. 4A, we found that treatment of rat cardiac fibroblasts with ATP for 24 hours actually induced S-nitrosylation of p65 subunit at Cys38. The promoter region of iNOS does not only contain NF- κ B binding sequences but also NFAT binding sequences (Fig. 4B). Treatment of cardiac fibroblasts with ATP induced iNOS expression, and the iNOS protein and p65 subunit were co-localized around the perinuclear regions (Fig. 4C). Immunoprecipitation study has revealed that ATP-induced iNOS forms a complex with p65. In contrast, treatment with IL-1 β strongly increased expression of iNOS protein but did not induce the interaction of iNOS with p65, because IL-1 β induces translocation of p65 from cytosol to the nucleus. We also found that flavin-binding domain of iNOS protein is a hot-spot in iNOS for the interaction with p65, and overexpression of iNOS fragment including flavin-binding domain completely suppressed the ATP-induced suppression of AT₁R signaling. Thus, the formation of signaling complex between iNOS and p65 around the perinuclear region may be essential for the ATP-induced AT₁R down-regulation.

6. Conclusion

Numerous studies have demonstrated that Cys modification by ROS and NO play important roles in agonist-induced signal transduction. Indeed, numerous proteins have been reported to be a target of NO [57]. We found that ROS and NO play important roles in the regulation of AT₁R gene expression. ROS increases AT₁R density through I κ B α degradation-dependent activation of NF- κ B, while NO decreases AT₁R density through direct S-nitrosylation of NF- κ B p65 subunit. As the physical

interaction between p65 and iNOS proteins are required for agonist-induced S-nitrosylation of p65, the location and the timing of ROS or NO production induced by agonist stimulation may be critical for the resultant regulation of AT₁R signaling. Identification of the target protein that induces I κ B α phosphorylation will be required for the understanding of the ROS-mediated signaling complex participating in AT₁R up-regulation. A growing body of evidence has suggested that ROS and NO work as second messengers in cellular signaling, and the emergence of Cys modification (thiol oxidation and S-nitrosylation) by ROS and RNS may presage a new era in cardiovascular biology. Unraveling the regulation of Ang II signaling by Cys modification will achieve new therapeutic targets with great potential to improve clinical outcomes.

References

- [1] L.T. Skeggs, F.E. Dorer, M. Levine, K.E. Lentz, J.R. Kahn, The biochemistry of the renin-angiotensin system, *Adv. Exp. Med. Biol.* 130 (1980) 1-27.
- [2] A.J. Turner, N.M. Hooper, The angiotensin-converting enzyme gene family: genomics and pharmacology, *Trends. Pharmacol. Sci.* 23 (2002) 177-183.
- [3] Y. Saito, B.C. Berk, Angiotensin II-mediated signal transduction pathways, *Curr. Hypertens. Rep.* 4 (2002) L167-L171.
- [4] R.M. Touyz, Reactive oxygen species as mediators of calcium signaling by Angiotensin II: implications in vascular physiology and pathophysiology, *Antioxid. Redox Signaling.* 7 (2005) 1302-1314.
- [5] L.A. Belova, Angiotensin II-generating enzymes, *Biochemistry (Mosc.)* 65 (2000) 1337-1345.
- [6] M. Miyazaki, S. Takai, Tissue Angiotensin II generating system by angiotensin-converting enzyme and chymase, *J. Pharmacol. Sci.* 100 (2006) 391-397.
- [7] S.R. Tipnis, N.M. Hooper, R. Hyde, E. Karran, G. Christie, A.J. Turner, A human homolog of angiotensin-converting enzyme. Cloning and functional expression as a captopril-insensitive carboxypeptidase, *J. Biol. Chem.* 275 (2000) 33238-33243.
- [8] R.A. Santos, F. Frezard, A.J. Ferreira, Angiotensin-(1-7): blood, heart, and blood vessels, *Curr. Med. Chem. Cardiovas. Hematol. Agents.* 3 (2005) 383-391.
- [9] C.M. Ferrario, Angiotensin-converting enzyme 2 and angiotensin-(1-7): an evolving story in cardiovascular regulation, *Hypertension* 47 (2006) 515-521.
- [10] M. de Gasparo, K.J. Catt, T. Inagami, J.W. Wright, Th. Unger, International union of pharmacology. XXIII. The angiotensin II receptors, *Pharmacol. Rev.* 52 (2000)

415-472.

- [11] C. Oro, H. Qian, W.G. Thomas, Type 1 angiotensin receptor pharmacology: signaling beyond G proteins, *Pharmacol. Ther.* 113 (2007) 210-226.
- [12] G.H. Gibbons, V.J. Dzau, The emerging concept of vascular remodeling, *N. Eng. J. Med.* 330 (1994) 1431-1438.
- [13] L. Hunyady, K.J. Catt, Pleiotropic AT1 receptor signaling pathways mediating physiological and pathogenic actions of Angiotensin II, *Mol. Endocrinol.* 20 (2006) 953-970.
- [14] J. Sadoshima, S. Izumo, The cellular and molecular response of cardiac myocytes to mechanical stress, *Annu. Rev. Physiol.* 59 (1997) 551-571.
- [15] J. Heineke, J.D. Molkentin, Regulation of cardiac hypertrophy by intracellular signaling pathways, *Nat. Rev. Mol. Cell. Biol.* 7 (2006) 589-600.
- [16] N. Onohara, M. Nishida, R. Inoue, H. Kobayashi, H. Sumimoto, Y. Sato, Y. Mori, T. Nagao, H. Kurose, TRPC3 and TRPC6 are essential for angiotensin II-induced cardiac hypertrophy, *EMBO J.* 25 (2006) 5305-5316.
- [17] S. Kiyonaka, K. Kato, M. Nishida, K. Mio, T. Numaga, Y. Sawaguchi, T. Yoshida, M. Wakamori, E. Mori, T. Numata, M. Ishii, H. Takemoto, A. Ojida, K. Watanabe, A. Uemura, H. Kurose, T. Morii, T. Kobayashi, Y. Sato, C. Sato, I. Hamachi, Y. Mori, Selective and direct inhibition of TRPC3 channels underlies biological activities of a pyrazole compound, *Proc. Natl. Acad. Sci. USA* 106 (2009) 5400-5405.
- [18] K. Lorenz, J.P. Schmitt, E.M. Schmitteckert, M.J. Lohse, A new type of ERK1/2 autophosphorylation causes cardiac hypertrophy, *Nat. Med.* 15 (2009) 75-83.
- [19] S. Rajagopalan, S. Kurz, T. Munzel, M. Tarpey, B.A. Freeman, K.K. Griendling, D.G. Harrison, Angiotensin II-mediated hypertension in the rat increases vascular superoxide

- production via membrane NADH/NADPH oxidase activation. Contribution to alterations of vasomotor tone, *J. Clin. Invest.* 97 (1996) 1916-1923.
- [20] M. Ushio-Fukai, R.W. Alexander, M. Akers, K.K. Griending, p38 mitogen-activated protein kinase is a critical component of the redox-sensitive signaling pathways activated by angiotensin II, *J. Biol. Chem.* 273 (1998) 15022-15029.
- [21] A.E. Loot, J.G. Schreiber, B. Fisslthaler, I. Fleming, Angiotensin II impairs endothelial function via tyrosine phosphorylation of the endothelial nitric oxide synthase, *J. Exp. Med.* 206 (2009) 2889-2896.
- [22] M. Nishida, S. Tanabe, Y. Maruyama, S. Mangmool, K. Urayama, Y. Nagamatsu, S. Takagahara, J.H. Turner, T. Kozasa, H. Kobayashi, Y. Sato, T. Kawanishi, R. Inoue, T. Nagao, H. Kurose, $G\alpha_{12/13}$ -and reactive oxygen species-dependent activation of c-Jun NH₂-terminal kinase and p38 MAPK by angiotensin receptor stimulation in rat neonatal cardiomyocytes, *J. Biol. Chem.* 280 (2005) 18434-18441.
- [23] T. Fujii, N. Onohara, Y. Maruyama, S. Tanabe, H. Kobayashi, M. Fukutomi, Y. Nagamatsu, N. Nishihara, R. Inoue, H. Sumimoto, F. Shibasaki, T. Nagao, M. Nishida, H. Kurose, $G\alpha_{12/13}$ -mediated production of reactive oxygen species is critical for angiotensin receptor-induced NFAT activation in cardiac fibroblasts, *J. Biol. Chem.* 280 (2005) 23041-23047.
- [24] M. Nishida, Y. Sato, A. Uemura, Y. Narita, H. Tozaki-Saitoh, M. Nakaya, T. Ide, T. Suzuki, K. Inoue, T. Nagao, H. Kurose, P2Y₆ Receptor- $G\alpha_{12/13}$ signaling in cardiomyocytes triggers pressure overload-induced cardiac fibrosis, *EMBO J.* 27 (2008) 3104-3115.
- [25] K. Sasaki, Y. Yamano, S. Bardhan, N. Iwai, J.J. Murray, M. Hasegawa, Y. Matsuda, T. Inagami, Cloning and expression of a complementary DNA encoding a bovine adrenal

- angiotensin II type-1 receptor, *Nature (Lond)* 351 (1991) 230-233.
- [26] S. Jayadev, R.D. Smith, G. Jagadeesh, A.J. Baukal, L. Hunyady, K.J. Catt, N-linked glycosylation is required for optimal AT_{1a} Angiotensin receptor expression in COS-7 cells. *Endocrinology* 140 (1999) 2010-2017.
- [27] T.S. Elton, M.M. Martin, Angiotensin II type1 receptor gene regulation. Transcriptional and posttranscriptional mechanisms, *Hypertension* 49 (2007) 953-961.
- [28] Y. Zou, H. Akazawa, Y. Qin, M. Sano, H. Takano, T. Minamino, N. Makita, K. Iwanaga, W. Zhu, S. Kudoh, H. Toko, K. Tamura, M. Kihara, T. Nagai, A. Fukamizu, S. Umemura, T. Iiri, T. Fujita, I. komuro, Mechanical stress activates angiotensin II type 1 receptor without the involvement of angiotensin II, *Nat. Cell Biol.* 6 (2004) 499-506.
- [29] N. Yasuda, S. Miura, H. Akazawa, T. Tanaka, Y. Qin, Y. Kiya, S. Imaizumi, M. Fujino, K. Ito, Y. Zou, S. Fukuhara, S. Kunimoto, K. Fukuzaki, T. Sato, J. Ge, N. Mochizuki, H. Nakaya, K. Saku, I. Komuro, Conformational switch of angiotensin II type 1 receptor underlying mechanical stress-induced activation, *EMBO Rep.* 9 (2008) 179-186.
- [30] K. Ohyama, Y. Yamano, T. Sano, Y. Nakagomi, T. Hamakubo, I. Morishima, T. Inagami, Disulfide bridges in extracellular domains of Angiotensin II receptor type IA, *Regul. Pept.* 57 (1995) 141-147.
- [31] J. Zhang, L. Villacorta, L. Chang, Z. Fan, M. Hamblin, T. Zhu, C.S. Chen, M.P. Cole, F.J. Schopfer, C.X. Deng, M.T. Garcia-Barrio, Y-H. Feng, B.A. Freeman, Y.E. Chen, Nitro-oleic acid inhibits angiotensin II-induced hypertension, *Circ. Res.* 107 (2010) 540-548.
- [32] A. Sugawara, K. Takeuchi, A. Uruno, Y. Ikeda, S. Arima, M. Kudo, K. Sato, Y. Taniyama, S. Ito, Transcriptional suppression of type 1 angiotensin II receptor gene expression by peroxisome proliferators-activated receptor- γ in vascular smooth muscle

- cells, *Endocrinology* 142 (2001) 3125-3134.
- [33] X. Zhao, M.M. Martin, T.S. Elton, Basal level transcriptional regulation of the human angiotensin II type 1 receptor gene, *Biochim. Biophys. Acta.* 1494 (2000) 181-184.
- [34] K.B. Beason, M.E. Steinhilber, T.S. Elton, An A/T-rich *cis*-element is essential for rat angiotensin II type 1A receptor transcription in vascular smooth muscle cells. *Biochim. Biophys. Acta.* 1444 (1999) 25-34.
- [35] T. Kubo, N. Kinjyo, A. Ikezawa, T. Kambe, R. Fukumori, Sp1 decoy oligodeoxynucleotide decreases angiotensin receptor expression and blood pressure in spontaneously hypertensive rats. *Brain Res.* 992 (2003) 1-8.
- [36] R.T. Cowling, D. Gurantz, J.F. Peng, W.H. Dillmann, B.H. Greenberg, Transcription factor NF- κ B is necessary for up-regulation of type 1 angiotensin II receptor mRNA in rat cardiac fibroblasts treated with tumor necrosis factor- α or interleukin-1 β , *J. Biol. Chem.* 277 (2002) 5719-5724.
- [37] M.M. Martin, X. Victor, X. Zhao, J.K. McDougall, T.S. Elton, Identification and characterization of functional angiotensin II type 1 receptors on immortalized human fetal aortic vascular smooth muscle cells, *Mol. Cell. Endocrin.* 183 (2001) 81-91.
- [38] R.T. Cowling, X. Zhang, V.C. Reese, M. Iwata, D. Gurantz, W.H. Dillmann, B.H. Greenberg, Effects of cytokine treatment on angiotensin II type 1A receptor transcription and splicing in rat cardiac fibroblasts, *Am. J. Physiol.* 289 (2005) H1176-H1183.
- [39] D. Gurantz, R.T. Cowling, J. Francisco, J. Villarreal, B.H. Greenberg, Tumor necrosis factor- α upregulates angiotensin II type 1 receptors on cardiac fibroblasts, *Circ.Res.* 89 (1999) 272-279.
- [40] Y. Ikeda, K. Takeuchi, T. Kato, Y. Taniyama, K. Sato, N. Takahashi, A. Sugawara, S.

- Ito, Transcriptional suppression of rat angiotensin AT_{1a} receptor gene expression by interferon-gamma in vascular smooth muscle cells, *Biochem. Biophys. Res. Commun.* 262 (1999) 494-498.
- [41] G. Nickening, T.J. Murphy, Down-regulation by growth factors of vascular smooth muscle angiotensin receptor gene expression, *Mol. Pharmacol.* 46 (1994) 653-659.
- [42] S. Wassmann, U. Laufs, A.T. Baumer, K. Muller, C. Konkol, H. Sauer, M. Bohm, G. Nickenig, Inhibition of geranylgeranylation reduces angiotensin II-mediated free radical production in vascular smooth muscle cells: involvement of angiotensin AT₁ receptor expression and Rac1 GTPase, *Mol. Pharmacol.* 59 (2001) 646-654.
- [43] T. Ichiki, M. Usui, M. Kato, Y. Funakoshi, K. Ito, K. Egashira, A. Takeshita, Downregulation of angiotensin II type 1 receptor gene transcription by nitric oxide, *Hypertension* 31 (1998) 342-348.
- [44] G. Nickenig, F. Michaelsen, C. Muller, A. Berger, T. Vogel, A. Sachinidis, H. Vetter, M. Bohm, Destabilization of AT₁ receptor mRNA by calreticulin, *Circ. Res.* 90 (2002) 53-58.
- [45] T. Ichiki, K. Takeda, T. Tokunou, Y. Funakoshi, K. Ito, N. Iino, A. Takeshita, Reactive oxygen species-mediated homologous downregulation of angiotensin II type 1 receptor mRNA by angiotensin II, *Hypertension* 37 (2001) 535-540.
- [46] M. Sundaresan, Z.X. Yu, V.J. Ferrans, D.J. Sulciner, J.S. Gutkind, K. Irani, P.J. Goldschmidt-Clermont, T. Finkel, Regulation of reactive-oxygen-species generation in fibroblasts by Rac1, *Biochem. J.* 318 (1996) 379-382.
- [47] M. Nishida, R. Suda, Y. Nagamatsu, S. Tanabe, N. Onohara, M. Nakaya, Y. Kanaho, T. Shibata, K. Uchida, H. Sumimoto, Y. Sato, H. Kurose, Pertussis toxin upregulates angiotensin type1 receptors through TLR4-mediated Rac activation, *J. Biol. Chem.* 285

(2010) 15268-15277.

- [48] Y. Lao, D.C. Chang, Mobilization of Ca^{2+} from endoplasmic reticulum to mitochondria plays a positive role in the early stage of UV- or TNF α -induced apoptosis, *Biochem. Biophys. Res. Commun.* 373 (2008) 42-47.
- [49] Y. Tsuchiya, T. Asano, K. Nakayama, T. Kato Jr, M. Karin, H. Kamata, Nuclear IKK β is an adaptor protein for I κ B α ubiquitination and degradation in UV-induced NF- κ B activation, *Mol. Cell* 39 (2010) 570-582.
- [50] Y.S. Bae, J.H. Lee, S.H. Choi, S. Kim, F. Almazan, J.L. Witztum, Y.I. Miller, Macrophages generate reactive oxygen species in response to minimally oxidized low-density lipoprotein: toll-like receptor 4- and spleen tyrosine kinase-dependent activation of NADPH oxidase 2. *Circ. Res.* 104 (2009) 210-218.
- [51] T.T. Chuang, L. Iacovelli, M. Sallese, A. De Blasi, G protein-coupled receptors: heterologous regulation of homologous desensitization and its implications, *Trends Pharmacol. Sci.* 17 (1996) 416-421.
- [52] Y. Cordeaux, S.J. Hill, Mechanisms of cross-talk between G-protein-coupled receptors, *Neurosignals* 11 (2002) 45-57.
- [53] K. Obasanjo-Blackshire, R. Mesquita, R.I. Jabr, J.D. Molkenin, S.L. Hart, M.S. Marber, Y. Xia, R.J. Heads, Calcineurin regulates NFAT-dependent iNOS expression and protection of cardiomyocytes: co-operation with Src tyrosine kinase, *Cardiovasc. Res.* 71 (2006) 672-683.
- [54] Z.T. Kelleher, A. Matsumoto, J.S. Stamler, H.E. Marshall, NOS2 regulation of NF- κ B by S-nitrosylation of p65, *J. Biol. Chem.* 282 (2007) 30667-30672.
- [55] H.E. Marshall, J.S. Stamler, Inhibition of NF- κ B by S-nitrosylation, *Biochemistry* 40 (2001) 1688-1693.

- [56] J.S. Stamler, E.J. Toone, S.A. Lipton, N.J. Sucher, (S)NO signals: translocation, regulation, and a consensus motif, *Neuron* 18 (1997) 691-696.
- [57] B. Lima, M.T. Forester, D.T. Hess, J.S. Stamler, S-nitroylation in cardiovascular signaling, *Circ. Res.* 106 (2010) 633-646.

Figure legends

Figure 1. AT_1R signaling pathways in the heart. ASK; apoptosis signal-regulating kinase, PTK; protein tyrosine kinase, VDCC; voltage-dependent Ca^{2+} channel.

Figure 2. Regulation of AT_1R transcription and translation by various factors. PPAR; peroxisome proliferator-activated receptor, LDL; low-density lipoprotein.

Figure 3. Hypothetical mechanism of ROS-mediated AT_1R up-regulation induced by *Pertussis* toxin in rat cardiac fibroblasts.

Figure 4. Mechanism of AT_1R down-regulation induced by NO. (A) principle of biotin-switch assay. (B) Potential binding sites of transcriptional factors existed in iNOS promoter region. (C) Changes in localization of iNOS and p65 proteins induced by ATP and IL-1 β . Cardiac fibroblasts were treated with ATP (100 μ M) or IL-1 β (1 ng/ml) for 24 hours. (D) Localization of p65 defines iNOS-dependent S-nitrosylation of p65 and resultant AT_1R down-regulation.




Figure 1. Nishida et al.




Figure 2. Nishida et al.




Figure 3. Nishida et al.




Figure 4. Nishida et al.